Fundamental Shifts of Central Hardwood Forests

Under Climate Change

Wu Ma¹, Jingjing Liang¹, Jonathan R. Cumming², Eungul Lee³, Amy B. Welsh¹, James V. Watson¹, Mo Zhou^{1*}

¹ School of Natural Resources, West Virginia University, USA

² Department of Biology, West Virginia University, USA

³ Department of Geology and Geography, West Virginia University, USA

Keywords: Climate change; fire regime; matrix growth model; forest dynamics; Monte Carlo simulations; Appalachians

***Corresponding author:**

Mo Zhou

PO Box 6125, School of Natural Resources

West Virginia University

Morgantown 26506, USA

Tel: (304) 293-5827; Fax: (304) 293-2441

Email: Mo.Zhou@mail.wvu.edu

Abstract

The Central Hardwood Region (CHR) of the United States constitutes one of the most diverse ecoregions in North America and the most extensive temperate deciduous forest in the world. Despite the economic and ecological significance of the CHR, the long-term effects of changes in climate and fire regime on forest structure remain largely unknown. In this study, we developed an integrated climate-sensitive matrix framework to synchronously couple (1) forest dynamics, (2) mean fire interval, (3) population density, and (4) future climate scenarios to study the community and population structure of CHR forests under climate change and associated changes of fire regimes. Using Monte Carlo simulations and coupled forest dynamics-disturbance models, we projected that the CHR would undergo a major shift in forest community structure from the present to year 2100. The fundamental changes would consist of a transition of dominant species from oak and hickory to maple species, reduced species diversity (9.6 – 11.5%), and substantial declines in stand basal area (55.1 – 62.0%) and stand volume (56.3 – 62.4%). These projected changes will have profound ecological and economic implications. Ecologically, changes in tree species diversity favoring maples would alter ecosystem processing of nutrients and subsequent nutrient flows to drainage waters within the region. Habitat change would alter the broad spectrum of organisms relying on the forest, leading to a redistribution of wildlife species, further heightening the risks for endangered species. Economically, the total stumpage value throughout the CHR would be reduced by 54.5 – 59.8% from approximately \$1,317 billion to \$529 – 599 billion. On the brink of these fundamental shifts, our study calls for ecologically- and economically-informed conservation and mitigation strategies to better prepare society for the associated changes in ecosystem services and economic benefits derived from CHR forests.

Introduction

The Central Hardwood Region (CHR) of the United States, covered by approximately 58 million ha of forests stretching from the upper southeast U.S. to Indiana and from Oklahoma to Pennsylvania (Fig. 1), constitutes the most extensive temperate deciduous forest in the world (Box & Fujiwara, 2015). The CHR is well known for its wide variety of oak-hickory forest resources that provide significant ecological and economic benefits to local, regional, and national communities. The CHR forms the headwaters for many major U.S. rivers and plays critical roles in improving and protecting soil and water resources as well as in mitigating flooding (Bernhardt & Palmer, 2011). It is home to a wide array of flora and fauna species (Schmidt & McWilliams, 2003) and is one of the most diverse ecoregions in North America (Mueller, 1996). The CHR supports ninety percent of hardwoods and one third of the total forest growing stock in the continental United States (Hicks, 1998). The high quality hardwood timber resources play a vital role in regional employment and wood related industries.

A growing body of evidence suggests that global climate change will have significant impacts on forest ecosystems by affecting the distribution and variation of key environmental factors, such as CO₂, humidity, and incoming solar radiation (e.g., Aber *et al.*, 2001, Allen *et al.*, 2010, Boisvenue & Running, 2006, Latta *et al.*, 2010, Lindner *et al.*, 2010, Schoene & Bernier, 2012). These changes will affect species composition and the productivity of forest ecosystems in this region (Aber *et al.*, 2001, Boisvenue & Running, 2006, Latta *et al.*, 2010, Shugart *et al.*, 2003, Smith *et al.*, 1995) as well as forest ecosystem processes through alterations in resource acquisition and resource utilization efficiency (Hansen & Dale, 2001, Hansen *et al.*, 2001, Helmick *et al.*, 2014, Juday *et al.*, 2005). Precipitation and temperature, and their seasonality and extremes, may change species' ranges, inter-species relationships, fire frequency, and other

ecosystem processes in the CHR that will have broad ecological and economic implications across the region and beyond (Alexander & Arthur, 2010, Lafon *et al.*, 2005, Parisen & Moritz, 2009). In addition, recent theoretical advances and empirical evidence (Cardinale *et al.*, 2012, Liang *et al.*, 2015, Naeem *et al.*, 2012, Tilman *et al.*, 1997) have revealed substantial impact of the loss of biodiversity on the functioning of ecosystems. The impact of climate change and biodiversity loss on CHR forests, however, has yet to be quantified.

Natural disturbances are a major factor affecting forest dynamics and composition (Fischer *et al.*, 2013), and influence the development of effective ecosystem restoration and management practices (Foster, 2000, Zhou & Buongiorno, 2006). Throughout the CHR, forest dynamics have been driven historically by high intensity stand-replacing fires necessary for the successful regeneration of shade intolerant species (Albrecht & Mccarthy, 2006, Brose *et al.*, 2013, McEwan *et al.*, 2011, McEwan *et al.*, 2007, Schuler *et al.*, 2012, Signell *et al.*, 2005). However, human intervention has reduced both the intensity and area of CHR forests burned since 1940, which has led to a widespread transition in dominant species from oak (*Quercus spp.*) and hickory (*Carya spp.*) to other early successional maple species (*Acer spp.*). This transition is termed the mesophication of the eastern hardwood forests (Fralish & McArdle, 2009, Nowacki & Abrams 2008). Accompanying this transition has been a reduction in area of bottomland hardwood forests and original oak savannas (Schmidt & McWilliams, 2003). In modeling fire impacts, fire regimes are typically based on the vegetation associations (Hann *et al.*, 2004, Keane *et al.*, 2002). Recent modeling efforts consider climate variables as predictors (Jiang *et al.*, 2012, Parisen & Moritz, 2009, Westerling *et al.*, 2006), and synthesize existing fire history information and mean fire intervals (MFI) based on physical mechanisms associated with dry climatic conditions (Guyette *et al.*, 2010). Temperature, precipitation, and their interaction prove to be the

most significant factors controlling fire frequencies and intensity in forest ecosystems (Morgan *et al.*, 2001), and these environmental factors are predicted to change in the future.

Reliable forest growth models are lacking for the CHR. One of the first forest growth models for the region was established by Perkey (1985). This whole-stand model simulated stand-level attributes, but did not specify population structure. Later, the distance-independent individual tree growth and yield system OAKSIM (Hilt, 1985) was developed for even-aged upland oak stands in southern Ohio and southeastern Kentucky, but this model has rarely been used due to limited applicability and a lack of stability (Brooks & Miller, 2011). Three more recent models for the region, namely SILVAH (Marquis & Ernst, 1992), the Forest Vegetation Simulator (FVS, Bush, 1995), and the Stand Damage Model (Colbert & Racin, 1995), have substantial prediction bias ranging from 20 to 140% of actual trees per acre (Brooks & Miller, 2011). Another whole-stand growth and yield model (Brooks, 2012) demonstrates better accuracy, but a lack of recruitment module renders this model inappropriate for long-term projections. More recently, LANDIS PRO has been developed to project forest successional trajectories and stand development patterns (Wang *et al.*, 2014). However, data are missing for several key tree species including yellow poplar, white ash, and black gum (He *et al.*, 2012); consequently, this model is susceptible to errors in terms of structured forest populations.

The primary objective of this study was to develop an empirical model of forest dynamics to study the successional patterns of CHR forests under future climate change. We then extended this modeling framework to account for climate-induced changes of fire regimes. We further employed Monte Carlo simulations in projecting future forest states and stumpage values under four future climate scenarios: RCP2.6, RCP4.5, RCP6.0 and RCP8.5. Ecological and economic

implications of the simulation results are discussed in relation to known ecosystem services provided by CHR forests.

Materials and Methods

Data

Forest and Stumpage Price Data

The CHR encompasses three ecoregions, namely the Blue Ridge Mountains, the historic Great Valley, and the ridges and valleys of the Allegheny Mountains (see Bailey, 2004 and references therein). Additionally, the CHR covers a wide range of stand states, community species composition, and climatic conditions (Fig. 1). Data used for model calibration consisted of 5,196 re-measured permanent sample plots (PSP's) from the Forest Inventory and Analysis (FIA) database (Sharon *et al.*, 2011) following three criteria. First, stand characteristics of the sample plots were closely monitored and re-measured. Second, stands had at least one live tree at the time of both measurements. Finally, plots were located in forests without any evidence of silvicultural treatments or any other forms of human interference, such as cutting and artificial regeneration. For validation purposes, we acquired an additional 1,107 plots randomly located throughout the region to test model accuracy.

For each sample plot, physical site attributes included geographic coordinates, slope, aspect, and elevation. Tree data, including species, diameter, and status (recruitment, live, or dead), were also collected on site. The research area is largely dominated by seven major species: *Quercus alba* (white oak), *Quercus velutina* (black oak), *Quercus rubra* (northern red oak), *Carya glabra* (pignut hickory), *Acer saccharum* (sugar maple), *Acer rubrum* (red maple), and *Liriodendron tulipifera* (yellow-poplar). White oak had the highest basal area of all the species

(13.9%), followed by black oak (10.9%), northern red oak (5.8%), pignut hickory (4.7%), sugar maple (4.5%), red maple (4.4%), and yellow-poplar (3.9%) (see Supplemental Information, Table S1). Due to the high diversity in this region, there also exist over 100 other tree species. For simplicity and due to computational constraints, we classified all tree species into seven species groups according to their taxonomic features: white oak species (*Quercus* – *Quercus*, QQ), red oak species (*Quercus* – *Lobatae*, QL), *Juglandaceae* (JD), *Sapindaceae* (SD), *Gymnosperms* (GS), *Fagus* (FG), and Other Angiosperms (OA) (see Supplemental Information, Table S1). Within each species group, trees were further categorized into seventeen diameter at breast height (DBH) classes of 5-cm increments, except for the first class (2.54 – 7 cm) and the last (82 cm and above) class.

Among all the variables studied (Table 1), the average recruitment (R) and total stem density (N) were the highest for *Sapindaceae* (SD) and lowest for *Fagus* (FG, see Supplemental Information, Table S2). The average interval between two inventories was roughly 6 years. At the individual tree level (see Supplemental Information, Table S3), the *Quercus* – *Lobatae* (QL) had the largest diameter at breast height (D) and the highest average annual diameter growth (g). *Juglandaceae* (JD) had the lowest mortality rate (m), and *Gymnosperms* (GS) had the highest.

For the analysis of stumpage values, we used the stumpage price data for the CHR in 2015 from the West Virginia Timber Market Report (<http://ahc.caf.wvu.edu/ahc-resources-mainmenu-45/timber-market-report-mainmenu-62>, accessed July 3, 2015) and assumed constant real prices in 2015 dollars in the region until 2100 to provide a current value estimate to understand implications in today's dollars.

Climate Data and Method

We assembled the historical records (30-year mean for 1981-2010) of the average annual temperature ($^{\circ}\text{C}$) and annual average of monthly mean precipitation (100 mm month^{-1}) from the downscaled 4-km resolution Parameter-Elevation Regression on Independent Slopes Model (PRISM) datasets (Daly *et al.*, 2008). PRISM calculates a climate-elevation regression for each digital elevation model grid cell, so the climate variables from PRISM, especially precipitation, are applicable for interpolation over mountain ranges (Daly *et al.*, 2008). The PRISM data set was shown to be a more accurate representation of spatial climatic patterns in the United States than the WorldClim and Daymet datasets (Daly *et al.*, 2008). For the prediction of future climatic changes in the CHR, we extracted the future relative changes (trend ratio) over the region (75°W - 95°W and 34.5°N - 42°N ; shown as the box in Figure 1) for the four scenarios of the IPCC AR5 report (Blyth *et al.*, 2007). The trend ratios of temperature and precipitation were obtained from the KNMI Climate Change Atlas (<http://climexp.knmi.nl/>, last accessed 10-August-2015). The four future climate scenarios of Representative Concentration Pathways (RCPs) were chosen to span a wide range of possible future conditions: RCP8.5 is a business-as-usual scenario with increasing greenhouse gas emissions over time, leading to high greenhouse gas concentration levels; RCP6.0 is a stabilization scenario in which emissions rise quickly until 2060 and then decrease; RCP4.5 assumes quicker action to limit greenhouse gas emissions with emissions peaking in 2040 and declining strongly until 2080; and RCP2.6 describes an all-out effort to limit global warming to below 2°C , with emissions decreasing sharply after 2020 and to zero emissions from 2080 onward (IPCC, 2013).

We estimated future temperature and precipitation changes in the study area in three steps. First, we extracted the 30-year (1981-2010) average annual temperature and annual average of monthly mean precipitation at all the plots (5,196 for calibration and 1,107 for validation plots)

using the PRISM data. Second, the trend ratios of temperature and precipitation over the CHR were obtained under the four RCP scenarios of future climate change (2010-2100) from the KNMI Climate Change Atlas. Third, we multiplied the extracted historical normal (30-year mean) temperature and precipitation by the future trend ratios to estimate the future temperature and precipitation changes, respectively, at the validation plot sites in the CHR during 2010-2100. Over the area represented by the validation plots, future annual temperature showed an increasing trend under all four RCP scenarios, and the annual average of monthly mean precipitation did not change significantly over time under any scenario, compared to 2010 climate (Figure 2). Average annual temperature and annual average of monthly mean precipitation were predicted to have the greatest changes of +3.6 °C under RCP8.5 and +14.3 mm month⁻¹ under RCP6.0, respectively, over the next 90 years. RCP2.6 has the smallest increase of temperature, +0.8 °C, and decline of precipitation, -0.6 mm month⁻¹ (Fig. 2a, b). For the study area, the future climate in general would be the warmest in RCP8.5, whereas precipitation on average would be the highest in RCP 6.0 (Fig. 2a, b). The precipitation in RCP4.5 had a similar trend as that in RCP6.0, but the temperature was lower in the second half of the century. In RCP2.6, the temperature appeared relatively stationary and stayed the lowest beyond 2050.

Population Density Data

We collected the historical records of mean human population density in 2000 from Gridded Population of the World (GPW), v3 database (<http://sedac.ciesin.columbia.edu/data/collection/gpw-v3>, last accessed 09-June-2015). The range of population density over the entire CHR was from 0 to 1023 km⁻² and the mean population density is 28.2 km⁻². Spatially specific future population density across the CHR was estimated

based on the current CHR population density and the overall population trend in the United States in year 2000 for 2025, 2050, 2075 and 2100 (see Supplemental Information, Fig. S1) (Bos *et al.*, 1994), assuming a constant CHR–nationwide population ratio.

Forest Dynamics

We studied forest dynamics and successional patterns using the matrix model, a type of ecological population model that uses transition matrices to estimate the dynamics of structured populations (e.g., Caswell, 2001, Fieberg & Ellner, 2001, Liang & Picard, 2013). Developed from ecological studies that date back to the 1940's (Leslie, 1945, Lewis, 1942), matrix models have been widely employed to study the dynamics of forest ecosystems all over the world due to their accuracy and robustness in depicting structured forest populations (see Liang & Picard, 2013 and references therein).

A conventional matrix model predicts the structured population dynamics of forest stands from time t to $t+1$:

$$\mathbf{y}_{t+1} = \mathbf{G} \cdot \mathbf{y}_t + \mathbf{R} + \varepsilon \quad (1)$$

where $\mathbf{y}_t = [y_{ijt}]$ is a column vector representing the number of live trees per hectare of species i ($i = 1, 2, \dots, m$) and diameter class j ($j = 1, 2, \dots, n$) at time t and ε is a random error. $\mathbf{G}$ is a state- and time-dependent transition matrix describing how trees grow or die between t and $t+1$.

In a Climate-Sensitive Matrix (CSMatrix) model, an extension of the conventional matrix model to control for the climate variability (Liang *et al.*, 2011), $\mathbf{G}$ and $\mathbf{R}$ are revised to be functions of mean annual temperature (T) and annual average of monthly mean precipitation (P). Thus, Eq. (1) was extended to control for climate-sensitivity by recognizing effects of temperature and precipitation on tree growth, mortality, and recruitment as follows:

$$\mathbf{y}_{t+1} = \mathbf{G}_t(T, P) \cdot \mathbf{y}_t + \mathbf{R}_t(T, P) + \varepsilon \quad (2)$$

For the CSMatrix model, the time increment unit—the interval in year between t and $t+1$ of Eq. (2)—was one year. The detailed matrix growth model is described in the Supplemental Information.

The dependent variables, i.e., the rates of upgrowth, mortality, and recruitment (see Supplemental Information for definitions), were estimated from tree and stand attributes for 17 size classes and seven species groups using repeated measurements of 5,196 FIA permanent sample plots. For parsimony and accuracy, we only retained variables that met three rigorous criteria: statistical significance, expected biological responses, and contribution to the model goodness-of-fit. To avoid compromised type-I error rates and severe artifacts commonly associated with model selection procedures (Mac Nally, 2000), explanatory variables were selected using hierarchical partitioning (HP) by the average independent contribution of each variable to the overall goodness-of-fit (Chevan & Sutherland, 1991). The HP analysis was conducted with the hier.part package of the R program (Mac Nally & Walsh, 2004).

Accuracy of the resulting model was evaluated through short-term prediction errors of the validation plots, defined as the difference between the observed and the predicted stand state variables at the second inventory. Predicted stand state variables were obtained by setting the first inventory as the initial condition and applying Eq. 2 iteratively over the elapsed period between the two inventories.

We further compared the stand state variables predicted by the present model with those predicted with the conventional matrix model and LANDIS PRO. For a fair comparison, an independent post-sample validation dataset (Fig. 1) was used for the simulations. For each model, the root mean squared errors (RMSE) (Wooldridge, 2012) were calculated based on the

difference between the predicted and observed basal area by species and diameter class as a comparable measure of accuracy.

Fire Disturbances

To take into account impacts of altered wildfire regimes induced by climate change, we adopted a mean fire interval (MFI) model by Guyette *et al.* (2010) specifically designed for the fire management plans in the eastern and southern United States. The historical MFI model is parameterized using mean maximum temperature, precipitation, their interaction, and estimated population density to account for climate change as a main driver of fire regimes. The detailed MFI model structure and definition of fire severity classes are explained in the Supplemental Information.

We followed the conventional assumption that fire occurrence had an exponential distribution with its probability at time t ,

$$p(t) = 1 - e^{-t / MFI} \quad (3)$$

In our simulations, $p(t)$ was drawn as a uniformly distributed random variable ranging from 0 to 1. Thus the year of fire occurrence, t , is calculated by $-\ln(1-p(t))MFI$. MFI was updated every decade based on the projected climate data and every 25 years based on projected population density data from 2010 to 2100 for all four climate scenarios, respectively. The following probabilities were assigned to fire severity classes **1** to **5** (see Supplemental Information, Table S6), respectively, based on the rationale that lower severity fires tended to happen more frequently: 40, 25, 20, 10, and 5%.

Integrated Framework

It is important to represent interactions among the components to comprehensively assess how landscapes in this region may respond to climate change. Such an integrated model has the

potential to provide resource managers the ability to better visualize potential future landscapes resulting from the interaction of biological and physical processes. In this study, we developed an integrated CSMatrix framework to couple (1) the model of forest dynamics, (2) the mean fire interval model, (3) the population density model, and (4) future climate scenarios. We synchronously coupled the three models and future climate data to evaluate the long-term effects of changes in climate and fire regime on forest structure in the CHR, represented as the average predicted values from Monte Carlo simulations (Fig. 3). The simulated changes in landscape structure have important implications for the management of natural resources in eastern United States.

Results

Forest Dynamics

Starting from a large amount of explanatory variables (see Supplemental Information, Equations S4-S7), we selected a subset to be in the final model (Table 2) based on statistical and biological significance and contribution to the goodness-of-fit. The primary control variables for the final matrix model, DBH (D), stand basal area (B), and stem density (N), were significant at the $\alpha \leq 0.05$ level and contributed highly to the overall goodness-of-fit (see Supplemental Information, Table S4). All other selected variables had independent contributions greater than the average and most were significant and had consistent signs over different species groups.

Trees grew significantly faster when they were larger for all species groups, except for Gymnosperms (GS). Fitted mortality equations indicated that the probability of dying declined with tree diameter for most species, consistent with previous results (Lin *et al.*, 1998, Schulte & Buongiorno, 2004, Zenner, 2005). Mortality rate decreased significantly with stand basal area.

The parameters in the recruitment equations from maximum-likelihood estimation showed that the recruitment of a species increased strongly with the density of that species in the stand. Total number of trees (N) was the most significant predictor of recruitment and its effect was consistent over most species groups.

For the 1,107 validation plots, the basal area by species and diameter class predicted by the final model fell within the 95% confidence interval of the observed values in all the species-diameter classes, demonstrating a high accuracy of the CSMatrix model (Fig. 4).

Compared with the conventional matrix model and LANDIS PRO, our final model had 9.2 – 16.3% lower RMSE than the conventional matrix model, and LANDIS PRO had the lowest short-term accuracy, with 49.1 – 266% higher RMSE than the CSMatrix model across all the species (Fig. 4).

DBH, square DBH, stand basal area, and site productivity were the most significant variables for the volume models (see Supplemental Information, Table S7). The explanatory variables presented here accounted for 93.8 – 97.0% of the variability in stem volume across all species.

Simulations with Fire and Climate Scenarios

In all four climate scenarios, mean fire intervals were predicted to decrease considerably in 90 years, relative to the one predicted with 2010 climate (Fig. 2c). RCP8.5 had the shortest interval, slightly over 4 years, while RCP2.6 the longest, ~6 years, as it approached 2100. RCP4.5 and RCP6.0 had the average intervals, ~5 years, over the next 90 years.

Absent of fire disturbance, climate change increased the total stand basal area but the effect appears to be limited (Fig. 5) and mixed for individual species groups (Supplemental Information Figs.S2 – S8). When both climate change and associated fire disturbances were

simultaneously accounted for, based on 100, 000 simulations, the projected forest states over the next 90 years displayed completely different patterns. Total stand basal area under the four RCP scenarios declined dramatically over the first 70 years by approximately 50 percent, and converged to around $14.9 \text{ m}^2 \text{ ha}^{-1}$ for RCP2.6, followed by $14.2 \text{ m}^2 \text{ ha}^{-1}$ for RCP4.5, $14.0 \text{ m}^2 \text{ ha}^{-1}$ for RCP6.0, and the lowest $13.0 \text{ m}^2 \text{ ha}^{-1}$ for RCP8.5 (Fig. 5). The current average total basal area throughout the CHR is $29.1 \text{ m}^2 \text{ ha}^{-1}$, but would decline by 55.1 – 62.0% to $11.1 - 12.7 \text{ m}^2 \text{ ha}^{-1}$ by 2100 given climate and fire predictions.

In terms of species composition in percentage basal area, current forests in the CHR are dominated by oak (QQ 22.5%, QL 20.5%) and hickory (JD 15.4%) species. Maple species (SD) account for only 8.9% of the total stocking (Fig. 6a). Under climate change and associated alterations of fire regimes, forests in the CHR in year 2100 would be dominated by maple species (SD 55.2 – 60.4%), whereas oak species (QQ, QL) would diminish to 25.5 – 31.5%. GS and JD together would only account for 3.0 – 3.4% of the total stand stocking (Fig. 6b). These potential changes were consistent across all four climate change scenarios (Fig. 6b). Tree size diversity would increase by 27.9 – 30.6% over the first 20 years, and gradually drop back to a level similar to the present (Fig. 7). In contrast, tree species diversity would monotonically decrease by 9.6 – 11.5% over the next 90 years (Fig. 7).

Similarly, total stand volume exhibited a steep decline during the first 50 years of the model run and, despite some fluctuations, converged to $330.4 \text{ m}^3 \text{ ha}^{-1}$ for RCP2.6, followed by $313.1 \text{ m}^3 \text{ ha}^{-1}$ for RCP4.5, $310.1 \text{ m}^3 \text{ ha}^{-1}$ for RCP6.0, and $307.5 \text{ m}^3 \text{ ha}^{-1}$ for RCP8.5 (Table 3). Total volume stocking would decline by 56.3 – 62.4% from $633.5 \text{ m}^3 \text{ ha}^{-1}$ currently to $257.6 - 276.8 \text{ m}^3 \text{ ha}^{-1}$ throughout the region. OA had the greatest volume stocking decline ($191.6 - 203.6 \text{ m}^3 \text{ ha}^{-1}$), followed by JD ($51.3 - 52.5 \text{ m}^3 \text{ ha}^{-1}$), QL ($45.6 - 47.8 \text{ m}^3 \text{ ha}^{-1}$), and QQ ($37.6 - 41.2 \text{ m}^3$

ha⁻¹), whereas SD (-11.9 – 15.3 m³ ha⁻¹) had the least reduction. Assuming constant stumpage prices in 2015 dollars, the total current stumpage value throughout the CHR is approximately \$1,317 billion. Our projected change of forest population structure would lead to a 54.5 – 59.8% reduction in the total value to \$529 – 599 billion in current dollars over the next 90 years.

Discussion

According to the post-sample validation, the CSMatrix model is the most accurate for gymnosperms (RMSE=0.89) and maple species (RMSE=1.87), and is the least accurate for beech (RMSE=3.57) and white oak species (RMSE=3.03). The results suggested that the growth of beech and white oak species may depend on some factors other than those already controlled for by the CSMatrix model and, upon the emergence of additional data, efforts to improve the current model should focus these two species groups. Nevertheless, the CSMatrix model was accurate over all the species and diameter classes because the predicted means were all within the 95% confidence interval of the observed means. In terms of RMSE, the CSMatrix model also demonstrated a higher accuracy than its predecessors.

Our study has addressed the direct and indirect effects of climate change on forest dynamics in the CHR. The direct effects refer to the influence of climate change on upgrowth, ingrowth, and mortality of trees. The indirect effects refer to the climate-induced changes in fire disturbance and consequent influences on forest dynamics. Contrary to the prevailing view that climate change will boost forest growth (Boisvenue & Running, 2006, Pretzsch *et al.*, 2014), we only found slight differences between total stand basal area predicted under constant climate and any of the four climate scenarios. This result seems more in line with research suggesting that warmer temperature and prolonged growing seasons may not have a strong positive effect on

forest growth in southeastern U.S. (Melillo *et al.*, 2014) and tropical regions (van der Sleen *et al.*, 2014). The indirect effect of fire was in fact much greater in magnitude and substantially decreased the total stand basal area (Fig. 5). This may be attributable to the smaller predicted changes in precipitation (-0.5% to 12.7%) and temperature (6.6% to 29.6%) than fire frequency (26.1% to 52.9%) relative to the baseline climate. In all, our study suggests that, for the CHR study area, climate-induced changes in natural disturbances may have greater impacts on forest dynamics than the direct effects of climate on tree growth. This is consistent with previous findings of the compounding effects of stochastic shocks on ecological processes and the crucial impact of the interaction between stochastic shocks and nonlinearity on shaping the ecosystem dynamics (Zhou & Buongiorno, 2004, Fortin & Langevin, 2012).

Contrary to an earlier projection of an increased tree species richness under climate change in the eastern U.S. (Iverson & Prasad, 2001), our projected 9.6 – 11.5% decline in tree species diversity in the CHR (Fig. 7) is due to the reduction of oak and hickory species caused by altered fire regimes (see Supplemental Information, Fig. S2 – S8). Historically, shade intolerant oak and hickory species in the region were maintained by high-intensity fires that reduced competition and promoted regeneration (Nowacki & Abrams, 2008, Ruffner, 2005). Projected shortening of mean fire intervals (Fig. 2c) indicates that high-intensity fires will be gradually replaced by more frequent low-intensity fires, diminishing advantages to oak and hickory species and leading to a gradual mesophication of the CHR forests and dominance of shade-tolerant maple species in the long term (Nowacki & Abrams, 2008). Our projections, consistent with the overall biodiversity trend observed throughout this region (Alexander & Arthur, 2010, Fralish & McArdle, 2009, Iverson *et al.*, 1997), present a great challenge from climate change to biological conservation in the region: changes in precipitation, temperature, and associated fire regimes will

lead to a decline in tree species richness that may translate to a widespread decline in overall plant and ecosystem diversity.

The projected replacement of dominant oak and hickory species by maple species in the CHR would have profound ecological and economic impacts (Nowacki & Abrams, 2008). The shift in tree dominance will alter fundamental ecosystem processes, as litter inputs and chemistry, throughfall quantity and quality, and soil nutrient cycling change in response to species replacement. Within forest soils, dominant tree species influence soil processes and nutrient cycling through their influence on microbial communities, litter quantity and quality, and direct weathering of soil minerals (Lambers *et al.*, 2009). The different soil symbiotic mycorrhizal fungi that form obligate associations with tree roots control many of these processes. Oaks and hickories form symbioses with ectomycorrhizal (ECM) fungi, whereas maples form arbuscular mycorrhizal (AM) associations. These associations differ extensively in their influences on soil nutrient cycling and scavenging processes (Phillips *et al.*, 2013), and these activities influence leaf litter quality and subsequent decomposition (Cornelissen *et al.*, 2001, Hobbie *et al.*, 2006). In the current case, oaks and their associated ECM fungi produce litter with high C:N ratio and low calcium (Ca) availability, which limits decomposition by soil microbes and invertebrates (Fox *et al.*, 2010, Hobbie *et al.*, 2006). In contrast, litter in sugar maple-influenced stands exhibits low C:N ratios and elevated Ca availability in soils relative to many other CHR species (Dijkstra, 2003, Dijkstra & Fitzhugh, 2003). This trend in localized N and Ca enrichment may function to enhance decomposition and nitrification, leading to elevated nutrient losses from watersheds draining the CHR (Christ *et al.*, 2002, Piatek *et al.*, 2010).

Fundamental shifts in tree species will alter habitat structure — canopy light, temperature, moisture, and litter and decaying wood—ultimately affecting broader community characteristics

(Alexander & Arthur, 2010, Martin *et al.*, 2011). Soil and arboreal invertebrate communities differ between oak and maple forests (Huebner *et al.*, 2012), and changes from oak to maple-dominated communities will affect not only nutrient cycling facilitated by invertebrates but changes in habitat and higher-level trophic interactions within the forest as well, resulting in distribution changes among wildlife dependent on these habitat characteristics (Fox *et al.*, 2010, Huebner *et al.*, 2012, Rodewald & Abrams, 2002, Summerville *et al.*, 2003, Wilhelm, 1991). Oak and hickory are keystone species that provide essential food and habitat for many wildlife species (Fralish, 2004). In the eastern United States, an estimated 44 wildlife species, the majority being birds, are highly dependent on oak and hickory trees, which provide more abundant food resources for wildlife relative to maple due to a higher amount of mast, i.e., nuts, buds, twigs (Fralish, 2004). For most terrestrial species, such as the Southern flying squirrel (*Glaucomys volans*), projected transition in dominant tree species would lead to declines in habitat availability from the loss of oak/hickory forests. This species' diet consists primarily of nuts and acorns (Helmick *et al.*, 2014, Thomas & Weigl, 1998), especially in the winter (Saunders, 1998). The transition would also affect avian species, as average abundance of neotropical migrants is higher in oak forests compared to mesic sites (Sierzega & Eichholz, 2014). However, the response may not be universal, as some species, such as the Kentucky warbler (*Oporornis formosus*), may instead increase in abundance in more mesic sites. Fire can also influence habitat choices by wildlife. The Indiana bat (*Myotis sodalis*), a federally-listed endangered species, prefers to roost in hickory trees. However, roosting preferences can change when fire occurs (Johnson *et al.*, 2010). In unburned forests, Indiana bats will roost in hickories, oaks, or maples. When a forest has been burned, however, the Indiana bat prefers to roost in fire-killed maples.

In addition to ecological changes accompanying keystone species shifts, climate change and associated changes in tree species composition and productivity will affect the long-term production of timber within the CHR. Such loss of timber production, especially among oak species, would have profound implications for forest management and economic vibrancy in the region. In order to reduce potentially large losses in the future, landowners and managers must adapt their management practices to the predicted changes. Alternate harvesting plans or rotation lengths would immediately be called for in response to the emergence of new dominant species (Spittlehouse & Stewart, 2004). Forest operations must take into account warmer and more variable climates with more frequent fires. Forest industries should prepare for changes in wood quality and size induced by species shifts. The predicted reduction in timber volume could presage a certain degree of transition of employment from traditional timber management and production to the provision of nontimber forest products, especially forest farming that produces high-value specialty crops for culinary and medicinal purposes, such as mushrooms and ginseng (<http://nac.unl.edu/practices/forestfarming.htm>, accessed September 15, 2015). As a result, the regional economy would likely be impacted significantly. Natural resource policies that promote active adaptation and risk management are thus in urgent need for this region. It remains a subject of debate if fire intervention and suppression is cost-effective and ecologically beneficial (Hand *et al.*, 2015), therefore caution must be used when making policy recommendations regarding wildland fire management.

The success of forest migration, or management plans to mitigate climate change impacts on forested ecosystems, will depend not only on tree species' capacity to move geographically but also on the suitability of soils and soil microbial communities for invasion. In the current study, the large change in MFI frequency within the CHR may potentially be constrained by

edaphic factors existing throughout this region. Maples have greater Ca requirement/demand than oaks and soil Ca is typically higher in soils associated with sugar maple (Christ *et al.*, 2002, Dijkstra & Fitzhugh, 2003). Soil acidification in the eastern regions of the CHR over the past half century has compromised the capacity of these areas to sustain sugar maple (Elias *et al.*, 2009, Fenn *et al.*, 2006, St. Clair *et al.*, 2008). Areas further west may be less impacted (Huntington, 2000), and the capacity of sugar maple to thrive throughout the CHR may depend on the soil base status of potentially invaded sites. Finally, differences in the reproductive systems between the ECM fungi and AM fungi may also limit the migration capacity of maple (Wilkinson 1998). Thus, the health and function of a future forest will depend on soil chemical and microbiological factors needed to support migrating species establishment within these evolving systems.

The CSMatrix model, like all other empirical models, has its inherent limitations, and our results should be interpreted in an appropriate context. In this study, we used the variance of temperature and precipitation across space as a surrogate for the temporal variance in climatic conditions to analyze the effects of climate change on CHR forests. Therefore, there would be an extrapolation bias should future climatic conditions move beyond the range of temperatures and precipitation used herein (see Supplemental Information, Table S2). Because the estimated future climatic conditions over the next 90 years (Fig. 2a, b) largely match the current range, our 90-year simulation is minimally affected by the extrapolation bias. However, any long-term projection results beyond 90 years using the CSMatrix model should be interpreted with caution. Although the CSMatrix model did not account for other natural disturbances except for wildfire, our results offer a striking illustration of forest population dynamics under climate change and

associated change of fire regimes, and may be of unprecedented value to policy making in the United States as well as many ongoing and upcoming ecological studies.

Conclusion

In summary, we developed a Climate-Sensitive Matrix model to estimate forest dynamics as a function of climate, fire disturbance, and shifting forest population structures in diverse ecosystems for the CHR of the U.S. Under four IPCC climate change scenarios and accompanying alterations of fire regimes, we projected that the total stand basal area would decline dramatically by approximately 60% over the first 70 years, and converge to around 14.9 m² ha⁻¹ for RCP2.6, followed by 14.2 m² ha⁻¹ for RCP4.5, 14.0 m² ha⁻¹ for RCP6.0, and the lowest 13.0 m² ha⁻¹ for RCP8.5. Similarly, total stand volume had a steep decline during the first 50 years and, despite some fluctuations, converged to 330 m³ ha⁻¹ for RCP2.6, followed by 313 m³ ha⁻¹ for RCP4.5, 310 m³ ha⁻¹ for RCP6.0, and 308 m³ ha⁻¹ for RCP8.5. Tree size diversity increased by 27.9 – 30.6% over the first 20 years, and then decreased to a level similar to the present. In contrast, tree species diversity would gradually decline by 9.6 – 11.5% over the next 90 years. These changes in forest structure within the CHR will have regional ecological and economic repercussions. Changes in diversity favoring maples will alter ecosystem processing of nutrients and subsequent nutrient flows to drainage waters within the region. Habitat change will alter the broad spectrum of organisms relying on the forest, with concomitant changes in ecosystem-wide biodiversity. Finally, changes in species composition, stocking, and productivity of CHR forests will reduce the direct and indirect economic benefits generated by timber in the CHR. One uncertainty to these projections is the capacity of soil chemical and microbiological factors needed to support migrating forest tree species establishment within these evolving

ecosystems. Such preferences and influences may function to structure communities in new unforeseen ways and may further influence forest ecosystem development under future climate regimes.

The challenge to traditional forestry as well as to the economy brought upon by climate change is by no means unique to the CHR. Globally, changes have been observed or predicted. Response to the expected changes and associated risks summons actions from both private and public sectors. Landowners and managers need to adjust and adapt their practices while public policies must promote and facilitate such adjustments and adaptations in order to sustain both the forests and their related ecosystem services in the future.

Acknowledgements

This study is supported in parts by the Davis College of Agriculture, Natural Resources & Design, West Virginia University, under the US Department of Agriculture (USDA) McIntire–Stennis Funds WVA00105. We thank the USDA Forest Service for providing FIA data, and J. Shanmugasundaram for assisting with climatic analysis.

REFERENCES

- Aber J, Neilson RP, McNulty S, Lenihan JM, Bachelet D, Drapek RJ (2001) Forest processes and global environmental change: predicting the effects of individual and multiple stressors. *BioScience*, **51**, 735–751.
- Albrecht MA, McCarthy BC (2006) Effects of prescribed fire and thinning on tree recruitment patterns in central hardwood forests. *Forest Ecology and Management*, **226**, 88–103.
- Alexander HD, Arthur MA (2010) Implications of a predicted shift from upland oaks to red maple on forest hydrology and nutrient availability. *Canadian Journal of Forest Research*, **40**, 716–726.
- Allen CD, Macalady AK, Chenchouni H *et al.* (2010) A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Forest Ecology and Management*, **259**, 660–684.
- Bailey RG (2004) Identifying ecoregion boundaries. *Environmental Management*, **34**, S14–S26.
- Bernhardt ES, Palmer MA (2011) The environmental costs of mountaintop mining valley fill operations for aquatic ecosystems of the Central Appalachians. *Annals of the New York Academy of Sciences*, **1223**, 39–57.
- Boisvenue C, Running SW (2006) Impacts of climate change on natural forest productivity - evidence since the middle of the 20th century. *Global Change Biology*, **12**, 862–882.
- Bos E, Vu M, Massiah E, Bulatao R (1994) World population projections 1994–95: estimates and projections with related demographic statistics. *New York: Johns Hopkins University Press*.
- Box EO, Fujiwara K (2015) Warm-temperate deciduous forests around the northern hemisphere, *Springer*.

550 Brooks JR (2012) Central Appalachian hardwood whole stand growth and yield models.
 551 *Unpublished manuscript, Appalachian Hardwood Forest Research Alliance Technical*
 552 *Report.*

553 Brooks JR, Miller GW (2011) An evaluation of three growth and yield simulators for even-aged
 554 hardwood forests of the mid-Appalachian region. In: *Proceedings of the 17th Central*
 555 *Hardwood Forest Conference GTR-NRS-P, 78, 23 p.*

556 Brose PH, Dey DC, Phillips RJ, Waldrop TA (2013) A meta-analysis of the fire-oak hypothesis:
 557 does prescribed burning promote oak reproduction in eastern North America? *Forest*
 558 *Science, 59, 322-334.*

559 Bush RR (1995) Northeast (NE) variant overview: forest vegetation simulator. *Washington, DC:*
 560 *U.S. Department of Agriculture, Forest Service, 37 p.*

561 Cardinale BJ, Duffy JE, Gonzalez A *et al.* (2012) Biodiversity loss and its impact on humanity.
 562 *Nature, 486, 59-67.*

563 Caswell H (2001) Matrix population models. *John Wiley & Sons, Ltd.*

564 Chevan A, Sutherland M (1991) Hierarchical partitioning. *American Statistician, 45, 90-96.*

565 Christ MJ, Peterjohn WT, Cumming JR, Adams MB (2002) Nitrification potentials and
 566 landscape, soil and vegetation characteristics in two Central Appalachian watersheds
 567 differing in NO₃⁻ export. *Forest Ecology and Management, 159, 145-158.*

568 Colbert JJ, Racin G (1995) User's guide to the stand damage model: a component of the gypsy
 569 moth life system model (Version 1.1). *General Technical Report NE-207. Radnor, PA:*
 570 *U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station,*
 571 *38 p.*

572 Cornelissen J, Aerts R, Cerabolini B, Werger M, Van Der Heijden M (2001) Carbon cycling
 573 traits of plant species are linked with mycorrhizal strategy. *Oecologia*, **129**, 611-619.
 574 Daly C, Halbleib M, Smith JI *et al.* (2008) Physiographically sensitive mapping of climatological
 575 temperature and precipitation across the conterminous United States. *International*
 576 *Journal of Climatology*, **28**, 2031-2064.
 577 Dijkstra FA (2003) Calcium mineralization in the forest floor and surface soil beneath different
 578 tree species in the northeastern US. *Forest Ecology and Management*, **175**, 185-194.
 579 Dijkstra FA, Fitzhugh RD (2003) Aluminum solubility and mobility in relation to organic carbon
 580 in surface soils affected by six tree species of the northeastern United States. *Geoderma*,
 581 **114**, 33-47.
 582 Elias PE, Burger J, Adams M (2009) Acid deposition effects on forest composition and growth
 583 on the Monongahela National Forest, West Virginia. *Forest Ecology and Management*,
 584 **258**, 2175-2182.
 585 Fenn ME, Huntington T, McLaughlin S, Eagar C, Gomez A, Cook R (2006) Status of soil
 586 acidification in North America. *Journal of Forest Science*, **52**, 3-13.
 587 Fieberg J, Ellner SP (2001) Stochastic matrix models for conservation and management: a
 588 comparative review of methods. *Ecology Letters*, **4**, 244-266.
 589 Fischer A, Marshall P, Camp A (2013) Disturbances in deciduous temperate forest ecosystems of
 590 the northern hemisphere: their effects on both recent and future forest development.
 591 *Biodiversity and Conservation*, **22**, 1863-1893.
 592 Foster DR (2000) From bobolinks to bears: interjecting geographical history into ecological
 593 studies, environmental interpretation, and conservation planning. *Journal of*
 594 *Biogeography*, **27**, 27-30.

595 Fox VL, Buehler CP, Byers CM, Drake SE (2010) Forest composition, leaf litter, and songbird
 596 communities in oak-vs. maple-dominated forests in the eastern United States. *Forest*
 597 *Ecology and Management*, **259**, 2426-2432.

598 Fortin M, Langevin L (2012) Stochastic or deterministic single-tree models: is there any
 599 difference in growth predictions?. *Annals of Forest Science*, **69**, 271-282.

600 Fralish JS (2004) The keystone role of oak and hickory in the central hardwood forest. In:
 601 Spetich, M.A. (ed) Upland oak ecology symposium: history, current conditions, and
 602 sustainability. *General Technical Reports*. SRS-73. Asheville, NC pp. 78-87.

603 Fralish JS, Mcardle TG (2009) Forest dynamics across three century-length disturbance regimes
 604 in the Illinois Ozark Hills. *The American Midland Naturalist*, **162**, 418-449.

605 Guyette RP, Stambaugh MC, Dey DC (2010) Developing and using fire scar histories in the
 606 southern and eastern United States. *Joint Fire Science Program Final Report, web site:*
 607 *www.firescience.gov*.

608 Hann W, Shlisky A, Havlina D *et al.* (2004) Interagency fire regime condition class guidebook.
 609 *Last update October 2007: Version 1.3. [Homepage of the Interagency and The Nature*
 610 *Conservancy fire regime condition class website, USDA Forest Service, U.S. Department*
 611 *of the Interior, The Nature Conservancy, and Systems for Environmental Management]*.
 612 *[Online]. Available: www.frcc.gov*.

613 Hansen A, Dale V (2001) Biodiversity in US forests under global climate change. *Ecosystems*, **4**,
 614 161-163.

615 Hansen AJ, Neilson RP, Dale VH, Flather C, Iverson L, Currie DJ, Bartlein P (2001) Global
 616 change in forests: responses of species, communities, and biomes. *BioScience*, **51**, 765-
 617 779.

618 Hand MS, Wibbenmeyer MJ, Calkin DE, Thompson MP (2015) Risk preferences, probability
619 weighting, and strategy tradeoffs in wildfire management. *Risk Analysis*, **35**, 1876-1891.

620 He HS, Wang WJ, Fraser JS, Shifley SR, Larsen DR (2012) LANDIS: a spatially explicit model
621 of forest landscape disturbance, management, and succession. *LANDIS PRO 7.0 USERS*
622 *GUIDE*.

623 Helmick KR, Barrett TL, Barrett GW (2014) Dietary resource preference of the Southern flying
624 squirrel (*Glaucomys volans*). *The American Midland Naturalist*, **171**, 371-374.

625 Hicks RR (1998) Ecology and management of central hardwood forests, *John Wiley & Sons*.

626 Hilt DE (1985) OAKSIM: an individual-tree growth and yield simulator for managed, even-aged,
627 upland oak stands. *Reserch Paper. NE-562. Broomall, PA: U.S. Department of*
628 *Agriculture, Forest Service, Northeastern Forest Experiment Station*, 21 p.

629 Hobbie SE, Reich PB, Oleksyn J, Ogdahl M, Zytowskiak R, Hale C, Karolewski P (2006) Tree
630 species effects on decomposition and forest floor dynamics in a common garden. *Ecology*,
631 **87**, 2288-2297.

632 Huebner K, Lindo Z, Lechowicz M (2012) Post-fire succession of collembolan communities in a
633 northern hardwood forest. *European Journal of Soil Biology*, **48**, 59-65.

634 Huntington TG (2000) The potential for calcium depletion in forest ecosystems of southeastern
635 United States: review and analysis. *Global Biogeochemical Cycles*, **14**, 623-638.

636 Blyth W, Yang M, Bradley R (2007) Climate policy uncertainty and investment risk. *OECD*
637 *Publishing*.

638 IPCC (2013) Summary for policymakers. In: climate change 2013: the physical science basis.
639 *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental*

Panel on Climate Change Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.

Iverson LR, Dale ME, Scott CT, Prasad A (1997) A GIS-derived integrated moisture index to predict forest composition and productivity of Ohio forests (USA). *Landscape Ecology*, **12**, 331-348.

Iverson LR, Prasad AM (2001) Potential changes in tree species richness and forest community types following climate change. *Ecosystems*, **4**, 186-199.

Johnson JB, Ford WM, Rodrigue JL, Edwards JW, Johnson CM (2010) Roost selection by male Indiana myotis following forest fires in central Appalachian hardwoods forests. *Journal of Fish and Wildlife Management*, **1**, 111-121.

Juday GP, Barber VA, Duffy P *et al.* (2005) Forests, land management, and agriculture. In: *Arctic Climate Impact Assessment*. (eds Symon C, Arris L, Heal B) pp. 781-862. New York, NY, Cambridge University Press.

Jiang Y, Zhuang Q, Mandallaz D (2012) Modeling large fire frequency and burned area in Canadian terrestrial ecosystems with Poisson models. *Environmental Modeling & Assessment*, **17**, 483-493.

Keane RE, Parsons R, Hessburg P (2002) Estimating historical range and variation of landscape patch dynamics: limitations of the simulation approach. *Ecological Modeling*, **151**, 29-49.

Lafon CW, Hoss JA, Grissino-Mayer HD (2005) The contemporary fire regime of the central Appalachian Mountains and its relation to climate. *Physical Geography*, **26**, 126-146.

Lambers H, Mougél C, Jaillard B, Hinsinger P (2009) Plant-microbe-soil interactions in the rhizosphere: an evolutionary perspective. *Plant and Soil*, **321**, 83-115.

662 Latta G, Temesgen H, Adams D, Barrett T (2010) Analysis of potential impacts of climate
663 change on forests of the United States Pacific Northwest. *Forest Ecology and*
664 *Management*, **259**, 720-729.

665 Leslie PH (1945) The use of matrices in certain population mathematics. *Biometrika*, **33**, 183-
666 212.

667 Lewis EG (1942) On the generation and growth of a population. *Sankhya*, **6**, 93-96.

668 Liang J, Picard N (2013) Matrix model of forest dynamics: an overview and outlook. *Forest*
669 *Science*, **59**, 359-378.

670 Liang J, Zhou M, Tobin PC, McGuire AD, Reich PB (2015) Biodiversity influences plant
671 productivity through niche–efficiency. *Proceedings of the National Academy of Sciences*,
672 **112**, 5738-5743.

673 Liang J, Zhou M, Verbyla D, Zhang L, Springsteen AL, Malone T (2011) Mapping forest
674 dynamics under climate change: a matrix model. *Forest Ecology and Management*, **262**,
675 2250-2262.

676 Lin CR, Buongiorno J, Prestemon J, Skog K (1998) Growth model for uneven-aged loblolly pine
677 stands. *USDA. Forest Service Reserch Paper. FPL-RP-569. Madison*.

678 Lindner M, Maroschek M, Netherer S *et al.* (2010) Climate change impacts, adaptive capacity,
679 and vulnerability of European forest ecosystems. *Forest Ecology and Management*, **259**,
680 698-709.

681 Mac Nally R (2000) Regression and model-building in conservation biology, biogeography and
682 ecology: the distinction between—and reconciliation of—‘predictive’ and ‘explanatory’
683 models. *Biodiversity and Conservation*, **9**, 655-671.

684 Mac Nally R, Walsh C (2004) Hierarchical partitioning public-domain software. *Biodiversity*
685 *Conservation*, **13**, 659-660.

686 Marquis DA, Ernst RL (1992) User's guide to SILVAH: stand analysis, prescription, and
687 management simulator program for hardwood stands of the Alleghenies. *General*
688 *Technical Report. NE-162. Radnor, PA: U.S. Department of Agriculture, Forest Service,*
689 *Northeastern Forest Experiment Station*, 124 p.

690 Martin KL, Hix DM, Goebel PC (2011) Coupling of vegetation layers and environmental
691 influences in a mature, second-growth central hardwood forest landscape. *Forest Ecology*
692 *and Management*, **261**, 720-729.

693 Mcewan RW, Dyer JM, Pederson N (2011) Multiple interacting ecosystem drivers: toward an
694 encompassing hypothesis of oak forest dynamics across eastern North America.
695 *Ecography*, **34**, 244-256.

696 Mcewan RW, Hutchinson TF, Long RP, Ford DR, McCarthy BC (2007) Temporal and spatial
697 patterns in fire occurrence during the establishment of mixed-oak forests in eastern North
698 America. *Journal of Vegetation Science*, **18**, 655-664.

699 Melillo JM, Richmond TC, Yohe GW (2014) Climate change impacts in the United States: the
700 third national climate assessment. *US Global Change Research Program*, 841.

701 Morgan P, Hardy CC, Swetnam TW, Rollins MG, Long DG (2001) Mapping fire regimes across
702 time and space: understanding coarse and fine-scale fire patterns. *International Journal of*
703 *Wildland Fire*, **10**, 329-342.

704 Mueller RF (1996) Biodiversity: central Appalachian plant distributions and forest types. *Wild*
705 *Earth*, **6**, 37-43.

706 Naeem S, Duffy JE, Zavaleta E (2012) The functions of biological diversity in an age of
 707 extinction. *Science*, **336**, 1401-1406.

708 Nowacki GJ, Abrams MD (2008) The demise of fire and “mesophication” of forests in the
 709 eastern United States. *Bioscience*, **58**, 123-138.

710 Parisen MA, Moritz MA (2009) Environmental controls on the distribution of wildfire at
 711 multiple spatial scales. *Ecological Monographs*, **79**, 127-154.

712 Perkey AW (1985) GROAK (Grow OAK) on a computer. *Northern Journal of Applied Forestry*,
 713 **2**, 99-100.

714 Phillips RP, Brzostek E, Midgley MG (2013) The mycorrhizal-associated nutrient economy: a
 715 new framework for predicting carbon–nutrient couplings in temperate forests. *New*
 716 *Phytologist*, **199**, 41-51.

717 Piatek KB, Munasinghe P, Peterjohn WT, Adams MB, Cumming JR (2010) A decrease in oak
 718 litter mass changes nutrient dynamics in the litter layer of a Central Hardwood Forest.
 719 *Northern Journal of Applied Forestry*, **27**, 97-104.

720 Pretzsch H, Biber P, Schütze G, Uhl E, Rötzer T (2014) Forest stand growth dynamics in Central
 721 Europe have accelerated since 1870. *Nature Communications*, **5**.

722 Rodewald AD, Abrams MD (2002) Floristics and avian community structure: implications for
 723 regional changes in eastern forest composition. *Forest Science*, **48**, 267-272.

724 Ruffner CM (2005) Understanding the evidence for historical fire across eastern forests. *Fire in*
 725 *Eastern Oak Forests: Delivering Science to Land Managers. Proceedings of a*
 726 *Conference. GTR-NRS-P-1*.

727 Saunders DA (1998) Adirondack mountains. *State University of New York, College of*
 728 *Environmental Science and Forestry*, 216 p.

729 Schmidt TL, McWilliams WH (2003) Shifts and future trends in the forest resources of the
 730 Central Hardwood region. *United States Department of Agriculture Forest Service*
 731 *General Technical Report NC*, pp 21-31.

732 Schoene DHF, Bernier PY (2012) Adapting forestry and forests to climate change: A challenge
 733 to change the paradigm. *Forest Policy and Economics*, **24**, 12-19.

734 Schuler TM, Thomas-Van Gundy M, Adams MB, Ford WM (2012) Analysis of two pre-
 735 shelterwood prescribed fires in a mesic mixed-oak forest in West Virginia. In:
 736 *Proceedings of the 18th Central Hardwood Forest Conference*. pp 26-28.

737 Schulte BJ, Buongiorno J (2004) A growth and yield model for naturally-regenerated mixed
 738 shortleaf pine forests in the southern United States of America. *International Forestry*
 739 *Review*, **6**, 19-29.

740 Sharon W, Barbara L, Barbara M *et al.* (2011) The forest inventory and analysis database:
 741 database description and users manual version 5.1 for phase 2.

742 Shugart H, Sedjo R, Sohngen B (2003) Forests and global climate change: potential impacts on
 743 U.S. forest resources. *Pew Center on Global Climate Change, Arlington, Virginia*. 52 p.

744 Sierzega KP, Kevin P, Eichholz MW (2014) Factors influencing avian habitat selection between
 745 oak-hickory and mesic forests in southern Illinois. In: *Proceedings of the 19th Central*
 746 *Hardwood Forest Conference, Groninger, John W., Holzmueller, Eric J, Nielsen,*
 747 *Clayton K., Dey, Daniel C., eds.* pp 102-103.

748 Signell SA, Abrams MD, Hovis JC, Henry SW (2005) Impact of multiple fires on stand structure
 749 and tree regeneration in central Appalachian oak forests. *Forest Ecology and*
 750 *Management*, **218**, 146-158.

751 Smith TM, Halpin PN, Shugart HH, Secrett CM (1995) 6 Global forests. In: As climate changes:
 752 international impacts and implications. (eds Strzepek KM, Smith JB) 146 p. *Cambridge,*
 753 *UK, Cambridge University Press.*

754 Spittlehouse DL, Stewart RB (2004) Adaptation to climate change in forest management.
 755 *Journal of Ecosystems and Management*, **4**, 1-11.

756 St. Clair SB, Sharpe WE, Lynch JP (2008) Key interactions between nutrient limitation and
 757 climatic factors in temperate forests: a synthesis of the sugar maple literature. *Canadian*
 758 *Journal of Forest Research*, **38**, 401-414.

759 Summerville KS, Crist TO, Kahn JK, Gering JC (2003) Community structure of arboreal
 760 caterpillars within and among four tree species of the eastern deciduous forest.
 761 *Ecological Entomology*, **28**, 747-757.

762 Thomas RB, Weigl PD (1998) Dynamic foraging behavior in the southern flying squirrel
 763 (*Glaucomys volans*): test of a model. *The American Midland Naturalist*, **140**, 264-270.

764 Tilman D, Lehman CL, Thomson KT (1997) Plant diversity and ecosystem productivity:
 765 theoretical considerations. *Proceedings of the National Academy of Sciences*, **94**, 1857-
 766 1861.

767 Van der Sleen P, Groenendijk P, Vlam M, Anten NP, Boom A, Bongers F, Pons TL, Terburg G,
 768 Zuidema PA (2014) No growth stimulation of tropical trees by 150 years of CO₂
 769 fertilization but water-use efficiency increased. *Nature Geoscience*, **8**, 24-28.

770 Wang WJ, He HS, Fraser JS, Iii FRT, Shifley SR, Spetich MA (2014) LANDIS PRO: a
 771 landscape model that predicts forest composition and structure changes at regional scales.
 772 *Ecography*, **37**, 226–230

773 Westerling AL, Hidalgo HG, Cayan DR, Swetnam TW (2006) Warming and earlier spring
 774 increase western U.S forest wildfire activity. *Science*, **313**, 940-943.
 775 Wilhelm GS (1991) Implications of changes in floristic composition of the Morton Arboretum's
 776 East Woods. In: *Proceedings of the Oak Woods Management Workshop. Eastern Illinois*
 777 *University, Charleston*. pp 31-54.
 778 Wooldridge JM (2012) Introductory econometrics: a modern approach. *Cengage Learning*.
 779 Zenner EK (2005) Development of tree size distributions in Douglas-fir forests under differing
 780 disturbance regimes. *Ecological Applications*, **15**, 701-714.
 781 Zhou M, Buongiorno J (2004) Nonlinearity and noise interaction in a model of forest
 782 growth. *Ecological Modelling*, 180, 291-304.
 783 Zhou M, Buongiorno J (2006) Forest landscape management in a stochastic environment, with
 784 an application to mixed loblolly pine-hardwood forests. *Forest Ecology and Management*,
 785 **223**, 170-182.

786 **Table 1.** Definitions and units of variables used in the CSMatrix model for CHR forests.

Variable	Unit	Definition/explanation
<i>B</i>	m ² ha ⁻¹	Total stand basal area
<i>C</i>	m ³ ha ⁻¹ yr ⁻¹	Site productivity
<i>D</i>	cm	Diameter at breast height
<i>g</i>	cm yr ⁻¹	Annual diameter growth
<i>E</i>	km	Plot elevation
<i>S</i>	degrees	Plot slope
<i>H_d</i>		Tree size diversity in Shannon's index
<i>H_s</i>		Tree species diversity in Shannon's index
<i>T</i>	°C	Mean annual temperature
<i>P</i>	100 mm month ⁻¹	Annual average of monthly mean precipitation
<i>m</i>	yr ⁻¹	Annual tree mortality
<i>N</i>	trees ha ⁻¹	Number of trees per hectare
<i>R</i>	trees ha ⁻¹ yr ⁻¹	Recruitment, the number of trees per hectare growing into the smallest diameter class (2.54-7cm) in a year
<i>V</i>	m ³ ha ⁻¹	Stem volume
<i>MFI</i>	year	Mean fire interval

787

788

789

790

791

792

793

794

795

796

797

798 **Table 2.** Estimated parameters of the CSMatrix model for CHR forests.

	QQ		QL		JD		SD		GS		FG		OA	
	Diameter growth													
Const	0.066	***	0.202	***	0.051	***	0.077	***	0.181	***	0.492	***	0.0619	***
D	0.006	***	0.007	***	0.009	***	0.004	*	-0.001	*	0.004	*	0.0194	***
D ²	-0.0001	***	-0.0001	***	-0.0001	***			-0.0001	*	-0.0001	***	-0.0004	***
H _d											-0.167	***		
P*D	0.003	***	0.001	*	0.001	*	0.002		-0.003	*	0.001	*	0.0026	**
P*D ²						*	-0.0001	*	0.0001	*			-0.0001	
P*B			0.018		-0.016									
T*D	0.0002	***	0.0001		0.00004		0.001	***	0.001	***	0.0003	*	-0.0006	***
T*D ²							-0.0001	*	-0.0001	*			0.00002	***
T*B			-0.002	*	0.001									
	Mortality													
Const	-0.418	***	-0.723	***	-0.570	***	-1.200	***	0.066	*	-1.465	***	-0.5454	***
D	-0.007		0.044	***	-0.051	***								
D ²	0.001	***			0.001	***								
B	-1.372	***	-1.631	***	-1.494	***	-0.953	***					-0.9055	***
H _s									-0.665	***				
H _d											-0.285			
S											0.018	*		
C											0.035	*		
P*D	-0.021	***	-0.039	***										
P*B	0.567	***	0.899	***	0.459	***	0.290	**					0.1885	***
P*H _s									0.079	*				
P*S											-0.002			
P*C														
T*D	-0.002	**												
T*B	0.036	*	0.025	**	0.030	**	0.029	*					0.0098	
T*H _s									-0.004	*				
T*S											-0.001	*		
	Recruitment													
Const	32.102		-79.636	***	-36.633	***	-80.766	***	-0.010	***	-0.012	***	0.1220	
N	0.186	***	0.313	***	0.121	***	0.166	***	0.189	**	0.212		0.1020	***
N ²	-0.046	***	-0.004		-0.022	***	-0.041	***			-0.463	*	0.0008	
H _d	-18.360	***												
H _s	-20.975				3.572									
P	-65.815	*											-	0.5110
P ²	6.071													0.0001
P*N	-0.035	*	-0.131	***	0.010	*	-0.021	*	-0.028	*	-0.011	*	-0.0114	
P*H _s	15.286													
P*H _d					-25.734	***								
T*N	-0.004	*	-0.004	*	-0.004	*	-0.002		-0.005		0.008		-0.0053	**
T*N ²			-0.004								0.023	*		
T*P													-0.8770	

Note: Significance levels: *≤0.05; **≤0.01; ***≤0.001.

Table 3. Total average volume (m³ ha⁻¹) for climate scenarios RCP2.6, RCP4.5, RCP6.0, RCP8.5 from year 2010-2100.

Year	RCP2.6	RCP4.5	RCP6.0	RCP8.5
<i>2010-2020</i>	461.1	468.7	448.5	443.6
<i>2020-2030</i>	394.6	392.1	367.2	380.9
<i>2030-2040</i>	369.6	355.4	339.4	352.4
<i>2040-2050</i>	352.2	320.2	327.4	325.0
<i>2050-2060</i>	330.4	313.1	316.7	307.5
<i>2060-2070</i>	316.1	294.0	296.8	286.2
<i>2070-2080</i>	307.0	283.6	297.8	270.5
<i>2080-2090</i>	292.7	270.3	269.7	257.8
<i>2090-2100</i>	276.8	257.6	261.7	238.2

1

2

3 **Figure 1.** Geographic distribution of the calibration (dots) and validation (+) plots, in the Central
 4 Hardwood Region (CHR). Inset shows the relative location of CHR in the contiguous United
 5 States. The box indicates the CHR region where the future relative changes of temperature and
 6 precipitation (trend ratio) were extracted for four future climate scenarios.

Figure 2. Temporal changes of mean annual temperature (°C), annual average of monthly mean precipitation (100 mm month⁻¹) and mean fire interval (year) across the 1107 sample plots estimated under four climate scenarios of RCP2.6, RCP4.5, RCP6.0, and RCP8.5, and the climate constant at year 2010.

Figure 3. Conceptual diagram showing how the CSMatrix framework integrated forest dynamics, climate data, and the Mean Fire Interval Model to project the population dynamics of CHR forests under future climate change scenarios.

Figure 4. Average predicted and observed stand states at the second inventory for the conventional matrix model (CON), climate-sensitive matrix model (CS), and LANDIS PRO 7.0 on the 1107 post-sample validation plots, with the 95% confidence interval of the observed mean.

Figure 5. Stand basal area under wildfire disturbance under four climate scenarios RCP2.6, RCP4.5, RCP6.0, and RCP8.5 from year 2010-2100.

Figure 6. Species composition in terms of basal area in year 2010 (a) and year 2100 under wildfire disturbance with four climate scenarios RCP2.6, RCP4.5, RCP6.0, and RCP8.5 (b). Vertical bars represent one standard error of the average predicted values from Monte Carlo simulations. The pie charts to the right show the corresponding percentage of basal area by species and climate scenario.

Figure 7. Tree species (a-d) and size diversity (e-h) under wildfire disturbance with 4 climate scenarios RCP2.6, RCP4.5, RCP6.0, and RCP8.5 from year 2010-2100.