Biotransformation of Tetracycline by a Novel Bacterial Strain *Stenotrophomonas maltophilia* DT1

Yifei Leng^{1, 4}, Jianguo Bao^{1,*}, Gaofeng Chang², Han Zheng¹, Xingxing Li¹, Jiangkun Du¹, Daniel Snow³, and Xu Li^{4,*}

Affiliations:

1. School of Environment Studies, China University of Geosciences, Wuhan 430074, PR China
2. Tianjin Environmental Protection Technical Development Center, Tianjin 300191, PR China
3. Water Sciences Laboratory, University of Nebraska-Lincoln, Lincoln, Nebraska, 68583, USA
4. Department of Civil Engineering, University of Nebraska-Lincoln, Lincoln, Nebraska, 68588, USA

***Corresponding Authors**

Jianguo Bao
bjianguo@cug.edu.cn

Xu Li
xuli@unl.edu

ABSTRACT

Although several abiotic processes have been reported that can transform antibiotics, little is known about whether and how microbiological processes may degrade antibiotics in the environment. This work isolated one tetracycline degrading bacterial strain, *Stenotrophomonas maltophilia* strain DT1, and characterized the biotransformation of tetracycline by DT1 under various environmental conditions. The biotransformation rate was the highest when the initial pH was 9 and the reaction temperature was at 30°C, and can be described using the Michaelis-Menten model under different initial tetracycline concentrations. When additional substrate was present, the substrate that caused increased biomass resulted in a decreased biotransformation rate of tetracycline. According to disk diffusion tests, the biotransformation products of tetracycline had lower antibiotic potency than the parent compound. Six possible biotransformation products were identified, and a potential biotransformation pathway was proposed that included sequential removal of N-methyl, carbonyl, and amine function groups. Results from this study can lead to better estimation of the fate and transport of antibiotics in the environment and has the potential to be utilized in designing engineering processes to remove tetracycline from water and soil.

Keywords: Tetracycline; biotransformation; hydrolysis; *Stenotrophomonas maltophilia*; transformation products.

1. INTRODUCTION

Tetracycline is a broad-spectrum antibiotic widely used for human and animal health [1-3]. More than 6,800 tons of tetracycline are used worldwide in animal agriculture [4, 5]. Because antibiotics cannot be absorbed or metabolized entirely by the animal, a reported 30 - 90% of antibiotics administered are excreted into the environment via urine and feces [6]. Residues of tetracycline have been detected in wastewater [7-9], surface water [10-12], groundwater [10, 11], river sediments[13], and soil [14, 15]. Tetracycline compounds and their transformation products can disrupt the structure and function of natural microbial communities [15, 16] and cause the emergence of antibiotic resistance among bacteria [4, 17].

Tetracycline compounds can be transformed through abiotic processes in the environment. Tetracycline can quickly hydrolyze in water [18], and the process is affected by temperature, pH, ionic strength, and the chelation with cations [5, 19]. Tetracycline is also sensitive to light, and some transformation products from photolysis are more toxic than the parent compound [20].

Biotransformation is an important yet poorly understood process. It can lead to the formation of transformation products different from those resulting from abiotic processes [5]. Earlier studies reported the removal of tetracycline from water by mixed microbial communities in river sediment [21] and in nitrifying granular sludge [22], a process often characterized as a combination of quick sorption and slow biotransformation [23]. Existing pure culture studies mostly focus on fungal species. For example, *Pleurotus ostreatus* mycelium is proven to be capable of converting oxytetracycline in soil into less toxic or

nontoxic substances [24]. Besides, laccase from the white rot fungus *Trametes versicolor* can transform tetracycline and eliminate its ecotoxicity [25]. The lignin and manganese peroxidase from the white rot fungus *Phanerochaete chrysosporium* can also effectively transform tetracyclines [26, 27], although the toxicity of the transformation products was unknown. Pure cultures of bacterial strains capable of degrading tetracycline compounds have also been reported [28, 29]; however, little is known about the kinetics or products of specific bacterial transformation of tetracycline compounds.

The objective of this study was to characterize the biotransformation of tetracycline by a novel bacterial strain *Stenotrophomonas maltophilia* strain DT1. In this study, the biotransformation kinetics of strain DT1 was determined under various temperature, pH, and background carbon conditions. Chemical and biotransformation products were identified and potential biotransformation pathways are proposed. Finally, the antimicrobial potency of the biotransformation products were determined. Results from this study can lead to better estimation of the fate and transport of antibiotics in the environment and has the potential to be utilized in designing engineering processes to remove tetracycline from water and soil.

2. MATERIALS AND METHODS

2.1. Chemicals and Solutions

All chemicals were purchased from Aladdin (Shanghai, China), including analytical grade tetracycline, and high performance liquid chromatography (HPLC) grade methanol and acetonitrile. Lysogeny broth (LB) medium was prepared using 10 g L⁻¹ tryptone, 5 g L⁻¹ yeast extract, and 5 g L⁻¹ NaCl. Mineral medium (MM) was made using 1.5 g L⁻¹ K₂HPO₄, 0.5 g L⁻¹

KH₂PO₄, 1.0 g L⁻¹ NaCl, and 0.2 g L⁻¹ MgSO₄·7H₂O (pH 7.0)[30]. MM amended with 50 mg L⁻¹ 0.22 µm nylon filter sterilized (Restek Corp., USA) tetracycline (MM-T), MM supplemented with 10 g L⁻¹ casein peptone (MM-P), and MM-P amended with 50 mg L⁻¹ 0.22 µm nylon filter sterilized tetracycline (MM-PT) were also prepared.

2.2. Isolation of Tetracycline Degrading Bacteria

In order to enrich tetracycline degrading bacteria in soil, 1 L water containing 400 mg L⁻¹ tetracycline was poured onto ~1 meter² grassland within the China University of Geosciences (Wuhan, China) once a week in Fall 2013. Four months later, 4 g of top soil was placed in 100 mL LB medium containing 50 mg L⁻¹ tetracycline. The solution was incubated at 30°C on a shaker set at 150 rpm in the dark. After three days of incubation, the enrichment culture were plated on MM-PT agar plates and grown at 30°C in the dark. Three days later eight colonies were further purified using the streak plate method on MM-PT agar for 1 day. Single colonies were used to inoculate fresh MM-PT solutions. The solutions were incubated at 30°C on a shaker set at 150 rpm in the dark, and the tetracycline concentration in the solution was monitored using HPLC.

2.3. Identification of the Strain

Among the eight isolates, strain DT1 exhibited the ability to biotransform tetracycline. Strain DT1 was identified according to the Bergey's Manual of Systematic Bacteriology [31] and the phylogenetic analysis using the 16S rRNA gene[32]. Genomic DNA of the strain was extracted using the Genomic DNA Extraction Kit (Axygen, USA) and amplified using a PCR protocol targeting the 16S rRNA gene [33]. The PCR amplicon was purified using the PCR Clean Up Kit (Shanghai, China), sequenced on a Applied Biosystems

3730-XL DNA analyzer, and deposited under the accession number KP996643 at GenBank.

Phylogenetically related sequences were downloaded from NCBI, and aligned with the sequence of DT1 using ClustalX 1.8.3 with default settings [34]. An unrooted phylogenetic tree was constructed using the Neighbor-Joining method with bootstrap values estimated from 1,000 replicated analyses [35] in MEGA version 5.10 [36].

2.4. Tetracycline Biotransformation

Biotransformation experiments were conducted in batch reactors. Strain DT1 was grown in LB medium containing 20 mg L⁻¹ tetracycline at 30°C on a shaker set at 150 rpm. After reaching the mid-exponential phase, cells were harvested and washed twice with sterilized physiological saline (0.9 % NaCl). Cells were adjusted to OD_{600nm} = 1.00 prior to being added to 50 mL MM-PT (1%, v/v) in a 250 mL flask[37]. Four series experiments were designed based on results from preliminary experiments. First, five temperatures (i.e., 20, 25, 30, 35, 40°C) were tested with an initial pH 9.0. Second, five initial pH values (i.e., 6.0, 7.0, 8.0, 9.0, 10.0) were tested at 30°C. Third, five initial tetracycline concentrations (i.e., 10, 25, 50, 75, 100, 200 mg L⁻¹) were tested with an initial pH 9.0 at 30°C. Finally, effects of four background substrates (i.e., glucose, glycerol, lactose, or sucrose, all at 10 g L⁻¹) on the transformation of tetracycline were assessed. These substrates were previously used in biotransformation studies on DDT [38], azo dyes [39] and iopromide [40]. For all four series, (1) the initial tetracycline concentration was 50 mg L⁻¹ unless stated otherwise; (2) all tests were performed in triplicates; (3) a blank control without bacteria was included in all tests; (4) all flasks were covered in aluminum foil to prevent photo degradation; and (5) liquid samples were collected daily for 7 days and the concentration of the parent compound was

measured using HPLC.

2.5. Quantification of Tetracycline using HPLC

To measure tetracycline, 1 mL solution from each batch reactor was centrifuged at 8,000 g at 4°C for 10 min. Mcilvaine-Na₂EDTA buffer was added to the sample supernatant in an equal volume fashion to chelate metal ions [41], and the mixture was filtered through 0.22 µm filters prior to being preserved at -20°C. Stored samples were loaded onto a Shimadzu LC-06A system. A C18 reversed-phase column (4.6×250 mm, 5 µm, Agilent Technologies) was operated at 40°C, with a 1 mL min⁻¹ mobile phase consisting of 67% (v/v) 0.1 M ammonium oxalate in high purity water, 22% (v/v) acetonitrile and 11% (v/v) methanol. The injection volume was 20 µL, and the column isocratic elution was monitored by a UV detector at 355 nm.

2.6. Modeling

In degradation experiments containing strain DT1, the decrease in tetracycline concentration was attributed to both biotransformation and hydrolysis. In control experiments containing no strain DT1, the decrease in tetracycline concentration was attributed to hydrolysis only. Both the overall degradation process (Equation A) and the hydrolysis process (Equation B) can be described using first-order kinetics. The change of tetracycline concentration due to biotransformation (Equation C) equals the observed overall degradation (experiments containing the bacterial strain DT1) minus the change of tetracycline concentration due to hydrolysis (control experiments containing no bacterial cells).

$$C_{HB} = C_0 - C_0 \times e^{-k_{HB} \times T} \quad (A)$$

$$C_H = C_0 - C_0 \times e^{-k_H \times T} \quad (B)$$

$$C_B = C_{HB} - C_H = C_0 \times e^{-k_H \times T} - C_0 \times e^{-k_{HB} \times T} \quad (C)$$

C_0 is the initial concentration of tetracycline (mg L^{-1}), k_{HB} is the first-order reaction rate constant of the overall degradation reaction (day^{-1}), k_H is the first-order reaction rate constant of hydrolysis (day^{-1}), T is the time (day), C_{HB} is the concentration change of tetracycline from the overall degradation (C_{HB} was measured from the reactors containing strain DT1), C_H is the concentration change of tetracycline due to hydrolysis only (C_H was measured from the abiotic control reactors), C_B is the concentration change of tetracycline due to biotransformation (mg L^{-1}). The values of k_H and k_{HB} were listed in Table S1. Because there was a 3-day delay in bacteria growth when the initial pH was 6 (Fig. S3), Equation D was modified from Equation C to describe the biotransformation kinetics for the experiment with an initial pH 6.

$$C_B = C_0 \times e^{-k_H \times (T-3)} - C_0 \times e^{-k_{HB} \times (T-3)} \quad (D)$$

After taking the derivative on both sides of Equation C over time, biotransformation rate (v_B , $\text{mg L}^{-1} \text{d}^{-1}$) can be calculated as

$$v_B = C_0 \times (k_H \times e^{-k_H \times T} - k_{HB} \times e^{-k_{HB} \times T}) \quad (E)$$

In the experiment with various initial tetracycline concentrations, because the biotransformation rate varied with time, the initial rates (i.e., the biotransformation rate at $T=0$) was used as representative biotransformation rates to see if the experimental data can fit the Michaelis-Menten model:

$$v = \frac{v_{\max} \times S}{K_m + S} \quad (F)$$

Where v is the initial biotransformation rate (d^{-1}), $v_{\max}$ is the maximum initial biotransformation rate (d^{-1}), K_m (mg L^{-1}) is half saturation constant, and S is the initial concentration of tetracycline (mg L^{-1}). Non-linear regression analysis was conducted using

the software Origin 9.0.

2.7. Antibacterial Activities of Tetracycline Transformation Products

The antibacterial activities of tetracycline and its transformation products were measured using the disk diffusion method [42]. *Escherichia coli* cells were grown in nutrient broth at 30°C to the stationary phase. Cells were diluted to OD_{600nm} = 1.00 and spread on nutrient agar plates. Wells were made on agar plate surface and filled with 300 µL solution collected from various phases of the treatment reactors containing DT1 and control reactors containing no bacterial cells. Plates were incubated at 30°C for 18 hr. Inhibition zones around the wells were measured in mm (including well diameter). The measurements were done for all three replicate transformation experiments.

2.8. Identification of the Transformation Products

Three sets of reaction conditions were evaluated: 1) MM-T solution containing strain DT1 (hydrolysis plus biotransformation), 2) MM-T solution containing no bacteria (hydrolysis), and 3) MM solution (no tetracycline). Solutions from each reaction condition were collected on Day 0 and Day 3, and the transformation products from the three conditions were identified using LC tandem mass spectrometry Q Exactive hybrid Quadrupole-Orbitrap mass spectrometer (Thermo scientific, Bremen, Germany). Aqueous samples were extracted using solid-phase extraction cartridges (Oasis HLB, 6cc/150mg, Waters) as described in previous studies [41, 43]. Separation was performed on a C18 column (4.6×250 mm, 5 µm, Agilent Technologies) with an injection volume of 5 µL. Flow rate was set at 0.3 mL min⁻¹ at the isocratic mode for 30min. Mobile phase consisted of 0.1% formic acid, acetonitrile, and methanol at 67:22:11. Mass spectra were processed using the Xcalibur

2.1 software (Thermo Scientific). The mass accuracy accepted for the experiments was estimated to be below 5 ppm. The MS analysis was performed with a heated electrospray ionization (HESI) source in positive mode with a spray voltage of 3.5 kV, S-lens RF level of 50%, and a capillary temperature of 300°C. The MS acquisition was performed in full scan mode 50-1000 Da with a mass resolution of 70000 [41, 43-45]. The molecular structure for tetracycline and their transformation products were tentatively proposed by the detection of predicted mass, changing in full scan ion intensity, and ring double bond equivalent number (RDB) from the reacted sample (Table 1).

3. RESULTS

3.1. Strain Isolation and Identification

A total of 8 isolates were obtained from the top soil that was exposed to tetracycline for 4 months. Strain DT1 was selected for further analyses due to its ability to biotransform tetracycline. Based on the 16S rRNA gene sequence, strain DT1 was identified as a *Stenotrophomonas maltophilia* strain (Fig. S1). *S. maltophilia* strain DT1 was determined to be aerobic and Gram-negative. Its colonies were round, shiny, light yellow on LB agar. It could grow on MM-T agar plate which contained 50 mg L⁻¹ tetracycline. Growth was observed over a temperature range of 20-40°C, a pH range of 6.0–10.0, and a salinity range of 0-5 % NaCl. It showed positive results in the methyl red test, in the catalase/oxidase/urease test, in indole production and nitrate reduction, and showed negative results in gelatin hydrolysis and Voges–Proskauer test.

3.2. Characteristics of Tetracycline Degradation

Because the biotransformation experiments were conducted in aqueous solution, hydrolysis was an intrinsic part of the experiment. Hence, biotransformation cannot be directly measured in this study. In order to assess biotransformation, for all of the experimental conditions tested we included a degradation experiment that contained strain DT1 and a control experiment that contained no bacterial cells. The differences in the tetracycline profiles between the two experiments were attributed to biotransformation (Fig. 1). Both the overall degradation and the hydrolysis followed first order reaction kinetics. The fastest overall degradation appeared at 30°C, where 44.43 mg L⁻¹ tetracycline was degraded in 4 days. The changes of tetracycline concentrations due to hydrolysis and due to combined hydrolysis and biotransformation at other temperatures are plotted in Fig. S2. Hydrolysis rate constants increased with temperature (Fig. 2A insert). The biotransformation rate at time zero, i.e., the initial biotransformation rate, was the highest (Fig. 2B). The initial biotransformation rate was the highest at 30°C with a value of 22.69 mg L⁻¹d⁻¹ (Fig. 2B insert). Specific biotransformation rates, i.e., biotransformation rates normalized to biomass, could not be calculated because strain DT1 formed biofilm on the inner wall of the reactors and made it impossible to quantify bacterial biomass.

The hydrolysis and the biotransformation of tetracycline were also affected by solution pH (Fig. S3 and Fig. 3). The fastest hydrolysis occurred when the initial pH was 10 (Fig. 3A). The hydrolysis rate constant increased with the increasing initial pH (Fig. 3A insert). The highest initial biotransformation rate was at pH 9 (Fig. 3B insert). At the end of the degradation experiment, the final pH of all five tests was close to 9 (Fig. S4). There was a 3-day lag phase in the biotransformation of tetracycline at initial pH 6 (Fig. 3B).

3.3. Effects of Initial Tetracycline Concentrations

The Michaelis-Menten model (Equation F) could satisfactorily describe the biotransformation kinetics of tetracycline by DT1 as a function of initial tetracycline concentration under 100 mg L^{-1} (Fig. 4). Based on non-linear regression analysis, $v_{\max}$ and K_m equal $62.14 \text{ mg L}^{-1}\text{d}^{-1}$ and 80.52 mg L^{-1} , respectively.

3.4. Effects of Background Carbon Substrates

In general, the addition of carbon substrates in the background increased cell density and reduced tetracycline biotransformation rate (Fig. 5). For example, when glucose was present, the cell density was more than twice the control and the biotransformation rate was much lower than the control. Glycerol was an exception. The presence of glycerol had no evident impact on either cell density or biotransformation rate (Fig. 5).

3.5. Transformation Products

The biotransformation products exhibited lower antimicrobial potency than the parent compound. In the treatment reactors where both hydrolysis and biotransformation occurred, residual tetracycline dropped from 50.00 mg L^{-1} to 2.23 mg L^{-1} by Day 7 (Fig. 6). The antimicrobial potency of the treatment reactors were contributed from residual tetracycline and transformation products from hydrolysis and biotransformation. The antimicrobial potency decreased, as shown in the decrease of the diameter of inhibitory zone from 19.0 mm to 10.1 mm (Fig. 6). In comparison, in the control reactors where only hydrolysis occurred, residual tetracycline dropped from 50.00 mg L^{-1} to 24.93 mg L^{-1} by Day 7. The antimicrobial potency of the control reactors were from residual tetracycline and transformation products from hydrolysis. The diameter of inhibitory zones only decreased slightly from 19.0 mm to

17.0 mm (Fig. 6).

A peak corresponding to either an epimer or isomer of tetracycline was identified (Fig. S5 and Table 1) in Day 3 of the experiment. Tetracycline eluted at 7.42 min with an m/z value of 445.1603 for its protonated form, while the peak for either 4-epi-tetracycline (ETC) or isotetracycline (ISO-TC) ^{5, 45}, had the same m/z value but an earlier retention time (i.e., 6.51 min). Previous work in aqueous systems at near neutral pH suggests that formation of ISO-TC is unusual [19, 46].

Six possible biotransformation products were also identified (Fig. S5, Fig. S6 and Table 1) and the mass spectra of the parent compound and proposed transformation products are shown in Fig. S7 to S14. A putative biotransformation pathway of tetracycline by *S. maltophilia* DT1 was proposed in Fig. 7. Formation of compound TP 431 could occur when the N-(CH₃)₂ group on tetracycline C-4 is de-methylated. Formation of TP 415 may occur via dehydrogenation. After losing a carbonyl group, TP 415 may be converted to compound TP 387. Finally, compound TP 370 may form when an amine group was further removed from compound TP 387 (Fig. 7). Biotransformation may also start with de-methylation at C-4 of ISO-TC and form compound ISO-TP 415. After losing a carbonyl group, compound ISO-TP 415 was converted to compound ISO-TP 387. It is noted that the intensity of ISO-TP 415 and ISO-TP 387 was lower than TP 415 and TP 387 (Table 1).

4. DISCUSSION

Different from sulfonamide compounds, whose degradation by pure bacterial cultures [47-49] and microbially mediated abiotic processes [50] have been reported, little is known

about how microbial processes may transform tetracycline compounds. Several *Stenotrophomonas* strains have demonstrated the ability to degrade organic pollutants, such as pesticide [51], insecticide [52, 53], BTEX [54], PAH [55], and steroid hormones [56]. These strains can transform toxic pollutants by opening their loop structure or by cutting functional groups. For example, *Stenotrophomonas sp.* THZ-XP can biotransform the pesticide acetamiprid by cutting its cyano substituent [52].

We noticed that although strain DT1 could form colonies on MM-T agar plates, they could not grow in MM-T liquid medium. This suggests that strain DT1 may not be able to use tetracycline as sole carbon and energy source. The growth of strain DT1 on MM-T agar plates might be due to the utilization of the trace amount of organic matters in agar. The biotransformation of tetracycline by strain DT1 might be attributed to the detoxification mechanism, which relies on the flavin-dependent monooxygenase encoded by *tet(X)* [57].

Temperature and pH can influence the hydrolysis of tetracycline. Similar to the findings from a previous study [19], this study demonstrated that high temperatures accelerated tetracycline hydrolysis. According to the collision theory, higher temperatures lead to higher reaction rates by causing more collisions between particles [58]. High pH also accelerated the hydrolysis of tetracycline. Tetracycline is an amphoteric molecule with three dissociation constants ($pK_a=3.3, 7.7, 9.7$) [59]. As pH increased from 6 to 10, different reactive species with various degrees of ionization might have dominated the solution and led to different hydrolysis rates [19].

For the temperature and pH tested (except pH 6), biotransformation increased in the first four days before leveling off (Fig. 2 and 3). Despite the buffer system in the solution,

strain DT1 changed the final pH of the solution to ~9 at the end of the experiment regardless the initial pH values (Fig. S4). Bacterial cells can break down peptone in the reaction medium and release alkaline compounds such as ammonia and amines, which can cause increase in pH [60].

Background organic compounds can provide microbes with additional sources of energy, nutrients, and growth factors, but may also trigger carbon catabolite repression. In this study, the background substrates that promoted biomass growth lowered the tetracycline biotransformation rate (Fig. 5). This may be explained by carbon catabolite repression, which is a general regulatory phenomenon. After bacteria adapt to a preferred (i.e., rapidly biodegradable) carbon and energy source, the synthesis of enzymes involved in utilizing other carbon and energy sources would be inhibited. For example, the presence of succinate or glucose lead to substantial biomass growth, but also caused decreased phenol biodegradation in *Pseudomonas putida* [61]. In the presence of these background organics, phenol hydroxylase was hardly induced, because the activation function of its activator protein PhlR was interfered [61]. Glycerol cannot be utilized by *S. maltophilia* [31], which can explain why the presence of glycerol had no effect on cell density and tetracycline biotransformation rate.

Previous studies showed that the tetracycline transformation products from photolysis had higher toxicity than the parent compound [20, 62]. In contrast, the tetracycline transformation products from fungal laccase had lower toxicity than the parent compound [25]. This study also showed that biotransformation products had lower antimicrobial potencies than the parent compound (Fig. 6). Hence, microbial transformation of tetracycline

may be more desirable from an environmental perspective than physicochemical processes as it reduces the biological activities of the antibiotic.

During hydrolysis, under the action of hydroxide ion from water the hydroxyl group on C6 may attack the carbonyl group on C11 and form iso-tetracycline (ISO-TC) irreversibly in alkaline pH range [63]. Also, tetracycline can undergo reversible epimerization on C4 under certain acidic conditions and form the corresponding 4-epi-tetracycline (ETC) with the participation of hydrogen ions from water at acidic pH conditions [64]. Moreover, tetracycline can turn into ISO-TC or ETC at neutral or weak alkaline conditions (pH =6.5–9), causing decrease in tetracycline concentrations during hydrolysis [5]. The intensity of subsequent transformation products ISO-TP 415 and ISO-TP 387 were nearly an order of magnitude lower than TP 415 and TP 387 (Table 1), suggesting the pathway through ISO-TC was not the dominant pathway.

The loss of the N-methyl group by the demethylation of the dimethylamino group at position C4 from the biotransformation in this study was also observed in previous studies where tetracycline compounds were transformed by potassium ferrate [65], chloramine [66], or manganese oxide [64]. Liu and co-workers reported that oxytetracycline underwent a cleavage at the C1-C12a bond and formed a diradical intermediate, then lost a carbonyl group and formed another diradical, which would close the ring [67]. The loss of the amide group at C2 was also observed in the degradation of tetracycline by laccase [68] and advanced oxidation [67]. The main structure of tetracycline often remains largely intact, because the four ring structure is very stable [13].

5. CONCLUSIONS

In this study, *S. maltophilia* strain DT1 capable of degrading tetracycline was isolated from a contaminated soil. Under pH 9 and 30°C, strain DT1 reached the highest tetracycline degradation kinetics. The biotransformation kinetics can be described by the difference between two first-order kinetics. The presence of easily biodegradable substrates increased the biomass significantly, but it also lowered the biotransformation rate of tetracycline. Compared to hydrolysis, biotransformation was more effective in lowering the antimicrobial potency of the parent compound. Six possible biotransformation products were identified and a potential degradation pathway was presented.

ACKNOWLEDGEMENTS

This study was supported by Natural Science Foundation of China (No. 41373083), and US National Science Foundation (CBET-1351676).

REFERENCES

- [1] R. Li, Y. Zhang, C.C. Lee, L. Liu, Y. Huang, Hydrophilic interaction chromatography separation mechanisms of tetracyclines on amino-bonded silica column, *J Sep Sci*, 34 (2011) 1508-1516.
- [2] X. Ding, S. Mou, Ion chromatographic analysis of tetracyclines using polymeric column and acidic eluent, *J Chromatogr A*, 897 (2000) 205-214.
- [3] P. Nielsen, N. Gyrd-Hansen, Bioavailability of oxytetracycline, tetracycline and chlortetracycline after oral administration to fed and fasted pigs, *J Vet Pharmacol Ther*, 19 (1996) 305-311.
- [4] R. Daghrir, P. Drogui, Tetracycline antibiotics in the environment: a review, *Environ Chem Lett*, 11 (2013) 209-227.
- [5] B. Halling-Sørensen, G. Sengeløv, J. Tjørnelund, Toxicity of tetracyclines and tetracycline degradation products to environmentally relevant bacteria, including selected tetracycline-resistant bacteria, *Arch Environ Con Tox*, 42 (2002) 263-271.
- [6] J. Bound, N. Voulvoulis, Pharmaceuticals in the aquatic environment—a comparison of risk assessment strategies, *Chemosphere*, 56 (2004) 1143-1155.
- [7] T.R. Burch, M.J. Sadowsky, T.M. LaPara, Fate of antibiotic resistance genes and class 1 integrons in soil microcosms following the application of treated residual municipal wastewater solids, *Environ Sci Technol*, 48 (2014) 5620-5627.
- [8] Y. Zhang, C. Zhang, D.B. Parker, D.D. Snow, Z. Zhou, X. Li, Occurrence of antimicrobials and antimicrobial resistance genes in beef cattle storage ponds and swine treatment lagoons, *Sci. Total Environ.*, 463-464C (2013) 631-638.
- [9] Y.P. Zhang, D.D. Snow, D. Parker, Z. Zhou, X. Li, Intracellular and Extracellular Antimicrobial Resistance Genes in the Sludge of Livestock Waste Management Structures, *Environ. Sci. Technol.*, 47 (2013) 10206-10213.
- [10] F. Hernández, J.V. Sancho, M. Ibáñez, C. Guerrero, Antibiotic residue determination in environmental waters by LC-MS, *Trac-trend Anal Chem*, 26 (2007) 466-485.
- [11] J.J. López Peñalver, C.V. Gómez Pacheco, M. Sánchez Polo, J. Rivera Utrilla, Degradation of tetracyclines in different water matrices by advanced oxidation/reduction processes based on gamma radiation, *J Chem Technol Biot*, 88 (2013) 1096-1108.
- [12] C. Fan, J. He, Proliferation of antibiotic resistance genes in microbial consortia of sequencing batch reactors (SBRs) upon exposure to trace erythromycin or erythromycin-H₂O, *Water Res*, 45 (2011) 3098-3106.
- [13] R. Delépée, D. Maume, B. Le Bizec, H. Pouliquen, Preliminary assays to elucidate the structure of oxytetracycline's degradation products in sediments: Determination of natural tetracyclines by high-performance liquid chromatography–fast atom bombardment mass spectrometry, *J Chromatogr B: Biomed Sci Appl*, 748 (2000) 369-381.
- [14] F. Liu, G.-G. Ying, R. Tao, J.-L. Zhao, J.-F. Yang, L.-F. Zhao, Effects of six selected antibiotics on plant growth and soil microbial and enzymatic activities, *Environ Pollut*, 157 (2009) 1636-1642.
- [15] S. Thiele-Bruhn, I.-C. Beck, Effects of sulfonamide and tetracycline antibiotics on soil microbial activity and microbial biomass, *Chemosphere*, 59 (2005) 457-465.

- [16] A.B. Caracciolo, E. Topp, P. Grenni, Pharmaceuticals in the environment: Biodegradation and effects on natural microbial communities. A review, *J Pharmaceut Biomed*, 106 (2015) 25-36.
- [17] E. Michalova, P. Novotna, J. Schlegelova, Tetracyclines in veterinary medicine and bacterial resistance to them. A review, *Vet Med-czech*, (2004).
- [18] Y. Wu, R. Fassihi, Stability of metronidazole, tetracycline HCl and famotidine alone and in combination, *Int J Pharm*, 290 (2005) 1-13.
- [19] K.A. Loftin, C.D. Adams, M.T. Meyer, R. Surampalli, Effects of ionic strength, temperature, and pH on degradation of selected antibiotics, *J Environ Qual*, 37 (2008) 378-386.
- [20] S. Jiao, S. Zheng, D. Yin, L. Wang, L. Chen, Aqueous photolysis of tetracycline and toxicity of photolytic products to luminescent bacteria, *Chemosphere*, 73 (2008) 377-382.
- [21] B.-V. Chang, Y.-L. Ren, Biodegradation of three tetracyclines in river sediment, *Ecol Eng*, 75 (2015) 272-277.
- [22] Y.-J. Shi, X.-H. Wang, Z. Qi, M.-H. Diao, M.-M. Gao, S.-F. Xing, S.-G. Wang, X.-C. Zhao, Sorption and biodegradation of tetracycline by nitrifying granules and the toxicity of tetracycline on granules, *J Hazard Mater*, 191 (2011) 103-109.
- [23] N. Prado, J. Ochoa, A. Amrane, Biodegradation and biosorption of tetracycline and tylosin antibiotics in activated sludge system, *Process Biochem.*, 44 (2009) 1302-1306.
- [24] L. Migliore, M. Fiori, A. Spadoni, E. Galli, Biodegradation of oxytetracycline by *Pleurotus ostreatus* mycelium: a mycoremediation technique, *J Hazard Mater*, 215 (2012) 227-232.
- [25] T. Suda, T. Hata, S. Kawai, H. Okamura, T. Nishida, Treatment of tetracycline antibiotics by laccase in the presence of 1-hydroxybenzotriazole, *Bioresour Technol*, 103 (2012) 498-501.
- [26] X. Wen, Y. Jia, J. Li, Degradation of tetracycline and oxytetracycline by crude lignin peroxidase prepared from *Phanerochaete chrysosporium*—a white rot fungus, *Chemosphere*, 75 (2009) 1003-1007.
- [27] X. Wen, Y. Jia, J. Li, Enzymatic degradation of tetracycline and oxytetracycline by crude manganese peroxidase prepared from *Phanerochaete chrysosporium*, *J Hazard Mater*, 177 (2010) 924-928.
- [28] S. Ghosh, M. Sadowsky, M. Roberts, J. Gralnick, T. LaPara, *Sphingobacterium* sp. strain PM2-P1-29 harbours a functional tet (X) gene encoding for the degradation of tetracycline, *J. Appl. Microbiol*, 106 (2009) 1336-1342.
- [29] G. Volkers, L. Schuldt, G.J. Palm, G.D. Wright, W. Hinrichs, Crystallization and preliminary X-ray crystallographic analysis of the tetracycline-degrading monooxygenase TetX2 from *Bacteroides thetaiotaomicron*, *Acta Crystallogr., Sect. F: Struct. Biol. Cryst. Commun*, 66 (2010) 611-614.
- [30] G. Wang, Y. Zhao, H. Gao, W. Yue, M. Xiong, F. Li, H. Zhang, W. Ge, Co-metabolic biodegradation of acetamiprid by *Pseudoxanthomonas* sp. AAP-7 isolated from a long-term acetamiprid-polluted soil, *Bioresour Technol*, 150 (2013) 259-265.
- [31] L. Mcclung, *Bergey's Manual of Systematic Bacteriology*, Volume 2, *Int J Syst Evol Micr*, 37 (1987) 91-91.

- [32] J.R. Marchesi, T. Sato, A.J. Weightman, T.A. Martin, J.C. Fry, S.J. Hiom, W.G. Wade, Design and evaluation of useful bacterium-specific PCR primers that amplify genes coding for bacterial 16S rRNA, *Appl Environ Microb*, 64 (1998) 795-799.
- [33] M. Goodfellow, E. Stackebrandt, *Nucleic acid techniques in bacterial systematics*, J. Wiley, 1991.
- [34] M.A. Larkin, G. Blackshields, N. Brown, R. Chenna, P.A. McGettigan, H. McWilliam, F. Valentin, I.M. Wallace, A. Wilm, R. Lopez, Clustal W and Clustal X version 2.0, *Bioinformatics*, 23 (2007) 2947-2948.
- [35] K.-H. Chu, L. Alvarez-Cohen, Trichloroethylene degradation by methane-oxidizing cultures grown with various nitrogen sources, *Water Environ Res*, 68 (1996) 76-82.
- [36] K. Tamura, J. Dudley, M. Nei, S. Kumar, MEGA4: molecular evolutionary genetics analysis (MEGA) software version 4.0, *Mol Biol Evol*, 24 (2007) 1596-1599.
- [37] W. Luo, Y. Zhao, H. Ding, X. Lin, H. Zheng, Co-metabolic degradation of bensulfuron-methyl in laboratory conditions, *J Hazard Mater*, 158 (2008) 208-214.
- [38] A.A. Sari, S. Tachibana, K. Itoh, Determination of co-metabolism for 1, 1, 1-trichloro-2, 2-bis (4-chlorophenyl) ethane (DDT) degradation with enzymes from *Trametes versicolor* U97, *J Biosci Bioeng*, 114 (2012) 176-181.
- [39] K. Chen, W. Huang, J. Wu, J. Houn, Microbial decolorization of azo dyes by *Proteus mirabilis*, *J Ind Microbiol Biot*, 23 (1999) 686-690.
- [40] B. Xu, P. Gao, Z. Liu, G. Xue, Y. Liu, F. Wu, Influence of Cosubstrates on Iopromide Degradation by *Pseudomonas* sp. I-24, *Water Air Soil Poll*, 225 (2014).
- [41] X. Wu, Y. Wei, J. Zheng, X. Zhao, W. Zhong, The behavior of tetracyclines and their degradation products during swine manure composting, *Bioresour Technol*, 102 (2011) 5924-5931.
- [42] A. Das, S. Datta, S. Mukherjee, S. Bose, S. Ghosh, P. Dhar, Evaluation of antioxidative, antibacterial and probiotic growth stimulatory activities of *Sesamum indicum* honey containing phenolic compounds and lignans, *LWT--Food Sci Technol*, 61 (2015) 244-250.
- [43] A. Jia, Y. Xiao, J. Hu, M. Asami, S. Kunikane, Simultaneous determination of tetracyclines and their degradation products in environmental waters by liquid chromatography-electrospray tandem mass spectrometry, *J Chromatogr A*, 1216 (2009) 4655-4662.
- [44] M. Llorca, S. Rodriguez-Mozaz, O. Couillerot, K. Panigoni, J. de Gunzburg, S. Bayer, R. Czaja, D. Barcelo, Identification of new transformation products during enzymatic treatment of tetracycline and erythromycin antibiotics at laboratory scale by an on-line turbulent flow liquid-chromatography coupled to a high resolution mass spectrometer LTQ-Orbitrap, *Chemosphere*, 119 (2015) 90-98.
- [45] C. Nebot, M. Guarddon, F. Seco, A. Iglesias, J.M. Miranda, C.M. Franco, A. Cepeda, Monitoring the presence of residues of tetracyclines in baby food samples by HPLC-MS/MS, *Food Control*, 46 (2014) 495-501.
- [46] H. Sanderson, F. Ingerslev, R.A. Brain, B. Halling-Sørensen, J.K. Bestari, C.J. Wilson, D.J. Johnson, K.R. Solomon, Dissipation of oxytetracycline, chlortetracycline, tetracycline and doxycycline using HPLC-UV and LC/MS/MS under aquatic semi-field microcosm conditions, *Chemosphere*, 60 (2005) 619-629.

- [47] E. Topp, R. Chapman, M. Devers-Lamrani, A. Hartmann, R. Marti, F. Martin-Laurent, L. Sabourin, A. Scott, M. Sumarah, Accelerated Biodegradation of Veterinary Antibiotics in Agricultural Soil following Long-Term Exposure, and Isolation of a Sulfamethazine-degrading sp, *J Environ Qual*, 42 (2013) 173-178.
- [48] W. Tappe, M. Herbst, D. Hofmann, S. Koepchen, S. Kummer, B. Thiele, J. Groeneweg, Degradation of sulfadiazine by *Microbacterium lacus* strain SDZm4, isolated from lysimeters previously manured with slurry from sulfadiazine-medicated pigs, *Appl Environ Microb*, 79 (2013) 2572-2577.
- [49] B. Ricken, P.F. Corvini, D. Cichocka, M. Parisi, M. Lenz, D. Wyss, P.M. Martínez-Lavanchy, J.A. Müller, P. Shahgaldian, L.G. Tulli, Ipso-hydroxylation and subsequent fragmentation: a novel microbial strategy to eliminate sulfonamide antibiotics, *Appl Environ Microb*, 79 (2013) 5550-5558.
- [50] J.L. Mohatt, L. Hu, K.T. Finneran, T.J. Strathmann, Microbially mediated abiotic transformation of the antimicrobial agent sulfamethoxazole under iron-reducing soil conditions, *Environ Sci Technol*, 45 (2011) 4793-4801.
- [51] Y.-j. Shen, P. Lu, H. Mei, H.-j. Yu, Q. Hong, S.-p. Li, Isolation of a methyl parathion-degrading strain *Stenotrophomonas* sp. SMSP-1 and cloning of the *ophc2* gene, *Biodegradation*, 21 (2010) 785-792.
- [52] H. Tang, J. Li, H. Hu, P. Xu, A newly isolated strain of *Stenotrophomonas* sp. hydrolyzes acetamiprid, a synthetic insecticide, *Process Biochem*, 47 (2012) 1820-1825.
- [53] Y.-J. Zhao, Y.-J. Dai, C.-G. Yu, J. Luo, W.-P. Xu, J.-P. Ni, S. Yuan, Hydroxylation of thiacloprid by bacterium *Stenotrophomonas maltophilia* CGMCC1. 1788, *Biodegradation*, 20 (2009) 761-768.
- [54] E.Y. Lee, Y.S. Jun, K.-S. Cho, H.W. Ryu, Degradation characteristics of toluene, benzene, ethylbenzene, and xylene by *Stenotrophomonas maltophilia* T3-c, *J Air Waste Manage*, 52 (2002) 400-406.
- [55] A.L. Juhasz, G. Stanley, M. Britz, Metabolite repression inhibits degradation of benzo [a] pyrene and dibenz [a, h] anthracene by *Stenotrophomonas maltophilia* VUN 10,003, *J Ind Microbiol Biot*, 28 (2002) 88-96.
- [56] Z. Li, R. Nandakumar, N. Madayiputhiya, X. Li, Proteomic analysis of 17 β -estradiol degradation by *Stenotrophomonas maltophilia*, *Environ Sci Technol*, 46 (2012) 5947-5955.
- [57] W. Yang, I.F. Moore, K.P. Koteva, D.C. Bareich, D.W. Hughes, G.D. Wright, TetX is a flavin-dependent monooxygenase conferring resistance to tetracycline antibiotics, *J Biol Chem*, 279 (2004) 52346-52352.
- [58] J.E. Leffler, E. Grunwald, Rates and equilibria of organic reactions: as treated by statistical, thermodynamic and extrathermodynamic methods, Courier Corporation, 2013.
- [59] Z. Qiang, C. Adams, Potentiometric determination of acid dissociation constants (pK_a) for human and veterinary antibiotics, *Water Res*, 38 (2004) 2874-2890.
- [60] T. Bergan, J.R. Norris, Methods in microbiology, Academic Press, 1979.
- [61] C. Müller, L. Petruschka, H. Cuypers, G. Burchhardt, H. Herrmann, Carbon catabolite repression of phenol degradation in *Pseudomonas putida* is mediated by the inhibition of the activator protein PhlR, *J Bacteriol*, 178 (1996) 2030-2036.

- [62] R.X. Guo, J.Q. Chen, Phytoplankton toxicity of the antibiotic chlortetracycline and its UV light degradation products, *Chemosphere*, 87 (2012) 1254-1259.
- [63] C. Waller, B. Hutchings, R. Broschard, A. Goldman, W. Stein, C. Wolf, J. Williams, DEGRADATION OF AUREOMYCIN. VII. 1 AUREOMYCIN AND ANHYDROAUREOMYCIN, *J Am Chem Soc*, 74 (1952) 4981-4982.
- [64] W.-R. Chen, C.-H. Huang, Transformation kinetics and pathways of tetracycline antibiotics with manganese oxide, *Environ Pollut*, 159 (2011) 1092-1100.
- [65] Y. Ma, N. Gao, C. Li, Degradation and pathway of tetracycline hydrochloride in aqueous solution by potassium ferrate, *Environ Eng Sci*, 29 (2012) 357-362.
- [66] Y. Wan, A. Jia, Z. Zhu, J. Hu, Transformation of tetracycline during chloramination: Kinetics, products and pathways, *Chemosphere*, 90 (2013) 1427-1434.
- [67] Y. Liu, X. He, Y. Fu, D.D. Dionysiou, Degradation kinetics and mechanism of oxytetracycline by hydroxyl radical-based advanced oxidation processes, *Chem Eng J*, 284 (2016) 1317-1327.
- [68] M. de Cazes, M.P. Belleville, E. Petit, M. Llorca, S. Rodríguez-Mozaz, J. de Gunzburg, D. Barceló, J. Sanchez-Marcano, Design and optimization of an enzymatic membrane reactor for tetracycline degradation, *Catal Today*, 236 (2014) 146-152.

Table 1. Characteristics of the parent compound and the degradation products from hydrolysis and/or biotransformation.

Retention Time (min)	Compound	Ion	Predicted mass (m/z)	Measured mass (m/z)	Elemental composition	Double-bond equivalents (RDB)	Intensity
6.51	ETC or ISO-TC	[M+H] ⁺	445.1605	445.1603	C ₂₂ H ₂₅ O ₈ N ₂	11.5	2.77E+07
7.42	TC*	[M+H] ⁺	445.1605	445.1603	C ₂₂ H ₂₅ O ₈ N ₂	11.5	4.79E+07
7.02	TP 431	[M+H-CH ₂] ⁺	431.1449	431.1445	C ₂₁ H ₂₃ O ₈ N ₂	11.5	4.29E+06
9.93	TP 415	[M+H-CH ₂ -CH ₂ -2H] ⁺	415.1136	415.1133	C ₂₀ H ₁₉ O ₈ N ₂	12.5	5.69E+06
11.66	ISO-TP 415	[M+H-CH ₂ -CH ₂ -2H] ⁺	415.1136	415.1136	C ₂₀ H ₁₉ O ₈ N ₂	12.5	1.09E+05
22.06	TP 387	[M+H-CO-CH ₂ -CH ₂ -2H] ⁺	387.1187	387.1184	C ₁₉ H ₁₉ O ₇ N ₂	11.5	5.72E+06
23.01	TP 370	[M+H-CO-NH ₃ -CH ₃ -CH ₃] ⁺	370.0994	370.0921	C ₁₉ H ₁₆ O ₇ N	12.5	1.37E+07
24.84	ISO-TP 387	[M+H-CO-CH ₂ -CH ₂ -2H] ⁺	387.1187	387.1184	C ₁₉ H ₁₉ O ₇ N ₂	11.5	6.59E+05

*TC=tetracycline

Figure 1. The residual concentrations of tetracycline in the presence and absence of strain DT1 at 30°C and initial pH 9. Error bars represent the standard deviation from triplicate experiments.

Figure 2. The temporal change of tetracycline concentration due to hydrolysis (A) and biotransformation (B) at different temperatures. Lines were modeled using Equations B and C. Error bars are standard deviation from triplicate experiments. The values of R^2 are 0.9279-0.9900 in (A) and 0.9443-0.9965 in (B).

A **B**

Figure 3. The temporal change of tetracycline concentration due to hydrolysis (A) and biotransformation (B) at various initial pHs. Lines were modeled using Equations B, C and D. Error bars were standard deviations from triplicate experiments. The values of R^2 are 0.9766-0.9977 in (A) and 0.9410-0.9967 in (B).

Figure 4. Michaelis-Menten simulation of the biotransformation rates of tetracycline by strain DT1. The value of R^2 is 0.9384. Error bars are standard deviations from triplicate experiments.

Figure 5. Effects of background carbon substrates on tetracycline biotransformation rate and on cell density.

Figure 6. The antibacterial potency of the degradation products as measured using inhibition zones and the concentration of residual tetracycline at 30°C and initial pH 9.0. The inner diameter of the wells was 6 mm in the disk diffusion test. The error bars are standard deviations from triplicate experiments.

Figure 7. Proposed biodegradation pathway of tetracycline by *S. maltophilia* strain TD1. Intermediates that were not detected are parentesize.

Tetracycline

Biotransformation

Main Biotransformation product

Biotransformation and hydrolysis

Decrease antibiotic potency