Characterization and photodegradation of dissolved organic matter (DOM) from a tropical lake and its dominant primary producer, the cyanobacteria *Microcystis aeruginosa*.

Thais B Bittar^{a,*}, Aron Stubbins^a, Armando A. H. Vieira^b, Kenneth Mopper^c

^aSkidaway Institute of Oceanography, Marine Sciences Department, University of Georgia. 10 Ocean Science Circle, Savannah, Georgia, 31411, U.S.A.

* Corresponding author: thais.bittar@skio.uga.edu, 912-598-2445 (office), 757-472-3499 (cell)

aron.stubbins@skio.uga.edu

^bLaboratory of Phycology, Department of Botany, Federal University of São Carlos. Via Washington Luis, km 235, São Carlos, São Paulo, 13565-905, Brazil.

ahvieira.ufscar@gmail.com

^cDepartment of Chemistry and Biochemistry, Old Dominion University. 4541 Hampton Boulevard, Norfolk, Virginia, 23529, U.S.A.

kmopper@odu.edu

22 ***Keywords***

23 Dissolved organic carbon, photodegradation, cyanobacteria, *Microcystis*, tropical lake,

24 freshwater, CDOM, fluorescence, FTICR-MS

25

26 *Abbreviations*

27 a , Napierian absorption coefficient; AI_{mod} , modified aromaticity index; ANOVA, analysis of
28 variance; BB, Barra Bonita; CDOM, chromophoric dissolved organic matter; CHO, molecular
29 formula containing Carbon, Hydrogen and Oxygen only; CHO+N, molecular formulae with C,
30 H, O and nitrogen; CHO+S/P, molecular formulae with C, H, O, P and/or S, and without N; Da,
31 Daltons; DOC, dissolved organic carbon; DOM, dissolved organic matter; EEM, excitation
32 emission matrix; ESI, electrospray ionization; FDOM, fluorescent dissolved organic matter;
33 FTIC-MS, Fourier transform ion cyclotron mass spectrometry; JSI, Jaccard similarity index; λ ,
34 decay constant; NPOC, non-purgeable organic carbon; PARAFAC, parallel factor analysis; r.u.,
35 Raman units; SUVA, specific UV absorption; UVB, ultraviolet B.

36

37

38

39

40

41

42

43

44

45

Abstract

This study investigates optical and high-resolution molecular signatures and photochemical degradation of DOM from the Barra Bonita Reservoir (BB-DOM), a tropical eutrophic lake, as well as from its dominant phytoplankton species, the cyanobacteria *Microcystis aeruginosa* (*Microcystis*-DOM). Consistent with a predominantly autotrophic source, BB-DOM and *Microcystis*-DOM exhibited high protein-like fluorescence and contained a large number of aliphatics. *Microcystis*-DOM was enriched in peptide-like formulae, while BB-DOM had higher chromophoric and fluorescent DOM (CDOM and FDOM) and was enriched in moderately unsaturated formulae, indicating additions of terrigenous DOM and/or in situ processing of autochthonous material in the lake. Consistent with its higher CDOM content, BB-DOM was more photoreactive than *Microcystis*-DOM. For both types of DOM, photodegradation resulted in loss of CDOM, FDOM, moderately unsaturated structures, high O/C and low H/C formulae, and preservation of aliphatics. The majority of photoproducts of 0.5 d irradiation were subsequently removed by day 7, and photoproducts represented a minor fraction of the photo-irradiated DOM. For BB-DOM, molecular formula photolability increased with increasing aromaticity index values, while for *Microcystis*-DOM, molecular formula photolability increased with molecular mass. Photodegradation increased the proportion of molecular formulae containing N (CHO+N) in BB-DOM, while the molecular mass and the proportion of CHO+N formulae decreased upon photo-irradiation of *Microcystis*-DOM. In concert, these molecular shifts due to photodegradation decreased the diversity of and increased the similarity between BB-DOM and *Microcystis*-DOM, suggesting the selective pressure exerted by photochemistry selects for the survival of similar compounds in both samples.

69 *Introduction*

70 Photochemical degradation is one of the most important pathways of removal and
71 transformation of dissolved organic matter (DOM) in aquatic systems (Mopper et al., 2015).
72 Complete photooxidation of dissolved organic carbon (DOC) to inorganic forms (CO₂, CO) and
73 photochemically-mediated alterations in DOM composition play significant roles in global C-
74 cycling (Mopper et al., 1991; Stubbins et al., 2011). Optical, elemental and structural changes in
75 DOM caused by light exposure have several implications for processes in the water column. For
76 instance, exposure to sunlight, particularly ultra-violet (UV) radiation, can affect chemical
77 composition (Helms et al., 2014; Minor et al., 2007) and biological availability of DOM,
78 increasing or decreasing biodegradation (Moran et al., 2000; Stedmon and Markager, 2005;
79 Tranvik and Bertilsson, 2001; Tranvik et al., 2000). UV exposure also causes loss of color (i.e.,
80 photobleaching) (Helms et al., 2013; Stubbins et al., 2012), increasing light penetration depths
81 and the exposure of microorganisms to photosynthetically active radiation and harmful UV
82 radiation, which affect both autotrophic and heterotrophic metabolism (Moran and Zepp, 1997;
83 Vodacek et al., 1997; Walsh et al., 2003).

84 Carbon fluxes in freshwater ecosystems play an important role in global biogeochemical
85 cycles (Aufdenkampe et al., 2011; Battin et al., 2009; Cole et al., 2007) and DOC concentration
86 is one of the parameters used to define the trophic status of lakes (Williamson et al., 1999). In
87 natural lakes and man-made reservoirs, DOM is comprised of a mixture of products from
88 indigenous phytoplankton primary productivity (autochthonous DOM), the surrounding
89 watershed (terrigenous DOM) (Williamson et al., 1999), and the biotic and abiotic reworking of
90 this DOM (Bittar et al., 2015). Generally, autochthonous DOM is more bio-labile, and less

chromophoric and photo-labile, while terrigenous DOM is less bio-labile, and more chromophoric and photo-labile (Benner and Kaiser, 2011).

High resolution Fourier transform ion cyclotron mass spectrometry (FTICR-MS) brings a major fraction of DOM into our analytical window, providing high accuracy molecular-level information regarding elemental and inferred structural composition that can be related to source (Dittmar and Stubbins, 2014; Kujawinski, 2002) and photochemical transformations of DOM (Gonsior et al., 2009; Kujawinski et al., 2004; Stubbins et al., 2010). The majority of DOM photodegradation studies from lakes are from temperate regions (Anesio and Granéli, 2003; Bertilsson and Tranvik, 2000; Boreen et al., 2008). There are few studies addressing the photochemical effects on DOM in tropical lakes, which are subject to intense solar radiation and warm temperatures throughout the year, favoring both high phytoplankton productivity and DOM photodegradation. The extent to which radiation-induced mechanisms in aquatic systems can be extrapolated from the temperate regions to the tropics is not clear (Lewis, 2000).

The Barra Bonita Reservoir (22°29' S, 48°34' W) is one of six cascade, man-made river-reservoirs created for power generation by constructing dams across the Tietê River in São Paulo State, one of the most populous areas of Brazil. The reservoir is located downstream from the confluence between the Tietê and the Piracicaba rivers, where the hydrology regime transitions from lotic to lentic (Tundisi et al., 2008). Seasonal patterns consist of a wet season (December through February) and a dry season (May to August), with only moderate variations in water temperature throughout the year (20 to 30°C) (Dellamano-Oliveira et al., 2008). The reservoir is a eutrophic (i.e., nutrient-rich) and polymictic (i.e., thermally unstratified) tropical ecosystem, that has undergone eutrophication as a result of urban, agricultural and industrial development in the State of São Paulo (Tundisi et al., 2008). Increasing eutrophication of freshwater

environments results in higher growth of cyanobacteria, such as *Microcystis* spp., *Anabaena* spp., and *Pseudo anabaena* spp. (Paerl and Huisman, 2008; Tien et al., 2002). Among these, those belonging to the genus *Microcystis* are of particular interest, as they often dominate massive and potentially toxic blooms in fresh and brackish waters around the world (Sivonen and Jones, 1999), producing significant amounts of DOM. While extensive research has focused upon *Microcystis* toxicity, the fate of DOM produced by this species is not well known (Paerl and Otten, 2013). In the Barra Bonita Reservoir, the most abundant species of phytoplankton is *M. aeruginosa*, which is present primarily as free cells (Calijuri et al., 2002; Dellamano-Oliveira et al., 2008). High tolerance to light stress, buoyancy regulation and low edibility are competitive advantages of *M. aeruginosa* (Brookes and Granf, 2001; Kluijver et al., 2012; Liu et al., 2004), which, combined with warm temperatures and light- and nutrient-replete conditions, lead to dominance of this species in the Barra Bonita Reservoir throughout the year (Dellamano-Oliveira et al. 2008). Consequently, *M. aeruginosa* is an important source of autochthonous DOM in the reservoir during both wet and dry seasons (Dellamano-Oliveira et al., 2008; Vieira et al., 2013). Terrigenous DOM, on the other hand, originates from watershed runoff and riverine input, particularly in the wet season. The catchment area is predominantly agricultural, interspersed with small industrial and urban areas. Watershed runoff is also the cause of high inflows of inorganic N and P, mainly of agricultural origin (Tundisi et al., 2008), which favors phytoplankton growth and release of DOM (Vieira et al., 2013; Welch and Lindell, 1992).

The goals of this study were, firstly, to characterize and compare the optical and molecular signatures of the DOM from the Barra Bonita Reservoir (BB-DOM) and from its dominant primary producer (laboratory cultured *Microcystis*; *Microcystis*-DOM). Secondly, the effects of photodegradation upon BB- and *Microcystis*-DOM were assessed in laboratory-

controlled UVB irradiation experiments. DOM was characterized and the photochemically-mediated alterations were identified using bulk (DOC concentration), optical (chromophoric and fluorescent properties) and molecular-level (FTICR-MS) analyses. In addition, given that the DOM in the reservoir had undergone some degree of photodegradation by being exposed to full sunlight prior to our sampling, we anticipated that this sample would be more likely composed of photodegraded *Microcystis*-DOM than of unaltered *Microcystis* exudates. In order to test this hypothesis, we assessed whether the FTICR-MS profile of the DOM from the reservoir was more similar to that of UVB-exposed *Microcystis*-DOM than to the non-irradiated sample.

Materials and methods

Barra Bonita DOM (BB-DOM) sampling

Samples were collected from the Barra Bonita Reservoir in an 800 m wide section of the reservoir, ~6 km upstream of the hydroelectric dam and ~20 km downstream from the point where the Piracicaba River joins the Tietê River (**Fig. 1**). Water depth was ~20 m at the sampling site. A depth-integrated 20 L water sample was collected on January 28, 2008, using a Niskin sampler and sterile (autoclaved) collection bottle and tubing. The depth integrated sample was generated by combining equal volumes of water collected from 0, 1, 3, 7, 10, 15 and 18 m. On the sampling day, a massive phytoplankton bloom was visible and microscopic analysis showed that the bloom was dominated by free cells of *Microcystis* sp. UVB radiation was measured at the sampling site using a portable radiometer (EPP 2000, StellarNet) during daytime on January 22, 2008, with maximum intensity of 2.7 W m^{-2} (**Fig. S1**).

The 20 L water sample was returned to the laboratory within 2 hours and filtered through pre-combusted $1.2 \mu\text{m}$ pore size filters (GF/C, Whatman) to remove large particles. The $<1.2 \mu\text{m}$ fraction was subsequently filtered using a tangential flow filtration system through a sterile 0.2

160 μm hollow fiber polysulfone cartridge (QuixStand Systems, GE) under low pressure (<5 psi). A
 161 subset of this filtrate containing the DOM ($<0.2 \mu\text{m}$) was frozen at -20°C and shipped in the dark
 162 to Old Dominion University for the irradiation experiments. The filtration system was pre-
 163 washed with 5% NaClO and rinsed thoroughly with ultrapure deionized water (MilliQ,
 164 Millipore). Total organic carbon concentration measured in the ultrapure water after running
 165 through the filtration unit as an operational blank was $<1 \mu\text{M-C}$, which is below the minimum
 166 detection limit reported for DOC analysis using a Shimadzu high temperature combustion system
 167 (e.g., $2.8 \pm 0.3 \mu\text{M-C}$) (Stubbins and Dittmar, 2012).

168 ***Microcystis aeruginosa* DOM (*Microcystis*-DOM)**

169 Axenic (bacteria-free) cultures of a *M. aeruginosa* strain isolated from the Barra Bonita
 170 Reservoir were used to obtain *Microcystis*-derived DOM. The strain is maintained in axenic
 171 conditions in the Culture Collection of Microalgae at the Botany Department, Federal University
 172 of São Carlos (World Data Center for Microorganisms No. 835). Axenic *M. aeruginosa* was
 173 cultured in batch mode in sterile (autoclaved) pH 7 Artificial Seawater McLachlan (ASM-1)
 174 medium containing inorganic nitrogen (NaNO_3 , 2 mM-N) and phosphorus (K_2HPO_4 and
 175 Na_2HPO_4 , 0.2 mM-P) in concentrations similar to that of the Barra Bonita Reservoir water
 176 (Dellamano-Oliveira et al., 2008; Gorham et al., 1964). In spite of the word seawater in the ASM
 177 acronym, this medium does not contain NaCl . DOC concentration of the medium was $<10 \mu\text{M-C}$
 178 prior to addition of *M. aeruginosa*. Cultures were lightly stirred and maintained in a temperature-
 179 controlled room ($23 \pm 1^{\circ}\text{C}$) under a light:dark cycle of 12:12 h and light intensity of 32.7 W m^{-2}
 180 (40 W fluorescent tubes, Philips TLT 40W/75 Extra Day Light). *M. aeruginosa* growth rates
 181 were calculated from in vivo chlorophyll-a fluorescence using a Turner 10-AU fluorometer
 182 calibrated with chlorophyll-a standards (Sigma Chemical Co.). *M. aeruginosa* cultures reached

the late exponential/early stationary growth phase in 30 d (data not shown), at which point the cells were removed by filtration through pre-combusted 1.2 μm pore size filters (GF/C, Whatman). DOM released by the cells that accumulated in the batch cultures was collected by tangential flow filtration through a sterile 0.2 μm pore size polypropylene cassette module (Pellicon Millipore) under low pressure (≤ 5 psi). Cleaning and handling of filtration apparatus were performed as described above for BB-DOM.

Simulated UVB irradiation

Prior to the irradiation experiments, the DOM samples were thawed and filtered once again through acid-washed, sterile (autoclaved) 0.22 μm pore size polycarbonate membrane filters (GTTP Isopore, Millipore) to remove any particles that could have formed during freezing. All glassware were previously autoclaved and combusted, and filtrations and handling of samples were performed in a sterile flow hood; the hood itself was sterilized with 70% ethanol solution and exposed to germicide UV bulbs for 1 h. In the sterile hood, DOM samples were transferred to 250 mL, pre-combusted quartz flasks and irradiated for 0, 0.5, 1, 3 and 7 d in a chamber equipped with 4 UVB bulbs (275 to 315 nm, TL40W-12-RS Philips) and kept at 20°C ($\pm 1^\circ\text{C}$). The UVB intensity reaching the flasks was 2.7 W m^{-2} in order to simulate UVB levels at the surface of the Barra Bonita Reservoir. In these experiments, UVB exposure for 0.5 d and 7 d was equivalent to 1 d and 14 d, respectively, of exposure to natural UVB at the surface of the reservoir, assuming a 12:12 h light:dark cycle. Due to a limited volume of samples, particularly BB-DOM, that could be transported from Brazil to the lab at Old Dominion University, typical dark controls were not run in these experiments. While this is not the ideal setup, great care was taken to avoid contamination of the DOM samples with bacteria, i.e., samples were filter-sterilized twice and materials that came in contact with the samples were pre-combusted (500°C,

overnight) and/or autoclaved. In addition, the UVB radiation used in these time-series experiments is likely to inhibit bacterial growth and activity during these relatively short term experiments.

DOM characterization

Irradiated samples were not filtered prior to analyses for DOM characterization. The absence of visible particles and the low absorbance coefficients of CDOM at wavelengths longer than 500 nm indicated that photo-flocculation did not occur during these irradiations. In addition, photo-flocculation has been shown to take place in DOM samples with levels of terrestrial material and iron significantly greater than those in this study (Helms et al., 2013). The lack of particles detectable by eye or via spectrophotometry provided further evidence that microbial growth did not occur during the irradiations.

Bulk concentrations of DOM are reported as non-purgeable DOC and were obtained by high temperature combustion (680 °C) and non-dispersive infrared detection (NDIR) using a total organic carbon analyzer (TOC-Vcph, Shimadzu). At least triplicate sample injections were made for each sample, with a standard deviation of 0.2. The instrument was calibrated with standard solutions of potassium hydrogen phthalate from 0.1 to 1 mM-C (Stubbins and Dittmar, 2012).

Chromophoric (absorbance) and fluorescent DOM (CDOM and FDOM) were characterized. Ultrapure water was used as blanks for CDOM and FDOM analyses. Absorbance spectra of the ASM-1 medium represented 15% of the absorbance coefficient at 254 nm obtained for non-irradiated *Microcystis*-DOM. The absorbance spectrum of the medium was not used as a blank for CDOM analysis because the medium composition changed during the 30 d of *Microcystis* culturing, i.e., the light-absorbing constituents (mainly nitrate) were consumed by

Microcystis. Measurements of total dissolved nitrogen (TDN) on day 0 and day 30 of *Microcystis* culturing showed that TDN was reduced to 10% of its initial concentration. Therefore, nitrate concentrations and the background absorbance of the medium was likely negligible after 30 d. For each DOM sample, duplicate absorption spectra of CDOM from 250 to 800 nm at 1 nm intervals were obtained using a UV-visible diode array spectrophotometer (Agilent 8453) and a 5 cm acid-washed quartz cuvette. Spectra were corrected for light scatter by subtracting absorption from 750 to 800 nm, and Napierian light absorption coefficients were calculated ($a \text{ m}^{-1}$) (Green and Blough, 1994). First and second derivative spectra were determined using linear regression over sliding 21 nm intervals. The second derivative spectra were used to accurately determine peaks that appeared as shoulders in the absorption coefficient spectra (Helms et al., 2013; Loiselle et al., 2009). Specific UV absorbance (SUVA_{254}) was calculated using the decadic light absorption coefficient at 254 nm (m^{-1}) normalized to DOC (mg-C L^{-1}) and used as a proxy for aromaticity (Weishaar et al., 2003).

Excitation-emission matrix (EEM) spectra for analysis of FDOM were determined using a spectrofluorometer (Fluoromax 2) and a 1 cm acid-washed quartz cuvette (Spencer et al., 2007). Each EEM was comprised of 37 emission spectra from 300 to 500 nm in 1 nm intervals, at excitation from 240 to 420 nm in 5 nm intervals and both emission and excitation slits were set to 5 nm. Blank EEMs were collected from ultrapure water and subtracted from sample EEMs. EEMs were corrected using instrument-specific excitation (xcorrect) and emission (mcorrect) correction files, and for inner-filter effects using Matlab version 7.14 (Cory et al., 2010). Corrected EEMs were normalized to Raman units and FDOM components were identified by parallel factor analysis (PARAFAC). A PARAFAC model was developed using EEMs for 122 samples from *Microcystis*-DOM and BB-DOM using the DOM Fluor toolbox in Matlab version

7.14 (Stedmon and Bro, 2008). The model was validated using random split-half analysis and the number of components was determined by visual examination of the residuals from models with gradually increasing number of components. Intensities of each component were calculated as the maximum fluorescent intensity of each component, i.e., using maxval values (Stedmon and Bro, 2008).

Three samples from each time series experiment (BB-DOM and *Microcystis*-DOM) were selected for analysis by negative mode electrospray ionization (ESI, Apollo II) Fourier transform ion cyclotron mass spectrometry (FTICR-MS) (Bruker Daltonics 12 Tesla Apex Qe), at the College of Sciences Major Instrumentation Cluster (COSMIC), Old Dominion University, Virginia, U.S.A. (Sleighter et al., 2008). The samples analyzed by FTICR-MS from each DOM were: 0 d (non-irradiated), 0.5 d and 7 d UVB irradiated samples. Whole samples (unfractionated) were prepared by dilution (50:50 v/v) with ultrapure methanol plus 0.1% ammonium hydroxide (pH 8) and infusion rates into the ESI ion source were 120 $\mu\text{L h}^{-1}$. Ion accumulation time in the hexapole was 1 s and 300 transients were collected with a 4M word time domain (Chen et al., 2011). Mass spectra of peaks with signal to noise ratio > 4 ranged from 200 to 800 m/z and were internally calibrated using data lists of fatty acids and a list of peaks common to all samples (Sleighter et al., 2008). Molecular formulae assignment was performed using a toolbox in Matlab version 7.14, based on exact masses (± 800 ppb) and published rules (Spencer et al., 2014; Stubbins et al., 2010) for atomic composition as follows: $C \leq 50$; $H/C \geq 0.3$; $O/C \leq 1$; $N \leq 4$; $S \leq 2$; $P \leq 1$. The relatively high number of N and S was allowed in the molecular assignment because DOM from *Microcystis* axenic cultures is likely to contain unaltered amino acids and peptides which may have multiple N and S atoms. Total number of peaks was 1581, 1634 and 1130 for BB-DOM 0 d, 0.5 d and 7 d, respectively; and 1280, 1404 and 1180 for

Microcystis-DOM 0 d, 0.5 d and 7 d, respectively. Number of single-charged and single-formula assigned peaks was 436, 426 and 150 for BB-DOM 0 d, 0.5 d and 7 d, respectively, and 309, 328 and 151, for *Microcystis*-DOM 0 d, 0.5 d and 7 d, respectively. Only single-charged peaks and unambiguously assigned molecular formulae were included in the final list of compounds comprising each DOM sample. The trends observed in total peaks and single-formulae assigned peaks were similar, with lesser peak diversity in the samples irradiated longer.

Assigned formulae were categorized as CHO-only: only C, H and O in the molecular formula; CHO+N: formulae with N, regardless of the presence of P or S (CHON, CHONS, CHONP and CHONSP); and CHO+P/S: formulae with either P or S, but without N (CHOP, CHOS and CHOPS). Peaks were also categorized into likely structural groups, based on double bond equivalents (DBE) (McLafferty and Turecek, 1993), modified aromaticity index ($AI_{mod} = [(1 + \#C - 0.5 * \#O - \#S - 0.5 * \#H) / (\#C - 0.5 * \#O - \#S - \#N - \#P)]$), where $\#C$ = number of carbon atoms, $\#O$ = number of oxygen atoms, $\#S$ = number of sulfur atoms, $\#H$ = number of hydrogen, $\#N$ = number of nitrogen atoms and $\#P$ = number of phosphorus atoms in a given formula, (Koch and Dittmar, 2006), the elemental ratios of H/C and O/C, and the number of N in the formula. The molecular criteria used to define the structural groups in this study were similar to those previously described by Stubbins et al. (2014). Structural groups were aromatics ($AI_{mod} \geq 0.5$), moderately unsaturated structures ($AI_{mod} < 0.5$, $O/C < 0.9$ and $H/C < 1.5$), unsaturated aliphatics ($N=0$, $O/C < 0.9$ and $2 \leq H/C \leq 1.5$), peptide-like ($N > 0$, $O/C < 0.9$, $H/C \geq 1.5$), saturated fatty acids ($N=0$, $O/C < 0.9$ and $H/C > 2$) and sugars ($O/C \geq 0.9$).

Interpretation of FTICR-MS data within time-series comparisons

A comparison of FTICR-MS derived formulae present at 0 d, 0.5 d and 7 d within each DOM time-series experiment revealed six photoreactivity-defined pools of DOM: (1) Highly

photolabile (present in 0 d, absent in 0.5 d, absent in 7 d); (2) Moderately photolabile (present in 0 d, present in 0.5 d, absent from 7 d); (3) Photoresistant (present in 0 d, present in 0.5 d, present in 7 d); (4) Photolabile photoproducts (absent from 0 d, present in 0.5 d, absent from 7 d); (5) Photoresistant photoproducts (absent from 0 d, present in 0.5 d, present in 7 d); (6) “Long-term” photoproducts (absent from 0 d, absent from 0.5 d, present in 7 d).

Statistical analyses

Changes in DOC concentration, SUVA₂₅₄ and fluorescence intensity of PARAFAC-derived FDOM components over the irradiation time-series were fit with the following three-parameter decay function: $y_t = y_0 \exp(-\lambda t) + b$, where y_0 = y-axis value at time 0 (0 d), representing the photoreactive DOM fraction at time 0; y_t = modeled y-axis value at a given time; λ = decay constant, or rate of decay; t = time (d); b = y-axis projected value at time equals infinity, representing the non-reactive DOM fraction; and exp = the base of the natural logarithm, using Sigma Plot v. 12.0 from Systat Software Inc. Decay constants (λ , d⁻¹) and standard errors are presented.

The Jaccard similarity index (JSI) (Jaccard, 1912) ranging from 0 to 1, where 0 = dissimilar and 1 = identical, was calculated as $JSI = \# \text{ of formulae common to both samples} / (\# \text{ formulae unique to BB-DOM} + \# \text{ formulae unique to } Microcystis\text{-DOM} + \# \text{ formulae common to both samples})$, to evaluate similarity between BB-DOM and *Microcystis*-DOM, before and after 0.5 d and 7 d irradiations. Frequency distributions of molecular formulae are presented for the variables AI_{mod} and molecular mass (Da) as number of formulae per class intervals (bins). The bins for each variable were mutually exclusive and defined arbitrarily as 0.1 intervals for AI_{mod} and 200 Da intervals for molecular mass. Unpaired, non-parametric Mann-Whitney tests were used to compare molecular formulae composition of BB- and *Microcystis*-DOM at a given

time point, and an unpaired non-parametric ANOVA (Kruskal-Wallis) test was used to compare molecular mass and AI_{mod} between time points for each DOM type.

Results

Characterization of BB-DOM and Microcystis-DOM before irradiation (0 d)

DOC concentration, and absolute absorbance and fluorescence intensities were greater for BB-DOM than for *Microcystis-DOM* (**Table 1, Figs. 2A-B and 3**). After DOC-normalization, absorbance ($SUVA_{254}$) and fluorescence remained greater for BB-DOM than *Microcystis-DOM* (**Table 1, Figs. 3 and 4**). The second derivative spectra revealed four CDOM peaks in BB-DOM, at 290 nm (a_{290}), ~315 nm (a_{315}), 362 nm (a_{362}) and 430 nm (a_{430}); and four CDOM peaks in *Microcystis-DOM*, at 282 nm (a_{282}), ~310 nm (a_{310}), 350 nm (a_{350}) and 382 nm (a_{382}) (**Fig. 2C-D, solid arrows**). The PARAFAC model identified 3 components (**Fig. S2**), which are described using classical nomenclature (Coble, 1996) as peak T (bio-labile, protein-like, with Ex/Em = 275/338); peak M (semi-bio-labile, marine or autochthonous humic-like, with Ex/Em = 315/388); and peak C+A (bio-refractory, terrigenous humic-like, with Ex/Em = 350/451 and 245/451, respectively). BB-DOM had similar proportion of fluorescence of peaks T, C+A and M, while peak T accounted for just over half of *Microcystis-DOM* fluorescence (**Table 1, Fig. 4**).

FTICR-MS showed differences in the chemical composition of BB- and *Microcystis-DOM* at 0 d (non-irradiated) (**Table 2**). BB-DOM molecular mass was 38 Da lower than *Microcystis-DOM* (Mann-Whitney, $p < 0.0001$). BB-DOM was dominated by CHO-only formulae (81%) and moderately unsaturated structures (56%), while *Microcystis-DOM* was mainly comprised of N-containing formulae (CHO+N, 55%) and the most common structures were moderately unsaturated (32%) and peptide-like formulae (30%). Unsaturated aliphatics

represented 30% and 24%, in BB- and *Microcystis*-DOM, respectively. BB- and *Microcystis*-DOM had 105 formulae in common before irradiation (0 d), and the Jaccard similarity index (JSI) between the two types of DOM at 0 d was 0.16 (**Table 2**). JSI calculated for each elemental and structural class of FTICR-MS formulae showed that saturated fatty acids (0.71), unsaturated aliphatics (0.39), and CHO+P/S (0.29) exhibited the highest similarity between BB- and *Microcystis*-DOM (**Table 2**).

UVB-degradation of BB-DOM and Microcystis-DOM

In BB-DOM, UVB irradiation caused a linear decrease in CDOM absorbance over time, with greater bleaching at shorter wavelengths (**Fig. 2A**). For *Microcystis*-DOM, UVB exposure caused a non-linear decrease in CDOM absorbance from 250 nm to 335 nm over time, and an increase in absorbance from 335 nm to 500 nm within 0.5 d followed by a decrease at all wavelengths from 0.5 d to 7 d of exposure (**Fig. 2B**). UVB exposure caused qualitative shifts in CDOM, as shown by the second derivative CDOM absorption spectra. In BB-DOM, peak *a*290 shifted by ~10 nm towards shorter wavelengths (*a*280), peak *a*315 did not shift significantly and peak *a*362 shifted 10 nm towards longer wavelengths (*a*372) over time, while peak *a*430 disappeared within 0.5 d (**Fig. 2C**). In *Microcystis*-DOM, CDOM peaks *a*282, *a*350 and *a*382 shifted towards shorter wavelengths by 10 nm (*a*272), 30 nm (*a*320) and 5 nm (*a*377), respectively, while peak *a*310 disappeared within 0.5 d (**Fig. 2D**).

Within the first 0.5 d of UVB exposure, BB-DOM absorbance at 254 (CDOM *a*254) was reduced by 41% and at the end of the experimental time-series (7 d) *a*254 was reduced by 80% (**Fig. 3A**). For *Microcystis*-DOM, *a*254 increased slightly within 0.5 d and was reduced by 63% at 7 d (**Fig. 3B**). Photochemical loss of DOC in BB-DOM was measurable from 0.5 d to 7 d of UVB exposure, while DOC decay in *Microcystis*-DOM was minimal until 1 d of exposure, but

measurable between 1 d and 7 d (**Fig. 3**). DOC decay constant values (λ , d⁻¹) over the irradiation time-series were similar for both types of DOM, while λ of SUVA₂₅₄ were significantly greater for BB-DOM than for *Microcystis*-DOM (**Fig. 3**). Total DOC loss over the time-series was 32% and 19% in BB-DOM and *Microcystis*-DOM, respectively. The more efficient removal of CDOM relative to bulk DOC resulted in reductions in SUVA₂₅₄ by 74% and 54% for BB- and *Microcystis*-DOM, respectively.

For the first 0.5 d of exposure, DOC-normalized fluorescence intensity of peaks T, M and C+A in BB-DOM decreased, respectively, by 65%, 58% and 26%, while in *Microcystis*-DOM the decrease was 91%, 87% and 49%, respectively (**Fig. 4**). λ of DOC-normalized fluorescence intensity of peaks T and M over the experimental time-series were smaller in BB-DOM than in *Microcystis*-DOM (**Fig. 4A-D**), while λ of peak C+A were the smallest among FDOM components, and similar between the two types of DOM (**Fig. 4E-F**).

Based on FTICR-MS data, shifts occurring within 0.5 d of irradiation were evident in BB-DOM when relative peak intensity, rather than number of formulae (i.e., presence/absence), of each structural class was considered (**Table 2**). Shifts in the number of molecular formulae, on the other hand, were more dramatic in 7 d-irradiated samples than in 0.5 d-irradiated samples, relative to non-irradiated samples (0 d) for both DOM types (**Table 2, Fig. 5**). At the end of the experimental time series (7 d), the total number of formulae was reduced by 65% in BB- and 51% in *Microcystis*-DOM. The variation in molecular mass (Da) was not significant in BB-DOM over time of UVB exposure (ANOVA, $p>0.05$), while in *Microcystis*-DOM, molecular mass decreased by 50 Da after 7 d (ANOVA, $p<0.0001$) (**Table 2**). In BB-DOM, UVB exposure decreased the relative content of CHO-only formulae (81% to 59%), and caused a relative enrichment in CHO+N formulae (7% to 17%), while in *Microcystis*-DOM, UVB exposure

resulted in relative increase in CHO-only (27% to 57%) and a decrease in CHO+N formulae (55% to 19%) at 7 d (**Table 2**). As a consequence of these opposite photochemically-mediated shifts, the elemental compositions and molecular masses of BB- and *Microcystis*-DOM were more similar to one another at 7 d than prior to irradiation (0 d), with CHO of 57-59%, CHO+N of 17-19% and a 15 Da difference in median molecular mass (**Table 2**). Structural changes caused by UVB irradiation were similar for both BB- and *Microcystis*-DOM, with preferential photochemical removal of moderately unsaturated structures, and photochemical enrichment of aliphatics (**Table 2, Fig. 5**). Photodegradation shifted both DOM types towards a less oxidized (lower O/C) and more saturated (higher H/C) state, particularly at 7 d of exposure (**Fig. 5**). In summation, the elemental, mass-related, structural and oxidation/saturation shifts over the irradiation time-series gradually increased the Jaccard similarity index (JSI) between BB-DOM and *Microcystis*-DOM from 0.16 at 0 d, to 0.19 at 0.5 d and 0.59 at 7 d (**Table 3**).

Photochemically produced compounds

Photoproducts accounted for a minor fraction of the total number of formulae in all irradiated samples: 15% in both 0.5 d and 7 d BB-DOM, and 17% and 32%, respectively, in 0.5 d and 7 d *Microcystis*-DOM. Photoproducts at the shortest irradiation time (0.5 d) were classified as photoresistant or photolabile depending on whether they were, respectively, present or absent in the 7 d irradiated sample (see section ***Interpretation of FTICR-MS data within time-series comparisons*** for descriptions). The photoproducts of 0.5 d irradiation of BB-DOM were more photolabile than those of *Microcystis*-DOM. In BB-DOM, 65 formulae appeared within 0.5 d of exposure (photoproducts), of which 50 (77%) were photolabile, and the other 15 (23%) were photoresistant (i.e. present at 7 d). For *Microcystis*-DOM, 55 photoproducts were detected at 0.5 d, 32 (58%) were photolabile and 23 (42%) were photoresistant until 7 d. For BB-DOM, the

photoresistant photoproducts had O/C <0.3 while most photolabile photoproducts had O/C spanning from 0.3 to 0.6 (**Fig. 5E**), suggesting continued photo-oxidation occurred from 0.5 d to 7 d. In *Microcystis*-DOM, most photoresistant photoproducts had H/C from 1.5 to 2, while photolabile photoproducts had H/C across the range from 1 to >2 (**Fig. 5F**). For both types of DOM, the majority of photoproducts had low AI_{mod} values (<0.2) and molecular mass between 300 and 500 Da and, thus, no clear patterns of photolability of photoproducts were detected regarding aromaticity/degree of molecular conjugation or molecular mass (data not shown). “Long-term” photoproducts accounted for 7 formulae in the 7 d-irradiated BB-DOM and 26 formulae in the 7 d-irradiated *Microcystis*-DOM, and the majority of these had high H/C (**Fig. 5E, 5F**).

Photochemical lability gradient

The distribution in van Krevelen space of each photolability defined group (i.e., photoresistant, moderately- and highly photolabile, see section ***Interpretation of FTICR-MS data within time-series comparisons*** for descriptions) showed that the highly photolabile formulae had O/C spanning from 0.2 to 0.4 and H/C between 1.2 and 1.5 for BB- and *Microcystis*-DOM (**Fig. 5A, 5B**). In the moderately photolabile group these ranges were expanded to greater O/C (up to 0.55 in BB and to 0.9 in *Microcystis*) as well as greater H/C (up to 1.6 in BB and 1.9 in *Microcystis*) (**Fig. 5A-D**). Photoresistant compounds had mostly low O/C (<0.2) and/or high H/C (between 1.75 and 2.5) (**Fig. 5A-F**).

The frequency of distribution of formulae in each photolability-defined group for the variables AI_{mod} (modified aromaticity index) and molecular mass (Da) is presented in **Fig. 6**. In BB-DOM, highly photolabile compounds had high values of AI_{mod} (0.3 to 0.39), moderately photolabile had intermediate AI_{mod} (0.1 to 0.29), and photoresistant compounds had the lowest

AI_{mod} (≤ 0.1) (**Fig. 6A**), indicating an association between photolability and the degree of molecular conjugation. There was no clear association between photolability and molecular mass in BB-DOM, since all three groups had frequency of distribution of formulae centered between 300 and 400 Da (**Fig. 6B**). These results are in agreement with the overall significant decrease in AI_{mod} (ANOVA, $p < 0.0001$) and not significant change in molecular mass (ANOVA, $p > 0.05$) in BB-DOM over the UVB irradiation time-series (**Table 2**). In *Microcystis*-DOM, all three photolability-defined groups had AI_{mod} < 0.1 (**Fig. 6C**) and, thus the relationship between photolability and molecular conjugation was less clear than for BB-DOM. Nonetheless, a number of moderately photolabile formulae with AI_{mod} of 0.2-0.29 (33 formulae) and 0.4-0.49 (35 formulae) resulted in overall decrease in AI_{mod} of *Microcystis*-DOM over the experimental time series (ANOVA, $p = 0.005$, **Table 2**). Highly and moderately photolabile *Microcystis*-DOM formulae had masses between 400 and 500 Da (**Fig. 6D**), while the photoresistant compounds were smaller (300 to 400 Da) (**Fig. 6D**), showing an association between photolability and molecular mass in *Microcystis*-DOM. Such association was reflected in a statistically significant 50 Da-decrease in molecular mass of *Microcystis*-DOM over the irradiation (ANOVA, $p < 0.0001$, **Table 2**).

Discussion

Signatures of BB-DOM and *Microcystis*-DOM

The exudation of DOM by live phytoplankton is a natural process influenced by physiological status, membrane permeability, and the intracellular excess of specific compounds (Fogg, 1983; Mague et al., 1980), among other factors. *Microcystis* is the dominant phytoplankton in Barra Bonita (Dellamano-Oliveira et al., 2008; Vieira et al., 2008), and thus is expected to contribute significantly with the autochthonous production of DOM in the reservoir.

Microcystis-DOM is highly biolabile (Bittar et al., 2015), which implies high biological alteration and turnover rates in the environment (Vieira et al., 2013). Other sources of DOM also exist for Barra Bonita. Therefore, BB-DOM and *Microcystis*-DOM may exhibit quite distinct molecular features.

DOC production rates in laboratory cultures do not accurately reflect the amount of DOC released by natural populations of *Microcystis* in Barra Bonita, given that phytoplankton DOC production is influenced by growth conditions (Fogg 1983). Therefore, the amount of BB-DOM that originates from *Microcystis* exudation could not be determined in this study. Optical and FTICR-MS provide information on DOM quality, including dominant sources and subsequent processing (Fellman et al., 2010; Dittmar and Stubbins, 2014). In this study, many characteristics of *Microcystis*-DOM were shared with BB-DOM. Given the non-quantitative character of these analyses, these similarities have to be interpreted with great care. For instance, 24% of the FTICR-MS formulae identified in BB-DOM had molecular features identical to axenic *Microcystis*-DOM (**Table 2**) and, unsaturated aliphatics and saturated fatty acids displayed high similarity indexes between BB- and *Microcystis*-DOM (0.39 and 0.71, respectively, **Table 1**), implying that *Microcystis* could be an important source of this material in the reservoir. However, identical molecular formulae can represent distinct isomers (i.e., completely distinct compounds, Sleighter and Hatcher, 2008) and the overlap between BB- and *Microcystis*-DOM does not rule out other sources (e.g., other phytoplankton) for these formulae. Similarly, BB-DOM and *Microcystis*-DOM had similar DOC-normalized peak T, which may indicate that this component in the reservoir is strongly influenced by the continuous release of protein-like material by *Microcystis*. Furthermore, BB-FDOM had high relative intensity of peak T when compared to terrestrially-influenced waters (Cory et al., 2007), which may simply indicate that

FDOM in the reservoir is highly influenced by phytoplankton-derived material. Nonetheless, other possible sources of FDOM must be considered. Besides phytoplankton activity, bacterial production (Boyd and Osburn, 2002; Cammack et al., 2004) and riverine discharge and/or watershed runoff of lignin-phenols from vascular plants are also sources of peak T (Maie et al., 2007).

BB-DOM also displayed features distinct from those of *Microcystis*-DOM. A greater content of CDOM (a_{254} and $SUVA_{254}$), more CHO-only formulae and the stronger signal of peaks M and C+A (**Table 1 and 2, Figs. 3 and 4**), for example, suggest the influence of terrigenous humic material input (Coble 1996, Minor et al., 2012), runoff from the agricultural surroundings (Borisover et al., 2009) and byproducts of microbial respiration (Coble, 1996) as potential sources of DOM in the reservoir. Another clear distinction between these types of DOM was the higher relative contribution of moderately unsaturated structures in BB-DOM versus a high relative content of peptide-like formulae in *Microcystis*-DOM (**Table 2**). Moderately unsaturated formulae can represent multiple biogeochemically distinct structural isomers, including carboxylic-rich alicyclic molecules (CRAM) (Hertkorn et al., 2006) and lignin (Dittmar and Stubbins, 2014; Sleighter and Hatcher, 2008; Stubbins et al., 2010), which is unique to vascular plants and thus used as a tracer of terrigenous DOM (Hedges and Mann, 1979; Hernes and Benner, 2006; Spencer et al., 2009). Either microbial reworking of phytoplankton DOM that results in formation of CRAM (Hertkorn et al., 2006), or riverine and runoff inputs of terrigenous DOM could be sources of moderately unsaturated structures in the reservoir. Both CRAM and lignin-like material are considered less biolabile than freshly produced phytoplankton DOM (Hertkorn et al., 2006, Ward et al., 2013) and thus are more likely

to accumulate over time in the reservoir than peptides/proteins from phytoplankton, which are more biolabile (Benner and Kaiser 2011).

Photodegradation

In spite of similar DOC decay rates, qualitative photo-alteration (e.g. loss of CDOM and total molecular formulae) of BB-DOM was more pronounced than for *Microcystis*-DOM. Decay of FDOM components also differed between BB- and *Microcystis* DOM, likely due to distinct starting compositions. For instance, peak T is likely comprised of protein-like material in *Microcystis*-DOM, but could be a mixture of protein- and lignin-like moieties in BB-DOM (Maie et al., 2007). The decay rates of FDOM components are also likely influenced by simultaneous underlying photoproduction of FDOM, particularly peaks C+A and peak M (Helms et al., 2014; Ishii and Boyer, 2012).

The decreases in CDOM absorbance, SUVA₂₅₄ values (**Figs. 2 and 3**), DOC-normalized FDOM, and AI_{mod} (**Table 2, Fig. 4**) suggested that photodegradation caused loss of aromatic, potentially terrestrial signatures, transforming BB-DOM into a more aliphatic-rich DOM pool, which is in agreement with previous reports (Helms et al., 2014; Sleighter and Hatcher, 2008; Stubbins et al., 2010). These results suggest that photodegradation has the potential to mask the allochthonous-DOM signature and overestimate the *Microcystis*-DOM signature in BB-DOM at the surface layer of the reservoir, given the typical water column stability of this water body and high insolation throughout the year.

Photoproducts comprised a minor portion of the photodegraded samples, in agreement with previous studies (Stubbins et al., 2010). While photoproduct are described here and in previous photochemistry studies (Stubbins et al., 2010; Rossel et al., 2013), the unequivocal identification of ‘new’ formulae can be complicated by shifts in charge competition in the

FTICR-MS ionization source due to photodegradation. Such shifts could have resulted in some molecules present in initial samples increasing in abundance (i.e., with higher S/N) in the mass spectra for the photodegraded sample due to the loss of DOC and a probable accompanying loss of the total number of molecules competing for charge. For *Microcystis*-DOM, photodegradation resulted in reduced molecular size, as suggested by blue-shifts of CDOM peaks (**Fig. 2C**) (Helms et al., 2013; Helms et al., 2008; Pavia et al., 1979) and the decrease in median of molecular mass (Da) detected by FTICR-MS (**Table 2, Fig. 6D**). Photodegradation affected the elemental composition of BB- and *Microcystis*-DOM in different ways (**Table 2**), with preferential removal of CHO-only formulae and moderately unsaturated compounds, potentially representative of lignin-phenols, in BB-DOM (Rossel et al., 2013; Stubbins et al., 2010), and preferential removal of CHO+N in *Microcystis*-DOM (Bittar et al., 2015). These results indicate that photodegradation has the potential to reduce the overall degree of molecular conjugation, size and N content of autochthonous DOM in high primary productivity waters. Such transformations have implications to downstream processes such as biological utilization of this less aromatic, lower molecular mass and N-depleted pool of DOM.

Effects of photodegradation on the similarity between BB-DOM and Microcystis-DOM

In spite of potential differences between DOM from lab-cultured and natural populations of *Microcystis*, the *Microcystis*-DOM examined in this study represents ‘pure’ phytoplankton-derived DOM, i.e., that has not undergone any degree of photochemical or bacterial degradation prior to the UVB experiments. Since the reservoir is continuously exposed to sunlight, BB-DOM has unequivocally undergone a degree of photodegradation prior to our sampling. Thus, we hypothesized that BB-DOM at the time of sampling (0 d) would more likely contain photodegraded *Microcystis*-DOM than unaltered *Microcystis*-DOM. If that was the case, the

FTICR-MS profile of initial BB-DOM (0 d) would be more similar to the photodegraded *Microcystis*-DOM (0.5 d and 7 d) than to non-degraded *Microcystis*-DOM (0 d). This trend was observed to a certain extent, as photodegraded *Microcystis*-DOM became more similar to BB-DOM at 0 d, with similarity indexes increasing from 0.16 to 0.18 at 0.5 d and to 0.24 at 7 d of exposure to UVB (**Table 3**). FTICR-MS analyses were not run in replicates, so the statistical significance of these JSI could not be assessed. At the same time, increasing similarity was observed between photodegraded BB-DOM (0.5 d and 7 d) and the non-irradiated *Microcystis*-DOM (0 d) (from 0.16 to 0.17 at 0.5 d, and to 0.23 at 7 d; **Table 3**). One could argue that photoreactivity of *Microcystis*-DOM was theoretically lower compared to photoreactivity of the terrigenous fraction of BB-DOM and, consequently, photodegraded BB-DOM lost its terrigenous features, becoming more similar to *Microcystis*-DOM (0 d). While similarity increased over time of UVB exposure in both directions (i.e., photodegraded BB-DOM was more similar to unaltered *Microcystis*-DOM and vice versa, **Table 3**), the similarity index between the two 7 d-photodegraded samples was 0.59, ~3-fold greater than the JSI of any other sample pair (**Table 3**). These results suggest that (1) BB-DOM is not ‘simply’ composed of photodegraded *Microcystis*-DOM, but by a more complex mixture of sources, and that (2) *Microcystis*-DOM, despite low aromaticity, is not photochemically inert. Most notably, observation of the highest JSI for BB- and *Microcystis*-DOM after 7 d of irradiation indicates that photodegradation increases the similarities between samples by decreasing the chemical diversity of DOM through the selective removal and production of chemically defined suites of molecules (Stubbins et al., 2010; Kellerman et al., 2014).

Final considerations

This study showed that DOM from the tropical eutrophic Barra Bonita Reservoir is likely influenced by autochthonous sources, such as *Microcystis* exudation, and that photodegradation has the potential to alter not only the optical and chemical properties of BB-DOM, but also those of “pure” phytoplankton DOM. Significantly more quantitative field data for light attenuation and for the apparent quantum yield spectra of reactions need to be experimentally determined (i.e. a wavelength dependent photo-by-photon accounting is required) in order to extrapolate these laboratory data on photochemical rates to the water column. The euphotic zone for primary production in the Barra Bonita Reservoir varies from 1 to ~12 m throughout the year (Dellamano-Oliveira et al., 2008), but due to high concentrations of CDOM, UVB radiation in the Barra Bonita Reservoir likely penetrates only the first few centimeters of the surface layer. While this represents a minor portion of the water column, DOM across the whole reservoir area of 325 km² is subject to photodegradation by UVB year-round. Due to water column stability and *Microcystis* buoyancy regulation capability, phytoplankton blooms dominated by this species typically remain at the water surface for days to weeks. Given that bacterial abundance and activity tend to be low at the water surface (Vieira et al., 2013), and that phytoplankton release of DOM is generally greater during daylight hours (Kaplan and Bott, 1982), UVB degradation of *Microcystis*-derived DOM could be a significant mechanism of DOM removal and transformation at the surface of the reservoir. Photodegradation of DOM is considered an important process of C cycling in temperate lakes, including those where the autochthonous DOM component is equally or more dominant relative to terrigenous sources (Bertilsson and Tranvik, 2000; Kellerman et al., 2014; Minor et al., 2012), and this study extends the potential relevance of this process into tropical lakes.

Acknowledgements

This work was funded by the National Science Foundation (OCE# 0728634) and Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP #05/51263-5). TBB thanks FAPESP for a PhD scholarship (FAPESP #05/57213-0). We thank Margie Mulholland, Andrew Gordon, Dave Burdige and Pat Hatcher at Old Dominion University for use of laboratory facilities and instruments. The authors wish to thank Hongmei Chen, John Helms and Pat Hatcher for valuable discussions during the development of this study.

References

- Anesio, A.M., Granéli, W., 2003. Increased photoreactivity of DOC by acidification: Implication for carbon cycle in humic lakes. *Limnol. Oceanogr.* 48, 735-744.
- Aufdenkampe, A.K., Mayorga, E., Raymond, P.A., Melack, J., Doney, S.C., Alin, S.R., Aalto, R.E., Yoo, K., 2011. Riverine coupling of biogeochemical cycles between land, oceans, and atmosphere. *Front. Ecol. Environ.* 9, 53-60.
- Battin, T.J., Luysaert, S., Kaplan, L.A., Aufdenkampe, A.K., Richter, A., Tranvik, L.J., 2009. The boundless carbon cycle. *Nature Geoscience* 2, 598-600.
- Benner, R., Kaiser, K., 2011. Biological and photochemical transformations of amino acids and lignin phenols in riverine dissolved organic matter. *Biogeochemistry* 102, 209-222.
- Bertilsson, S., Tranvik, L.J., 2000. Photochemical transformation of dissolved organic matter in lakes. *Limnol. Oceanogr.* 49, 753-762.
- Bittar, T.B., Vieira, A.A.H., Stubbins, A., Mopper, K., 2015. Competition between photochemical and biological degradation of dissolved organic matter from the cyanobacteria *Microcystis aeruginosa*. *Limnol. Oceanogr.* Doi: 10.1002-lno.10090.

- 619 Boreen, A.L., Edhlund, B.L., Cotner, J.B., McNeill, K., 2008. Indirect photodegradation of
620 dissolved free amino acids: The contribution of singlet oxygen and the differential
621 reactivity of DOM from various sources. *Environ. Sci. Technol.* 42, 5492-5498.
- 622 Borisover, M., Laor, Y., Parparov, A., Bukhanovsky, N., Lado, M., 2009. Spatial and seasonal
623 patterns of fluorescent organic matter in lake Kinneret (Sea of Galilee) and its catchment
624 basin. *Water Res.* 43, 3104-3116.
- 625 Boyd, T.J., Osburn, C.L., 2002. Changes in CDOM fluorescence from allochthonous and
626 autochthonous sources during tidal mixing and bacterial degradation in two coastal
627 estuaries, Ocean Sciences Meeting 2002. Elsevier, Honolulu, pp. 189-210.
- 628 Brookes, J.D., Granf, G.G., 2001. Variations in the buoyancy response of *Microcystis*
629 *aeruginosa* to nitrogen, phosphorus and light. *J. Plankton Res.* 23, 1399-1411.
- 630 Calijuri, M.C., Dos Santos, A.C.A., Jati, S., 2002. Temporal changes in the phytoplankton
631 community structure in a tropical and eutrophic reservoir (Barra Bonita, S.P., Brazil). *J.*
632 *Plankton Res.* 24, 617-634
- 633 Cammack, W.K.L., Kalff, J., Prairie, Y.T., Smith, E.M., 2004. Fluorescent dissolved organic
634 matter in lakes: Relationships with heterotrophic metabolism. *Limnol. Oceanogr.* 49,
635 2034-2045.
- 636 Chen, H., Stubbins, A., Hatcher, P.G., 2011. A mini-electrodialysis system for desalting small
637 volume saline samples for Fourier transform ion cyclotron resonance mass spectrometry.
638 *Limnol. Oceanogr. Methods* 9, 582-592.
- 639 Coble, P.G., 1996. Characterization of marine and terrestrial DOM in seawater using excitation-
640 emission matrix spectroscopy. *Mar. Chem.* 51, 325-346.

- 641 Cole, J.J., Prairie, Y.T., Caraco, N.F., McDowell, W.H., Tranvik, L.J., Striegl, R.G., Duarte,
642 C.M., Kortelainen, P., Downing, J.A., Middelburg, J.J., Melack, J., 2007. Plumbing the
643 global carbon cycle: Integrating inland waters into the terrestrial carbon budget.
644 *Ecosystems* 10, 171-184.
- 645 Cory, R., Miller, M.P., McKnight, D.M., Guerard, J.J., Miller, P.L., 2010. Effect of instrument-
646 specific response on the analysis of fulvic acid fluorescence spectra. *Limnol. Oceanogr.*
647 *Methods* 8, 67-78.
- 648 Cory, R.M., McKnight, D.M., Chin, Y.-P., Miller, P., Jaros, C.L., 2007. Chemical characteristics
649 of fulvic acids from arctic surface waters: Microbial contributions and photochemical
650 transformations. *J. Geophys. Res.* 112, G04S51, 10.1029/2006jg000343.
- 651 Dellamano-Oliveira, M.J., Vieira, A.A.H., Rocha, O., Colombo-Corbi, V., Sant'anna, C.L., 2008.
652 Phytoplankton taxonomic composition and temporal changes in a tropical reservoir.
653 *Fundam. Appl. Limnol.* 171, 27-38.
- 654 Dittmar, T., Stubbins, A., 2014. Dissolved organic matter in aquatic systems, in: Birrer, B.,
655 Falkowski, P., Freeman, K. (Eds.), *Treatise of Geochemistry*. Elsevier, pp. 125-156.
- 656 Fellman, J.B., Hood, E., Spencer, R.G.M., 2010. Fluorescence spectroscopy opens new windows
657 into dissolved organic matter dynamics in freshwater ecosystems: A review. *Limnol.*
658 *Oceanogr.* 55, 2452-2462.
- 659 Fogg, G.E., 1983. The ecological significance of extracellular products of phytoplankton
660 photosynthesis. *Botanica Marina* 26, 3-14.
- 661 Gonsior, M., Peake, B.M., Cooper, W.T., Podgorski, D., D'andrilli, J., Cooper, W.J., 2009.
662 Photochemically induced changes in dissolved organic matter identified by ultrahigh

663 resolution Fourier transform ion cyclotron resonance mass spectrometry. Environ. Sci.
664 Technol. 43, 698-703.

665 Gorham, P.R., McLachlan, J., Hammer, U.T., 1964. Isolation and culture of toxic strains of
666 *Anabaena flos-aquae*. Breb. Int. Ver. Theor. Angew. Limnol. Verh. 19, 796-804.

667 Green, S.A., Blough, N.V., 1994. Optical absorption and fluorescence properties of
668 chromophoric dissolved organic matter in natural waters. Limnol. Oceanogr. 39, 1903-
669 1916.

670 Hedges, J.I., Mann, D.C., 1979. The characterization of plant tissues by their lignin oxidation
671 products. Geochim. Cosmochim. Acta 43, 1803-1807.

672 Helms, J.R., Mao, J., Stubbins, A., Schmidt-Rohr, K., Spencer, R.G.M., Hernes, P.J., Mopper,
673 K., 2014. Loss of optical and molecular indicators of terrigenous dissolved organic matter
674 during long-term photobleaching. Aquat. Sci. 76, 353-373.

675 Helms, J.R., Stubbins, A., Perdue, E.M., Green, N.W., Chen, H., Mopper, K., 2013.
676 Photochemical bleaching of oceanic dissolved organic matter and its effect on absorption
677 spectral slope and fluorescence. Mar. Chem. 155, 81-91.

678 Helms, J.R., Stubbins, A., Ritchie, J.D., Minor, E.C., Kieber, D.J., Mopper, K., 2008. Absorption
679 spectral slopes and slope ratios as indicators of molecular weight, source, and
680 photobleaching of chromophoric dissolved organic matter. Limnol. Oceanogr. 53, 955-
681 969.

682 Hernes, P.J., Benner, R., 2006. Terrigenous organic matter sources and reactivity in the North
683 Atlantic Ocean and a comparison to the Arctic and Pacific oceans. Mar. Chem. 100, 66-
684 79.

- 685 Hertkorn, N., Benner, R., Frommberger, M., Schmitt-Kopplin, P., Witt, M., Kaiser, K., Kettrup,
686 A., Hedges, J.I., 2006. Characterization of a major refractory component of marine
687 dissolved organic matter. *Geochim. Cosmochim. Acta* 70, 2990-3010.
- 688 Ishii, S.K., Boyer, T.H., 2012. Behavior of reoccurring PARAFAC components in fluorescent
689 dissolved organic matter in natural and engineered systems: A critical review. *Environ.*
690 *Sci. Technol.* 46, 2006-2017.
- 691 Jaccard, P., 1912. The distribution of the flora in the alpine zone. *New Phytol.* 9(2), 37-50.
- 692 Kaplan, L.A., Bott, T.L., 1982. Diel fluctuations of DOC generated by algae in a piedmont
693 stream. *Limnol. Oceanogr.* 27, 1091-1100.
- 694 Kellerman, A.M., Dittmar, T., Kothawala, D.N., Tranvik, L.J., 2014. Chemodiversity of
695 dissolved organic matter in lakes driven by climate and hydrology. *Nat. Commun.* Doi:
696 10.1038/ncomms4804.
- 697 Kluijver, A., Yu, J., Houtekamer, M., Middelburg, J.J., Liu, Z., 2012. Cyanobacteria as a carbon
698 source for zooplankton in eutrophic lake Taihu, China, measured by ¹³C labeling and
699 fatty acid biomarkers. *Limnol. Oceanogr.* 57, 1245-1254.
- 700 Koch, B.P., Dittmar, T., 2006. From mass to structure: An aromaticity index for high-resolution
701 mass data of natural organic matter. *Rapid Commun. Mass Spectrom.* 20, 926-932.
- 702 Kujawinski, E.B., 2002. Electrospray ionization Fourier transform ion cyclotron mass
703 spectrometry (ESI-FT-ICR-MS): Characterization of complex environmental mixtures.
704 *Environ. Forensics* 3, 207-216.
- 705 Kujawinski, E.B., Del Vecchio, R., Blough, N.V., Klein, G.C., Marshall, A.G., 2004. Probing
706 molecular-level transformations of dissolved organic matter: Insights on photochemical

707 degradation and protozoan modification of DOM from electrospray ionization Fourier
708 transform ion cyclotron resonance mass spectrometry. Mar. Chem. 92, 23-37.

709 Lewis, W.M.J., 2000. Basis for the protection and management of tropical lakes. Lakes Reserv.
710 Res. Manage. 5, 35-48.

711 Liu, Z., Hader, D.P., Sommaruga, R., 2004. Occurrence of mycosporine-like amino acids
712 (MAAs) in the bloom-forming cyanobacterium *Microcystis aeruginosa*. J. Plankton Res.
713 26, 963-966.

714 Loiselle, S.A., Bracchini, L., Dattilo, A.M., Rici, M., Tognazzi, A., Cozar, A., Rossi, C., 2009.
715 Optical characterization of chromophoric dissolved organic matter using wavelength
716 distribution of absorption spectral slopes. Limnol. Oceanogr. 52, 590-597.

717 Mague, T.H., Friberg, E., Hughes, D.J. , Morris, I., 1980. Extracellular release of carbon by
718 marine phytoplankton: A physiological approach. Limnol. Oceanogr. 25, 262-279.

719 Maie, N., Scully, N.M., Pisani, O., Jaffe, R., 2007. Composition of a protein-like fluorophore of
720 dissolved organic matter in coastal wetland and estuarine ecosystems. Water Res. 41,
721 563-570.

722 McLafferty, F.W., Turecek, F., 1993. Interpretation of mass spectra. University Science Books,
723 Mill Valley, California, USA.

724 Minor, E.C., Dalzell, B.J., Stubbins, A., Mopper, K., 2007. Evaluating the photoalteration of
725 estuarine dissolved organic matter using direct temperature-resolved mass spectrometry
726 and UV-visible spectroscopy. Aquat. Sci. 69, 440-455.

727 Mopper, K., Kieber, D.J., Stubbins, A., 2015. Marine photochemistry of organic matter:
728 Processes and impacts, in: Hansell, D.A., Carlson, C.A. (Eds.), Marine Photochemistry of
729 Organic Matter. Elsevier, pp. 389-450.

- 730 Mopper, K., Zhou, X., Kieber, R.J., Kieber, D.J., Sikorski, R.J., Jones, R.D., 1991.
731 Photochemical degradation of dissolved organic carbon and its impact on the oceanic
732 carbon cycle. *Science* 353, 60-62.
- 733 Moran, M.A., Sheldon, W.M.J., Zepp, R.G., 2000. Carbon loss and optical property changes
734 during long-term photochemical and biological degradation of estuarine dissolved
735 organic matter. *Limnol. Oceanogr.* 45, 1254-1264.
- 736 Moran, M.A., Zepp, R.G., 1997. Role of photoreactions in the formation of biologically labile
737 compounds from dissolved organic matter. *Limnol. Oceanogr.* 42, 1307-1316.
- 738 Morris, D.P., Zagarese, H., Williamson, C.E., Balseiro, E.G., Hargreaves, B.R., Modenutti, B.,
739 Moeller, R., Queimalinos, C., 1995. The attenuation of solar UV radiation in lakes and
740 the role of dissolved organic carbon. *Limnol. Oceanogr.* 40, 1381-1391.
- 741 Paerl, H.W., Huisman, J., 2008. Blooms like it hot. *Science* 320, 57-58.
- 742 Paerl, H.W., Otten, T.G., 2013. Blooms bite the hand that feeds them. *Science* 342, 433-434.
- 743 Pavia, D.L., Lampman, G.M., Kriz, G.S., 1979. Introduction to spectroscopy: A guide for students
744 of organic chemistry. Saunders.
- 745 Rossel, P.E., Vähätalo, A.V., Witt, M., Dittmar, T., 2013. Molecular composition of dissolved
746 organic matter from a wetland plant (*Juncus effusus*) after photochemical and microbial
747 decomposition (1.25 yr): Common features with deep sea dissolved organic matter. *Org.*
748 *Geochem.* 60, 62-71.
- 749 Sleighter, R.L., Hatcher, P.G., 2008. Molecular characterization of dissolved organic matter
750 (DOM) along a river to ocean transect of the lower Chesapeake Bay by ultrahigh
751 resolution electrospray ionization Fourier transform ion cyclotron resonance mass
752 spectrometry. *Mar. Chem.* 110, 140-152.

- 753 Sleighter, R.L., McKee, G., Liu, Z., Hatcher, P.G., 2008. Naturally present fatty acids as internal
754 calibrants for Fourier transform mass spectra of dissolved organic matter. *Limnol.*
755 *Oceanogr. Methods* 6, 246-253.
- 756 Spencer, R.G., Baker, A., Ahad, J.M., Cowie, G.L., Ganeshram, R., Upstill-Goddard, R.C., Uher,
757 G., 2007. Discriminatory classification of natural and anthropogenic waters in two U.K.
758 Estuaries. *Sci. Total Environ.* 373, 305-323.
- 759 Spencer, R.G.M., Guo, W., Raymond, P.A., Dittmar, T., Hood, E., Fellman, J. , Stubbins, A.,
760 2014. Source and biolability of ancient dissolved organic matter in glacier and lake
761 ecosystems on the Tibetan plateau. *Geochim. Cosmochim. Acta* 142, 64-74.
- 762 Spencer, R.G.M., Stubbins, A., Hernes, P.J., Baker, A., Mopper, K., Aufdenkampe, A.K., Dyda,
763 R.Y., Mwamba, V.L., Mangangu, A.M., Wabakanghanzi, J.N., Six, J., 2009.
764 Photochemical degradation of dissolved organic matter and dissolved lignin phenols from
765 the congo river. *J. Geophys. Res.* 114, G03010, 10.1029/2009jg000968.
- 766 Stedmon, C.A., Bro, R., 2008. Characterizing dissolved organic matter fluorescence with parallel
767 factor analysis: A tutorial. *Limnol. Oceanogr. Methods* 6, 572-279.
- 768 Stedmon, C.A., Markager, S., 2005. Tracing the production and degradation of autochthonous
769 fractions of dissolved organic matter by fluorescence analysis. *Limnol. Oceanogr.* 50,
770 1415-1426.
- 771 Stubbins, A., Dittmar, T., 2012. Low volume quantification of dissolved organic carbon and
772 dissolved nitrogen. *Limnol. Oceanogr. Methods* 10, 347-352.
- 773 Stubbins, A., Lapierre, J.F., Berggren, M., Prairie, Y., Dittmar, T., Del Giorgio, P.A., 2014.
774 What's in an EEM? Molecular signatures associated with dissolved organic fluorescence
775 in boreal Canada. *Environ. Sci. Technol.* 48, 10598–10606.

- 776 Stubbins, A., Law, C.S., Uher, G., Upstill-Goddard, R.C., 2011. Carbon monoxide apparent
777 quantum yields and photoproduction in the Tyne Estuary. *Biogeosciences* 8, 703-713.
- 778 Stubbins, A., Niggemann, J., Dittmar, T., 2012. Photo-lability of deep ocean dissolved black
779 carbon. *Biogeosciences* 9, 1661-1670.
- 780 Stubbins, A., Spencer, R.G.M., Chen, H.M., Hatcher, P.G., Mopper, K., Hernes, P.J., Mwamba,
781 V.L., Mangangu, A.M., Wabakanghanzi, J.N., Six, J., 2010. Illuminated darkness:
782 Molecular signatures of Congo River dissolved organic matter and its photochemical
783 alteration as revealed by ultrahigh precision mass spectrometry. *Limnol. Oceanogr.* 55,
784 1467-1477.
- 785 Sugiyama, Y., Hatcher, P.G., Sleighter, R.L., Suzuki, T., Wada, C., Kumagai, T., Mitamura, O.,
786 Katano, T., Nakano, S., Tanaka, Y., Drucker, V.V., Fialkov, V.A., Sugiyama, M. 2014
787 Developing an understanding of dissolved organic matter dynamics in the giant Lake
788 Baikal by ultrahigh resolution mass spectrometry *Limnology*. 15, 127-139.
- 789 Tien, C.J., Krivtsov, V., Levado, E., Sigee, D.C., White, K.N., 2002. Occurrence of cell-
790 associated mucilage and soluble extracellular polysaccharides in Rostherne Mere and
791 their possible significanc. *Hydrobiologia* 485, 245-252.
- 792 Tranvik, L.J., Bertilsson, S., 2001. Contrasting effects of solar UV radiation on dissolved organic
793 sources for bacterial growth. *Ecol. Lett.* 4, 458-463.
- 794 Tranvik, L.J., Olofsson, H., Bertilsson, S., 2000. Photochemical effects on bacterial degradation
795 of dissolved organic matter in lake water, in: Bell, C.R., Brylinsky, M., Johnson-Green,
796 P.C. (Eds.), *Microbial Biosystems: New Frontiers. Proc. 8th Int. Symp. Microb. Ecol.*,
797 Halifax, Canada, pp. 193-200.

- 798 Tundisi, J.G., Matsumura-Tundisi, T., Abe, D.S., 2008. The ecological dynamics of Barra Bonita
799 (Tietê River, S.P., Brazil) Reservoir: Implications for its biodiversity. *Braz. J. Biol.* 68,
800 1079-1098.
- 801 Vieira, A.A.H., Colombo-Corbi, V., Bagatini, I.L., Dellamano-Oliveira, M.J., Tessarolli, L.P.,
802 2013. Rhamnose and hydrolysis of muf- α -L- rhamnopyranoside coupled with producers
803 of rhamnose-rich extracellular polysaccharides in a hypereutrophic reservoir. *Aquat.*
804 *Microb. Ecol.* 69, 169-181.
- 805 Vieira, A.A.H., Ortolano, P.H.C., Girolodo, D., Dellamano-Oliveira, M.J., Bittar, T.B., Lombardi,
806 A.T., Sartori, A.L., Paulsen, B.S., 2008. Role of hydrophobic extracellular polysaccharide
807 of *Aulacoseira granulata* (Bacillariophyceae) on aggregate formation in a turbulent and
808 hypereutrophic reservoir. *Limnol. Oceanogr.* 53, 1887-1899.
- 809 Vodacek, A., Blough, N.V., DeGrandpre, M.D., Peltzer, E.T., Nelson, R.K., 1997. Seasonal
810 variation of CDOM and DOC in the Middle Atlantic Bight: Terrestrial inputs and
811 photooxidation. *Limnol. Oceanogr.* 42, 674-686.
- 812 Walsh, J.J., Weisberg, R.H., Dieterle, D.A., He, R., Darrow, B.P., Jolliff, J.K., Lester, K.M.,
813 Vargo, G.A., Kirkpatrick, G.J., Fanning, K.A., Sutton, T.T., Jochens, A.E., Biggs, D.C.,
814 Nababan, B., Hu, C., Muller-Karger, F.E., 2003. Phytoplankton response to intrusions of
815 slope water on the West Florida Shelf: Models and observations. *J. Geophys. Res.* 108,
816 C6, 3190, 10.1029/2002JC001406.
- 817 Ward, N.D., Keil, R.G., Medeiros, P.M., Brito, D.C., Cunha, A.C., Dittmar, T., Yager, P.L.,
818 Krusche, A.V., Richey, J.E., 2013. Degradation of terrestrially derived macromolecules
819 in the Amazon River. *Nature Geoscience* 6, 530-533.

820 Weishaar, J.L., Aiken, G., Bergamaschi, B.A., Fram, M.S., Fujii, R. , Mopper, K., 2003.
821 Evaluation of specific ultraviolet absorbance as an indicator of the chemical composition
822 and reactivity of dissolved organic carbon. *Environ. Sci. Technol.* 37, 4702-4708.

823 Welch, E.B., Lindell, T., 1992. Ecological effects of wastewater: Applied limnology and
824 pollution effects. E & FN Spon, London.

825 Williamson, C.E., Morris, D.P., Pace, M.L., Olson, O.G., 1999. Dissolved organic carbon and
826 nutrients as regulators of lake ecosystems: Resurrection of a more integrated paradigm.
827 *Limnol. Oceanogr.* 44, 795-803.

828

829

830

831

832

833

834

835

836

837

838

839 *Figure Captions*

840 **Figure 1.** Location of the Barra Bonita Reservoir, Tietê River and Piracicaba River and this
 841 study's sampling site (A, B, C) and a bloom of *Microcystis aeruginosa* in the reservoir (D, E).
 842 Sources: (A) Wikimedia Commons, http://pt.wikipedia.org/wiki/S%C3%A3o_Paulo, (B) Google
 843 maps, (C) <http://www.brazilinvestmentguide.com/state-guides/sao-paulo/> (C, D), personal
 844 records.

845 **Figure 2.** CDOM absorbance spectra (A, B) and second derivative CDOM absorption spectra (C,
 846 D) of non-irradiated (0 d) and UVB irradiated (0.5, 1, 3, 7 d) DOM from the Barra Bonita
 847 Reservoir (A, C) and axenically-cultured *Microcystis* (B, D). Solid arrows and dashed arrows
 848 represent peaks and shoulders in the CDOM spectra of non-irradiated (0 d) and UVB irradiated
 849 DOM, respectively.

850 **Figure 3.** Loss of dissolved organic carbon (DOC, mM-C, crosses, short-dashed lines), CDOM
 851 absorbance at 254 nm (a_{254} , m^{-1} , circles, long-dashed lines) and specific UV absorbance
 852 (SUVA_{254} , $\text{mgC L}^{-1} \text{m}^{-1}$, triangles, solid lines) over UVB irradiation time series of DOM from
 853 Barra Bonita Reservoir (A) and axenically-cultured *Microcystis* (B). Decay rates (λ , d^{-1}) ($\pm$
 854 standard error); adjusted r^2 and p value of modelled data are presented.

855 **Figure 4.** Decay of DOC-normalized fluorescence intensity of peak T (A, B), peak M (C, D),
 856 and peak C+A (E, F) over UVB irradiation time series of DOM from Barra Bonita Reservoir (A,
 857 C, E) and axenically-cultured *Microcystis* (B, D, F). Decay rates (λ , d^{-1}) ($\pm$ standard error);
 858 adjusted r^2 and p value of modelled data are presented.

859 **Figure 5.** Van Krevelen diagrams of FTICR-MS formulae in each photolability defined group:
 860 photoresistant (black dots), highly photolabile (white dots), moderately photolabile (blue crosses)

that comprise samples at 0 d (non-irradiated) for BB-DOM (A) and *Microcystis*-DOM (B); photoresistant, moderately photolabile, photoresistant photoproducts (red triangles) and photolabile photoproducts (open squares) that comprise samples irradiated for 0.5 d for BB-DOM (C) and *Microcystis*-DOM (D); and photoresistant, photoresistant photoproducts and long-term photoproducts (blue triangles) that comprise samples irradiated for 7 d for BB-DOM (E) and *Microcystis*-DOM (F).

Figure 6. Frequency of distribution of number of formulae classified as photoresistant (black bars), moderately photolabile (blue bars) and highly photolabile (white bars) in each range interval (bin) of modified aromaticity index (AI_{mod}) (A, C) and molecular mass (Da) (B, D) for BB-DOM (A, B) and *Microcystis*-DOM (C, D).

Figure S1. UVB radiation intensity ($W\ m^{-2}$) in the Barra Bonita Reservoir (São Paulo State, Brazil) (A) during daylight hours and (B) throughout the year 2007.

Figure S2. PARAFAC-derived FDOM components and spectral loadings.

881 Table 1
 882 DOC concentrations, CDOM (a_{254} and $SUVA_{254}$) and FDOM absolute, relative (in brackets, %)
 883 and DOC-normalized fluorescence intensity measurements of DOM from the Barra Bonita
 884 Reservoir (BB-DOM) and from axenic cultures of *Microcystis aeruginosa* (*Microcystis*-DOM)
 885 before irradiation (0 d).

	BB-DOM	<i>Microcystis</i> -DOM
DOC (mM-C)	1.60	0.97
a_{254} (m ⁻¹)	93	17
$SUVA_{254}$ (mgC L ⁻¹ m ⁻¹)	2.10	0.64
FDOM total (r.u.; %)	4.7 (100)	1.9 (100)
Peak T	1.6 (33)	1.0 (51)
Peak M	1.4 (29)	0.3 (17)
Peak C+A	1.8 (38)	0.6 (32)
FDOM total (r.u. mM-C ⁻¹)	3.0	2.0
Peak T	1.0	1.0
Peak M	0.9	0.3
Peak C+A	1.1	0.6

893 Table 2. Molecular formulae characterization of BB-DOM and *Microcystis*-DOM at each irradiation time (0 d, 0.5 d and 7 d), and
 894 common formulae ('Common 0 d') and Jaccard similarity index (JSI) of BB-DOM and *Microcystis*-DOM prior to irradiation (0 d).
 895 Number (#) of singly-assigned formulae, median of molecular mass (Da), median of modified aromaticity index (AI_{mod}), elemental
 896 composition (number of CHO-only, CHO+P/S and no N, and CHO+N formulae) and structural composition (number of formulae [#]
 897 and relative peak intensity [rel. int.]). Numbers in brackets are percentages (%).

		BB 0 d	BB 0.5 d	BB 7 d	<i>Microcystis</i> 0 d	<i>Microcystis</i> 0.5 d	<i>Microcystis</i> 7 d	Common 0d	JSI
All	#	436	426	150	309	328	151	105	0.16
Molecular mass	median	365	362	368	403	413	353	326	--
AI_{mod}	median	0.19	0.18	0.09	0.13	0.10	0.05	0.07	--
CHO	# (%)	353 (81)	351 (82)	89 (59)	82 (27)	90 (27)	86 (57)	67	0.18
	Rel. int.	0.79	0.84	0.64	0.29	0.31	0.73		
CHO+P/S	# (%)	53 (12)	44 (10)	36 (24)	58 (19)	72 (22)	36 (24)	26	0.29
	Rel. int.	0.15	0.09	0.26	0.21	0.22	0.18		
CHO+N	# (%)	30 (7)	31 (7)	25 (17)	169 (55)	166 (51)	29 (19)	12	0.06
	Rel. int.	0.06	0.06	0.10	0.51	0.47	0.09		
aromatics	# (%)	6 (1)	11 (3)	11 (7)	19 (6)	25 (8)	8 (5)	1	0.04
	rel. int.	0.01	0.02	0.05	0.11	0.10	0.02		
moderately unsaturated	# (%)	246 (56)	214 (50)	32 (21)	100 (32)	97 (30)	19 (13)	19	0.06
unsaturated	rel. int.	0.41	0.29	0.13	0.26	0.24	0.09		
aliphatics	# (%)	143 (33)	165 (39)	67 (45)	73 (24)	84 (26)	76 (50)	61	0.39
saturated	rel. int.	0.48	0.60	0.58	0.29	0.32	0.67		
fatty acids	# (%)	23 (5)	17 (4)	23 (15)	18 (6)	25 (8)	27 (18)	17	0.71
peptide-like	rel. int.	0.06	0.04	0.16	0.03	0.06	0.12		
	# (%)	16 (4)	16 (4)	13 (9)	94 (30)	93 (28)	20 (13)	7	0.07
sugars	rel. int.	0.04	0.04	0.06	0.28	0.26	0.10		
	# (%)	2 (0)	3 (1)	4 (3)	5 (2)	4 (1)	1 (1)	0	0.00
	rel. int.	0.00	0.00	0.02	0.03	0.03	0.00		

898

	BB-DOM 0 d	BB-DOM 0.5 d	BB-DOM 7 d
<i>Microcystis</i> -DOM 0 d	0.16	0.17	0.23
<i>Microcystis</i> -DOM 0.5 d	0.18	0.19	0.27
<i>Microcystis</i> -DOM 7 d	0.24	0.26	0.59

899

900

901

902

903

904

905

906

907

908

909

910

911

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6