Carbon isotope measurements reveal unexpected cycling of dissolved organic matter in the deep Mediterranean Sea

Chiara Santinelli (1), Christopher Follett (2, 3), Simona Retelletti Brogi (1), Li Xu (4), and Daniel Repeta (2)

1. Istituto di Biofisica, Via Moruzzi 1, Pisa, Italy

2. Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, MA, 02543. USA.

3. Lorenz Center, Department of Earth Atmospheric, and Planetary Sciences, Massachusetts Institute Technology Cambridge, MA 02139

4. Department of Marine Geology and Geophysics, Woods Hole Oceanographic Institution, Woods Hole, MA, 02543. USA.

Follett@mit.edu, Lxu@whoi.edu, simona.retelletti@pi.ibf.cnr.it, drepeta@whoi.edu

ABSTRACT

One of the most intriguing aspects of dissolved organic carbon (DOC) dynamics in the Mediterranean Sea (Med Sea) is that in the intermediate and deep waters, DOC concentrations are equal to the lowest values found in the deep Atlantic and Pacific (36-42 μM). The very low DOC values in the deep Med Sea were unexpected since the renewal time of deep waters in the basin is only 20-126 years. Over this short timescale, we expected just a very small, nearly undetectable, fraction of refractory DOC (RDOC) to be removed. The first DOC isotope data show that DOC in Med Sea deep water is more depleted in both $\Delta^{14}\text{C}$ and $\delta^{13}\text{C}$ than in the deep Atlantic Ocean, with an estimated age of 4500-5100 years. These data suggest that at least 10%, and up to 45%, of the Atlantic RDOC entering the Med Sea is removed and replaced by isotopically lighter DOC in less than 126 years. Potential allochthonous sources include: fossil methane and methane derived DOC seeps from sediments, anthropogenic combustion products and terrestrial organic matter delivered by the atmosphere, rivers, and groundwaters. Using current information, it is not possible to quantify the relative contribution of these potential sources. Based on estimated flux and isotopic value, atmospheric input of soluble organic carbon from soils or combustion products as well as DOC from groundwater are the most likely

sources of allochthonous DOC. Our results suggest that DOC cycling in the deep Med Sea is dynamic and support the idea that a substantial fraction (up to 45%) of what has traditionally been defined as “refractory” DOC imported from the Atlantic Ocean, can be removed on temporal scales of <126 y, thereby opening intriguing questions about deep sea DOC cycling.

Key words

Dissolved organic carbon, carbon cycle, radiocarbon, refractory carbon, Mediterranean Sea

1. Introduction

In the global ocean, dissolved organic carbon (DOC) concentrations range between 34 μM in the deep North Pacific and $\sim 80 \mu\text{M}$ in oligotrophic surface waters of mid ocean gyres. DOC concentrations are generally high in the surface layer, where most of the production processes occur, undergo a marked reduction in the mesopelagic layer, due to net microbial mineralization, and have low values in the deep ocean ($> 1000 \text{ m}$). The fraction of deep ocean DOC remaining when concentrations reach values $< 42 \mu\text{M}$, is considered “refractory” (RDOC), indicating little potential for additional removal by microbial respiration on short time scales (lifetime of $\sim 16,000$ years; Hansell, 2013). DOC stable carbon isotopes values ($\text{DOC}\delta^{13}\text{C}$) are $\sim -21\text{‰}$ to -22‰ throughout the whole water column, indicating an autochthonous microbial source for most marine DOC (Druffel et al.,

1992). DOC concentrations decrease along the pathway of abyssal circulation, from ~ 48 μM in the deep North Atlantic (Carlson et al., 2010), to ~ 40 μM in the Southern Ocean (Hansell and Carlson, 2013) and ~ 34-38 μM in the deep North Pacific (Hansell and Carlson, 1998; Hansell and Carlson, 2013). The decrease in DOC concentration along the trajectory of abyssal circulation suggests slow, continuous removal of RDOC in the deep ocean, of the order ~3 nM C y^{-1} (Hansell, 2013). However, looking at the deep RDOC distribution more in detail, it appears that the removal of RDOC may not be a continuous process. Localized sinks for RDOC have been proposed in the deep, far North Pacific and at mid depth in the subtropical South Pacific, while localized sources may be present in the Indian Ocean, the Arabian Sea (Hansell and Carlson, 1998) and at the Equator (Hansell and Carlson, 2013).

Radiocarbon measurements support the concept of a very old pool of DOC in the deep ocean. Radiocarbon values for RDOC are highly depleted (~ -400 to -530‰), compared to DIC (-80 to -230‰), indicating that RDOC is markedly older than DIC (4000-6000 vs. 100-1000 y). RDOC becomes further depleted in $\Delta^{14}\text{C}$ as water ages and moves from the deep North Atlantic (~ -390‰) to the North Pacific (-530‰), in good agreement with the depletion in $\text{DIC}\Delta^{14}\text{C}$ (from -80 to -230‰) (Druffel et al., 1992). Due to its long apparent residence time in the ocean, RDOC should be well mixed throughout the water column.

In surface waters of the N. Atlantic, $\text{DOC}\Delta^{14}\text{C}$ is strongly depleted relative to $\text{DIC}\Delta^{14}\text{C}$ (Druffel et al., 1992). The $\text{DOC}\Delta^{14}\text{C}$ depth profile, with ~ 70-80 μM DOC of $\Delta^{14}\text{C}$ -230‰ in surface water and ~ 40 μM DOC of $\Delta^{14}\text{C}$ ~ -390‰ in deep water, is therefore well described by a two-component model (TCM) of 36-42 μM RDOC (~ -390‰) mixed throughout the entire water column, and a second, semi-labile component (SLDOC) with a modern radiocarbon age that is cycled within the upper ocean (< 1000 m) (Williams and Druffel, 1987)

Recently, Follett et al. (2014) proposed that the TCM is not adequate to fully describe the complexity and richness of marine DOC cycle. Follett et al. (2014) used stepwise oxidation of DOC to estimate the isotopic distribution of components within DOC in order to uncover carbon fluxes that may be obscured by the TCM. Their results suggest that a significant fraction of deep DOC is modern and supported by a 1 $\text{Pg}\cdot\text{y}^{-1}$ carbon flux from sinking particles. These results open new and intriguing hypothesis about DOC cycle.

The Mediterranean Sea (Med Sea) is a natural laboratory where these hypotheses may be tested. The major physical processes occurring in the oceans (i.e.: thermohaline

1 circulation, deep water formation, seasonal cycle of stratification, intense meso and sub-
2 mesoscale activity) all occur within the basin and Med Sea DOC concentrations and vertical
3 profiles are similar to the oceanic ones (Santinelli, 2015). It is important to remember that
4 the Med Sea is a small semi-enclosed basin characterized by low primary production
5 (Lazzari et al., 2012) and low sinking particle fluxes; only ~ 10% of open ocean values
6 (Speicher et al., 2006). It is ultra-oligotrophic and shows an anomalous N/P ratio (N:P > 25
7 in the eastern Med Sea; N:P ~ 20 in the western Med Sea; Krom et al., 2005; Ribera
8 d'Alcalà et al., 2003). Its deep waters are ~ 10°C warmer than deep waters in the ocean
9 and they are characterized by (i) fast ventilation rates and short residence times; (ii)
10 enhanced respiration (Christensen et al., 1989) and DOC mineralization rates (Santinelli et
11 al., 2010). The Med Sea thermohaline circulation is driven by the inflow of the Atlantic
12 Water (AW) at the Gibraltar strait. The AW increases its salinity during its west to east
13 transit through the basin, as a result of intense evaporation that is characteristic of the
14 region. Due to its increased salinity, AW sinks in the Levantine Basin forming the
15 Levantine Intermediate Water (LIW). LIW crosses the entire basin at 200-500 m moving
16 from east to west and finally exits through the Gibraltar Strait at depths > 100 m. Deep
17 water formation occurs every winter in both the western (West Med) and eastern (East
18 Med) basins by a mechanism analogous to the polar Atlantic deep convection.

19 An intriguing observation is that in the intermediate and deep waters of the Med
20 Sea, DOC concentrations are equal to the lowest values found in the deep Atlantic and
21 Pacific (36-42 μM ; Santinelli, 2015 and literature therein). The lowest DOC values in the
22 deep Med Sea (~ 36 μM) are unexpected since the renewal time of deep waters in the
23 basin is only 20-126 years (Andrieu and Merlivat, 1988; Schiltzer et al., 1991). Applying
24 removal rates of RDOM characteristic of the deep global ocean (~3 nM y^{-1} ; Hansell, 2013),
25 we expected that just a very small (~ 0.06-0.38 μM), nearly undetectable fraction of RDOC
26 should be removed over the residence time of deep water in the basin. The Med Sea is a
27 concentration basin (evaporation is higher than the input of freshwater) with significant
28 influxes of terrestrial carbon from Europe and North Africa. As a consequence, the
29 concentration of DOC in the deep Med Sea should be slightly higher than in the deep
30 Atlantic Ocean (~ 42 μM). The surprisingly low DOC concentrations measured in the
31 intermediate and deep waters can be explained by different hypothesis:

32 1) The processes that remove RDOC are much faster in the deep Med Sea than in
33 the deep ocean, as suggested by Hansell and Carlson (2013). The net removal of ~ 8 μM
34 RDOC takes less than 100 years in the Med Sea compared to ~ 1000 years in the deep

ocean. Faster rates of RDOC removal could be explained by (i) the higher (~ 13-14°C) temperature, characteristic of the deep Med Sea; (ii) different communities of RDOC-degrading microbes; and/or (iii) priming of bathypelagic RDOC-degrading microbes by the input of labile DOC to depth after wintertime deep mixing events. The faster removal of RDOC should not affect its radiocarbon value, we therefore expect radiocarbon values to be similar for DOC in the deep Med Sea and deep Atlantic Ocean.

2) The concentration of RDOC is < 36 µM in both the oceans and the Med Sea, but deep Atlantic oceanic DOC values are elevated due to the input of semi-labile DOC (SLDOC) from sinking particles (Follett et al., 2014). Here we define SLDOC as any carbon that accumulates in the ocean but is removed on timescales of decades or less. Our definition does not distinguish between carbon removed on seasonal and decadal timescales, and includes the semi-labile (seasonal removal) and semi-refractory (decadal removal) DOC as defined by Hansell (2013). In this scenario, a fraction of deep DOC that has been traditionally defined as RDOC (< 42 µM; lifetime of ~ 16,000 y) is not refractory with respect to the timescale of its removal, but is semi-labile, and in steady-state through supply by dissolution of sinking particles and removal by as yet unquantified processes, but most likely through microbial oxidation. The generally lower fluxes of sinking particles characteristic of the Med Sea (Speicher et al., 2006) would reduce the supply of SLDOC to depth, resulting in lower steady state concentrations of DOC. A reduction in the supply of autochthonous, surface derived, semi-labile, modern radiocarbon fraction from deep DOC will lower the radiocarbon value of the remaining organic matter. We therefore expect radiocarbon values to be more depleted (older age) in the deep Med Sea than in the deep Atlantic Ocean.

These two hypotheses represent different scenarios for deep sea DOC cycling that are not easily tested in the global ocean. The hypotheses are not mutually exclusive, and higher degradation rates for RDOC could be made more visible by lower supply of SLDOC at depth. Here we present the first stable and radioisotope data for DOC in the West Med, these data will help elucidate sources, sinks and cycling pathways for DOC in the deep ocean.

2. Material and methods

2.1. Study area and sampling stations.

Seawater samples for DOC, $\text{DOC}\Delta^{14}\text{C}$ and $\text{DOC}\delta^{13}\text{C}$ were collected in three stations located in the Tyrrhenian Sea (St. A), Sardinian Channel (St. B) and Sardinian Sea (St. C) during the oceanographic cruise WMED-BIOOPT 2012, carried out on the *R/V Urania*, from March 22nd to April 10th 2012 (Fig. 1, Table 1). Samples for DICA^{14}C were collected in the Tyrrhenian Sea during the oceanographic cruise WMED-BIOOPT 2013, carried out on the *R/V Minerva II*, April 14th 2013 (Fig. 1, Table 3). Pressure, conductivity, temperature and dissolved oxygen concentration were measured with a SBE 911 plus CTD, equipped with a rosette sampler fitted with 10 L Niskin bottles. CTD-oxygen was calibrated by Winkler titration of select discrete samples. CTD data were kindly provided by the Rome section of the Institute of Atmospheric Sciences and Climate (ISAC), Italian National Research Council (CNR) (Dr. F. Bignami).

2.2. Sample collection and storage

2.2.1 DOC

DOC samples were collected at 5, 10, 25, 50, 75, 100, 150, 200, 300, 400, 500 m and every 250 m from 500 m to the bottom. Water was collected directly from the Niskin bottles into amber glass sample bottles that were preconditioned with filtered open-sea deep water and rinsed three times with the sample before its collection. Samples were immediately filtered through sterile 0.2 μm nylon membrane filters (Minisart Nylon 25 mm, 17845 ACK) under low pressure by high-purity air. The first ~ 200 ml of the filtrate was discarded. Filtered samples were stored at 4°C in the dark until analysis (within 3 months). DOC was also measured directly from the samples used for radiocarbon analysis.

2.2.2. $\text{DOC}\Delta^{14}\text{C}$ and $\text{DOC}\delta^{13}\text{C}$

All radio and stable isotope samples were collected and processed using protocols recommended by the National Ocean Sciences Accelerator Mass Spectrometry (NOSAMS) facility in Woods Hole, MA. The protocols were developed to minimize the potential for contamination by radiocarbon used in shipboard ^{14}C tracer measurements. No contamination of shipboard work areas or Niskin sampling bottles was detected by swipe test. Samples were returned to Woods Hole for processing in a radiocarbon clean facility.

$\text{DOC } \Delta^{14}\text{C}/\delta^{13}\text{C}$ samples were collected in new, 2-10l polycarbonate bottles cleaned by soaking in 10% bleach (1 week) and 1% HCl (> 1 week) followed by > 5 rinses with high purity (> 18 M Ω) water. Samples were drawn directly from the Niskin bottles and filtered in-

line through Whatman Polycap 36 TC capsule filters (0.2 μm , cleaned by soaking in 1% HCl overnight, followed by 5l high purity water rinse and 1l sample rinse). Samples were capped, triple-bagged and frozen immediately after collection.

2.2.3 DIC $\Delta^{14}\text{C}$

Samples for dissolved inorganic carbon were collected in 500 ml glass bottles supplied by NOSAMS. Bottles were filled to overflow directly from the Niskin bottles. A small amount of the sample was removed, the bottle was poisoned with 100 μl of a saturated solution of HgCl_2 , sealed, and stored in the dark at room temperature. Further processing was done at NOSAMS using established protocols (McNichol et al., 1994).

2.3 Sample measurement

2.3.1 DOC

The DOC measurements were carried out in the laboratory at the Pisa section of the Institute of Biophysics (IBF), CNR, with a Shimadzu TOC-VCSN, equipped with a quartz combustion column filled with 1.2% Pt on alumina pillows of ~ 2 mm diameter. Samples were acidified with 2N high purity HCl and sparged for 3 min with CO_2 -free ultra-high purity air in order to remove inorganic carbon. 150 μl of the sample were injected in the furnace after a three-fold rinsing with the sample to be analyzed. From 3 to 5 replicate injections were performed until the analytical precision was lower than 1%. A four-point calibration curve was done with standard solutions of potassium hydrogen phthalate in the same concentration range as the samples. The system blank was measured every day at the beginning and the end of analyses using low-carbon water (2-3 μM C) produced by a Milli-Q system. Measurement reliability was assessed twice daily by comparison of data with DOC Consensus Reference Waters (CRM) (Hansell, 2005) (batch #12, measured concentration: 41.7 ± 1.2 μM , standard error 0.10 μM , $n = 28$). This quality assurance procedure ensures the integrity of the calibration curve (Sharp et al., 2002).

2.3.2 DOC $\Delta^{14}\text{C}$ and DOC $\delta^{13}\text{C}$

Samples for DOC $\Delta^{14}\text{C}/\delta^{13}\text{C}$ were processed at NOSAMS using protocols detailed in Beaupré et al. (2007). Briefly, approximately 1L of sample was transferred to an all-quartz reaction flask and acidified to pH ~ 2 with high purity phosphoric acid. The flask was placed in a gas extraction apparatus and sparged with helium to remove carbon dioxide.

DOC was oxidized by UV irradiation, and the resulting carbon dioxide stripped with a helium sparge and cryogenically trapped. Further details are provided on the NOSAMS website at http://www.whoi.edu/nosams/Submitting_Guidelines.

3. Results

3.1. Hydrologic characteristics

Potential temperature/salinity (θ/S) diagrams allow for the identification of the major water masses occurring in the study area: Atlantic Water (AW), Levantine Intermediate Water (LIW), Western Mediterranean Deep Water (WMDW) and Tyrrhenian Deep Water (TDW) (Fig. 2). The AW ($S < 37.2$, $\theta = 15.0$ - 16.5 °C), occupies the surface layer of station B, the marked increase in S at stations A and C ($S = 37.9$ - 38.1) indicates that these stations are affected by a modified vein of AW, probably coming from the East Med. The thermocline is not very strong, with surface temperatures (0-10 m) between 15.0 - 16.8 °C at all three stations (Fig. 2a and 3).

The LIW is easily recognizable by its salinity maximum (38.6 - 38.74). Its core becomes more mixed (less salty and colder) moving from station A (330-600 m; $S = 38.70$ - 38.74 , $\theta = 13.9$ - 14.1 °C) to station B (280-600 m; $S = 38.65$ - 38.70 , $\theta = 13.7$ - 14.0 °C) and station C (250-600 m, $S = 38.55$ - 38.61 , $\theta = 13.25$ - 13.73 °C), where it is still visible (Fig. 2b and 3). The vertical profiles of oxygen clearly show a minimum (< 183 μM) in the core of LIW (200-500 m), with a ~ 10 μM decrease moving from station A to C (Fig. 3).

The WMDW ($S \sim 38.47$; $\theta = 12.89$ - 12.92 °C; $\text{DO} = 194$ - 198 μM) can be observed at stations B and C (Fig. 2c and 3). This water mass forms during intense, wintertime cooling events in the Gulf of Lions and spreads throughout the deep western basin. At both stations a very dense ($\sigma_\theta > 29.11$ Kg m^{-3}), salty ($S > 38.47$) tongue of WMDW is clearly visible. WMDW is slightly warmer at station B (Fig. 2c). This water mass is most likely the residual of the “anomalous” deep water formation event in 2005-2006 (López-Jurado et al., 2005; Schroeder et al., 2006). Changes in the physical properties of the WMDW were related either to the severity of weather conditions at the time of formation in winter 2004/2005 or to the transfer of the Eastern Mediterranean Transient (EMT) (Malanotte-Rizzoli et al., 1999; Roether et al., 1996) signal to the West Med (López-Jurado et al., 2005; Schroeder et al., 2006). At station A, deep water is warmer (12.99 - 13.04 °C), saltier (38.49 - 38.51) and lower in oxygen (186 - 188 μM) than the WMDW (Fig. 2c and 3). These characteristics are typical of the TDW, formed by the mixing between the LIW and the

WMDW. The very low oxygen values indicate the absence of recently ventilated deep water in the Tyrrhenian Sea (Fig. 3).

Apparent Oxygen Utilization (AOU) values (Table 1) are in the range of those reported for the TDW (57-71 μM) and the WMDW (52-73 μM) in 2008 (Santinelli et al., 2010), while they are higher than those observed in the WMDW in 2005 (43-56 μM), confirming the exceptional characteristics of the WMDW formed in winter 2005 and the removal of oxygen with time due to the mixing with the resident waters and the microbial respiration.

3.2. DOC vertical profiles

DOC vertical profiles show the classical shape with the highest concentrations in the surface water (60-65 μM), similar to values in the North Atlantic subtropical gyre, and a gradual decrease to 39-40 μM at ~ 300 m. Below 500 m, the values remain very low (38-42 μM) until the bottom (Fig. 4). DOC concentration and vertical distribution is in agreement with the data reported for the Tyrrhenian Sea (Santinelli et al., 2013) and the West Med (Santinelli et al., 2010).

Focusing on the deep water, it is noteworthy that DOC concentrations are the same as those reported in 2008 in the region, but they are markedly lower than those observed in 2005 in the WMDW (56 ± 9 μM , Santinelli et al., 2010). The 14 μM (56-42 μM) decrease is in agreement with the AOU increase and suggests that a fraction of DOC, exported to depth during the exceptional event of deep water formation in winter 2005, was removed on the short temporal scale.

3.3. Carbon Isotopes

Samples for radiocarbon measurements were collected in the surface layer (2 m), in the core of LIW (480 m, maximum of S) at station B, in the WMDW (the deepest samples at stations B and C), in the TDW (the deepest sample at station A), and at intermediate depths (1500 m at stations A and C; 2500 m at station A).

In the surface layer, $\text{DOC}\Delta^{14}\text{C}$ ranges between -237 to -286‰, suggesting a radiocarbon age of 2120-2640 years. $\text{DOC}\delta^{13}\text{C}$ values range from -22.0 to -22.7‰ (Table 2 and Fig. 4). Just one sample was collected in the LIW with $\text{DOC}\Delta^{14}\text{C}$ of -425‰ and $\text{DOC}\delta^{13}\text{C}$ of -22.8‰, indicating the presence of DOC that is older than in the surface layer,

but younger than in the deep waters (Table 2 and Fig. 4). $\text{DOC}\Delta^{14}\text{C}$ in the deep waters (1500-bottom) have values between -445 to -475‰, indicating a radiocarbon age of 4670-5120 years old. $\text{DOC}\delta^{13}\text{C}$ ranges between -23.0 and -24.2‰ (Table 2 and Fig. 4).

$\text{DIC}\Delta^{14}\text{C}$ values for the Tyrrhenian Sea (Station A, Table 3) decrease from +31.3‰ in the surface to +18.6‰ in the core of the LIW at 400 m, reach a minimum of -34.6‰ at 2500 m, before rebounding slightly to -22.6‰ in the deepest sample collected at 3390 m. These values confirm that the Med Sea waters are renewed in less than 126 years (post-bomb values).

4. Discussion

4.1. $\text{DIC}\Delta^{14}\text{C}$ and ventilation time of Med Sea waters

The vertical pattern of $\text{DIC}\Delta^{14}\text{C}$ values is similar to an earlier report by Broecker and Gerard (1969) for samples collected in the West Med and in the Tyrrhenian Sea in summer 1956 and 1958, before significant bomb radiocarbon had penetrated the basin (Fig. 1 and Table 3). In the West Med Broecker and Gerard (1969) reported a value of -48‰ for surface water, -60‰ for LIW (320 m), and -69‰ for deep water (2700 m) (Table 3). A fourth sample collected from the deep (3390 m) Tyrrhenian Sea, close to our station A (Fig. 1), was significantly older, with a $\text{DIC}\Delta^{14}\text{C}$ value of -83‰. The difference in radiocarbon values from the deep West Med and the Tyrrhenian Sea reflects more active deep water formation in the West Med, as indicated by higher oxygen concentrations and lower values for AOU at our stations B and C compared to station A (Table 1). The ~+80‰ increase in surface water $\text{DIC}\Delta^{14}\text{C}$ values, and ~40-50‰ increase in intermediate and deep water values are indicative of the rapid penetration of bomb radiocarbon to all depths in the basin, in agreement with Broecker and Gerard's earlier assessment of deep Mediterranean ventilation times of < 100 y for the West Med and < 200 y for the Tyrrhenian Sea.

4.2 Radiocarbon in surface water, source and age of SLDOC in the Med Sea

As expected from past studies on radiocarbon at open ocean sites, $\text{DOC}\Delta^{14}\text{C}$ values are more highly depleted than $\text{DIC}\Delta^{14}\text{C}$ throughout the entire water column, reflecting the longer residence time of organic carbon in seawater. In the euphotic zone,

photosynthesis fixes carbon with $\Delta^{14}\text{C} \sim \text{DIC}\Delta^{14}\text{C}$ (+31‰) into SLDOC that is slowly degraded by microbial cycling. The depletion in surface water $\text{DOC}\Delta^{14}\text{C}$ (~ -237 to -286‰, Table 2) relative to $\text{DIC}\Delta^{14}\text{C}$ (+31‰, Table 3) can be explained by a two-component mixture of modern radiocarbon age SLDOC and old radiocarbon age DOC. Since no strong upwelling regions occur in the Med Sea (excluding the Gulf of Lions) the source of the old DOC in the surface layer is most likely the AW inflow. Surface carbon isotope values for both the $\text{DOC}\delta^{13}\text{C}$ (-21.9 to -22.7‰) and the $\text{DOC}\Delta^{14}\text{C}$ (-237 to -286‰) are slightly lower (more negative) than those measured in the surface Atlantic Ocean ($\text{DOC}\delta^{13}\text{C} = -21.0$ to -21.3 ‰, $\text{DOC}\Delta^{14}\text{C} -210$ to -253 ‰) (Druffel et al., 1992) even though DOC values are quite similar (59-65 μM for the North Atlantic vs 60-65 μM for the Mediterranean Sea) (Table 2 and Fig. 4). For radiocarbon, these differences could arise from changes in the isotopic value of SLDOC due to differences in location and time of sampling. Bomb radiocarbon reached maximal values in ocean subtropical gyres in the mid to late 1970s (McNichol and Aluwihare, 2007) and has been decreasing since that time. $\text{DIC}\Delta^{14}\text{C}$ values measured by Druffel et al. (1992) in North Atlantic surface waters (< 100 m) in 1991 were ~ +115‰, considerably higher than values we measured in the Mediterranean Sea 21-22 years later (+31‰). Assuming the RDOC in AW inflow has an isotopic value similar to RDOC in deep Atlantic Ocean water (> 1000 m, $\text{DOC} = 43$ μM ; $\text{DOC}\Delta^{14}\text{C} = -394$ ‰; Druffel et al., 1992), we can calculate that in the Med Sea surface waters (mean $\text{DOC} = 62$ μM ; mean $\text{DOC}\Delta^{14}\text{C} = -267$ ‰, Fig. 4 and Table 2), $\text{SLDOC}\Delta^{14}\text{C}$ would have a value of ~ +20‰, similar to the $\text{DIC}\Delta^{14}\text{C}$, using the following equation.

(equation 1)

$$(\text{DOC}\Delta^{14}\text{C})_{\text{surfMed}} \cdot [\text{DOC}]_{\text{surfMed}} = (\text{DOC}\Delta^{14}\text{C})_{\text{deepNA}} \cdot [\text{DOC}]_{\text{deepNA}} + (\text{SLDOC}\Delta^{14}\text{C}) \cdot ([\text{DOC}]_{\text{surfMed}} - [\text{DOC}]_{\text{deepNA}})$$

where surfMed is the average of the surface (2 m) Med Sea samples (Fig. 4 and Table 2) and deepNA is the average of samples collected in the deep (> 1000 m) North Atlantic.

4.3. Radiocarbon in deep water, source and age of RDOC in the Med Sea

The small number of samples and spatial placement of our study sites does not allow for meaningful interpretation of differences in $\text{DOC}\Delta^{14}\text{C}$ between deep water samples. For the purposes of the following discussion, we therefore take a deep water average value of -460 ± 12 ‰ (> 1000 m, $n = 6$) to be representative of deep Med Sea

1 DOC $\Delta^{14}\text{C}$. This value is lower than the average RDOC $\Delta^{14}\text{C}$ value of $-394 \pm 13\text{‰}$ for the
2 deep ($> 1000\text{ m}$, $n = 9$) North Atlantic reported by Druffel et al (1992), even though DOC
3 concentrations in the two studies were comparable ($43 \pm 1\text{ }\mu\text{M}$ in the deep North Atlantic,
4 $42 \pm 2\text{ }\mu\text{M}$ in the deep Mediterranean Sea). Flerus et al. (2012) reported DOC $\Delta^{14}\text{C}$ values
5 for a fraction of RDOC that was recovered by solid phase extraction onto a hydrophobic
6 resin along a North-South cruise track that passed just west of the Gibraltar Strait. Solid
7 phase extractable RDOC $\Delta^{14}\text{C}$ values ranged from ~ -225 to -370‰ in surface ($< 100\text{ m}$)
8 water and $\sim -450\text{‰}$ in deep ($> 500\text{ m}$) water. Druffel et al. (1992) found similar values ($-$
9 329 to -402‰ surface, -454 to -587‰ deep) for humic substances adsorbed onto
10 hydrophobic solid phases (XAD) from Sargasso seawater. In all cases, the $\Delta^{14}\text{C}$ value of
11 adsorbed DOC in deep water was more depleted (older) than in whole DOC by -40 to $-$
12 173‰ . These studies demonstrate that within deep North Atlantic DOC, there is a fraction
13 (at least 20-25%) with radiocarbon values as depleted as deep Med Sea DOC $\Delta^{14}\text{C}$.

14 Atlantic DOC enters the Med Sea through Gibraltar in the AW inflow. During AW
15 transit into the basin and its transformation into Med Sea intermediate and deep waters,
16 Atlantic SLDOC is removed and replaced by SLDOC released in the surface layer of the
17 Med Sea. However, Atlantic RDOC, which makes up $\sim 70\%$ of the total DOC, should
18 remain. Due to the comparable concentrations of DOC in the deep Atlantic and Med Sea
19 and the relatively short ventilation time of the Med Sea, a similar age for DOC was
20 expected. The $\sim 66\text{‰}$ depletion of deep Med Sea DOC $\Delta^{14}\text{C}$ (-460‰), compared to deep
21 North Atlantic DOC $\Delta^{14}\text{C}$ (-394‰) was therefore surprising. The depletion in DOC $\Delta^{14}\text{C}$
22 cannot be due to radioactive decay of ^{14}C , since renewal of water in the Med Sea is too
23 fast ($< 126\text{ y}$) compared to the half-life of ^{14}C (5730 y). The depletion must therefore arise
24 from the rapid replacement of a fraction of “refractory” DOC coming from the Atlantic
25 (RDOC_{Atl}) with Med Sea DOC having an older radiocarbon age (DOC_{Med}).

26 The minimum amount of RDOC_{Atl} replacement needed to explain the older deep
27 Mediterranean values can be calculated if we assume DOC_{Med} to be fully depleted in ^{14}C ($-$
28 1000‰). In this case, a $4\text{ }\mu\text{M}$ ($\sim 10\%$) replacement of RDOC_{Atl} with DOC_{Med} will yield the
29 observed deep DOC $\Delta^{14}\text{C}$ value. Carbon autochthonous to Med Sea, produced by
30 microbes, cannot be older than 126 y (the renewal time of Med Sea waters) and therefore
31 should have $\Delta^{14}\text{C}$ values similar to DIC (-34 to $+31\text{‰}$, Table 3). As a consequence, to
32 lower the $\Delta^{14}\text{C}$ value of deep Med Sea DOC, DOC_{Med} must have a source external to the
33 basin.

The average deep Mediterranean $\text{DOC}\delta^{13}\text{C}$ value is -23.7‰ , which is significantly lighter than $\text{DOC}\delta^{13}\text{C}$ in the deep Atlantic or Pacific (-20.8‰ and -21.1‰ respectively), but similar to $\text{DOC}\delta^{13}\text{C}$ in the Canada Basin (Griffith et al., 2012). The more negative $\text{DOC}\delta^{13}\text{C}$ value can be explained by the replacement of a fraction of RDOC_{Atl} by isotopically light DOC_{Med} . Assuming that the minimum $4\text{ }\mu\text{M}$ DOC_{Med} needed to account for low $\text{DOC}\Delta^{14}\text{C}$ is the same carbon that lowers $\text{DOC}\delta^{13}\text{C}$, DOC_{Med} should have a $\delta^{13}\text{C}$ value of $\sim -48\text{‰}$, and an annual flux of $1.9\text{-}2.5 \times 10^{+12}\text{ g C y}^{-1}$, obtained by multiplying the amount of carbon replaced ($4\text{ }\mu\text{M}$) by the inflow at the Gibraltar Strait ($1.28\text{-}1.68\text{ Sv}$; Hopkins 1999; Bethoux and Gentili, 1999). Isotopically light carbon with annual fluxes of this magnitude are not characteristic of most sources allochthonous to the Med Sea (Table 4).

Potential sources of DOC_{Med} that need to be evaluated for their flux and isotopic value include (1) fossil methane and methane derived DOC seeps from sediments, (2) anthropogenic combustion products and terrestrial organic matter delivered by the atmosphere, rivers, and groundwaters.

4.4 Potential sources of DOC_{Med}

4.4.1 Methane

Methane arises from the microbial reduction of sedimentary organic matter and carbon dioxide during diagenesis (biogenic methane) or from thermal cracking of sedimentary carbon, usually deep in the geosphere under elevated temperatures over long periods of time (thermogenic methane). Biogenic processes yield methane that is isotopically lighter (typically -50‰ to $< -80\text{‰}$) than thermogenic methane (typically -40‰ to -55‰), although there is some overlap (Whiticar et al., 1986). Transformation of methane into biomass by methanotrophs leads to further fractionation of $\sim -30\text{‰}$ (Summons et al., 1994). The radiocarbon value of biogenic methane can range from fully modern to highly depleted, depending on the radiocarbon content of the organic matter or carbon dioxide used as substrate, while thermogenic methane is thought to be fully depleted in radiocarbon ($\sim -1000\text{‰}$). We are not aware of any isotopic measurements of methane in the Med Sea. A few measurements of $^{13}\text{C}\text{-CH}_4$ from pore waters of the Adriatic Sea and Gulf of Lions sediments suggest a biogenic origin for shallow gas in the Adriatic Sea (^{13}C -70 to -92‰ ; Garcia-Garcia et al., 2007) and perhaps mixed biogenic/thermogenic origin in the Gulf of Lions (^{13}C -53 to -82‰ ; Garcia-Garcia et al., 2006).

1 The total flux of methane from sediments places an upper bound on the potential
2 contribution of methane to DOC_{Med} . Biogenic and thermogenic methane enters the Med
3 Sea either by direct injection of methane charged muds (mud volcanoes), venting from
4 pockmarks and fissures (gas bubbles and flares) or through diffusive fluxes from
5 subsurface sediments with high methane concentrations (Boetius and Wenzhöfer, 2013,
6 Romer et al, 2014). The large number of potential input mechanisms, and the spatial and
7 temporal variability of these inputs makes a basin-wide estimate of methane flux difficult.
8 Using hydroacoustic profiling of methane flares in the water column, sediment core
9 incubations, and benthic chambers, Romer et al. (2014) estimated the annual flux of
10 methane to the water column to range from 0.4 to $94 \times 10^{+9} \text{ g C y}^{-1}$ in a 225 km^2 section of
11 the Nile deep sea fan. If this flux is scaled to the entire Med Sea, the annual flux of
12 methane from sediments would range from 0.4 to $104 \times 10^{+13} \text{ g C y}^{-1}$. This value most
13 likely overestimates the flux of methane into the deep Med Sea by a significant amount,
14 since many areas of the Med Sea floor lie outside the methane hydrate stability zone and
15 are characterized by carbon-lean sediments, with little potential for methane generation or
16 storage.

17 As a second approach, we weighted the estimated global sediment to seawater flux
18 of methane ($0.03\text{-}0.05 \text{ Pg C y}^{-1}$, with an additional 0.03 Pg C y^{-1} from deep sea mud
19 volcanoes) (Kvenvolden et al., 2001, Milkov et al, 2003) to the areal extent of sediments
20 that lie within the methane hydrate stability zone in the Med Sea. Using this approach the
21 basin-wide annual methane flux is markedly lower $\sim 3\text{-}4 \times 10^{+11} \text{ g C y}^{-1}$.

22 In the water column, methane can escape to the atmosphere, be oxidized to CO_2 ,
23 or fixed into microbial biomass with a fraction subsequently released as DOC. We are
24 unaware of any measurements of the conversion efficiency of methane to DOC by either
25 aerobic or anaerobic methane oxidizers. If, from estimates of bacterial growth efficiency on
26 marine DOC, we assume 10-30% of methane is fixed into biomass by bacteria, that 10%
27 of the fixed carbon is released as DOC (Carlson, 2002), and that $\sim 10\%$ of this DOC is
28 refractory (Ogawa et al., 2001), methane could potentially contribute to DOC_{Med}

29 Alternatively, we might assume that the conversion of methane to DOC has an
30 efficiency similar to the conversion of net marine primary production to RDOC. Taking into
31 account that less than 1% of the net annual production in the open ocean is required to
32 maintain the marine reservoir of RDOC at steady state with a radiocarbon age of 16,000 y
33 (Hansell, 2013), then the contribution of fossil methane carbon to DOC_{Med} is probably less
34 than $3 \times 10^{+9}$ to $1 \times 10^{+13} \text{ g C y}^{-1}$.

Methanotrophy in surface sediments may add isotopically depleted DOC directly into the water column (Pohlman et al. 2011; Boetius and Wenzhöfer, 2013) further increasing the flux of methane sourced carbon to DOC_{Med}. We are not aware of any current estimates of direct input of methane derived DOC from sediments at any site, and we cannot estimate the magnitude of this additional flux. Uncertainties in the annual flux of fossil methane into the deep Med Sea, and the amount of methane carbon that is transformed into RDOC are so large that it is unclear if fossil methane contributes significantly to DOC_{Med}. However, unless future estimates of methane flux for the region or the methane to DOC conversion efficiency by methanotrophs are revised substantially upwards, it seems unlikely that methane contributes more than a fraction to DOC_{Med}, and other sources of aged carbon are needed to explain the isotopic depletion of deep Med Sea DOC.

4.4.2 Rivers and atmosphere

Assuming that DOC in Med Sea deep water is a mixture of marine DOC ($\delta^{13}\text{C} \sim -21\text{‰}$) coming from the Atlantic Ocean and terrestrial DOC ($\delta^{13}\text{C} \sim -27\text{‰}$) coming from rivers, atmosphere and marine groundwaters surrounding the Med Sea, we can calculate that 55% ($\sim 23 \mu\text{M}$) of deep Med Sea DOC is marine, and 45% ($\sim 19 \mu\text{M}$) is of terrestrial origin. If the $23 \mu\text{M}$ of marine DOC had a $\Delta^{14}\text{C}$ value equal to deep North Atlantic DOC (-394‰) then the $\Delta^{14}\text{C}$ value of the terrestrial component would be $\sim -540\text{‰}$ and it should have an annual flux of $9\text{--}12 \times 10^{+12} \text{ g C y}^{-1}$, estimated multiplying the Atlantic inflow at Gibraltar ($1.28\text{--}1.68 \text{ Sv}$) by $19 \mu\text{M}$. The total estimated terrestrial input of DOC to the Med Sea is $1\text{--}13 \times 10^{+12} \text{ g C y}^{-1}$, including both river ($0.6\text{--}0.7 \times 10^{+12} \text{ g C y}^{-1}$) and atmospheric ($0.4\text{--}13 \times 10^{+12} \text{ g C y}^{-1}$) DOC inputs (Santinelli, 2015 and literature therein). Additionally, rivers deliver between $0.7\text{--}1.0 \times 10^{+12} \text{ g C y}^{-1}$ of particulate organic carbon to the Med Sea, some fraction of which may become solubilized as DOC after delivery.

River input was estimated considering all the available data that include the major Med Sea rivers (Rhône, Po and Ebro) and some of the minor ones. DOC concentrations are from multi-year observations in the Rhône river (Sempéré *et al.*, 2000; Panagiotopoulos *et al.*, 2012), Po river (Pettine *et al.*, 1998) and Arno river (Retelletti *et al.*, 2015). In contrast, DOC data are taken sporadically in the other rivers. If we consider the discharge (Ludwig *et al.*, 2009), DOC data are available for 50% of the rivers (81% in the West Med and 34% in the East Med). DOC concentrations ranged between 133 and 366 μM in the West Med and between 92 and 292 μM in the East Med. The lack of DOC data

1 in some rivers was overcome by using the mean DOC concentration computed by all the
2 available data (176-225 μM in the West Med and 157 μM in the East Med) and multiplying
3 this value by the discharge of the rivers without DOC data (for further details refer to
4 Santinelli, 2015).

5 Data on wet and dry DOC depositions from the atmosphere are scarce for the Med
6 Sea. DOC fluxes from the atmosphere were therefore estimated by using the largest range
7 calculated referring to three papers. Pulido-Villena *et al.*, (2008) report direct DOC
8 concentration on total atmospheric deposition, collected at the Cap Ferrat (West Med Sea)
9 continuously from December 2005 to December 2006. These authors observed an
10 atmospheric DOC input ranging from 0.04 to 1.2 $\text{mmol m}^{-2}\text{d}^{-1}$, with the highest values
11 associated with Saharan dust deposition events. Economou and Mihalopoulos, (2002)
12 measured TOC in 47 rain samples collected at Heraklion (Creta Island, East Med sea)
13 from September 1999 to May 2000. They observed concentrations ranging from 16.6 to
14 358 μM , with a volume weighted mean of 100 μM . An annual wet deposition for TOC of
15 0.14 $\text{mmol m}^{-2} \text{d}^{-1}$ was calculated multiplying 100 μM by an annual precipitation rate of 0.5
16 m. De Vicente *et al.*, (2012) measured the concentration of water-soluble organic carbon
17 (WSOC), in both dry and wet deposition, collected in 2004 and 2005 by a passive sampler
18 deployed in three reservoirs located in Southern Spain within an area that experienced
19 intense dust deposition. The annual average concentrations of WSOC, considering both
20 dry and wet depositions, ranged from 0.18 to 0.42 $\text{mmol m}^{-2} \text{d}^{-1}$.

21 The minimum and maximum of the fluxes reported in these three papers were
22 multiplied by the Med Sea area ($2.5 \times 10^6 \text{ km}^2$), in order to have a regional estimate of
23 atmospheric input. This estimate is affected by large uncertainties, since it is based on the
24 assumption that this range is valid for the whole basin and for the whole year. The flux
25 could be underestimated since Saharan dust deposition, which seems to play a major role
26 on DOC input, is probably more intense and frequent in the southern part of the West Med,
27 where no data are available. On the other hand, Saharan dust deposition doesn't occur
28 every day, leading to an overestimation of the fluxes if the highest values are taken into
29 account. The most reasonable range is the one based on annual average (0.14 to 0.42
30 $\text{mmol m}^{-2} \text{d}^{-1}$) and it suggests an average DOC flux of $1.3\text{-}4.6 \times 10^{+12} \text{ g C y}^{-1}$.

31 Recently, Total Dissolved Nitrogen (TDN) and Phosphorus (TDP) fluxes from the
32 atmosphere were studied in bulk deposition samples simultaneously collected at 10
33 locations, around the Med Sea, during one year period (June 2001–May 2002) (Markaki *et al.*
34 *et al.*, 2010). These data are more representative of integrated fluxes to the basin, but no

DOC data are available in this study. The authors observed dissolved inorganic nitrogen (DIN) fluxes ranging from 33.2 to 36.6 mmol m⁻² y⁻¹ and dissolved organic nitrogen (DON) to be on average 32% of TDN, DON fluxes should therefore be 15.6-17.2 mmol m⁻² y⁻¹. Assuming a DOC/DON ratio in atmospheric dissolved organic matter (DOM) ranging between 10 and 18, typical ranges of Med Sea DOM (Pujo Pay et al., 2011; Santinelli et al. 2012), atmospheric DOC input should range from 4.7 to 9.3 x 10⁺¹² g C y⁻¹. These values are slightly higher than those estimated by taking into consideration direct DOC measurements on atmospheric deposition.

No DOCΔ¹⁴C data is available for Med Sea rivers, however we assume that the values are similar to the range (+364 to -158‰) reported by Raymond and Bauer, (2001) for temperate and tropical rivers that drain a wide variety of terrestrial watershed types, including large population centers (Potomac, Washington D.C.; Susquehanna, Philadelphia; Hudson, New York; all USA). Wastewater treatment associated with population centers may contribute a substantial fraction of petroleum derived carbon (Griffith et al., 2009).

Atmospheric DOCΔ¹⁴C spans a slightly wider range of values from +49 to -522‰, reflecting inputs from modern terrestrial biomass to fossil fuel combustion in urban areas (Table 4).

Estimates of carbon flux and isotopic range for terrestrial DOC inputs are based on only a small number of measurements, and have large uncertainties when extrapolated to the entire basin over decadal timescales. However, the low flux, enriched isotopic values, and lability of riverine DOC (5-50% removed in months) (Holmes et al., 2008; Lønborg et al., 2009; Retelletti et al., 2015) suggest rivers contribute only a minor fraction of DOC_{Med}, although processing of riverine particulate organic matter (0.7-1.0 x 10⁺¹² g C y⁻¹) may further increase this contribution. It is noteworthy that most of the carbon in the Rhone River continental shelf is much older (POCΔ¹⁴C = -310±66‰) than would be expected if it had been derived from contemporarily synthesized sources. The authors explained these heavily depleted values with the presence of aged organic carbon, such as kerogens, graphite, petroleum residues or black carbon (Cathalot et al., 2013). The flux and isotopic value of DOC delivered from the atmosphere, however, could be sufficient to supply DOC_{Med}, even if only a fraction of this carbon is refractory.

4.4.3 Submarine groundwater

Submarine groundwater may represent another crucial source of pre-aged terrestrial DOC and DIC to the Med Sea (Moore, 2010). Recently, using ²²⁸Ra mass

balance, Rodellas et al. (2015) estimated that submarine groundwater discharge (SGD) to the Med Sea is $0.3\text{-}4.8 \cdot 10^{+12} \text{ m}^3 \text{ y}^{-1}$. This flux is equal to or up to 16 times larger than river input ($0.3 \cdot 10^{+12} \text{ m}^3 \text{ y}^{-1}$) and it represents a significant source of nutrients (N, P, Si) to the Med Sea (DIN: $(20\text{-}1500) \cdot 10^{+9} \text{ mol y}^{-1}$; DIP: $(0.1\text{-}4.6) \cdot 10^{+9} \text{ mol y}^{-1}$; DSi $(20\text{-}650) \cdot 10^{+9} \text{ mol y}^{-1}$) (Rodellas et al., 2015). The only paper reporting total organic carbon concentration (TOC) in Mediterranean SGD refers to the coast of the island of Majorca (Basterretxea et al., 2010). These authors measured a TOC concentration ranging between 80 and 169 μM in 4 embayments. Assuming these values are representative of SGD to the Med Sea SGD could account for a TOC flux of $0.3\text{-}9.7 \times 10^{+12} \text{ g C y}^{-1}$. This estimate includes uncertainties, but suggests that the magnitude of SGD can be relevant and cannot be overlooked. It is also important to remember that submarine groundwaters are an important source of aged DIC (Gramling et al., 2003), and that if recycled through bacteria to DOC, SGD could contribute to the synthesis of pre-aged DOC.

4.5 Refractory DOC cycling

Stable and radiocarbon isotope values of deep Med Sea DOC suggest at least 10% and up to 45% of the Atlantic RDOC entering the Med Sea is removed and replaced by isotopically depleted DOC in less than 126 years. This finding can be explained by two hypothesis:

1. Unique conditions of the deep Med Sea allow for the removal of a substantial fraction (up to 45%) of “refractory” DOC imported from the Atlantic. Carbon that is refractory in the global ocean becomes semi-labile in the Med Sea.
2. Not all of the $\sim 40 \mu\text{M}$ RDOC entering to the Med Sea is refractory (cycles on timescales of 1000s of years). A fraction of RDOC is in fact semi-labile since it can be removed in < 126 years, and the true “refractory” fraction of DOC_{Atl} is only $\sim 20\text{-}25 \mu\text{M}$. This hypothesis posits that the deep DOC inventory is at a steady state. A fraction is continuously removed and replaced by semi-labile DOC released by sinking POC or produced by chemoautotrophy and/or other processes (Follett et al., 2014). In the Med Sea primary production and the flux of sinking autochthonous POC are both low (Lazzari et al., 2012; Speicher et al., 2006), and some RDOC is replaced by older and lighter DOC from allochthonous sources. Using current information, it is not possible to clearly identify the relative contribution of potential

allochthonous sources. However, based on estimated flux and isotopic value, atmospheric inputs of soluble organic carbon from soils or combustion products and DOC from submarine groundwaters are the most likely sources of DOC_{Med}. Atmospheric soluble organic carbon and DOC by SGD can be then transported to depth by deep water formation or by sorption onto/desorption off of sinking particles.

We can think of no reason or evidence to suggest that the Med Sea should be different from the global ocean with respect to the lability of RDOC (hypothesis 1). If RDOC persists in the ocean for several millennia, then it spends significant time in the surface ocean where temperature, salinity, and nutrient concentrations all bracket conditions found in the deep Med Sea. Exposure to elevated temperatures in surface waters should draw down RDOC values to the values ~ 20-25 μM that we find in the deep Med Sea. However, the high fidelity with which the two-component model (TCM) replicates the observed distribution of DOC radiocarbon values in the surface ocean (Williams and Druffel, 1987; Druffel et al., 1992; Beaupré and Aluwihare, 2010) suggests RDOC is not substantially degraded. Additionally, Griffith et al. (2012) report DOC $\delta^{13}\text{C}$ (~ -23 to -24 ‰) and DOC $\Delta^{14}\text{C}$ values (-400 to -500 ‰) for deep sea DOC in the Canadian Basin of the Arctic Ocean, similar to values we observe in the deep Med Sea. DOC concentration values in Griffith's samples are ~ 40-42 μM . The DOC $\delta^{13}\text{C}$ values in particular suggest that replacement of 30 % RDOC has occurred, even though temperatures are 0 °C to -1 °C throughout the deep basin.

Hypothesis 2 is consistent with a growing body of radiocarbon measurements made in the deep sea (Druffel and Bauer, 2000; Griffith et al, 2012; Follett et al, 2014) that are otherwise difficult to explain. In this scenario, replacement of SLDOC upwelled from the deep ocean with SLDOC in the surface ocean would not impact its radiocarbon value, and the distribution of radiocarbon predicted by the TCM would be unaffected.

Our calculations assume that the RDOC_{Atl} has an isotopic value equal to DOC in the deep North Atlantic (43 μM ; $\delta^{13}\text{C}$ = -21‰; $\Delta^{14}\text{C}$ = -394‰). Recently, Follett et al (2014) used serial oxidation of deep sea DOC from the North Pacific to measure its isotopic diversity and distribution. The results show a continuum of isotopic values from -1000‰ to +150‰ indicating that deep Pacific DOC includes organic matter from post-bomb to fossil carbon sources. The authors divided deep Pacific DOC into three fractions: refractory carbon (~ 24 μM) with $\Delta^{14}\text{C}$ between -1000‰ and -600‰, semi-labile carbon (~ 9 μM) with a modern (+50‰) radiocarbon value, probably supplied by sinking particles, and an aged

1 carbon fraction ($\sim 7 \mu\text{M}$, $\sim -275\text{‰}$, similar to deep sea $\text{DIC}\Delta^{14}\text{C}$) that could represent
2 carbon derived from meso- and bathypelagic chemoautotrophy or from entrainment and
3 advection of semi-labile DOC. If the concentration of RDOC in the deep North Atlantic is
4 equal to the North Pacific, and the radiocarbon values of these fractions are adjusted for
5 aging by the difference in $\text{DIC}\Delta^{14}\text{C}$ value (-146‰) between the deep North Atlantic (-84‰)
6 and Pacific (-230‰), we can calculate the amount and radiocarbon value of RDOC in the
7 deep N. Atlantic as $\sim 24 \mu\text{M}$ at -654‰ and $\sim 7 \mu\text{M}$ at -129‰ (Fig. 5). By this scenario, only
8 $12 \mu\text{M}$ of labile, modern ($\Delta^{14}\text{C} \sim -28\text{‰}$) carbon would need to be replaced by $11 \mu\text{M}$ carbon
9 with $\Delta^{14}\text{C}$ of -247‰ and $\delta^{13}\text{C}$ of -31‰ to achieve the deep Med Sea $\text{DOC}\Delta^{14}\text{C}$ value of $-$
10 460‰ and $\delta^{13}\text{C}$ value of -23.7‰ (Fig. 5b). An annual flux of $5.5\text{--}7.0 \times 10^{+12} \text{ g C y}^{-1}$ would
11 be needed to support this replacement. Alternatively, if the $7 \mu\text{M}$ of aged carbon is
12 supplied by chemoautotrophy and is included in the semi-labile fraction, then $19 \mu\text{M}$ of
13 carbon could be replaced by $18 \mu\text{M}$ DOC_{Med} with isotopic values of -201‰ and -27‰ , in
14 the range of $\text{DOC}\Delta^{14}\text{C}$ and $\text{DOC}\delta^{13}\text{C}$ values for riverine and atmospheric input (Fig. 5c).

15 The two hypotheses are not mutually exclusive. They represent different scenarios
16 for deep sea DOC cycling that can be addressed with further measurement and modeling.
17 If hypothesis 2 is correct, then this would necessitate a re-evaluation of traditional
18 definitions and inventories of refractory marine DOC in the deep ocean. Further, the
19 hypothesis more intimately ties the distribution of deep sea DOC to surface production in
20 the surface and the to the flux and dissolution of sinking particulate organic matter. It
21 suggests that a significant fraction of accumulated DOC in the deep sea is ultimately
22 utilized by the microbial community.

23

24

25 5. Conclusions

26 The first radiocarbon data on DOC in the Med Sea indicate that at least 10% and
27 up to 45% of the Atlantic RDOC entering the Med Sea is removed and replaced by
28 isotopically lighter and older DOC in less than 126 years. Large uncertainties are
29 associated to atmospheric, riverine, ground water and methane fluxes, limiting an
30 estimate of their relative importance. However, based on measured fluxes and isotopic
31 values, atmospheric inputs of soluble organic carbon from soils or combustion products
32 as well as DOC from groundwaters are the most likely sources of allochthonous DOC.
33 These data suggest that the true “refractory” fraction of DOC (the fraction that cycles

on millennial timescales) could be as low as ~ 20-25 μM . The Med Sea can be viewed as a reactor, where marine DOC imported from the Atlantic is replaced by terrestrial DOC, with the release of $0.5\text{-}0.6 \cdot 10^{+12} \text{ g CO}_2 \text{ y}^{-1}$ to the atmosphere. The outflow of deep Med Sea water through Gibraltar may be a source of terrestrial DOC to the ocean, with a flux of $8.6\text{-}11.5 \cdot 10^{+12} \text{ g C y}^{-1}$, estimated multiplying 19 μM by the outflow of Med waters (1.2-1.6 Sv) (Hopkins 1999; Bethoux and Gentili, 1999). By this account, the Med Sea would add about ~ 1-2% of total DOC as terrestrial carbon to the deep N. Atlantic. Although this amount would become rapidly undetectable in the Atlantic after mixing, it may be possible to detect this source at stations close to Gibraltar through measurements of $\text{DOC}\delta^{13}\text{C}$.

The mechanisms that lead to RDOC replacement are unknown. If replacement is common in the deep ocean (Follett et al., 2014), then replacement by isotopically lighter terrestrial carbon in the Med Sea and Arctic Ocean only makes the process in these basins more visible than in the global ocean. If however, unique processes or microbial communities in the deep Med Sea and Arctic Ocean convert RDOC into semi-labile DOC, then the mechanisms involved are of great interest.

Our results highlight the need for additional data on radiocarbon in both the West and East Med as well as additional information about radiocarbon in the external sources to the basin (atmosphere, rivers, groundwaters) and further studies on DOC cycling in this peculiar basin. The idea that the concentration of refractory DOC could be lower than 36-42 μM further strengthens the importance of DOC in the global carbon cycle, and elevates the possibility that DOC could represent a key element in maintaining dark ocean metabolism.

Acknowledgments

We thank Francesco Bignami and Rosalia Santoleri (CNR-ISAC) who provided us dissolved oxygen and CTD data and invited us to participate in their cruise. We further thank Luciano Nannicini, Stefano Vestri and Andrea Paluselli for their support in the sampling procedure during the cruises. Particular thanks are due to the captains and crews of the CNR RV 'Urania' and RV 'Minerva I', for their assistance and continuous support during the cruises. We also thank the staff of the National Ocean Sciences Accelerator Mass Spectrometry facility for their generous advice and assistance with isotope measurements.

This research was supported by PERSEUS project, funded by the European Commission under Seventh Framework Program Theme "Oceans of Tomorrow" OCEAN.2011-3, Grant Agreement No. 287600; the Italian Flagship project RITMARE funded by the Italian Ministry of Research and University. We acknowledge support from

the National Science Foundation (NSF) Science and Technology Center for Microbial Oceanography Research and Education (C-MORE) Grant CFF-424599, from the Gordon and Betty Moore Foundation through grant GBMF3298 and GBMF1711 to DJR, by the WHOI-Gori Fellowship program for SRB.

6. References

- Andrie, C., Merlivat, L., 1988. Tritium in the western Mediterranean Sea during 1981 Phycemed cruise. *Deep-Sea Res. I* 35, 247-267.
- Avery, G.B.Jr., Willey, J.D., Kieber, R.J., 2006. Carbon isotopic characterization of dissolved organic carbon in rainwater: Terrestrial and marine influences. *Atmospheric Environment* 40, 7539-7545.
- Basterretx, G., Tovar-Sanchez, A., Beck, A.J., Masque, P., Bokuniewicz, H.J., Coffey, R., Duarte, C.M., Garcia-Orellana, J., Garcia-Solsona, E., Martinez-Ribes, L., Vaquer-Sunyer, R., 2010. Submarine Groundwater Discharge to the Coastal Environment of a Mediterranean Island (Majorca, Spain): Ecosystem and Biogeochemical Significance. *Ecosystems* 13, 629-643 DOI: 10.1007/s10021-010-9334-5.
- Beaupré, S.R., Druffel, E.R.M., Griffin, S., 2007. A low-blank photochemical extraction system for concentration and isotopic analyses of marine dissolved organic carbon. *Limnology and Oceanography: Methods* 5(6), 174-184.
- Beaupré, S.R. and Aluwihare, L.I., 2010. Constraining the 2-component model of marine dissolved organic radiocarbon. *Deep Sea Res. II* 57, 1494-1503.
- Bernardoni, V., Calzolari, G., Chiari, M., Fedi, M., Lucarelli, F., Nava, S., Piazzalunga, A., Riccobono, F., Valli, G., Vecchi, R., 2013. Radiocarbon analysis on organic and elemental carbon in aerosol samples and source apportionment at an urban site in Northern Italy. *Journal of aerosol science* 56, 88-99.
- Bethoux, J.P., Gentili B., 1999. Functioning of the Mediterranean Sea: past and present changes related to fresh water input and climatic changes. *J. Mar. Sys.* 20, 33-47.
- Boetius, A., Wenzhöfer, F., 2013. Seafloor oxygen consumption fuelled by methane from cold seeps. *Nature Geoscience* 6, 725-734.
- Boldrin, A., Langone, L., Miserocchi, S., Turchetto, M., Acri, F., 2005. Po River plume on the Adriatic continental shelf: Dispersion and sedimentation of dissolved and suspended matter during different river discharge rates. *Marine Geology* 222–223, 135-158.

- 1 Broecker, W.S., Gerard, R., 1969. Natural Radiocarbon in the Mediterranean Sea.
2 Limnol. Oceanogr. 14, 883-8.
- 3 Carlson, C.A., 2002. Production and removal processes. In Hansell, D.A., Carlson,
4 C.A. (Eds.) Biogeochemistry of Dissolved Organic Matter, 1st Ed. Elsevier,
5 London, pp. 91-151.
- 6 Carlson, C.A., Hansell, D.A., Nelson, N.B., Siegel, D.A., Smethie, W.M., Khatiwala, S.,
7 Meyers, M.M., Wallner, E., 2010. Dissolved organic carbon export and
8 subsequent remineralization in the mesopelagic and bathypelagic realms of the
9 North Atlantic basin. Deep Sea Res. II 57, 1433-1445.
- 10 Cathalot, C., Rabouille, C., Tisnerat-Laborde, N., Toussaint, F., Kerherve, P., Buscail,
11 R., Loftis, K., Sund, M.-Y., Tronczynski, J., Azoury, S., Lansard, B, Treignier, C.,
12 Pastor, I., Tesi, T., 2013. The fate of river organic carbon in coastal areas: A
13 study in the Rhone River delta using multiple isotopic ($\delta^{13}\text{C}$, $\Delta^{14}\text{C}$) and organic
14 tracers. Geochimica et Cosmochimica Acta 118, 33–55.
- 15 Christensen, J.P., Packard, T.T., Dortch, F.Q., Minas, H.J., Gascard, J.C., Richez, C.,
16 Garfield, P.C., 1989. Carbon oxidation in the deep Mediterranean Sea: evidence
17 for dissolved organic carbon source. Global Biogeochem. Cycles 3, 315-335.
- 18 De Vicente, I., Ortega-Retuerta, E., Morales-Baquero, R., Reche, I., 2012. Contribution
19 of dust inputs to dissolved organic carbon and water transparency in
20 Mediterranean reservoirs. Biogeosciences 9, 5049-5060.
- 21 Druffel, E., Williams, P., Bauer, J., Ertel, J., 1992. Cycling of dissolved and particulate
22 organicmatter in the open ocean. J Geophys Res 97(C10), 15639-15659.
- 23 Druffel, E.R.M., Bauer, J.E., 2000. Radiocarbon distributions in Southern Ocean
24 dissolved and particulate organic matter. Geophysical Research Letters 27(2),
25 1495-1498.
- 26 Dusek, U., ten Brink, H.M., Meijer, H.A.J., Kos, G., Mrozek, D., Röckmann, T.,
27 Holzinger, R., Weijers, E.P., 2013. The contribution of fossil sources to the
28 organic aerosol in the Netherlands. Atmospheric Environment 74, 169-176.
- 29 Economou, C., Mihalopoulos, N., 2002. Formaldehyde in the rainwater in the eastern
30 Mediterranean: occurrence, deposition and contribution to organic carbon
31 budget. Atmos. Environ. 36, 1337-1347.
- 32 Flerus, R., Lechtenfeld, O.J., Koch, B.P., McCallister, S.L., Schmitt-Kopplin P., Benner,
33 R., Kaiser, K., Kattner, G., 2012. A molecular perspective on the ageing of
34 marine dissolved organic matter. Biogeosciences 9, 1935-1955.

- 1 Follett, C.L., Repeta, D.J., Rothman, D.H., Xu, L., Santinelli, C., 2014. Hidden cycle of
2 dissolved organic carbon in the deep ocean. *PNAS* 111(47), 16706-11.
- 3 García-García, A., Orange, D., Lorenson, T., Radakovitch, O., Tesi, T., Miserocchi, S.,
4 Berné, S., Friend, P.L., Nittrouer, C., Normand, A., 2006. Shallow gas off the
5 Rhône prodelta, Gulf of Lions. *Marine Geology* 234, 215-231.
- 6 García-García, A., Orange, D., Miserocchi, S., Correggiari, A., Langone, L., Lorenson,
7 T.D., Trincardi, F., Nittrouer, C.A., 2007. What controls the distribution of shallow
8 gas in the Western Adriatic Sea? *Continental Shelf Research* 27, 359-374.
- 9 Gramling, C.M., McCorkle, D.C. , Mulligan, A. E., Woods, T. L., 2003. A carbon isotope
10 method to quantify groundwater discharge at the land-sea interface. *Limnol.*
11 *Oceanogr.* 48(3), 957–970.
- 12 Griffith D.R., Barnes, R.T., Raymond, P.A., 2009. Inputs of Fossil Carbon from
13 Wastewater Treatment Plants to U.S. Rivers and Oceans. *Environmental*
14 *Science & Technoogy* 43(15), 5647–5651.
- 15 Griffith, D.R., McNichol, A.P., Xu, L., McLaughlin, F.A., Macdonald, R.W., Brown, K.A.,
16 Eglinton, T.I., 2012. Carbon dynamics in the western Arctic Ocean: insights from
17 full-depth carbon isotope profiles of DIC, DOC, and POC. *Biogeosciences* 9,
18 1217-1224.
- 19 Hansell, D.A., Carlson, C.A, 1998. Deep ocean gradients in the concentration of
20 dissolved organic carbon. *Nature* 395, 263-266.
- 21 Hansell, D.A., 2005. Dissolved organic carbon reference material program. *EOS Trans.*
22 *Amer. Geophys. Union* 86(35), 318.
- 23 Hansell, D.A., 2013. Recalcitrant dissolved organic carbon fractions. *Ann. Rev. Mar.*
24 *Sci.* 5, 3.1-3.25.
- 25 Hansell, D.A., Carlson, C.A., 2013. Localized Refractory Dissolved Organic Carbon
26 Sinks in the Deep Ocean. *Global Biogeochem. Cycles* 27, 705-710.
- 27 Harmelin-Vivien, M., Dierking, J., Bănar, D., Fontaine, M.F., Arlhac, D., 2010.
28 Seasonal variation in stable C and N isotope ratios of the Rhone River inputs to
29 the Mediterranean Sea (2004–2005). *Biogeochemistry* 100(1-3), 139-150.
- 30 Hopkins, T.S., 1999. The thermohaline forcing of the Gibraltar exchange. *J. Mar. Syst.*
31 20, 1-31.
- 32 Holmes R.M., McClelland, J.W., Raymond, P.A., Frazer, B.B., Peterson, B.J., Stieglitz,
33 M., 2008. Lability of DOC transported by Alaskan rivers to the Arctic Ocean.
34 *Geophysical Research Letters* 35, L03402, doi:10.1029/2007GL032837.

- Krom, M.D., Woodward, E.M.S., Herut, B., Kress, N., Carbo, P., Mantoura, R.F.C., Spyrese, G., Thingstad, T.F., Wassmann, P., Wexels-Riserf, C., Kitidis, V., Law, C.S., Zodiatis, G., 2005. Nutrient cycling in the south east Levantine basin of the eastern Mediterranean: Results from a phosphorus starved system. *Deep Sea Res. II* 52, 2879-2896.
- Kvenvolden, K.A., Lorenson, T.D., Reeburgh, W., 2001. Attention turns to naturally occurring methane seepage. *EOS* 82, 457.
- Lazzari, P., Solidoro, C., Ibello, V., Salon, S., Teruzzi, A., Béranger, K., Colella, S., Crise, A., 2012. Seasonal and inter-annual variability of plankton chlorophyll and primary production in the Mediterranean Sea: a modelling approach. *Biogeosciences* 9, 217–233.
- Lønborg C., Davidson, K., Álvarez-Salgado, X.A., Miller, A.E.J., 2009. Bioavailability and bacterial degradation rates of dissolved organic matter in a temperate coastal area during an annual cycle. *Marine Chemistry* 113, 219–226.
- López-Jurado, J.L., González-Pola, C., Vélez-Belchí, P., 2005. Observation of an abrupt disruption of the long-term warming trend at the Balearic Sea, western Mediterranean Sea, in summer 2005. *Geophysical Research Letters* 32, L24606.
- Ludwig, W., Dumont, E., Meybeck, M., Heussner, S., 2009. River discharges of water and nutrients to the Mediterranean and Black Sea: Major drivers for ecosystem changes during past and future decades? *Prog. Oceanogr.* 80, 199-217.
- Malanotte-Rizzoli, P., Manca, B.B., Ribera D'alcalà, M., Theocharis, A., Brenner, S., Budillon, G., Ozsoy, E., 1999. The Eastern Mediterranean in the 80s and in the 90s: the big transition in the intermediate and deep circulations. *Dynam. Atmos. Ocean* 29, 365-395.
- Markaki, Z., Loýe-Pilot, M.D., Violaki, K., Benyahya, L., Mihalopoulos, N., 2010. Variability of atmospheric deposition of dissolved nitrogen and phosphorus in the Mediterranean and possible link to the anomalous seawater N/P ratio. *Mar. Chem.* 120, 187-194.
- McNichol, A.P., Osborne, E.A., Gagnon, A.R., Fry, B., Jones, G.A., 1994. TIC, TOC, DIC, DOC, PIC, POC -Unique aspects in the preparation of oceanographic samples for ¹⁴C-AMS: *Nuclear Instruments and Methods in Physics Research B* 92, 162-165.

- McNichol, A.P., Aluwihare, L.I., 2007. The power of radiocarbon in biogeochemical studies of the marine carbon cycle: Insights from studies of dissolved and particulate organic carbon (DOC and POC). *Chemical Reviews* 107, 4423-466.
- Milkov, A.V., Sassen, R., Apanasovich, T.V., Dadashev, F.G., 2003. Global gas flux from mud volcanoes: A significant source of fossil methane in the atmosphere and the ocean. *Geophysical Research Letters* 30(2), 1037.
- Moore, W.S., 2010. The Effect of Submarine Groundwater Discharge on the Ocean. *Annual review of marine science* 2, 59-88.
- Ogawa, H., Amagau, Y., Koike, I., Kaiser, K., Benner, R., 2001. Production of refractory dissolved organic matter by bacteria. *Science* 292, 917-920.
- Panagiotopoulos, C., Sempéré, R., Para, J., Raimbault, P., Rabouille, C., Charrière, B., 2012. The composition and flux of particulate and dissolved carbohydrates from the Rhone River into the Mediterranean Sea. *Biogeosciences* 9, 1827-1844.
- Pettine, M., Patrolecco, L., Manganelli, M., Capri, S., Farrace, M.G., 1999. Seasonal variations of dissolved organic matter in the northern Adriatic Sea. *Mar. Chem.* 64, 153-169.
- Pohlman, J.W., Bauer, J.E., Waite, W.F., Osburn, C.L., Chapman, N.R., 2011. Methane hydrate-bearing seeps as a source of aged dissolved organic carbon to the oceans. *Nature Geoscience* 4, 37-41.
- Pujo-Pay, M., Conan, P., Oriol, L., Cornet-Barthaux, V., Falco, C., Ghiglione, J.-F., Goyet, C., Moutin, T., Prieur, L., 2011. Integrated survey of elemental stoichiometry (C, N, P) from the Western to Eastern Mediterranean Sea. *Biogeosciences* 8, 883-899.
- Pulido-Villena, E., Wagener, T., Guieu, C., 2008. Bacterial response to dust pulses in the western Mediterranean: Implications for carbon cycling in the oligotrophic ocean. *Global Biogeochem. Cycles* 22, GB1020.
- Raymond, P.A., Bauer, J.E., 2001. Use of ^{14}C and ^{13}C natural abundances for evaluating riverine, estuarine, and coastal DOC and POC sources and cycling: A review and synthesis. *Org Geochem* 32(4), 469-485.
- Retelletti Brogi, S., Gonnelli, M., Vestri, S., Santinelli, C., 2015. Biophysical processes affecting DOM dynamics at the Arno River mouth (Tyrrhenian Sea). *Biophysical Chemistry* 197, 1-9.

- Ribera d'Alcalà, M., Civitarese, G., Conversano, F., Lavezza, R., 2003. Nutrient ratios and fluxes hint at overlooked processes in the Mediterranean Sea. *J. Geophys. Res.* 108(C9), 8106.
- Rodellas, V., Garcia-Orellana, J., Masqué, P., Feldman, M., Weinstein, Y., 2015. Submarine groundwater discharge as a major source of nutrients to the Mediterranean Sea *PNAS*, 3926–3930.
- Roether, W., Manca, B.B., Klein, B., Bregant, D., Georgopoulos, D., Beitzel, V., Kovacevic, D.V., Luchetta, A., 1996. Recent changes in eastern Mediterranean deep waters. *Science* 271, 333-335.
- Römer, M., Sahling, H., Pape, T., dos Santos Ferreira, C., Wenzhöfer, F., Boetius, A., Bohrmann, G., 2014. Methane fluxes and carbonate deposits at a cold seep area of the Central Nile Deep Sea Fan, Eastern Mediterranean Sea. *Marine Geology* 347, 27-42.
- Santinelli, C., Nannicini, L., Seritti, A., 2010. DOC dynamics in the meso and bathypelagic layers of the Mediterranean Sea. *Deep Sea Res. II* 57, 1446-1459.
- Santinelli, C., Ibello, V., Lavezza, R., Civitarese, G., Seritti, A., 2012. New insights into C, N and P stoichiometry in the Mediterranean Sea: The Adriatic Sea case. *Cont. Shelf Res.* 44, 83-93.
- Santinelli, C., Hansell, D.A., Ribera d'Alcalà, M., 2013. Influence of stratification on marine dissolved organic carbon (DOC) dynamics: the Mediterranean Sea case. *Prog. Oceanogr.* 119, 68-77.
- Santinelli, C., 2015. DOC in the Mediterranean Sea. In: Hansell, D.A., Carlson C.A. (Eds.) *Biogeochemistry of Marine Dissolved Organic Matter*, 2nd Ed. Elsevier, pp. 579-608. Burlington: Academic Press.
- Schlitzer, R., Roether, W., Oster, H., Junghansh, H., Hausmann, M., Johannsen, H., Michelato, A., 1991. Chlorofluoromethane and oxygen in the Eastern Mediterranean. *Deep Sea Res.* 38, 1531-1551.
- Schroeder, K., Gasparini, G.P., Tangherlini, M., Astraldi, M., 2006. Deep and intermediate water in the western Mediterranean under the influence of the Eastern Mediterranean Transient. *Geophysical Research Letters* 33, L21607.
- Sempéré, R., Charrière, B., Van Wambeke, F., Cauwet, G., 2000. Carbon inputs of the Rhône River to the Mediterranean Sea: Biogeochemical implications. *Global Biogeochem. Cycles* 14, 669-681.

- 1 Sharp, J.H., Carlson, C.A., Peltzer, E.T., Castle-Ward, D.M., Savidge, K.B., Rinker,
2 K.R., 2002. Final dissolved organic carbon broad community intercalibration and
3 preliminary use of DOC reference materials. *Mar. Chem.* 77, 239-253.
- 4 Speicher, E.A., Moran, S.B., Burd, A.B., Delfanti, R., Kaberi, H., Kelly, R.P., Papucci,
5 C., Smith, J.N., Stavrakakis, S., Torricelli, L., Zervakis, V., 2006. Particulate
6 organic carbon export fluxes and size-fractionated POC/²³⁴Th ratios in the
7 Ligurian, Tyrrhenian and Aegean Seas. *Deep Sea Res. I* 53(11), 1810-1830.
- 8 Summons, R.E., Jahnke, L.L., Roksandic, Z., 1994. Carbon isotopic fractionation in
9 lipids from methanotrophic bacteria: Relevance for interpretation of the
10 geochemical record of biomarkers. *Geochimica et Cosmochimica Acta* 58(13),
11 2853-2863.
- 12 Whiticar, M.J., Faber, E., Schoell, M., 1986. Biogenic methane formation in marine and
13 freshwater environments: CO₂ reduction vs. acetate fermentation-isotope
14 evidence. *Geochimica et Cosmochimica Acta* 50(5), 693-709.
- 15 Williams, P.M., Druffel, E.R.M., 1987. Radiocarbon in dissolved organic matter in the
16 central North Pacific Ocean. *Nature* 330, 246-248.
- 17
18
19

Figure captions

Figure 1: Study area and sampling stations. The position of the stations reported in Broecker and Gerarld (1969) is also indicated (yellow squares).

Figure 2: Potential temperature/Salinity (θ/S) diagrams for the DOC sampling stations. The Levantine Intermediate Water (LIW, maximum of salinity) is clearly visible at the three stations. AW: Atlantic Water; WMDW: Western Mediterranean Deep Water; TDW: Tyrrhenian Deep Water. The black rectangles indicate the area that is enlarged and referred to in the subsequent plots for intermediate (b) and deep (c) water.

Figure 3: Vertical profiles of salinity, potential temperature and oxygen at the three stations.

Figure 4: Vertical profiles of DOC, with radiocarbon values and ages superimposed. DOC concentration refers to those measured in the bottle used for radiocarbon analysis. Cyan boxes: surface samples, Red boxes: LIW samples; Yellow boxes deep water samples (> 1000 m)

Figure 5: Vertical profiles of DOC for the Sargasso sea (a) (data from Druffel et al., 1992) and the average DOC profile for the West Med (b, c) (this study), with the hypothetical distribution of different radiocarbon fractions superimposed. In b we assume that the $7 \mu\text{M}$ of aged DOC at -129‰ is refractory; in c we assume this DOC is SLDOC from chemosynthesis, so it can be easily replaced.

Table 1: Site information for the three stations and physical chemical characteristics at the sampling depths

Station	Latitude °N	Longitude °E	Date	Bottom Depth m	Area	Sampling Depth	Pot Temp °C	Salinity	σ/θ	DO μM	AOU μM
A	38.50	13.50	Apr-7-2012	3440	Tyrrhenian Sea	2	16.79	37.90	27.79	244.7	-
						1500	13.20	38.55	29.11	186.5	71.2
						2500	13.04	38.51	29.10	188.0	70.6
						3429	12.99	38.49	29.10	188.5	70.4
B	38.33	8.57	Apr-1-2012	1894	Sardinian Channel	2	16.57	37.02	27.17	248.7	-
						480 (LIW)	13.84	38.70	29.08	177.9	76.2
						1851	12.89	38.47	29.11	197.2	62.2
C	40.16	7.36	Apr-4-2012	2702	Sardinian Sea	2	16.01	37.84	27.93	249.7	-
						1500	12.89	38.46	29.10	193.7	65.8
						2680	12.89	38.47	29.11	196.7	62.8

Table 2: DOC concentrations and isotopic values at the three stations.

Station	Depth (m)	DOC in bottle (μM) ^a	DOC NOSAMS (μM) ^b	DOC (μM) ^c	F Modern	Fm Err	$\delta^{13}\text{C}$ ‰	$\Delta^{14}\text{C}$ ‰	Age years	Age Err years
A	1	61.6	59.2	58.3	0.768	0.002	-22.7	-237	2120	20
	1500	41.1	45.6	40.1	0.559	0.002	-23.0	-445	4670	35
	2500	41.5	39.2	39.3	0.550	0.002	-24.1	-454	4810	30
	3429	-	-	41.8	0.538	0.002	-23.7	-464	4938	38
B	1	60.0	60.0	60.2	0.726	0.004	-21.9	-279	2570	40
	480 (LIW)	39.7	42.5	41.7	0.579	0.002	-22.8	-425	4390	25
	1851	41.6	46.0	40.3	0.529	0.001	-24.2	-475	5120	20
C	1	65.1	67.0	59.1	0.719	0.003	-22.0	-286	2640	35
	1500	38.7	42.0	41.3	0.534	0.002	-23.4	-470	5040	30
	2680	44.8	44.8	41.4	0.548	0.001	-23.6	-450	4762	20

^a DOC concentration in the bottle used for carbon isotopes measurements

^b DOC concentration estimated by the CO_2 produced during sample processing for carbon isotopes measurements.

^c DOC concentration in the replicate collected in amber glass bottle (see the Material and Methods section for further details)

Table 3: DIC $\Delta^{14}\text{C}$ values in the Med Sea compared to values reported in the literature. The position of the stations is reported in Fig. 1.

Station	Latitude °N	Longitude °E	Date	Area	Depth m	DIC Conc (mmol/kg)	$\delta^{13}\text{C}$ ‰	$\Delta^{14}\text{C}$ ‰	Age years	Age Err years	reference
A	40.07	12.70	Apr 2013	Tyrrhenian Sea	2	2.19	0.98	31.3	>Modern	-	This study
					400	2.32	0.75	18.6	>Modern	-	
					2500	2.31	0.99	-34.6	220	20	
					3390	2.31	0.99	-22.6	120	20	
367N	40.31	12.73	Aug 1956	Tyrrhenian Sea	3390	n.a.	0.70	-83±6	n.a.	n.a.	Broecker & Gerard, 1969
492X	39.2	6.17	Jul 1958	West Med	1	n.a.	n.a.	-48±10	n.a.	n.a.	
492W	39.12	7.60	Jul 1958	West Med	320	n.a.	n.a.	-60±10	n.a.	n.a.	
492U	38.73	7.00	Jul 1958	West Med	2700	n.a.	n.a.	-69±7	n.a.	n.a.	

n.a.: not available

Table 4: Literature stable and radiocarbon isotopes values for organic carbon measured in the possible allochthonous sources to the Med Sea

Source		DOC [μM]	DOC Flux (10^{12} gCy^{-1}) ^a	DOC $\Delta^{14}\text{C}$	DOC $\delta^{13}\text{C}$	References
Atlantic	Western North Atlantic, 1987 (0-100 m)	62	30-39	-238	-21.2	Druffel et al 1992
Atmosphere	TC in Aereosol in Milan urban area		0.4-13	-354 to -522		Bernardoni et al., 2013
	Organic carbon in PM10 in Amsterdam urban area			-325		Dusek et al., 2013
	Organic carbon in PM10 in a coastal location of Petten.			-177		
	Organic carbon in PM1-PM10 in various locations in Europe.			-118 to -453		Dusek et al., 2013 (Table 2 and literature therein)
	Rainwater North Carolina	132-540		49 to -102	-21.8 to -28.2	Avery et al. 2006
Rivers	Rhone River 2004-2005	8 to 62.5 (POC)	0.6-0.7		-25.46 to -28.55	Harmelin Vivien et al., 2010
	Po River 2000-2001	16-171 (POC)			-21.3 to 28.9	Boldrin et al., 2005
	Amazon (Brazil), Hudson (New York, USA), York (Virginia, USA) and Parker (Massachusetts, USA) rivers			364 to -158	-	Raymond & Bauer, 2001
Submarine groundwater	Island of Majorca (Spain)	80-169	0.3-9.7			Basterretxea et al., 2010
Methane	Adriatic Sea		0.003-10	-	-70 to -92‰	García-García et al., 2007
	Gulf of Lions			-	-53 to -82‰;	García-García et al., 2006

^a Fluxes estimated in this work

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5