1 Solid phase extraction method for the study of black carbon cycling in dissolved organic carbon
2 using radiocarbon
3

4 Alysha I. Coppola, Brett D. Walker, and Ellen R.M. Druffel

5 Department of Earth System Science, 2212 Croul Hall, University of California Irvine, Irvine,
6 CA 92617

7 corresponding author: acoppola@uci.edu
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

Abstract:

Radiocarbon analysis is a powerful tool for understanding the cycling of individual components within carbon pools, such as black carbon (BC) in dissolved organic carbon (DOC). Radiocarbon ($\Delta^{14}\text{C}$) measurements of BC provide insight into one source of aged, recalcitrant DOC. We report a modified solid phase extraction (SPE) method to concentrate $43\pm6\%$ of DOC (SPE-DOC) from seawater. We used the Benzene Polycarboxylic Acid (BPCA) method to isolate BC from SPE-DOC for subsequent ^{14}C analysis. We report SPE-BC $\Delta^{14}\text{C}$ values, SPE-BC concentrations, and the relative BPCA distributions from Milli-Q water process blanks, two riverine reference standards, as well as a coastal and an open ocean surface water sample. The composition of BC is less aromatic in the ocean samples than those in the river standards. We find higher BC $\Delta^{14}\text{C}$ values in the river standards (+148 to -462‰) than BC in the ocean samples (-592 to -712‰), suggesting that BC ages within oceanic DOC. We report that BC is $4.2 \pm 1.0\%$ of SPE-DOC in the open ocean surface sample, or $1.4\pm0.1 \mu\text{M C}$. This work provides the methodological basis by which global BC concentrations, compositions (eg. relative abundances of BPCA marker compounds) and $\Delta^{14}\text{C}$ values can be assessed.

Keywords: solid phase extraction, dissolved organic carbon (DOC), black carbon (BC), radiocarbon (^{14}C), BPCA, recalcitrant DOC (RDOC), Station M, radiocarbon analysis, preparative capillary gas chromatography, Suwannee River

1. Introduction

Black carbon (BC) is formed from incomplete combustion of fossil fuels and biomass. BC plays a strong role in the climate system, as it is considered second to CO₂ as the most important human emission based on its radiative forcing (Bond et al., 2013; Husain et al., 2007; Novakov and Rosen, 2013). BC also is hypothesized to be a long term carbon sink (Masiello and Druffel, 1998; Dai et al., 2005; Kuhlbusch and Crutzen, 1995), because its structure is composed of condensed aromatic rings making it stable and resistant to biological degradation (Goldberg, 1985; Forbes et al., 2006). After a fire, large amounts of charcoal in soils are oxidized and transported to river sheds (Myers-Pigg et al., 2015; Kim et al., 2004; Hockaday et al., 2007; Mannino and Harvey, 2004; Preston and Schmidt, 2006). BC is transported to the ocean by rivers, and is ubiquitous in the water column and sediments (Jaffe et al., 2013; Dittmar and Paeng, 2009; Stubbins et al., 2012a; Ziolkowski and Druffel, 2010; Suman et al., 1997; Masiello and Druffel, 1998; Coppola et al., 2014).

Once BC enters the ocean, it contributes to one of Earth's major organic carbon reservoirs, dissolved organic carbon (DOC, passes a 0.2–1.0 µm filter). While most DOC is believed to be produced by phytoplankton in the surface ocean, marine DOC is surprisingly thousands of ¹⁴C years old (Williams and Druffel, 1987). The presence of BC in the marine DOC pool may explain this mystery (Ziolkowski and Druffel, 2010; Masiello and Druffel, 1998) and was the motivation of this work. The structural composition and Δ¹⁴C values of SPE-BC within DOC provides insight into the composition of recalcitrant DOC, which escapes biological decomposition for thousands of years (Hansell et al., 2012; Ziolkowski and Druffel, 2010; Jiao et al. 2010; Stubbins et al., 2012). Two studies have identified BC in solid phase extracted DOC (SPE-DOC) with concentrations ranging from 0.6 to 0.8 µM (1 to 2% of total DOC) in the

Southern Ocean (Dittmar and Paeng, 2009) and $1.0 \pm 0.2 \mu\text{M}$ BC in surface North Atlantic Deep Water at the Bermuda Atlantic Time Series site (Stubbins et al., 2012a).

Ziolkowski and Druffel (2010) made the first BC $\Delta^{14}\text{C}$ measurements in ultrafiltered DOC (UDOC). They found open ocean BC had a range of ages from 15,680 to 20,100 ^{14}C yrs, providing evidence for BC stability on $>10^4$ year time scales. BC $\Delta^{14}\text{C}$ measurements in UDOC have limitations for evaluating the total DOC pool, because UDOC is only ~25% of DOC and selectively collects high molecular weight compounds (>1000 Da) (Amon and Benner, 1996; Walker et al., 2011; Benner, 2002; Aluwihare et al., 2002; Repeta et al., 2002; Benner et al., 1997). Studies of chemical composition and ^{14}C age of marine organic carbon show that the age of organic carbon increases with smaller sized molecules (Benner and Amon, 2015; Santschi et al., 1995; Walker et al., 2011). Thus, the presumably oldest, low molecular weight BC is not captured in UDOC.

We measured $\Delta^{14}\text{C}$ of BC in more of the DOC pool using SPE, which does not have an inherent size bias. The SPE method concentrates DOC based on chemical composition retaining polar and less-polar DOC compounds. This BC extraction method is an improvement over UDOC, because SPE recovers approximately half of the total DOC pool based on the sample type (Louchouart et al., 2010; Dittmar, 2008), without the time required for laborious UDOC isolation.

Our primary motivation is to use the SPE method to measure BC in DOC by determining the concentration and ^{14}C age of BC in ocean water. We use a styrene divinyl benzene copolymer resin to concentrate mostly hydrophilic DOC. We measure the mass and $\Delta^{14}\text{C}$ value of SPE-DOC and SPE-BC from a variety of standards and seawater. We assess blanks, reproducibility and the robustness of this SPE-DOC method. Comparisons of SPE-DOC to total

UV-oxidized DOC are used to evaluate whether SPE-DOC $\Delta^{14}\text{C}$ values are representative of total DOC $\Delta^{14}\text{C}$.

2. Materials and Methods:

2.1. Sample Collection

Coastal seawater DOC samples were collected from Newport Beach Pier (NBP) (33°60.70'N, 117°92.89'W) on August 8, 2013 at 1 m depth in 1-gallon bottles. Open ocean seawater DOC samples were collected from 20 m depth at Station M located on the Pacific abyssal plain (34°50'N, 123°00'W) on the *R/V New Horizon* in November 2004 using Niskin bottles. NBP samples were filtered through pre-baked 1.0 μm Whatman GFF filters and Station M DOC samples were filtered using pre-baked Whatman Polycap AS filter capsules (0.2 μm). All glassware in this study was soaked in 10% HCl, rinsed with deionized water and combusted at 550°C for 2 hrs. All seawater samples were frozen (at -20°C) until analysis. Seawater samples were separately analyzed for total DOC $\Delta^{14}\text{C}$ and [DOC] values using UV-oxidation techniques (Beaupré et al., 2007).

Seawater samples collected from Station M are from a long-term abyssal study site (4100m), located ~220 km west of Point Conception, CA in the NE Pacific (Smith and Druffel, 1998). The California current flows southward at Station M and has well-developed spring blooms. NBP is a coastal site, impacted by urban river discharge from the Santa Ana River (5km north of NBP) during discharge events in southern California. There were no river discharge events RECORDED BY THE GAUGE ON THE SANTA ANA RIVER IN THE CITY OF SANTA ANA ON AUGUST 8, 2013, NOR A MONTH PRIOR THAT MAY HAVE ADDED LARGE AMOUNTS OF RUNOFF DURING THE SAMPLE COLLECTION (AVAILABLE ON THE USGS SITE, SUPPLEMENTARY FIGURE 1).

2.2. Suwannee River Natural Organic Matter Standards

We used two freshwater, organic matter reference samples (IHSS 1R101N termed SR NOM I and IHSS 2R101N termed SR NOM II) purchased from the International Humics Substances Society (<https://ihss.humicsubstances.org/>) that were collected in May 1999 and May 2012, respectively. The Suwannee River drains the Okefenokee Swamp in southeastern Georgia, located at 30°48'14"N, 82°25'03"W, and has high DOC concentrations (82.7 mg C/L) with low concentrations of inorganic solutes (Serkiz and Perdue, 1990; Green et al., 2015). To create these standards, large volumes of water (~36,000L) from the Suwannee River were concentrated by reverse osmosis, desalted by cation exchange, freeze-dried and homogenized. When SR NOM I was sampled, two dams (built in 1950-1960) in the Suwannee River sill retained a high water level in the swamp and the site was relatively pristine and automobile vehicle access was prohibited (Green et al., 2015). The average flow rate of the river was 1.96 m³/s. In 2012, one of the two dams had been removed and it was no longer pristine (Green et al., 2015). The water level during 2012 was lower and the flow rate was much lower (0.46 m³/s). Both SR NOM I and SR NOM II were dissolved in Milli-Q water at DOC concentrations of 75–85 µM for SPE-BC analysis. These natural organic matter standards were also processed without SPE extraction, using only the BPCA method, and are called “total BC.”

2.3. Solid Phase Extraction of DOC (SPE-DOC)

We used a styrene divinyl benzene copolymer sorbent (Sigma Aldrich Diaion 13605, HP-20, pore size 200 Å), first used by De Jesus and Aluwihare, (2008), for extraction of DOC from seawater. To minimize the resin carbon blank, extensive cleaning (~1 week) was performed

using a glass soxhlet extractor with a pyrex glass insert to hold the resin during washings of methanol, acetone, ethyl acetate and dichloromethane at 65°C, 56°C, 77°C and 40°C, respectively. Each soxlet solvent washing lasted 24 hours and was performed sequentially in order of decreasing solvent polarity. According to the HP-20 manufacturers guidelines, sample water was acidified to pH 2 with hydrochloric acid (Fluka Traceselect 84415-500 ml) to increase extraction efficiency of organic components that contained exchangeable protons (Louchouart et al., 2010).

Figure 1 summarizes the SPE and BC methods. Briefly, large-volume frozen water samples (10 – 15 L) were defrosted, homogenized by shaking, acidified and loaded onto the SPE column. Samples were siphoned through 15 ml of resin in a glass Kontes column using 6 mm diameter pyrex tubing with acid cleaned silicone tubing, at a slow loading rate of 16 bed volumes per hour (240 ml/hr for surface samples). This flow rate was monitored and maintained over the course of the sample loading. The sample was loaded onto the resin three times to maximize interaction of DOC with the resin bed and increase DOC recovery.

In preparation for DOC elution, two bed volumes (30 ml) of Milli-Q water were passed through the column at 30 ml/hr to remove salts, and this fraction was discarded. SPE-DOC was eluted with solvents, each at a flow rate of 30 ml/hr using two bed volumes (30 ml). Solvents with different polarities were used to elute SPE-DOC into individual, glass vials in the following order: methanol, acetone, ethyl acetate and dichloromethane. These SPE-DOC fractions were dried under a stream of ultra-high purity (UHP) nitrogen gas. The SPE-DOC fractions were dissolved in a known volume of solvent, sub-sampled (5-10% of the volume), dried, then lyophilized for 24 hours. The SPE-DOC sub-samples were combusted separately to determine the percent yield of DOC (SPE-DOC/total DOC x 100) and SPE-DOC $\Delta^{14}\text{C}$ analyses.

2.4. Total BC and BC in Solid Phase Extracted DOC (SPE-BC)

The SPE-DOC methanol fractions were used to isolate BC using the Benzene Polycarboxylic Acid method, which oxidizes BC into marker compounds (BPCAs) for radiocarbon analysis (Ziolkowski et al., 2011; Schneider et al., 2011). The relative abundances of BPCAs produced during the oxidation step also provide qualitative BC structural information, because the more substituted BPCAs are derived from a more condensed aromatic BC network (Glaser et al. 1998; Ziolkowski et al., 2011; Coppola et al., 2013). The BPCA method characterizes only aromatic carbon as BC, and represents the most refractory aromatic constituents of pyrogenic carbon, but not the entire pool released to DOC (Masiello and Louchouart, 2013; Norwood et al., 2014; Myers-Pigg et al., 2015).

Briefly, eluted SPE-DOC extracts in glass vials were dried and lyophilized for 24 hrs. Total BC in SR NOM I and II was measured by adding the standard into quartz vials without SPE-processing. Concentrated nitric acid was added to vials (SPE-BC and total BC sample types) and placed in a pressure digestion chamber at 170°C for 8 hours to produce BPCAs (Coppola et al., 2013; Ziolkowski et al., 2011). The carbon in the carboxylic acid groups of the BPCA compounds is derived from adjacent aromatic groups of BC (B3CA, substituted with three carboxylic acids through B6CAs, those substituted with six carboxylic acids). After digestion, the solution was filtered, lyophilized and re-dissolved in methanol. Samples were derivatized using (trimethylsilyl) diazomethane in 2.0 M diethyl ether to convert carboxylic acid groups to methyl esters and an internal standard was added (500 µml of diphenic acid). BPCAs were collected on the preparative capillary gas chromatograph (PCGC), along with other BC standards (Hammes et al., 2007; Wiedemeier et al., 2015) using previously published techniques (Ziolkowski and Druffel, 2009; Coppola et al., 2013).

BC concentrations were determined from PCGC chromatographs. Peak areas of each BPCA relative to the internal standard were determined. Calibration curves (in mg C per unit mass) from commercially available BPCAs were applied for an estimate of BPCA mass. The BPCA method requires a conversion factor to convert the BPCAs formed from the nitric acid digestion into an estimate of the original BC mass. The conversion factor $24.2 \pm 1.6\%$ was used to convert BPCAs into BC concentrations (Ziolkowski and Druffel, 2010). This conversion factor was determined from oxidizing compounds with known chemical formulas (polycyclic aromatic hydrocarbons) to calibrate the BPCA method for carbon yield. Previously used conversion factors were based on yields from activated charcoal (with an unknown structure) (Glaser et al., 1998; Brodowski et al., 2005).

In preparation for ^{14}C analysis, B3CA through B6CA marker compounds (including nitrated B3CA and B4CAs) were collected in U-traps (-20°C) in the PCGC fraction collector. The B2CA marker compounds were not collected, because they are derived from lignins, a non-combusted material. The BPCAs in the U-trap were transferred using dichloromethane to quartz tubes, and dried under UHP nitrogen.

2.5. Radiocarbon Analyses

For $\Delta^{14}\text{C}$ analysis, samples were sealed in quartz tubes under vacuum with cupric oxide and silver wire and combusted to CO_2 at 850°C for 2 hrs. The CO_2 gas produced was cryogenically purified, and quantified manometrically, and used to calculate % DOC recovery. In some cases when sample sizes were too small for individual $\Delta^{14}\text{C}$ measurement (e.g., ethyl acetate and dichloromethane), fractions were combined and so-labeled. The resultant CO_2 was converted to graphite using the hydrogen reduction method on iron catalyst for ^{14}C analysis at

the Keck Carbon Cycle Accelerator Mass Spectrometry (AMS) Laboratory at U.C. Irvine (Santos et al., 2007). Radiocarbon measurements were reported as $\Delta^{14}\text{C}$ for geochemical samples without known age correction (Stuiver and Polach, 1977).

2.6. Carbon Blanks: Resin Blank and Process Blank Assessments

Three types of blanks were evaluated. First, resin blanks were processed to assess the carbon blank from each solvent that was passed through the resin before loading a sample. Resin blanks ranged from 2 to 6 $\mu\text{g C}$ each. Second, process blanks contained Milli-Q water (10 L each) and were subjected to the same processing steps as the samples. Process blanks were used to correct SPE-DOC samples for the $\Delta^{14}\text{C}$ value and mass of carbon blank added during processing, using a mass balance approach in Equation 1:

$$\Delta^{14}\text{C}_{\text{SPE-DOC}} = \frac{\Delta^{14}\text{C}_{\text{measured}}C_{\text{measured}} - \Delta^{14}\text{C}_{\text{Milli-Q}}C_{\text{Milli-Q}}}{C_{\text{measured}} - C_{\text{Milli-Q}}} \quad (\text{Equation 1})$$

where $\Delta^{14}\text{C}_{\text{SPE-DOC}}$ is the $\Delta^{14}\text{C}$ value corrected for SPE processing, $\Delta^{14}\text{C}_{\text{measured}}$ and C_{measured} are the measured $\Delta^{14}\text{C}$ value and mass of the sample, respectively, uncorrected for SPE processing, and $\Delta^{14}\text{C}_{\text{Milli-Q}}$ and $C_{\text{Milli-Q}}$ are the average $\Delta^{14}\text{C}$ value and mass of the Milli-Q water SPE-DOC, respectively (Table 1).

Third, for the assessment of blank carbon during the BPCA method, process blanks were generated using the Milli-Q water described above. The SPE-BC samples are corrected using an indirect approach as previously reported by Coppola et al., (2013) and Ziolkowski and Druffel (2009). Several BC standards from the carbon ring trial described by Hammes et al., (2007) were run alongside SPE-BC samples to correct for blank carbon. We found that approximately 6 – 10 μg dead blank C and 0.5 $\mu\text{g C}$ of modern carbon was added during BC processing of water samples. **3. Results**

3.1. Carbon Blank assessments

The mass of resin blanks from each of the four solvents ranged from 2 to 8 $\mu\text{g C}$, and their $\Delta^{14}\text{C}$ values ranged from $-499\pm 14\text{‰}$ to $-873\pm 2\text{‰}$ (Supplementary Table 1). The $\Delta^{14}\text{C}$ values of Milli-Q process blanks ranged from -660‰ to -1000‰ , and had masses that ranged from 2 – 10 $\mu\text{g C}$ (Table 1). Using an isotopic mass balance, we determined that Milli-Q water SPE-DOC $\Delta^{14}\text{C}$ values ranged from $-721\pm 77\text{‰}$ to $-927\pm 54\text{‰}$ (Supplementary Table 1). The Milli-Q water total DOC $\Delta^{14}\text{C}$ values obtained using UV-oxidation techniques ranged from -612 ± 57 to $-927\pm 54\text{‰}$, which is equal to the range in SPE-DOC Milli-Q $\Delta^{14}\text{C}$ values.

3.2. SPE-DOC recoveries and $\Delta^{14}\text{C}$ values

Percent SPE-DOC recoveries in all elutants for SR NOM I, SR NOM II, NBP and Station M samples were $62\pm 5\%$, $61\pm 4\%$, $43\pm 6\%$ and $44\pm 3\%$, respectively (Figure 2, Table 2). Percent SPE-DOC recoveries for ocean samples are comparable to those reported by Dittmar (2008) ($42\pm 8\%$) for surface marine DOC using a similar method (pre-made PPL cartridges with a smaller pore size of 150 $\text{\AA}$ and methanol elution).

For all samples, most of the DOC ($>89\%$) was eluted in the methanol fraction (Table 2, Figure 3). SR NOM I and SR NOM II standards had a very small amount of DOC ($\sim 1\%$) that eluted in the acetone fraction, and no DOC eluted in the ethyl acetate and dichloromethane fractions. For NBP, although the majority ($92\pm 4\%$) of coastal SPE-DOC was present in the methanol fraction, the less-polar (acetone, ethyl acetate and dichloromethane) fractions constitute 8% of the SPE-DOC. In the Station M sample, the methanol fraction was $89\pm 2\%$,

dichloromethane was $9\pm 2\%$ and there was no insignificant DOC in the acetone and ethyl acetate fractions ($1\pm 1\%$).

All SPE-DOC $\Delta^{14}\text{C}$ values were corrected using results from the process blanks (Equation 1). The methanol SPE-DOC $\Delta^{14}\text{C}$ values for SRNOMI are identical within 1 sigma ($+110\pm 7\%$, $+105\pm 10\%$, $+114\pm 8\%$) (Figure 2). The acetone SPE-DOC fractions in SRNOMI are $-286\pm 81\%$ and $-124\pm 68\%$, significantly lower than those of the methanol fractions. The $\Delta^{14}\text{C}$ value of total combusted SRNOMI is $+152\pm 3\%$ (Table 2), which is significantly higher than those of the methanol fractions, indicating that the SPE-DOC fraction is selectively isolating older DOC from the bulk material.

Duplicate methanol SPE-DOC $\Delta^{14}\text{C}$ values in SR NOM II are identical ($+20\pm 16\%$, $+20\pm 18\%$) (Table 2). The acetone fraction $\Delta^{14}\text{C}$ values ($-329\pm 57\%$ and $-480\pm 75\%$) are significantly lower than those of the methanol fractions. The total combusted SR NOM II $\Delta^{14}\text{C}$ value ($+45\pm 3\%$) is within 2 sigma of the SPE-DOC $\Delta^{14}\text{C}$ values measured in the methanol fractions (Table 2).

Duplicate methanol SPE-DOC $\Delta^{14}\text{C}$ values in the NBP samples are identical ($-263\pm 10\%$, $-255\pm 12\%$) (Figure 2). The acetone SPE-DOC $\Delta^{14}\text{C}$ values are $-717\pm 24\%$ and $-764\pm 10\%$, which were much lower than those of the methanol fractions. The $\Delta^{14}\text{C}$ values of the combined ethyl acetate/dichloromethane fractions ($-543\pm 41\%$ to $-691\pm 95\%$ Table 2) are intermediate to the other fractions. The NBP total DOC $\Delta^{14}\text{C}$ values measured by UV-oxidation are $-246\pm 3\%$, which is also equal to the methanol fraction $\Delta^{14}\text{C}$ values.

For Station M seawater, the methanol SPE-DOC $\Delta^{14}\text{C}$ values are $-318\pm 11\%$ and

-335±7‰ (Figure 2). The other fractions have lower $\Delta^{14}\text{C}$ values (acetone -678±25‰, ethyl acetate/dichloromethane -958±5‰) (Table 2). The total Station M DOC $\Delta^{14}\text{C}$ value is -299±3‰, which was significantly higher than one of the methanol SPE-DOC $\Delta^{14}\text{C}$ values.

3.3. BC concentrations, and relative BPCA distributions

The average BC/OC% values in SR NOM I, SR NOM II, NBP and Station M seawater samples are 8.2±1.0%, 11.9±2.0%, 20.5±2.0% and 4.2±1.0%, respectively (Table 3). The BC concentrations at in the NBP and Station M samples are 5.6±0.4 µM and 1.4±0.1 µM, respectively, in the SPE-DOC (Supplementary Table 2).

The relative BPCA abundances (i.e. B3CA, B4CA, B5CA, B6CA) in the methanol fractions of SR NOM I and II have low B3CAs and B4CAs abundances relative to those for B5CAs and B6CAs. However, there is a greater abundance of B6CAs in SR NOM II than those in SRNOMI. In contrast, the NBP samples have similar abundances of B3CAs, B4CAs and B5CAs and a low abundance of B6CAs (7±3%). We obtained BPCA abundances for the less polar SPE-BC fractions for the NBP samples only (Figure 4b), and the only significant difference between methanol with other less-polar fractions is the absence of B6CA marker compounds. The BPCA abundances in the Station M samples (Figure 4a) reveal a greater proportion of B3CAs than at NBP.

3.4 SPE-BC $\Delta^{14}\text{C}$ values

All SPE-BC $\Delta^{14}\text{C}$ values were corrected using an indirect approach (Table 3, Figure 4a). The methanol SPE-BC $\Delta^{14}\text{C}$ value of SR NOM I was +148±17‰. The total BC $\Delta^{14}\text{C}$ value for SR NOM I (without SPE) was significantly lower than that for the methanol SPE-BC (-49±33‰).

In contrast, duplicates of the methanol SPE-BC of SR NOM II ($-462 \pm 40\text{‰}$ and $-346 \pm 70\text{‰}$ within 2 sigma) were lower than that for the total BC ($-159 \pm 55\text{‰}$).

The methanol SPE-BC $\Delta^{14}\text{C}$ values in NBP seawater are $-712 \pm 36\text{‰}$ and $-678 \pm 19\text{‰}$ (within 2 sigma). The SPE-BC $\Delta^{14}\text{C}$ values of the acetone fractions are higher ($-587 \pm 47\text{‰}$ and $-615 \pm 20\text{‰}$). The NBP SPE-BC $\Delta^{14}\text{C}$ value of the ethyl acetate/dichloromethane mixture was equal to the acetone fractions of NBP SPE-BC fractions ($-593 \pm 15\text{‰}$) (Figure 4b).

Duplicates of the methanol SPE-BC $\Delta^{14}\text{C}$ values in the Station M samples are $-592 \pm 201\text{‰}$ and $-649 \pm 121\text{‰}$ (within 2 sigma), equal to the NBP methanol values (Figure 4a). No results were obtained for the acetone, dichloromethane or ethyl acetate fractions, due to their small masses.

4. Discussion

In the first section of the discussion, we assess the SPE-DOC method, by addressing the reproducibility of concentration and isotopic measurements in replicate standards, blanks and samples. In section 4.2, we discuss the variability of $\Delta^{14}\text{C}$ values obtained for SPE-DOC and total DOC analyses on the same samples. In section 4.3, we compare $\Delta^{14}\text{C}$ values obtained for SPE-BC and total BC for SR NOM I and SRNOMII samples. We also compare our results to those available for a deep UDOC BC sample in the Northeast Pacific and a surface sample in the North Central Pacific (NCP) (Ziolkowski and Druffel, 2010). In the final section, we discuss SPE-DOC and BC composition and cycling in these aqueous environments.

4.1 Methods assessment: Recoveries, SPE-DOC carbon blanks and reproducibility

Replicate measurements of SPE-DOC and SPE-BC recoveries and $\Delta^{14}\text{C}$ values agree within 2 sigma. This demonstrates that use of the method produces reproducible $\Delta^{14}\text{C}$ results

(Table 2, Table 3, Figure 2). Recoveries are higher for freshwater samples ($62\pm5\%$), likely due to the absence of salt in the sample. Recoveries were also consistent and reproducible for each sample type (Figure 2).

We report that masses of resin blanks acquired prior to sample loading, and Milli-Q process blanks are equal within error (Table 1, Supplementary Table 1), indicating that there is no additional carbon blank introduced to the large volume water samples. The SPE-DOC $\Delta^{14}\text{C}$ values of Milli-Q process blanks are equal within error to the $\Delta^{14}\text{C}$ values obtained for multiple 1-L UV oxidized Milli-Q blanks run from 2012-2014 in our laboratory (Supplementary Table 1). The SPE-Milli-Q process blank correction changed $\Delta^{14}\text{C}$ values by less than 10‰.

The SPE-BC process blanks are similar in size and $\Delta^{14}\text{C}$ value to that of resin and Milli-Q process blanks (Table 3). The SPE-BC process blanks are also comparable to those reported by Coppola et al., (2013). Together, these results suggest that no additional carbon blank is introduced by the SPE method and this method has low, consistent carbon blanks. Other studies quantify blanks by a direct approach (Dittmar, 2008; Flerus et al., 2012). Our blank assessment uses large sample volumes, standards and matrices to evaluate the $\Delta^{14}\text{C}$ value of blank carbon added indirectly during processing.

4.2 Comparison of $\Delta^{14}\text{C}$ values of SPE-DOC and total DOC

We present another comparison of $\Delta^{14}\text{C}$ values between SPE-DOC and (UV oxidized,) total DOC from splits of the same sample water (Table 2) (Druffel et al., 1992). Generally, we find that all methanol SPE-DOC $\Delta^{14}\text{C}$ values are equal to or lower ($\sim 25\pm 10\%$) than their corresponding total DOC $\Delta^{14}\text{C}$ values (Figure 2). Similarly, the $\Delta^{14}\text{C}$ values of SPE-DOC from SR NOM I and II standards are significantly lower ($\sim 35\pm 10\%$) than total DOC $\Delta^{14}\text{C}$ values

(Figure 2). The polarity-driven SPE method and its lower extraction efficiency for dissolved organic nitrogen compared to organic carbon may be responsible for the difference between total DOC and SPE-DOC $\Delta^{14}\text{C}$ values (Flerus et al., 2012). Another possible explanation for this $\Delta^{14}\text{C}$ offset could be selective concentration of younger humic materials that were precipitated out of solution at pH 2 during the collection of the standards (a step that we use in our SPE method). Our results suggest that SPE-DOC selectively isolates older DOC from the riverine standards, and to a lesser or no extent for seawater samples.

Using a mass balance between the SPE-DOC that is recovered ($43\pm 6\%$) and total DOC, we find the non-adsorbed DOC ($57\pm 5\%$) is more modern than their SPE-DOC counterpart by 67‰, 38‰, 9‰ and 17‰ for SR NOM I, SR NOM II, NBP and Station M, respectively (Supplementary Figure 2). For freshwater systems, the non-adsorbed DOC $\Delta^{14}\text{C}$ value is significantly different from the SPE-DOC $\Delta^{14}\text{C}$ value. For marine samples, the non-adsorbed DOC $\Delta^{14}\text{C}$ value is within error to the total DOC and SPE-DOC $\Delta^{14}\text{C}$ values (Supplementary Figure 2). While our sample set is limited, these results suggest that SPE-DOC $\Delta^{14}\text{C}$ results may be within error of the total DOC pool.

4.3 Comparison of SPE-BC, total BC and UDOC BC $\Delta^{14}\text{C}$ values and BPCA distributions

In contrast to SPE-DOC and bulk DOC, comparisons between SPE-BC and total BC of SR NOM I and SR NOM II standards show that BPCA distributions and $\Delta^{14}\text{C}$ values are dissimilar to one another (Table 3). This may indicate different sources, or differential cycling of BC within DOC. For SR NOM I and II, SPE-BC had a more aromatically condensed structure (greater abundance of B6CAs) than that for total BC (Table 3). Therefore, SR NOM I and SR NOM II BC that was not absorbed by SPE must have a greater proportion of less

aromatic BPCA marker compounds (B3CAs) to result in an overall lower B5CA and B6CA abundance in total BC.

The BPCA relative abundances revealed that SPE-BC isolates are more aromatically condensed than BC from the total BC pool. Additionally, the SPE-BC ^{14}C value of SR NOM I is higher than that of total BC, while that for SR NOM II is lower than that of the total BC. This may be due to variations in the environmental conditions present during sample collection of the two standards, or by differential cycling between SPE-BC and total BC. Ultrafiltration concentrates DOC based on size, whereas SPE chemically sorbs DOC based on polarity. We show that a more aromatically condensed BC structure (higher relative abundance of B6CAs) and higher $\Delta^{14}\text{C}$ values were found for the SPE-BC (Figure 4a) than for the UDOC (Ziolkowski and Druffel, 2010). This contrasted to expectation that the higher molecular weight (UDOC) fraction of BC might also be younger than the SPE-BC if BC, given the assumption that BC would follow the size reactivity relationships observed in DOC (see review by Benner and Amon, 2015). BC concentration has been found to correlate with DOC concentration (Myers-Pigg et al., 2015; Jaffe et al., 2013) in river waters. This suggests that with a larger DOC sample pool (43% SPE-DOC compared to 25% for UDOC), total BC may not be as aged as previously thought, or that there is discrete cycling of BC within these DOC fractions. Nonetheless, a larger sample set and comparisons of BC isolated by UDOM and SPE are needed to explain the relationship of the BC DOC fractions. BC concentration, structure and $\Delta^{14}\text{C}$ isolated by SPE and UDOC are not comparable.

4.3. SPE-DOC and BC cycling

Here, we investigate the different polarities of the SPE-DOC eluant fractions in the context of understanding DOC composition and cycling within these pools. By comparing the recovery of SPE-DOC in each of the polar and less-polar fractions, we found that the riverine SR NOM I and SR NOM II standards were exclusively polar with no significant differences in SPE-DOC composition between the two collection periods (Figure 3, Table 2). Methanol SPE-DOC fractions from both standards are composed of younger (perhaps humic, and/or carbohydrate-like) materials (Nwosu and Cook, 2014; Kuhn et al., 2015).

In contrast, NBP and Station M seawater samples have higher proportions of SPE-DOC eluting in the less polar fractions (Figure 3). Approximately 10% of oceanic SPE-DOC at Station M eluted with less-polar solvents. This agrees with previous studies that showed individual DOC fractions can have distinct compositions and low $\Delta^{14}\text{C}$ values in the ocean, such as total extractable lipids or acid insoluble fractions in DOC and particles (Loh et al., 2006; Wang et al., 2006; Hwang and Druffel, 2003).

Total DOC $\Delta^{14}\text{C}$ values and BC $\Delta^{14}\text{C}$ values in SR NOM I and SR NOM II are dissimilar. SR NOM I and II may represent different amounts of anthropogenic carbon. WE OBSERVE SIMILAR BC CONCENTRATIONS AND DIFFERENT BC $\Delta^{14}\text{C}$ VALUES IN THESE RIVER STANDARDS. SR NOM I BC HAS HIGH $\Delta^{14}\text{C}$ VALUES, PERHAPS INDICATIVE OF THE EXTENSIVE FIRES IN THE SWAMP DURING THE MID-1950s (GREEN ET AL., 2015). BOTH DOC AND BC $\Delta^{14}\text{C}$ VALUES WERE HIGHER IN 1999 (SR NOM I) THAN IN 2012 (SR NOM II) (FIGURE 2), SUGGESTING AN OLDER DOC AND BC SOURCE CONTRIBUTION IN SR NOM II. THE LOWER BC $\Delta^{14}\text{C}$ VALUE OF SR NOM II MAY SUGGEST ANTHROPOGENIC BC INPUTS, AS THE SUWANNEE RIVER SAMPLING SITE HAD NO LONGER BEEN CONSIDERED PRISTINE (GREEN ET AL., 2015) (FIGURE 4A). HYDROPHOBIC ACID

ISOLATES FROM REVERSE OSMOSIS OF SR NOM II COLLECTED IN 2012 HAVE A HIGHER AVERAGE MOLECULAR WEIGHT, AND ^{13}C NMR STUDIES REVEALED MORE AROMATIC FUNCTIONAL GROUPS THAN FOR SR NOM I (KRELLER ET AL., 2015; NWOSU AND COOK, 2014). This suggests that riverine BC may be influenced by aerosol anthropogenic input (Stubbins et al., 2012b; Spencer et al., 2014).

Our oceanic BC samples appear chemically distinct from the riverine standards we studied. Using the relative BPCA abundance as an indicator of BC aromaticity, we observed a smaller contribution of B3CAs in SPE-BC of SR NOM I and SR NOM II versus marine SPE-BC (Figure 4a). Our oceanic sample results also provide us with a glimpse of DOC-BC cycling at coastal and open ocean sites. Coastal NBP is the only sample with significant amounts of SPE-BC in all four solvent fractions ($\text{SPE-BC} = 5.6 \pm 0.4 \mu\text{M C}$). Given the high BC content, we believe this site is influenced by fossil fuels or local aged sources. Similarly, total suspended solids for a nearby river (~75 miles north, Santa Clara River, CA) had 44.3% BC/OC%, suggesting a large BC riverine flux from the Ventura Basin, with a low $\Delta^{14}\text{C}$ value of $-644 \pm 58\text{‰}$ (Masiello and Druffel, 2001). This may explain the high BC/OC% we observe in NBP samples ($20.5 \pm 2.0\%$). Seasonal hydrology plays a role in the quality of exported BC (Wagner et al., 2015). Based on this riverine study in the Colorado Rocky Mountains, we expect a riverine source of smaller BC structures (B3CAs) during low flow ($\ll 500 \text{ ft}^3/\text{s}$) periods. However, variations of BC structure during high and low flow river period and its impact on the ocean have yet to be investigated in urban environments.

Although the SPE-BC $\Delta^{14}\text{C}$ values at NBP and Station M are within error, their relative BPCA abundances are different, and may reveal reprocessing of the oceanic BC pool. We assume BC at Station M is representative of open ocean BC (with small contributions from

riverine anthropogenic sources) (Coppola et al., 2014; Masiello and Druffel, 1998). UV oxidation plays a strong role in degrading highly aromatic BC in the land-to-ocean transit and may remove these structures upon entering the open ocean (Stubbins et al., 2012; Masiello and Louchouart, 2013; Myers-Pigg et al., 2015). Coastal BC may be influenced by anthropogenic riverine BC more directly. The structure of BC at Station M has a greater relative abundance of B3CAs than that at NBP (Figure 4a) and a lower BC concentration. This may indicate degradation processes that remove aromatic BC with transport to the open ocean (e.g. photo-oxidation; Stubbins et al., 2012a).

Despite the limited number of BC $\Delta^{14}\text{C}$ studies, it is interesting to note the disparity between $\Delta^{14}\text{C}$ values as isolated by SPE and UDOC. At Station M, SPE-BC is about 12,000 ^{14}C yrs younger than BC isolated from UDOC ($20,100 \pm 3,000$ ^{14}C yrs; Ziolkowski and Druffel, 2010). This may suggest differential cycling of BC in chemical versus size fractionated DOC pools (Wagner and Jaffe, 2015). The discrepancy in BC $\Delta^{14}\text{C}$ values highlights the need for additional DOC BC measurements in order to constrain BC sources, cycling and the ^{14}C ages in the global ocean.

5. Summary and Implications

We report a modified, low-blank SPE method to isolate DOC from marine water samples. We report that SPE-DOC constitutes nearly half of the DOC pool ($43 \pm 6\%$). We present $\Delta^{14}\text{C}$ blank assessments of the SPE-DOC method from a limited set of solid river standards and surface ocean water samples. Our results suggest that SPE-DOC selectively isolates slightly older DOC from the riverine standards, and to a lesser or no extent for seawater samples. We

observe similar SPE-BC $\Delta^{14}\text{C}$ values in our coastal and open ocean sites, but different BPCA relative abundances.

Further measurements are needed to determine BC $\Delta^{14}\text{C}$ and structure (using relative BPCA distributions) to better understand the sources, sinks, transformations from rivers and cycling of BC in aquatic environments. One limitation of these analyses was the sample sizes of less-polar SPE-DOC fractions, which prevented analysis of relative BPCA abundances and $\Delta^{14}\text{C}$ values of less-polar BC at our open ocean site. The use of larger water samples would allow for less-polar BC to be evaluated and placed into the context of total and SPE-DOC ^{14}C ages. Chemical characterizations of SPE-DOC fractions at the molecular level (e.g nuclear magnetic resonance and Fourier transform ion cyclotron resonance mass spectrometry, FT-ICRMS) will reveal compositional relationships between SPE-DOC and SPE-BC.

481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499

500

501

502

Acknowledgements

We would like to thank Lihini Aluihware for her advice and assistance on modifications of their SPE methods; Steve Beaupré for collecting Station M DOC water samples, the technicians and crew of the *R/V New Horizon* (November 2004); Sheila Griffin for her technical support and assistance; Dachun Zhang for his help with the PCGC instrument and guidance; Xiaomei Xu for her help with ¹⁴C corrections; Danielle Glynn and Christopher Glynn for laboratory help; and John Southon and Keck Carbon Cycle AMS Lab at UC Irvine colleagues for support. We also thank Boris Koch for his advice and two anonymous reviews for their comments, which greatly improved the manuscript. We acknowledge support from NSF [OCE-0961980, OCE-1022716 and OCE-1458941]. We thank the Fred Kavli Foundation for funding.

503

504 References

- 505 Ahn, J.H, Grant, S.B., Surbeck, C.Q., Digiacomio, P.M., Nezlin, N.P., Jiang, S. 2005. Coastal
506 Water Quality Impact of Stormwater Runoff from an Urban
507 Watershed in Southern California. *Environ. Sci. Technol.*, 39,5940-5953.
- 508 Aluwihare, L.I., Repeta., D.J., Chen, R.F. 2002. Chemical composition and cycling of dissolved
509 organic matter in the Mid-Atlantic Bight. *Deep Sea Res. Part II: Trop. Stud. Oceanog.* 49,
510 4421–4437.
- 511 Amon, R.M.W, Benner, R. 1996. Bacterial utilization of different size classes of dissolved
512 organic matter. *Limnol. Oceanog.* 41, 41-51.
- 513 Beaupré, S.R., Griffin, S., Druffel, E.R.M. 2007. A low-blank photochemical extraction system
514 for concentration and isotopic analyses of marine dissolved organic carbon. *Limnol. and*
515 *Ocean. Methods* 5, 174–184.
- 516 Benner, R., Amon, R.M.W. 2015. The size-reactivity continuum of major bioelements in the
517 ocean. *Annu. Rev. Mar. Sci.* 7, 185-205.
- 518 Benner, R., Biddanda, B., Black, B., Mccarthy, M. 1997. Abundance, size distribution, and
519 stable carbon and nitrogen isotopic compositions of marine organic matter isolated by
520 tangential-flow ultrafiltration. *Mar. Chem* 57, 243-263.
- 521 Benner R. 2002. Chemical composition and reactivity In: D. Hansell and C. Carlson, Ed.,
522 *Biogeochemistry of Marine Dissolved Organic Matter*, Academic Press, 59–90.

523 Bond, T. C., S. J. Doherty, D. W. Fahey, P. M. Forster, T. Berntsen, B. J. DeAngelo, M. G.
 524 Flanner, et al., 2013. Bounding the Role of BC in the Climate System: A Scientific
 525 Assessment. *J. Geophys. Res. Atmos.*, 118, 1–173.

526 Coppola, A.I., Ziolkowski L.A., Druffel, E.R.M., 2013. Extraneous carbon assessments in
 527 radiocarbon measurements of BC in environmental matrices. *Radiocarbon*, 55,1631-1640.

528 Coppola, A.I., Ziolkowski, L.A., Masiello, C.A., Druffel, E.R.M. 2014. Aged black carbon in
 529 marine sediments and sinking particles. *Geophys. Res. Lett.* 41, 2147-2433.

530 Dai, X., Boutton, T.W., Glaser, B., Ansley, R.J., Zeck, W. 2005. BC in a temperate
 531 mixed-grass savanna. *Soil Bio. Biochem.* 10, 1879-1881.

532 De Jesus, R.P., Aluwihare, L.I. 2008. Natural abundance radiocarbon studies of dissolved organic
 533 carbon in the marine environment, Ph.D.thesis, Oceanogr., Univ. of Calif., San Diego.

534 Dittmar, T. 2008. The molecular level determination of BC in marine dissolved
 535 organic matter. *Org. Geochem.* 39, 396–407.

536 Dittmar, T., Koch, B., Hertkorn, N., Gerhard, K. 2008. A simple and efficient method for
 537 the solid-phase extraction of dissolved organic matter (SPE-DOM) from seawater.
 538 *Limnol. Oceanogr.: Methods* 6, 230-235.

539 Dittmar, T., Paeng, J. 2009. A heat-induced molecular signature in marine dissolved organic
 540 matter. *Nat. Geosci.* 2, 175-179.

541 Druffel, E.R.M., Williams, P., Bauer, J., Ertel, J.R. 1992. Cycling of Dissolved and Particulate
 542 Organic Matter in the Open Ocean. *Journ. Geophys. Res.*, 92:15, 639-659.

543 Dwight, R.H., Semenza, J.C., Baker, D.B., Olson, B.H. 2002. Association of urban runoff with
 544 coastal water quality in Orange County, California. *Water Envir. Res.*, 74(1), 82-90.

545 Glaser, B., L. Haumaier, G. Guggenberger, Zech W., 1998. BC in soils: the use of

546 benzenecarboxylic acids as specific markers. *Org. Geochem.*, 811-819.

547 Green, N.W., Perdue, E.M., Aiken, G.R., Butler, K.D., Chen, H., Dittmar, T., Niggemann, J.,

548 Stubbins, A. 2014. An intercomparison of three methods for the large-scale isolation of

549 oceanic dissolved organic matter. *Mar. Chem.*, 161, 14-19.

550 GREEN, N.S., MCINNIS, D., HERTKORN, N., MAURICE, P.A., PERDUE, E.M. 2015.

551 SUWANNEE RIVER

552 NATURAL ORGANIC MATTER: ISOLATION OF THE 2R101N REFERENCE SAMPLE BY

553 REVERSE OSMOSIS.

554 ENVIRON. ENG. SCI. 32, 38.

555 Hammes, K., et al. 2007. Comparison of quantification methods to measure fire-derived

556 (black/elemental) carbon in soils and sediments using reference materials from soil, water,

557 sediment and the atmosphere. *Glob. Biogeochem. Cyc* 21, GB002914

558 Husain, L., Dutkiewicz, V. A., Khan, A., and Ghauri, B. M., 2004. Characterization of

559 carbonaceous aerosols in urban air. *Atmos. Environ.*, 41, 6872–6883.

560 Hwang, J., Druffel, E.R.M. 2003. Lipid-like material as the source of the uncharacterized organic

561 carbon in the ocean? *Science* 299,881-884.

562 Jaffé, R., Ding, Y., Niggemann, J., Vähätalo, A.J., Stubbins, A., Spencer, R.G.M, Campbell, J.,

563 Dittmar, T. 2013. Global charcoal mobilization from soils via dissolution and riverine

564 transport to the oceans. *Science* 340, 345-347.

565 Kuhn, K.M., Neubauer, E., Hofmann, T., von der Kammer, F., Aiken, G.R., and Maurice, P.A.

566 (2015). Concentrations and distributions of metals associated with dissolved organic matter

567 from the Suwannee River (GA, USA). *Environ. Eng. Sci.* 32, 54-65.

568 Flerus, R., Lechtenfeld, O., Koch, B., McCallister, S., Schmitt-Kopplin, P., Benner, R., Kaiser,
 569 K., Kattner, G. 2012. A molecular perspective on the ageing of marine dissolved organic
 570 matter. *Biogeosci.* 9,6, 1935–1955.

571 Forbes, M., Raison, R., Skjemstad, J. O. 2006. Formation, transformation and transport of BC
 572 (charcoal) in terrestrial and aquatic ecosystems. *Sci. Total. Environ.*, 370, 190–
 573 206.

574 Goldberg, E. D.: BC in the Environment, Wiley, New York, New York, USA, 1985.

575 Jiao, N., Herndl, G.J., Hansell, D.A., Benner, R., Kattner, G., Wilhelm, S.W., Kirchman, D.L.,
 576 Weinbauer, M.G. 2010. Carbon Storage in the Global Ocean. *Nature* 8, 593–599.

577 Hockaday, W., Grannas, A., Kim, S., Hatcher, P. 2007. The transformation and mobility of
 578 charcoal in a fire impacted watershed. *Geochim. Cosmochim. Acta*, 71, 3432–3445,

579 Kim, S., Kaplan, L., Benner, R., Hatcher, P. 2004. Hydrogen deficient molecules in natural
 580 riverine water samples—evidence for the existence of BC in DOM. *Mar. Chem.*,
 581 92, 225–234, doi:10.1016/j.marchem. 2004.06.042.

582 Khosh, M. S., Xu, X., Trumbore, S.E. 2010. Small-mass graphite preparation by sealed tube zinc
 583 reduction method for AMS 14C measurements. *Nuc. Instrum. Methods Phys. Res. Sec. B*
 584 268, 7-8.

585 KRELLER, D.I., WU, Y., SUTTON, S., FURIO, A. 2015. ADSORPTION AND FRACTIONATION
 586 OF SUWANNEE
 587 RIVER ORGANIC MATTER ON METAL (Fe, Al) OXIDE COATED QUARTZ: A LIQUID
 588 CHROMATO-
 589 GRAPHIC INVESTIGATION. *ENVIRON. ENG. SCI.* 32, 45

590 Kuhlbusch, T., Crutzen, T. 1995. Toward a global estimate of black carbon in residues of

591 vegetation fires representing a sink of atmospheric CO₂ and a source of O₂. Global
592 Biogeochem. Cycles 9, 491-501.

593 Loh, A.N., Bauer, J.E., Druffel, E.R.M. 2004. Variable ageing and storage of dissolved organic
594 matter. Limnol. Oceanogr. 49, 735-740.

595 Louchouart. P., Amon, R.M.W., Duan, S., Pondell, C., Seward, S.M., White, N. 2010. Analysis
596 of lignin-derived phenols in standard reference materials and ocean dissolved organic
597 matter by gas chromatography/tandem mass spectrometry. Mar. Chem., 118, 85-97.

598 Mannino, A., Harvey, H. 2004. Black carbon in estuarine and coastal ocean dissolved organic
599 matter. Limnol. Oceanogr., 49, 735–740.

600 Masiello, C., Druffel E.R.M., 1998. Black carbon in deep-sea sediments. Science 280,
601 1911-1913.

602 Masiello, C.A., Druffel, E.R.M. 2001. Carbon isotope geochemistry of the Santa Clara River.
603 Global Biogeochem. Cyc., 15, 407-416.

604 Masiello, C., 2004. New directions in black carbon organic geochemistry. Mar. Chem., 92, 201–
605 213.

606 Masiello, C. A., and P. Louchouart (2013), Fire in the Ocean, Science, 340, 287–288,
607 doi:10.1126/science.1237688.

608 Middelburg, J., J. Nieuwenhuize, and P. Can-Breuge (1999), Black carbon in marine sediments,
609 Mar. Chem., 65, 245–252.

610 Myers-Pigg, A.N. et al., 2015. Labile pyrogenic dissolved organic carbon in major Siberian
611 Arctic rivers: Implications for wildfire-stream metabolic linkages. *Geophysical Research*
612 *Letters*, 42(2), 377-385.

613 Novakov, T., Rosen, H. 2013. The BC story: early history and new perspectives,
 614 *Ambio*, 42, 840–851.

615 NWOSU, U., COOK, R. 2014. A ¹³C NUCLEAR MAGNETIC RESONANCE AND ELECTRON
 616 PARAMAGNETIC
 617 SPECTROSCOPIC COMPARISON OF HYDROPHOBIC ACID, TRANSPHILIC ACID, AND
 618 REVERSE OSMOSIS
 619 MAY 2012 ISOLATES OF ORGANIC MATTER FROM THE SUWANNEE RIVER.
 620 *ENVIRON. ENG. SCI.*
 621 32, 14

622 Repeta, D.J., Quan, T.M., Aluwihare, L.I., Accardi, A. 2002. Chemical characterization of high
 623 molecular weight dissolved organic matter in fresh and marine waters. *Geochim. et.*
 624 *Cosmochim. Acta* 66, 955–962.

625 Preston, C.M., Schmidt M.W.I., 2006. Black (pyrogenic) carbon: a synthesis of current
 626 knowledge and uncertainties with special consideration of boreal regions. *Biogeosci.*
 627 3:397-420.

628 Santschi, P.H., Guo, L., Baskaran, M., Trumbore, S., Southon, J., Bianchi, T.S., Honeyman, B,
 629 Cifuentes, L. 1995. Isotopic evidence for the contemporary origin of high-molecular
 630 weight organic matter in oceanic environments. *Geochim. Cosmochim. Acta*, 59, 625–
 631 631.

632 Santos, G.M., Southon, J.R., Beaupre, S.R., Druffel, E.R.M. 2007. Ultra small-mass AMS ¹⁴C
 633 sample preparation and analyses at KCCAMS/UCI Facility. *Nucl. Instr. And Meth. B.*
 634 259, 293-302.

635 Schneider, M.P.W., Hilf, M., Vogt, U. F., Schmidt, M.W.I. 2010. The benzene polycarboxylic

636 acid (BPCA) pattern of wood pyrolyzed between 200°C and 1000°C. *Org. Geochem.* 41,
 637 1082-1088.

638 Schmidt, M. W., Noack, A. G. 2008. BC in soils and sediments: Analysis, distribution,
 639 implications, and current challenges. *Global Biogeochem. Cy.*, 14, 777–793.

640 Serkiz, S. M., Perdue, E. M. 1990. Isolation of dissolved organic matter from the Suwannee
 641 River using reverse osmosis. *Water Res.*, 24(7), 911–916.

642 Smith, K.L., Druffel, E.R.M. 1998. Long time-series monitoring of an abyssal site in the NE
 643 Pacific: an introduction. *Deep-Sea Res. Part II*, 45 (4-5):657-586.

644 Spencer, R. G. M., G. Weidong, P. Raymond, T. Dittmar, E. Hood, J. Fellman, Stubbins, A.
 645 2014. Source and biolability of ancient dissolved organic matter in glacier and lake
 646 ecosystems on the Tibetan Plateau. *Geochim. et Cosmochim. Acta.*, 142:64-74. doi:
 647 10.1016/j.gca.2014.08.006

648 Stubbins, A., Niggemann, J., Dittmar, T. 2012a. Photo-lability of deep ocean dissolved black
 649 Carbon. *Biogeo. Disc.*, 9, 485-505.

650 Stubbins, A., Hood, E., Raymond, P.A., Aiken, G.R., Sleighters, R.L., Hernes, P.J., Butman, D.,
 651 Hatcher, P.G., Striegl, R.G., Schuster, P. 2012b. Anthropogenic aerosols as a source of
 652 ancient dissolved organic matter in glaciers. *Nat. Geo.*, 5, 198-201.

653 Stuiver, M., and H.A. Polach, 1977. Reporting of ¹⁴C data. *Radiocarbon* 19, 355–363.

654 Wagner, S., Cawley, K.M., Rosario-Ortiz, F.L., Jaffe, R., 2015. In-stream sources and links
 655 between particulate and dissolved black carbon following a wildfire. *Biogeochem. DOI*
 656 10.1007/s10533-015-0088-1

657 Wagner, S., and R. Jaffe, 2015. Effect of photodegradation on molecular size distribution and
 658 quality of dissolved black carbon. *Org. Geochem* (85), 1-4.

659 Walker, B.D., Beaupre, S.R., Guilderson, T.P., Druffel, E.R.M., Mccarthy, M.D., 2011. Large-
 660 volume ultrafiltration for the study of radiocarbon signatures and size vs. Age
 661 relationships in marine dissolved organic matter. *Geochim. Cosmochim. Acta* 75, 5187-
 662 5202.

663 Wiedemeier, D. B, Abiven, S., Hockaday, W. C., Keiluweit, M., Kleber, M., Masiello, C. A.,
 664 McBeath, A. V., Nico, P. S., Pyle, L. A., Schneider, M. P. W., Smernik, R. J.,
 665 Wiesenberg, G. L. B. Schmidt, M. W. I. 2015. Aromaticity and degree of aromatic
 666 condensation of char. *Org. Geochem.*, 78:135-143.

667 Williams, P.M., Druffel, E.R.M., 1987. Radiocarbon in dissolved organic matter in the central
 668 north Pacific Ocean. *Nature* 330, 246-248.

669 Xu, X., Trumbore, S.E., Zheng, S., Southon, J.R., McDuffee, K.E., Luttgen, M., Liu, J.C. 2007.
 670 Modifying a sealed tube zinc reduction method for preparation of AMS graphite targets:
 671 Reducing background and attaining high precision. *Nuc. Instrum. Methods Phys. Res. Sect.*
 672 *B*, 259, 320–329.

673 Ziolkowski, L., Druffel E.R.M., 2009. Quantification of Extraneous Carbon during Compound
 674 Specific Radiocarbon Analysis of Black Carbon. *Anal. Chem* 81, 10158-10161.

675 Ziolkowski, L., Druffel E.R.M. 2010. Aged BC identified in marine dissolved organic carbon.
 676 *Geophys. Res. Lett.*, 37, L16601.

677 Ziolkowski, L., Chamberlin, A.R., Greaves, J., Druffel, E.R.M. 2011. Quantification of black
 678 carbon in marine systems using the benzene polycarboxylic acid method: a mechanistic
 679 and yield study. *Limnol. Oceanogr.: Methods*, 9, 140–149.

680

Table 1. Carbon blank mass and $\Delta^{14}\text{C}$ values of process blanks obtained from processing 10 L of Milli-Q water for each assessment. Averages are weighted averages. Milli-Q DOC $\Delta^{14}\text{C}$ values range from -926 ± 54 to $-612\pm57\text{‰}$ (n=60, S. Griffin and B. Walker).

SPE-DOC fraction	Sample Group	UCID lab code #	Blank size ($\mu\text{g C}$)	$\Delta^{14}\text{C}$ (‰) Blank corrected	$\pm$
Methanol	MQ I	17010	4	-891	23
	MQ III	17173	2	-877	76
	MQ IV	17185	10	nd	
	MQ IV	17187	8	nd	
	MQ IV	17175	6	-837	66
	MQ V	18649	7	-708	25
	MQ VI	18842	10	-747	34
	Average $\pm$SD:		6$\pm$3	-785	80
Acetone	MQ I	17026	2	-911	30
	MQ I	17028	2	-1000	4
	MQ I	17029	2	-1000	18
	MQ II	17137	2	-815	64
	MQ III	17176	4	-858	53
	MQ IV	17190	6	nd	
	MQ IV	17178	6	-815	65
	MQ V	18650	6	-836	67
	MQ VI	18843	10	-766	13
	Average $\pm$SD:		4$\pm$3	-836	87
Ethyl Acetate	MQ II	17138	2	-1000	6
	MQ III	17179	6	-912	56
	MQ IV	17191	6	nd	
	MQ IV	17192	6	nd	
	MQ IV	17184	6	-918	62
	MQ V	18651	2	nd	
	MQ VI	18844	4	-743	34
	Average $\pm$SD:		4$\pm$2	-866	107
Dichloromethane	MQ II	17030	2	-965	3
	MQ II	17032	2	-1000	4
	MQ II	17033	2	-1000	6
	MQ III	17179	2	-858	52
	MQ IV	17194	6	-801	63
	MQ IV	17183	2	-859	61
	MQ IV	17184	6	-862	53
	MQ V	18652	2	-1000	5
	MQ VI	18845	3	-660	50
	Average $\pm$SD:		3$\pm$2	-800	114

Table 2. Total [DOC] and SPE-DOC $\Delta^{14}\text{C}$ values for Suwannee River standards, NBP and Station M water samples. SPE-DOC $\Delta^{14}\text{C}$ values are corrected using mass balance techniques using Milli-Q SPE-DOC blanks (see text for details).

Total DOC				SPE-DOC					
Sample type	UCID lab code #	$\Delta^{14}\text{C}$ (‰)	$\pm$	SPE-DOC fraction (% recovery) ¹	UCID lab code #	Size (μg C)	$\Delta^{14}\text{C}$ (‰)	$\pm$ (‰)	Recovery (%)
Suwannee River standard 1R101N (1999)	17901- 17910; (n=10)	+152	3	Methanol	18619	690	+110	7	62 $\pm$ 4
				(99 $\pm$ 1%)	17848	590	+105	10	58 $\pm$ 4
					17582	789	+114	8	62 $\pm$ 4
				Acetone	17583	10	-286	81	
				(1 $\pm$ 1%)	17849	10	-124	68	
Suwannee River standard 2R101N (2012)	18796- 18799; (n=4)	+45	3	Methanol	18846	380	+20	16	58 $\pm$ 4
				(99 $\pm$ 1%)	18851	390	+20	18	62 $\pm$ 4
				Acetone	18847	38	-329	57	
				(1 $\pm$ 1%)	18852	39	-480	75	
NBP	17616 18270 18276 18253	-246	3	Methanol	18806	762	-263	10	49 $\pm$ 4
				(92 $\pm$ 4%)	18002	690	-255	12	44 $\pm$ 5
				Acetone	18007	20	-717	24	
				(4 $\pm$ 2%)	18003	57	-764	10	
				Ethyl acetate/ Dichloromethane	18008	6	-543	41	
				(2 $\pm$ 1%)	18804	6	-691	95	
Station M	8253 ² 8270 8271 8276	-299	3	Methanol	18267	401	-318	11	43 $\pm$ 3
				(89 $\pm$ 2%)	18243	514	-335	7	45 $\pm$ 3
				Acetone	18268	8	nd		
				(1 $\pm$ 1%)	18244	8	-678	25	
				Ethyl acetate/ Dichloromethane	18245	32	-958	5	
				(0 $\pm$ 1%/10 $\pm$ 2%)					

¹ Recovery in each SPE-DOC fraction calculated as the percentage of the total mass in all fractions.

² Station M UV-oxidation total DOC value is reported by Beaupre and Druffel (2009).

Table 3. SPE-BC $\Delta^{14}\text{C}$, relative BPCA abundance and total BC/OC percentages for Suwannee River standards, and NBP and Station M seawater samples.

Sample type	Fraction	UCID lab code #	Size ($\mu\text{g C}$)	SPE- BC $\Delta^{14}\text{C}$ (‰)	$\pm$	BPCA relative abundance ¹ B3CA:B4CA:B5CA:B6CA	Total BC/OC (%)
SR NOM I (1R101N)	Methanol	17586	112	+148	17	SPE-BC 9:11:53:27 total 16:22:38:21 ²	8 $\pm$ 2
SR NOM II (2R101N)	Methanol	18623 18850	31 20	-462 -346	40 70	SPE-BC 6:10:53:30 SPE-BC 6:12:51:31 total 11:33:37:19 ²	12 $\pm$ 2
NBP	Methanol	17940 17719	43 25	-678 -712	19 36	18:39:38:4 29:29:34:7	
	Acetone	17720 17941	16 42	-587 -615	47 20	23:30:47:0 27:35:37:0	21 $\pm$ 2
	Ethyl acetate/ Dichloromethane	17942 nd	62	-593	15	26:33:39:0 27:35:36:0	
Station M	Methanol	18247 18271	12 15	-592 -649	201 121	56:14:18:12 58:16:18:8	4 $\pm$ 1
Process blanks		17590 17448 17490 17261	8 6 6 10	-867 -864 -767 -917	2 3 4 2		

¹ Errors for the BPCA relative abundance determined by peak areas in FID trace is $\pm 3\%$

for B3CA, $\pm 4\%$ for B4CA, $\pm 3\%$ for B5CA and $\pm 2\%$ for B6CA

²total BPCA relative abundances were determined by SR NOM through the BPCA method

Figure 1. Flow chart of methods used to isolate and analyze SPE-DOC and SPE-BC.

Resin blanks are listed in Supplementary Table 1. Chemical structure of HP-20 resin from Aldrich Chemical catalog.

Figure 2 (a.) SPE-DOC $\Delta^{14}\text{C}$ values of methanol fractions for freshwater SR NOM samples, NBP and Station M seawater (open circles) with comparisons to total DOC $\Delta^{14}\text{C}$ values (black dash). (b.) Corresponding percent SPE-DOC recoveries per individual sample are shown as grey columns.

Figure 3. SPE-DOC recoveries for two sample types, riverine and marine DOC. Fraction compositions from methanol fraction (more polar, grey), acetone, ethyl acetate (EtOAc) (dark grey) to dichloromethane (less polar, black) are shown in a.) SR NOM I and II b.) NBP, and c.) Station M sites. There is no significant difference between SR NOM I and II with regard to SPE-DOC composition.

Figure 4 (a.) The relative BPCA abundance and $\Delta^{14}\text{C}$ values (white circles) of the methanol SPE-BC fractions for SR NOM I, SR NOM II, NBP, Station M SPE-BC. The inset shows UDOC relative BPCA abundances from Ziolkowski and Druffel, (2010) at Station M (1000m) and the North Central Pacific (50 m) (located 800km north of Oahu, Hawaii). Surface [BC] in UDOC-BC at NCP is $0.09\ \mu\text{M}$, and is $0.33\pm0.05\mu\text{M}$ in the deep at Station M. (b.) The relative BPCA abundance and $\Delta^{14}\text{C}$ values of all SPE-BC (white circles) fractions at NBP.

Figure 1.

Figure 2.

Figure 3.

Figure 4 a.,b.