Increase of transcription factor EB (TFEB) and lysosomes in rat DRG neurons and their transportation to the central nerve terminal in dorsal horn after nerve injury

Junyang Jung^{1,4*}, Noriko Uesugi^{2,4*}, Na Young Jeong^{3*}, Byung Sun Park¹, Hiroyuki Konishi^{2,4}, Hiroshi Kiyama^{2,4#}

*: These authors contributed equally to this work.

¹Department of Anatomy and Neurobiology, School of Medicine, Kyung Hee University, Heogi-Dong 1, Dongdaemun-Gu, Seoul 130-701, Republic of Korea.

²Department of Functional Anatomy & Neuroscience, Nagoya University, Graduate School of Medicine, 65 Tsurumaicho, Showa-ku, Nagoya 466-8550 Japan.

³Department of Anatomy and Cell Biology and Mitochondria, Hub Regulation Center, Dong-A University College of Medicine, 3-1 Dongdaesin-dong, Seo-gu, Busan 602-714, Republic of Korea

⁴Core Research for Evolutional Science and Technology (CREST) of the Japan Science and Technology Agency, Saitama, Japan.

#Corresponding author: Prof. Hiroshi Kiyama, Department of Functional Anatomy & Neuroscience, Nagoya University, Graduate School of Medicine, 65 Tsurumaicho, Showa-ku, Nagoya 466-8550, Japan. Tel: +81-52-744-2015, Fax: +81-52-744-2020, E-mail address: kiyama@med.nagoya-u.ac.jp

Number of figures: 10

Number of tables: 0

Number of pages: 25

Total number of words: 3,949

Keywords: lysosomal exocytosis, dorsal horn, spinal nerve injury, dorsal root ganglion neurons, microglia

Abstract

In the spinal dorsal horn (DH), nerve injury activates microglia and induces neuropathic pain. Several studies clarified an involvement of adenosine triphosphate (ATP) in the microglial activation. However, the origin of ATP together with the release mechanism is unclear. Recent *in vitro* study revealed that an ATP marker, quinacrine, in lysosomes was released from neurite terminal of dorsal root ganglion (DRG) neurons to extracellular space via lysosomal exocytosis. Here, we demonstrate a possibility that the lysosomal ingredient including ATP released from DRG neurons by lysosomal-exocytosis is an additional source of the glial activator in DH after nerve injury. After rat L5 spinal nerve ligation, mRNA for transcription factor EB (TFEB), a transcription factor controlling lysosomal activation and exocytosis, was induced in the DRG. Simultaneously both lysosomal protein, LAMP1- and vesicular nuclear transporter (VNUT)-positive vesicles were increased in L5 DRG neurons and ipsilateral DH. The increased quinacrine staining in DH was co-localized with LAMP1 immunoreactivity after nerve injury. In DH, LAMP1-positive vesicles were also co-localized with a peripheral nerve marker, IB4 lectin. Injection of the adenovirus encoding mCherry-LAMP1 into DRG showed that mCherry positive lysosomes are transported to the central nerve terminal in DH. These findings suggest that activation of lysosome synthesis including ATP packaging in DRG in response to nerve injury, the central transportation of the lysosome and subsequent its exocytosis from the central nerve terminal of DRG neurons could be a partial mechanism for activation of microglia in DH. This lysosome-mediated microglia activation mechanism may provide another clue to control nociception and pain.

Introduction

Adenosine triphosphate (ATP) was established as the source of free energy involved in biochemical pathways. However, ATP is now recognized as a key molecule both of an intracellular energy source and an intercellular signaling (Fields and Stevens, 2000). It has been demonstrated that ATP is released from axons *in vitro* (Holton, 1959; Soeda et al., 1997; Vizi et al., 1997; Jung et al., 2013) and involved in synaptic transmission (Engelman et al., 2004), and currently the role of ATP is established as a transmitter of relevant purinergic signaling (Pankratov et al., 2002; Zhang et al., 2003). ATP in the central nervous system (CNS) could be a crucial neurotransmitter because ATP-mediated activation of P2X and P2Y receptors in postsynaptic neuron, microglia, and astrocyte can trigger significant Ca^{2+} entry into cytoplasm and influence many metabolisms within cells (Lalo et al., 1998; Abbracchio et al., 2009). A recent study revealed that microglia express several ATP receptors after nerve injury and that released ATP contributed to the activation of microglia (Inoue et al., 2007). In addition, ATP-induced microglial activation in the spinal dorsal horn (DH) caused the neuropathic pain (Tsuda et al., 2003). Thus, the regulation of ATP release in DH may be crucial for the treatment of neuropathic pain associated with microglial activation. However, the mechanisms underlying ATP release to the extracellular space in DH are poorly understood.

Proposed pathways of ATP release from cells are vesicular exocytosis and diffusion via transmembrane pores (Pankratov et al., 2006; Imura et al., 2013). Previous reports have shown that ATP was stored in presynaptic terminals and it released mainly through vesicular exocytosis (Iijima, 1983; Morel and Meunier, 1981; Sawynok et al., 1993; Terrian et al., 1989). In our previous *in vitro* study, we suggested that ATP in lysosomes could be released from primary cultured dorsal root ganglion (DRG) neurons via lysosomal exocytosis (Jung et al., 2013). Lysosomes are acidified, enzyme-containing intracellular organelles to break down phagocytosed materials, cell debris and waste materials (Holtzman, 1989). Recently, it was identified that lysosomes have an additional property for regulatory exocytosis (secretory lysosomes) (Blott and Griffiths, 2002). The exocytotic process of mature lysosomes can be triggered by an increase of intracellular free Ca^{2+} (Blott and Griffiths, 2002; Andrews, 2000). Then, a microtubule-dependent step provides the movement of

exocytic lysosomes towards the plasma membrane (Andrews, 2000). The lysosomal exocytosis also can be triggered by chemicals that cause alkalization of lysosomes (Sundler, 1997). The lysosomal ingredients such as enzymes, degraded molecules, ions and ATP, which are all released by lysosomal exocytosis, could be mediators for the activation of microglia.

In the present study, we addressed a possibility of lysosomal exocytose is from central nerve terminal of DRG neurons into DH after peripheral nerve injury (Tsuda et al., 2003; Kim and Chung, 1992). We here demonstrated that peripheral nerve injury induced the transcription factor EB (TFEB) mRNA expression in DRG and subsequent increase of vesicular-nuclear-transporter (VNUT)-positive lysosomal vesicles in both DRG neurons and DH suggesting up-regulation of lysosomal exocytosis in DRG neurons. In addition the transportation of lysosomes from DRG to DH was also confirmed by injecting adenovirus encoding fluorescence-labeled lysosomal protein. The results suggested that lysosomal exocytosis in DH after peripheral nerve injury may be partly associated with microglial activation in DH.

Materials and Methods

Materials

The primary antibodies against LAMP1 (Sana Cruz Biotechnology, Sana Cruz, CA, USA, Cat# sc8098, RRID: AB_2134494; Cat# sc71489, RRID: AB 2265605) were used for immunostaining or western blotting. Phospho-p44/42 MAPK (p-ERK1/2, Cat# 9101S, RRID: AB 331046) and p44/42 MAPK (ERK1/2, Cat# 9102S, RRID: AB 10695746) were obtained from Cell signaling (Beverly, USA). VNUT and Iba-1 were obtained from MBL Co., Ltd (Woburn, MA, USA, Cat# BMP079, RRID: AB 10597575) and Wako (Osaka, Japan, Cat# 019-19741, RRID: AB_839504), respectively. Isolectin B4 (IB4) and all other reagents were purchased from Sigma-Aldrich (St. Luis, MO, USA).

Animals and surgical procedure for nerve injury

All animal experiments have been carried out in accordance with EU Directive 2010/63/EU for animal experiments. Also all animal experiments have been carried out in accordance with the University Animal Committee “Guidelines for the Care and Use of Laboratory Animals”, and were approved by the Nagoya University Institutional Animal Care and Use Committee. All possible efforts were made to minimize suffering. Adult male Wistar rats (7 weeks rats) were obtained from SLC (Hamamatsu, Japan). Animals were housed with a 12h light/dark cycle (8:00/20:00) at a constant room temperature (RT) of 23 ± 2 °C and humidity of 45~65%. After induction of anesthesia by intraperitoneal (i.p.) injection of pentobarbital (40mg/kg), the L5 nerve was tightly ligated with a 5-0 silk and cut distal to the ligature as previously described (Tozaki-Saitoh et al., 2008).

Tissue processing

Seven days after L5 spinal nerve ligation (SNL), the L5 spinal cord were collected and stored as appropriate for each experiment. For immunohistochemistry (IHC), tissues were perfused with 4% paraformaldehyde (PFA) in 0.1 M phosphate buffer (PB) and then excised. The samples were post-fixed by the same solution overnight at 4°C and treated by with 30% sucrose in 0.1-M PB for 3 days. They were then embedded in OCT and frozen immediately on dry ice. Frozen sections (16 μ m) were

cut from tissues and put on the polylysine-coated slide glasses (Fisher Scientific, Houston, TX, USA).

For the analysis of protein and RNA, the spinal cord was isolated from animals immediately after sacrifice. The spinal cord was separated into DHs at the L5 segment. Each sample was homogenized in a modified radioimmunoprecipitation assay buffer (RIPA; 50mmol/L Tris-HCl pH 7.4, 150mmol/L NaCl, 0.5% deoxycholic acid, 0.5% Triton X-100, 1 mmol/L phenylmethylsulfonyl fluoride, 1 mmol/L sodium o-vanadate, and protease inhibitor cocktail [Roche Molecular Biochemicals, Nutley, NJ, USA]). Protein concentrations were quantified by a Bradford assay following standard protocols. Total RNA was extracted from each sample by the conventional acid guanidineisothiocyanate/phenol/chloroform extraction (AGPC) method (Chomczynski and Sacchi, 2006).

Immunohistochemistry

Before staining, sample slides were fixed in 4% PFA for 10min. Following three washes in phosphate-buffered saline (PBS), the samples were permeabilized in PBS containing 0.3% Triton X-100 (PBST) and blocked with 5% bovine serum albumin (BSA) and 5% fetal bovine serum overnight at 4°C. Primary antibodies for each molecule were placed on the slides, which were incubated for 16 h at 4°C. Following three washes in PBS, slides were incubated for 2h at RT with Alexa Fluor 594 (Invitrogen, Carlsbad, CA, USA), Alexa Fluor 488 (Invitrogen, Carlsbad, CA, USA) or Alex Flour 647 (Invitrogen, Carlsbad, CA, USA) secondary antibodies. The slides were then washed three times with PBS, slide-mounted and subsequently cover-slipped. The immunolabeling was analyzed using an LSM700 imaging system(Carl Zeiss, Oberkochen, Germany).

Western blotting

Western blotting analysis was performed with the tissues described above Protein extracts were separated using 10% sodium dodecyl sulfate polyacrylamide gel electrophoresis and transferred to a nitrocellulose membrane (GE Healthcare, Waukesha, WI, USA). The membrane were blocked with 5% non-fat milk in Tris-buffered saline (TBS) containing 0.05% Tween 20 (TBST) overnight at 4°C and incubated with the following primary antibodies for 1 h at RT; mouse anti-beta actin (1:5000),

goatanti-LAMP1 (1:1000), p-ERK1/2 (1:1000) and ERK1/2 (1:1000). After three 10-min washes in TBST, the blots were incubated for 1 h at RT with the appropriate horseradish peroxidase-conjugated secondary antibodies (1:2000; Invitrogen; Carlsbad, CA, USA), developed in enhanced chemiluminescence reaction (Amersham; Buckinghamshire, England) for 1 min and film-detected (AGFA, Mortsel, Belgium).

Semi-quantitative reverse transcriptase polymerase chain reaction (RT-PCR)

RT-PCR analysis was performed with the tissues described above. In briefly, isolated RNA was obtained from L4-L6 spinal cord and then reverse-transcribed with random 6-mer oligo dT using SuperScript II (Invitrogen Corporation, Carlsbad, CA, USA). RT-PCR was performed using the specific primers as follows: TFEB, forward, 5'-GGTGCAGTCCTACCTGGAGA-3'; reverse, 5'-CTTTCTTCTGCCGTTCTTG-3', Neu-1, forward, 5'-TCGGCTCCGTAGACACTTT-3'; reverse, 5'-CGTGGTCATCACTGAGGAGA-3', VNUT, forward, 5'-GCT TCATCACTG TCACCACA-3'; reverse, 5'-CCAGGACAAGGTCTTTCTCA-3', LAMP1, forward, 5'-AACCCAGTGTGTCCAAGTA-3'; reverse, 5'-GCTGACAAAGATGTGCTCCT-3', GAPDH, forward, 5'-CAGCAATGCATCCTGCACC-3'; reverse, 5'-TGGACTGTGGTCATGAGCCC-3'. RT-PCR was performed using 30 PCR cycles depending on the target gene, with annealing temperatures of 60 °C. PCR products were visualized by 1% agarose gel electrophoresis followed by GelRed (Biotium, Hayward, CA, USA) staining.

ATP staining

For ATP staining *in vivo*, we used quinacrine dihydrochloride as a fluorescent dye. Quinacrine staining was performed as described in previous report (Lee et al., 2013). Briefly, ATP staining was performed by intraperitoneal injection with quinacrine (50 mg/kg) in PBS to rats three times per day (Belai and Burnstock 1994). Slide preparation was performed as described above.

Adenovirus production and its delivery *in vivo*

Cosmid pAxCALNLmLAMP1-mCherry and pAxCAWCre were constructed by the insertion of mouse LAMP1-mCherry or Cre recombinase sequence into the *Swa*I cloning site of pAxCALNLw or pAxCAWt respectively. Using a Cell Pfect transfection kit (GE Healthcare, Cat# 27-9268-01), recombinant adenovirus of AxCALNLmLAMP1-mCherry and AxCAWCre was generated by transfecting 293 cells with DNA-TPC (Takara Shuzo, Japan) and pAxCALNLmLAMP1-mCherry or pAxCAWCre. When these two viral vectors co-infected cells, Cre recombinase expressed under the CAG promoter activates the stuffer PolyA through the Cre/LoxP system. Recombinant adenovirus particle titer (VP) was determined spectrophotometrically. Each adenoviral vector was used at the concentration of 1.5×10^{10} VP/ μ L (AxCALNLmLAMP1-mCherry) or 5.0×10^9 VP/ μ L (AxCAWCre). The viral solutions were injected into L5 DRG (total, 1.0 μ L) concurrently when L5 nerve was injured.

Statistical analysis

Differences between groups were statistically analyzed using analysis of variance followed by Bonferroni's post hoc test. Data were assessed as mean $\pm$ SEM. Values were considered to be statistically significant at $p < 0.01$ after the analysis of four independent experiments.

Result

Increase of LAMP1 expression and TFEB mRNA in DRG neurons after nerve injury

To confirm whether the nerve injury increases lysosomal vesicles in DRG neurons *in vivo* as shown in the previous *in vitro* result (Jung et al., 2013), we immunostained injured L5 DRG with anti-LAMP1 antibody. Activating transcription factor 3 (ATF3) was used for a neuron-injury marker (Takeda et al., 2000; Tsujino et al., 2000; Ohba et al., 2003). A significant increase of LAMP1 immunoreactivity was observed in the ATF3-positive DRG neurons (Fig. 1A-C). Next, we examined a possibility that the ATP stores are associated with lysosomal vesicles in a DRG neuron using quinacrine staining and immunohistochemistry with anti-LAMP1 antibody. Fig. 1D showed clear merged signals of LAMP1 and quinacrine in a DRG neuron cell body. These data indicate that ATP could be localized in lysosomal vesicles in DRG neurons *in vivo* as well as DRG *in vitro* (Jung et al., 2013). The increase of

LAMP1 mRNA in *in vitro* DRG neurons was validated independently by semi-quantitative RT-PCR analysis. In the ipsilateral side of DRG neurons, mRNA for LAMP1 was increased compared with that of the contralateral side of DRG neurons (Fig.2A, B).

Recent studies have reported that TFEB activates lysosomal exocytosis (Medina et al., 2011; Settembre et al., 2011), but neuramidase-1 (Neu-1) negatively regulates lysosomal exocytosis (Yogalingam et al., 2008). To confirm induction of lysosomal synthesis and exocytosis in injured DRG neurons, we performed RT-PCR analysis with specific primers for TFEB and Neu-1 mRNAs. We found that TFEB mRNA level was significantly increased in the ipsilateral DRG neurons compared with contralateral DRG neurons, but no alteration of Neu-1 mRNA expression level in the injured DRG (Fig.2A, B). These results indicate that mRNAs for TFEB and LAMP1 as well as the number of lysosomes are concomitantly increased in the injured DRG neurons, suggesting that induced TFEB expression activates lysosomal synthesis and possibly lysosomal exocytosis in DRG neurons after nerve injury.

Increase of VNUT expression in DRG neurons after nerve injury and co-localization with LAMP1

Previous studies have revealed that VNUT plays a crucial role in transporting ATP into vesicles in the several cell types (Jung et al., 2013; Shehab et al., 2004; Swada et al., 2008). We therefore immunostained DRG sections with anti-VNUT and anti-LAMP1 antibodies. The VNUT immunoreactivity was significantly increased in the injured side compared with uninjured side of L5 DRG (Fig.3A, B). The increased VNUT immunoreactivities were also overlapped with LAMP1-positive profiles in the injured side (Fig.3C). RT-PCR analysis also indicated the increased mRNA expression level of VNUT in the injured DRG neurons compared with the control DRG neurons (Fig.3D, E). Thus, these findings suggest lysosomal vesicles are capable of storing ATP through VNUT in DRG neurons.

Increase of LAMP1 immunoreactivity in the ipsilateral DH after nerve injury

To determine the alteration of lysosomal localization in the spinal DH after sciatic nerve injury, we performed immunostaining using sections from the L5 level of spinal cord with anti-LAMP1 antibody. We found that LAMP1 immunoreactivity was significantly increased in the ipsilateral side of spinal DH (Fig.4A). Previous studies have reported that spinal nerve injury induces microglial activation in the injured side of spinal DH (Tsuda et al., 2005; Watkins and Maier, 2003). We confirmed the microglial activation and increase in the lysosome-increased ipsilateral DH using Iba-1 antibody (Fig.4A).

In addition, the increase of LAMP1 immunoreactivity was confirmed by Western blot analysis. LAMP1 expression in DH at day 1 after nerve injury was significantly increased and the increase was maintained at day 7 after nerve injury (Fig.4B, C). The increase of LAMP1 in DH after nerve injury had continued for 3 weeks (data not shown). A previous study showed that pERK1/2 activation occurs in the spinal DH after nerve injury (Zhang et al., 2007b). We used pERK1/2 as a positive control and ERK1/2 as a loading control. To assess whether the increase of LAMP1 immunoreactivity is in microglia or not, we performed simultaneous visualization of LAMP1 and Iba-1 immunoreactivities. Many large LAMP1 positive puncta were observed in ipsilateral DH. Many of LAMP1 immunoreactivities did not co-localized with Iba-1 immunostaining in the ipsilateral side of DH (Fig.4D). These results indicate that the increased large lysosomes mainly exist out of microglia after SNL.

Next, we accessed whether the ATP stores are associated with lysosomal vesicles in DH using quinacrine staining and immunohistochemistry with anti-LAMP1 antibody. In 6 day after SNL, rats were injected with quinacrine and, the next day, spinal cords were sampled for analyses. In L5 spinal cord sections, quinacrine staining was apparently more detectable in the ipsilateral dorsal horn compared to the contralateral side of the dorsal horn after SNL (Fig. 5A). In addition, the most prominent quinacrine-positive staining was found in the laminae I and II of the ipsilateral dorsal horn (Fig.5A arrows). Low basal constitutive quinacrine staining was found in the contralateral region of DH as well as in cell bodies of motor neurons in the ventral horn (Fig. 5A).

In superficial layers of DH, the quinacrine staining was co-labeled with LAMP1 (Fig.5B). High-

magnification images also showed that the most of intense quinacrine staining co-localized with LAMP1 (the lowest panel in Fig.5B). Although all LAMP1 immunoreactivities were not co-localized with the quinacrine staining, most of quinacrine staining showed co-localization with LAMP1 immunoreactivity, suggesting that there exist various types of lysosomes with abundant to less ATP. A quantitative analysis indicated that LAMP1-positive signals (Fig.5C) and quinacrine-positive signals (Fig.5D) were significantly increased in the ipsilateral DH compared with the contralateral side after SNL. Percentage of quinacrine/LAMP1-double positive signals in the total of LAMP1-positive signals indicated no significant difference between the ipsilateral and contralateral side of DH after SNL (Fig.5E). Taken together, these results suggested that number of ATP containing vesicles are increased in the injured side of DH after nerve injury and the increased ATP storage would be lysosomal vesicles in DH.

Increase of VNUT immunoreactivity in the injured DH and co-localization with LAMP1 immunoreactivity

Since higher rate of co-localization of VNUT and LAMP1 immunoreactivities were observed in DRG neurons after nerve injury, we also examined their co-localization in DH. The increases of mRNAs for LAMP1 and VNUT in the ipsilateral side of DH were identified 7days after nerve injury by RT-PCR (Fig. 6A, B). Next, we immunostained both VNUT and LAMP1 in the spinal cord sections. The VNUT immunostaining was significantly increased in the ipsilateral side compared with contralateral side of spinal DH (Fig.6C), and VNUT immunoreactivities were co-localized with LAMP1-positive profiles significantly in the ipsilateral side compared with the contralateral side of DH (Fig.6D,E). These findings suggest that lysosomal vesicles in DH are capable of storing ATP through VNUT.

Co-localizations of quinacrine/IB4in the nociceptive central afferent terminals *in vivo*

We previously reported that lysosomal activation is increased in the axon terminal of primary cultured DRG neurons (Jung et al., 2013). We therefore examined whether the LAMP1 positive vesicles locate in the central terminals from DRG neurons. To examine a possibility that the increased LAMP1

immunoreactivity seen in DH after nerve injury is in the central branch of DRG neurons, we compared LAMP1 immunoreactivity with IB4. We used IB4 as a maker of C-fiber nociceptor central terminals. Our IB4 staining existed in the afferent central terminal of lamina II (Munglani et al., 1995; Shehab et al., 2004) and the most prominent LAMP1 immunoreactivity was found in the laminae II of the ipsilateral dorsal horn within IB4 binding region (Fig. 7A, B). Next, we compared quinacrine staining with IB4 binding. We found that quinacrine staining overlapped with IB4 staining in the ipsilateral dorsal horn (Fig. 7C). Quantitative result showed that quinacrine/axon marker double-positive signals were increased in the ipsilateral side of L5 spinal dorsal horn (Fig. 7D). Thus, these findings indicate that the increase of quinacrine staining and lysosomal vesicles may occur in the axon terminals originated from the injured DRG neurons.

Localization of DRG-originated LAMP1 in dorsal horn

To evaluate the transportation of lysosomal vesicles to DH from DRG neurons more directly, we constructed adenovirus vector expressing LAMP1-mCherry. 7 days after infecting L5 DRG neurons with the LAMP1-mCherry-expressing adenovirus, a few L5 DRG neurons expressed the mCherry fluorescence (Fig. 8A), suggesting a minor DRG neurons were infected and expressed the mCherry-LAMP1. In the ipsilateral DH, the mCherry-positive fine nerve fibers were observed in both superficial and deeper layers, but not in the contralateral DH (Fig. 8B, C). In the ipsilateral DH, mCherry was co-localized with IB4 (Fig. 9 A-C arrows), but not with Iba1 (a marker for microglia, Fig. 9 D-F). This indicated that mCherry-labeled lysosomes synthesized in IB4 positive DRG neurons were transported to DH. Obviously the mCherry-labeled lysosomes were considered as DRG originated lysosomes. Collectively, these findings support an idea that the increase of quinacrine staining and lysosomal vesicles in DH after nerve injury would be partly due to the transportation of ATP containing lysosomes from DRG neurons.

Discussion

In this study, we demonstrated that nerve injury activated lysosomal biosynthesis in DRG neurons and

VNUT-positive lysosomes were transported to the primary afferent nerve terminal in DH (Fig. 10). We found that the injury of peripheral process of DRG neurons induced the expression of TFEB mRNA (Fig. 2A, B). TFEB is a master transcriptional regulator of lysosomal biogenesis, and the TFEB transcriptionally induces both the docking and fusion of lysosomes with the plasma membrane by regulating the expression of certain genes (Medina et al., 2011; Settembre et al., 2011, 2013). This suggests that the increase of lysosomal vesicles in DRG neurons could be due to the induction of TFEB in DRG neurons. Intriguingly, together with the increase of lysosomes in DRG neurons, VNUT mRNA was induced in the same DRG neurons, and prominent co-localization of VNUT and LAMP1 was observed on vesicles. This would suggest that ATP storage in lysosomes is activated in nerve-injured neurons for some reason.

What are the functional consequences of the lysosome increase in injured DRG neurons? One most likely explanation would be an involvement of lysosomes in membrane repair at injured peripheral nerve process (Settembre et al., 2013; Appelqvist et al., 2013). Previous studies have reported that lysosomal vesicles are involved in the process of membrane repair through Ca^{2+} -dependent synaptotagmin VII in the damaged cell by a way to fuse with the plasma membrane (Settembre et al., 2013; Appelqvist et al., 2013; Gerasimenko et al., 2001; Idone et al., 2008; McNeil, 2002). The present nerve injury might induce the influx of Ca^{2+} into DRG neurons and subsequently induces the number of lysosomal vesicles and lysosomal exocytosis under the activation of TFEB. The increased fusion of lysosomes with plasma membrane could function to repair the injured neurite of DRG neurons. The second possibility would be an intracellular clearance of molecules and organelles by lysosomal activation and exocytosis. Under pathological conditions, TFEB mediated activation of lysosomal biogenesis and exocytosis would be advantageous for removing intracellular toxic materials and injured organelle to outside of cell membrane (Medina et al., 2011; Settembre et al., 2013). Nerve injury elicits various types of stresses including oxidative stress and ER stress in neurons and these stresses would damage organelle and induce accumulation of unfolded proteins. For the removal of those unnecessary or even toxic contents, lysosomal degradation and clearance including autophagy

would be necessary in nerve-injured neurons. Although both the lysosome-mediated membrane repair and intracellular clearance are critical in injured peripheral process and soma of DRG neurons, it is likely that the increased lysosome plays some role in the central process as well. DRG neurons have both peripheral and central processes and it appears that no directional preferences to be transported exist for some neurotransmitters such as substance P and organelle such as mitochondria at least (Harman and Keen, 2010). In fact, the present adenovirus mediated expression of mCherry-LAMP1 clearly demonstrated the transport of lysosomes in the central branch of DRG neurons. Thus, in response to nerve injury, ATP-containing lysosomes would be delivered to both central and peripheral terminal of DRG neurons and whereby the ingredients including ATP might be released.

Recently lysosomes are considered as one of storage vesicles for ATP in neuronal and glial cells, and the lysosomal exocytosis possibly releases ATP together with other ingredients into the extracellular space (Jung et al., 2013; Shin et al., 2012; Zhang et al., 2007a; Pryazhnikov and Khiroug, 2008). The present co-localization study of LAMP1 and quinacrine in the same vesicle suggested that the increased lysosomal vesicles in the ipsilateral DH after nerve injury also contain ATP (Fig.5). Because the existence of ATP in lysosomes and ATP release by lysosomal exocytosis were demonstrated in cultured DRG neurons (Jung et al., 2013), it is likely that the lysosomal exocytosis is a mechanism underlying ATP release in DH after nerve injury. Substantial increase of lysosomal protein, LAMP1, was demonstrated in the injured DH by immunohistochemistry and Western blot analysis for LAMP1 (Fig.4 and 5). Thus, these findings suggest that the lysosomal ATP may be associated with nociceptive pathophysiology because the activation of ATP containing lysosomes occurred in the area where the central terminals of nociceptive c-fiber from DRG exist (Fig.7-9).

Several types of axons and processes including the central process of DRG neurons locate in DH. Because quinacrine- and LAMP1-positive vesicles were highly increased in DRG neurons after nerve injury as shown in Fig. 1, 2 and 3 and our previous study (Jung et al., 2013), it is very likely that ATP containing lysosomes is transported to DH in response to peripheral nerve injury. To clarify this possibility, peripheral nerve marker IB4 was used and LAMP1-positive structures existed jointly with

IB4 immunoreactivity in the ipsilateral DH with adenoviral infection (Fig. 8 and 9). In addition, we showed the increase of VNUT expression in the ipsilateral DH and the injured DRG neurons and its *in vivo* expression in lysosomal vesicles (Fig.3 and 6) as shown in our previous *in vitro* study (Jung et al., 2013). These findings together with our previous study (Jung et al., 2013) would support our hypothesis that peripheral nerve injury induces the transportation of ATP-containing lysosomes to the central branch of DRG neurons and suggests a possibility that ATP is released from axon terminals of DRG neurons through lysosomal exocytosis in addition to releases from astrocytes and microglia (Zhang et al., 2007b; Dou et al., 2012).

In conclusion, we demonstrated the transcriptional activation of lysosomal exocytosis activator TFEB in DRG neurons in response to nerve injury and prominent increase of lysosomes was also observed both in both DRG and DH of the spinal cord where DRG neurons project. Since it becomes evident that the transportation of lysosomes from DRG to DH through the central branch of primary afferent neurons, it is likely that ATP in addition to some additional ingredients are released by the lysosomal exocytosis at central terminal of DRG neurons. Although we could not demonstrate direct release of ATP from the primary afferent nerve terminal in DH via lysosomal exocytosis, previous *in vitro* study supported the possibility (Jung et al 2013). Thus, a regulation of lysosomal exocytosis at the axon terminals of DRG neurons may provide an alternative clue to control nociception and pain.

Acknowledgments: This study was partially supported by CREST, JST, Japan and a Grant-in-Aid for Scientific Research from the ministry of Education, Science, Sports and Culture of Japan (MEXT) and Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT and future Planning (J. Jung, 2015R1A2A2A01002735; N. Y. Jeong, 2015R1C1A1A02036863).

We are grateful to Profs. Inoue K and Tsuda M (Kyushu Univ., Japan) for their kind advices, Ms. Y. Tabata and N. Tawarayama for their technical assistance, and Ms. A. Asano for secretarial assistance.

References

- Abbracchio MP, Burnstock G, Verkhratsky A, Zimmermann H (2009) Purinergic signalling in the nervous system: an overview. *Trends Neurosci* 32:19-29.
- Andrews NW (2000) Regulated secretion of conventional lysosomes. *Trends Cell Biol* 10:316-321.
- Appelqvist H, Wäster P, Kågedal K, Öllinger K (2013) The lysosome: from waste bag to potential therapeutic target. *J Mol Cell Biol* 5:214-226.
- Belai A, Burnstock G (1994) Evidence for coexistence of ATP and nitric oxide in non-adrenergic, non-cholinergic (NANC) inhibitory neurons in the rat ileum, colon and anococcygeus muscle. *Cell Tissue Res* 278:197-200.
- Blott EJ, Griffiths GM (2002) Secretory lysosomes. *Nat Rev Mol Cell Biol* 3:122-131.
- Chomczynski P, Sacchi N (2006) The single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction: twenty-something years on. *Nat Protoc* 1:581-585.
- Dou Y1, Wu HJ, Li HQ, Qin S, Wang YE, Li J, Lou HF, Chen Z, Li XM, Luo QM, Duan S (2012) Microglial migration mediated by ATP-induced ATP release from lysosomes. *Cell Res* 22:1022-1033.
- Engelman HS, Mac Dermott AB (2004) Presynaptic ionotropic receptors and control of transmitter release. *Nat Rev Neurosci* 5:135-145.
- Fields RD, Stevens B (2000) ATP: an extracellular signaling molecule between neurons and glia. *Trends Neurosci* 23:625-633.
- Gerasimenko JV, Gerasimenko OV, Petersen OH (2001) Membrane repair: Ca(2+)-elicited lysosomal exocytosis. *Curr Biol* 11: R971-R974.
- Harmar A, Keen P (1982) Synthesis, and central and peripheral axonal transport of substance P in a dorsal root ganglion-nerve preparation in vitro. *Brain Res* 231:379-385.
- Holtzman E (1989) Autophagy and related phenomena, in *Lysosomes*, (Siekevitz P, eds), pp 243–316 New York Plenum Press.
- Holton, P (1959) The liberation of adenosine triphosphate on antidromic stimulation of sensory nerves. *J Physiol* 145:494-504.

- Idone V, Tam C, Andrews NW (2008) Two-way traffic on the road to plasma membrane repair. *Trends Cell Biol* 18:552-559.
- Iijima T (1983) Quinacrine-induced degeneration of non-adrenergic, non-cholinergic autonomic nerves in the rat anococcygeus muscle. *Cell Tissue Res* 230:639-648.
- Imura Y, Morizawa Y, Komatsu R, Shibata K, Shinozaki Y, Kasai H, Moriishi K, Moriyama Y, Koizumi S (2013) Microglia release ATP by exocytosis. *Glia* 61:1320-1330.
- Inoue K, Tsuda M, Tozaki-Saitoh H (2007) Modification of neuropathic pain sensation through microglial ATP receptors. *Purinergic Signal* 3:311-316.
- Jung J, Shin YH, Konishi H, Lee SJ., Kiyama H (2013) Possible ATP release through lysosomal exocytosis from primary sensory neurons. *Biochem. Biophys. Res. Commun.* 430, 488-493.
- Kim SH, Chung JM (1992) An experimental model for peripheral neuropathy produced by segmental spinal nerve ligation in the rat. *Pain* 50:355-363.
- Lalo U, Voitenko N, Kostyuk P (1998) Iono- and metabo-tropically induced purinergic calcium signalling in rat neocortical neurons. *Brain Res* 799:285-291.
- Lee SJ, Rao AS, Shin YH, Chung HJ, Huh Y, Ahn KH, Jung J (2013) A novel method using an acedan-based Zn(DPA) probe to monitor ATP localization in an in vivo system. *J Mol Histol* 44:241-247.
- Maeda M, Tsuda M, Tozaki-Saitoh H, Inoue K, Kiyama H (2010) Nerve injury-activated microglia engulf myelinated axons in a P2Y₁₂ signaling-dependent manner in the dorsal horn, *Glia* 58:1838-1846
- Mc Neil PL, (2002) Repairing a torn cell surface: make way, lysosomes to the rescue. *J Cell Sci* 115: 873-879.
- Medina DL, Fraldi A, Bouche V, Annunziata F, Mansueto G, Spampanato C, Puri C, Pignata A, Martina JA, Sardiello M, Palmieri M, Polishchuk R, Puertollano R, Ballabio A (2011) Transcriptional activation of lysosomal exocytosis promotes cellular clearance. *Dev Cell* 21:421-430.
- Morel N, Meunier FM (1981) Simultaneous release of acetylcholine and ATP from stimulated

- cholinergic synaptosomes. *J. Neurochem* 36:1766-1773.
- Munglani R., Bond A, Smith GD, Smith GD, Harrison SM, Elliot PJ, Birch PJ, Hunt SP (1995) Changes in neuronal markers in a mononeuropathic rat model relationship between neuropeptide Y, pre-emptive drug treatment and long-term mechanical hyperalgesia. *Pain* 63:21-31.
- Ohba N, Maeda M, Nakagomi S, Muraoka M, Kiyama H (2003) Biphasic expression of activating transcription factor-3 in neurons after cerebral infarction. *Brain Res Mol Brain Res* 115:147-156.
- Pankratov Y, Lalo U, Krishtal O, Verkhratsky A (2002) Ionotropic P2X purinoreceptors mediate synaptic transmission in rat pyramidal neurons of layer II/III of somato-sensory cortex. *J Physiol* 542:529-536.
- Pankratov Y, Lalo U, Verkhratsky A, North RA (2006) Vesicular release of ATP at central synapses. *Pflugers Arch* 452:589-597.
- Pryazhnikov E, Khiroug L (2008) Sub-micromolar increase in $[Ca^{2+}]_i$ triggers delayed exocytosis of ATP in cultured astrocytes. *Glia* 56:38-49.
- Sawynok J, Downie JW, Reid AR, Cahill CM, White TD (1993) ATP release from dorsal spinal cord synaptosomes: characterization and neuronal origin. *Brain Res* 610:32-38.
- Settembre C, Di Malta C, Polito, V.A., Garcia Arencibia, M., Vetrini, F., Erdin, S., et al, 2011. TFEB links autophagy to lysosomal biogenesis. *Science* 332, 1429-1433.
- Settembre C, Fraldi A, Medina DL, Ballabio A (2013) Signals from the lysosome: a control centre for cellular clearance and energy metabolism. *Nat Rev Mol Cell Biol* 14:283-296.
- Shehab SA, Spike RC, Todd AJ (2004) Do central terminals of intact myelinated primary afferents sprout into the superficial dorsal horn of rat spinal cord after injury to a neighboring peripheral nerve? *J. Comp. Neurol.* 474, 427-437.
- Shin YH, Lee SJ, Jung J (2012) Secretion of ATP from Schwann cells through lysosomal exocytosis during Wallerian degeneration. *Biochem Biophys Res Commun* 429:163-167.
- Soeda, H, Tatsumi H, Katayama Y (1997) Neurotransmitter release from growth cones of rat dorsal root ganglion neurons in culture. *Neuroscience* 77:1187-1199.
- Sundler, R (1997) Lysosomal and cytosolic pH as regulators of exocytosis in mouse macrophages.

- Acta Physiol Scand 161:553-556.
- Sawada K, Echigo N, Juge N, Miyaji T, Otsuka M, Omote H, Yamamoto A, Moriyama Y (2008) Identification of a vesicular nucleotide transporter. *Proc Natl Acad Sci USA* 105:5683-5686.
- Takeda M, Kato H, Takamiya A, Yoshida A, Kiyama H (2000) Injury-specific expression of activating transcription factor-3 in retinal ganglion cells and its colocalized expression with phosphorylated c-Jun. *Invest Ophthalmol Vis Sci* 41:2412-2421.
- Terrian DM, Hernandez PG, Rea MA, Peters RI, (1989) ATP release, adenosine formation, and modulation of dynorphin and glutamic acid release by adenosine analogues in rat hippocampal mossy fiber synaptosomes. *J Neurochem* 53:1390-1399.
- Tozaki-Saitoh H, Tsuda M, Miyata H, Ueda K, Kohsaka S, Inoue K, (2008) P2Y₁₂ receptors in spinal microglia are required for neuropathic pain after peripheral nerve injury. *J Neurosci* 28:4949-4956.
- Tsuda M, Shigemoto-Mogami Y, Koizumi S, Mizokoshi A, Kohsaka S, Salter MW, Inoue K (2003) P2X₄ receptors induced in spinal microglia gate tactile allodynia after nerve injury. *Nature* 424:778-783.
- Tsuda M, Inoue K, Salter MW (2005) Neuropathic pain and spinal microglia: a big problem from molecules in "small" glia. *Trends Neurosci* 28:101-107.
- Tsujino H, Kondo E, Fukuoka T, Dai Y, Tokunaga A, Miki K, Yonenobu K, Ochi T, Noguchi K (2000) Activating transcription factor 3 (ATF3) induction by axotomy in sensory and motoneurons: A novel neuronal marker of nerve injury. *Mol Cell Neurosci* 15:170-182.
- Vizi ES, Liang SD, Sperlágh B, Kittel A, Jurányi Z, (1997) Studies on the release and extracellular metabolism of endogenous ATP in rat superior cervical ganglion: support for neurotransmitter role of ATP. *Neuroscience* 79:893-903.
- Watkins LR, Maier SF (2003) Glia: a novel drug discovery target for clinical pain. *Nat Rev Drug Discov* 2:973-985.
- Yogalingam G, Bonten EJ, van de Vlekkert D, Hu H, Moshiah S, Connell SA, d'Azzo A (2008) Neuraminidase 1 is a negative regulator of lysosomal exocytosis. *Dev Cell* 15:74-86.

- Zhang JM, Wang HK, Ye CQ, Ge W, Chen Y, Jiang ZL, Wu CP, Poo MM, Duan S (2003) ATP released by astrocytes mediates glutamatergic activity-dependent heterosynaptic suppression. *Neuron* 40: 971-982.
- Zhang X, Chen Y, Wang C, Huang LY (2007a) Neuronal somatic ATP release triggers neuron-satellite glial cell communication in dorsal root ganglia. *Proc Natl Acad Sci USA* 104:9864-9869.
- Zhang Z, Chen G, Zhou W, Song A, Xu T, Luo Q, Wang W, Gu XS, Duan, S (2007b) Regulated ATP release from astrocytes through lysosome exocytosis. *Nat Cell Biol* 9:945-953.

Figure legends

Fig. 1. LAMP1 activation in the injured DRG neuronal cell bodies

(A) Confocal images showed ATF3 (green) activation in the injured DRG neurons. Scale bar, 200 μm . (B) High magnification images showed LAMP1 (red) activation in ATF3 (green)-positive DRG neuronal cell bodies (arrows). Scale bar, 50 μm . (C) Quantitative analysis of immunofluorescent staining against LAMP1 and ATF3. The number of LAMP1-activated ATF3 neuronal cell bodies in DH (200 μm x 200 μm) was calculated. $**p < 0.01$ (n=4). (D) Quinacrine staining (green) and immunostaining (red) with anti-LAMP1 antibody in a DRG neuron. Scale bar, 5 μm . Arrows show the co-localization of LAMP1-immunoreactivity and quinacrine.

Fig. 2. Alteration of LAMP1 and TFEB mRNAs expressions in the injured DRG

(A) RT-PCR analysis of LAMP1, TFEB and Neu-1 mRNA in the total RNA extract from the rat DRG on day 7 after peripheral nerve injury. Neuramidase 1 (Neu1) is a negative regulator for lysosomal exocytosis (Yogalingamet al., 2008). GAPDH is loading control. Cont, contralateral; Ipsi, ipsilateral. (B) Quantification of the relative intensity of bands derived by RT-PCR. $**p < 0.01$ (n=3).

Fig. 3. Increases of VNUT immunoreactivity and its mRNA in nerve injured DRG

(A) Confocal images showed that the vesicular nucleotide transporter (VNUT, green) immunoreactivity was increased in injured DRG neurons compared with uninjured DRG neurons after L5 SNL (upper left panel). Scale bar, 200 μm . DIC indicated that amounts of contralateral DRG neuron cell bodies are similar to ipsilateral DRG. DIC, Differential interference contrast. (B) Quantitative analysis showed that nerve injury induced an increase in the relative signal intensity of VNUT-positive profiles in DRGs (200 μm x 200 μm). $**p < 0.01$ (n=4). (C) Upper right panels (Scale bar, 30 μm) and lower panel (Scale bar, 15 μm) are high magnification images of injured DRG neuronal cell bodies (Fig. 2A). Arrows indicate VNUT (green) activation in LAMP1 (red)-positive DRG neuronal cell bodies. (D) Alterations in VNUT mRNA expression in DRG neurons after L5 SNL (n=4). (E) Quantification of the relative intensity of mRNA expression of VNUT in DRG after nerve

injury (n=4).

Fig. 4. Increase of LAMP1 immunoreactivity in DH after peripheral nerve injury

(A) Immunohistochemistry reveals increases in LAMP1 (red) level in the ipsilateral spinal horn (L5) 7 days after peripheral nerve injury. NeuN (blue) and Iba-1 (green) indicated the expression control and injured side, respectively. Scale bar, 200 μ m. (B) Protein extracts (10 μ g) from spinal dorsal horn were analyzed by western blotting. ERK1/2 was loading control. (C) Quantitative analysis of Western blotting for LAMP1 expression from three independent experiments. $**p<0.01$ (n=4). (D) Simultaneous demonstration of LAMP1 (red) and Iba1 (green). LAMP1 immunoreactivity is co-localized mainly out of Iba-1 in the ipsilateral side after nerve injury. The right hand side photo demonstrates higher magnification of the white square region seen in left. Scale bar, 50 μ m (left) and 10 μ m (right).

Fig. 5. Co-localization of LAMP1 immunoreactivity and quinacrine staining in the spinal dorsal horn

(A) Confocal images showing a ATP marker quinacrine (white) in the L5 spinal cord on day 7 after peripheral nerve injury, Scale bar, 500 μ m. Higher magnification images (lower panel) showed accumulation of quinacrine staining in the laminae II (arrows) of the ipsilateral dorsal horn. Scale bar, 200 μ m. (B) LAMP1 immunoreactivity (red) within the ipsilateral dorsal horn of the spinal cord 7 day after peripheral nerve injury co-localized with quinacrine staining (green) (arrows). Quin, quinacrine. Scale bar, 50 μ m. 3 panels in the bottom are higher magnification images of middle 3 panels. Scale bar, 20 μ m. (C) Quantitative analysis of immunofluorescent staining against LAMP1 (n=4). The number of LAMP1-positive profiles in DH (200 μ m x 200 μ m) was calculated. $**p<0.01$ (n=4). (D) Quantitative analysis of quinacrine fluorescence staining. The number of quinacrine-positive profiles in DH (200 μ m x 200 μ m) was calculated. Ipsi, ipsilateral. Cont, contralateral. $**p<0.01$ (n=5). (E) Percentage of quinacrine/LAMP1-double positive signals in the total LAMP1-positive signals.

Fig. 6. Increases of VNUT and LAMP1 mRNAs and their immunoreactivities in the injured DH

(A) RT-PCR analysis of LAMP1 and VNUT mRNAs in the spinal dorsal horn on day 7 after L5 SNL. GAPDH is loading control. Cont, contralateral; Ipsi, ipsilateral. (B) Quantification of the relative intensity of bands derived by RT-PCR. $**p < 0.01$ (n=3). (C) Confocal images showing that the vesicular nucleotide transporter (VNUT, green) immunoreactivity was increased in the ipsilateral side compared with contralateral side of DH after L5 SNL. Scale bar, 200 μm . (D) High magnification images of VNUT (green) and LAMP1 (red) immunostaining in DH. 20 μm . (E) Quantitative analysis showed that nerve injury induced VNUT-immunoreactivity profiles co-labelled with LAMP1-immunoreactivity per 100 μm^2 in the ipsilateral DH after nerve injury. $**p < 0.01$ (n=3).

Fig. 7. Intense LAMP1 immunoreactivity localizes in IB4 enriched layers

(A) IB4 (green) and LAMP1 (red) immunoreactivity in DH. Scale bar = 200 μm . (B) Higher magnification images of A. Scale bar = 200 μm . (C) quinacrine-positive puncta (green) is colocalized with IB4 (red) in the ipsilateral DH (arrows). Scale bar, 10 μm . (D) Quantitative analysis showed that nerve injury induced the increase of IB4-positive quinacrine puncta per 100 μm^2 in the ipsilateral dorsal horn after nerve injury. $**p < 0.01$ (n=4).

Fig. 8. Vesicles expressing LAMP1-mCherry in DRG neurons are anterogradely transported to dorsal horn.

(A) A representative image of L5 DRG which is injected the adeno-virus encoding LAMP1-mCherry. Arrows indicate the neurons expressing LAMP1-mCherry in DRG. (B) Contralateral dorsal horn of L5 spinal cord (Cont) 7 days after peripheral nerve injury and adeno-virus injection. (C) Ipsilateral side (Ipsi) of identical section with B. Many LAMP1-mCherry positive fibers are seen in the ipsilateral dorsal horn. Scale bars indicate 100 μm .

Fig. 9. Vesicles expressing LAMP1-mCherry are transported in IB4 positive fibers.

Higher magnification images showing LAMP1-mCherry (red) containing fibers in the dorsal horn of

L5 spinal cord (**A and D**). These are co-stained with IB4 (green in **B**) or Iba1 (green in **E**) respectively. (**C**) The merged images of **A** and **B**, (**F**) The merged image of **D** and **E**. Arrows in **A-C** indicate the LAMP1-mCherry and IB4 double-positive fibers. None of the LAMP1-mCherry positive structure was colocalized with Iba1-positive cells. Scale bars indicate 25 μm .

Fig. 10. Schematic demonstration of the present results

Red large dots, lysosomes; green small dots, ATP

Fig. 1

Fig. 2

A

B

Fig. 3

Fig. 4

A

B

C

D

Fig. 5

Fig. 6

A

B

C

D

E

Fig. 7

FIGURE 8

FIGURE 9

Fig. 10