1 Cool Bands: Wing bands decrease rate of heating, but not equilibrium temperature in
2 *Anartia fatima*

3

4 Jake Brashears^a, Annette Aiello^b, Brett M. Seymoure^{b,c*}

5 ^a *Department of Life Sciences, San Diego City College, San Diego 92101, United States*

6 ^b *Smithsonian Tropical Research Institute, Apartado 0843-03092 Balboa, Ancon, Rep.*
7 *Panama*

8 ^c *School of Life Sciences, Arizona State University, Tempe 85287, United States*

9 *Email address: jbrashea@sdccd.edu (J. Brashears), aielloa@si.edu (A. Aiello),*

10 **brett.seymoure@asu.edu (B.M. Seymoure)*

11 **Corresponding Author. Address: School of Life Sciences, Arizona State University,*

12 *Tempe, AZ. 85287. Tel: + (269) 501-8761*

13

14

15 *Running Title: Wing bands decrease rate of heating in Anartia fatima*

16

17

18 *Abstract*

19 Butterflies regulate their internal thoracic temperature in order to optimize
20 performance activities (e.g. flight, foraging). Previous research has shown that butterfly
21 wings, particularly the innermost portions, play a role in thermoregulation. We
22 investigated to see whether a lightly colored wing band would alter the thermal properties
23 of the banded peacock butterfly (*Anartia fatima*) with two within subject experiments in a

laboratory setting: 1) band color manipulation in which euthanized individuals were heated to thermal equilibrium with the band unaltered and then again with the wing darkened; 2) wing ablation in which individuals already run through experiment 1 were heated to equilibrium two more times; once with the outer portion of the wing including the band removed and then with the entire wing removed. Individuals were spread so that the dorsal surface of the wing was exposed to illumination from a lamp suspended above. Twelve *Anartia fatima* males were collected in Panama and were run through experiment one. Four individuals were run through experiment two. We found no effect of darkening the band on the internal thoracic equilibrium temperature, but the darkened band did increase the rate of heating. The wing ablation experiment revealed that wing removal lowered the internal thoracic equilibrium temperature but did not affect the heating rate. Therefore we show that butterfly bands may be important in butterfly thermoregulation and we discuss the importance of the wing band on thermoregulatory abilities in *Anartia fatima* with respect to the butterfly's natural history. We conclude that the wing band may allow butterflies to reduce heat stress induced by their warm environments.

Keywords: Butterfly Coloration, Radiation, Reflectance, Thermoregulation

Introduction

Thermoregulation is an important component of butterfly biology. Although the physical structures, such as wing size and thorax, ultimately limit thermoregulatory capacity, behavioral changes (e.g., basking, body postures) can greatly affect heat gain and loss and allow butterflies to keep their body temperature close to the optima for flight,

foraging, etc. (Heinrich, 1972; Kingsolver and Moffat, 1982; Kingsolver and Wiernasz, 1991; Kleckova et al., 2014; Rawlins, 1980; Rutowski et al., 1994). Through basking and adjusting body posture, butterflies increase internal temperature by absorbing solar radiation or by conducting heat through body parts placed onto warmer substrates (Heinrich, 1993; Huey et al., 2003; Kingsolver, 1987). Wing flapping or increased exposure to wind decreases body temperatures through convective cooling. Because butterflies are not able to change their coloration, wing color is thermally limiting in that darker coloration absorbs more solar radiation than lighter coloration (Berwaerts et al., 2001; Watt, 1968). Here, we revisit the question of how wing coloration affects thermoregulatory properties in a butterfly with a wing band.

Seminal research has demonstrated the effects of wing coloration on both the rate of heating and the internal thoracic temperature (Kammer and Bracchi, 1973; Schmitz, 1994; Wasserthal, 1975). Larger and darker wings (i.e., wings with more melanin) absorb more energy, thus increasing heat gain (Kammer and Bracchi 1973; Wasserthal 1975; Kingsolver 1987; Schmitz 1994). Although wing coloration is important, Wasserthal (1975) demonstrated that only the 15% of the wing surface area closest to the thorax (that is, coloration within 2-3mm of the thorax) affects thoracic temperature, the main target for butterfly thermoregulation (Kingsolver, 1985; May, 1979). However, these trends may not be applicable to all butterflies. Recent studies have shown a more complicated relationship between wing coloration and thoracic temperature.

Wing coloration can be produced by pigments, structurally (e.g. photonic crystals), or by a combination of both (Nijhout, 1991). Iridescent structural coloration absorbs more solar radiation and has a greater effect on internal temperatures than pigmentary

coloration (Bosi et al., 2008). Furthermore, De Keyser et al. (2015) found that color had less of an effect on thoracic temperature in small Lycaenid butterflies. Therefore, although there is a basic understanding of the role of coloration in butterfly thermoregulation, our understanding is based on a few species and further research is warranted. No research has explicitly tested the thermoregulatory function of wing bands, even though many banded butterflies occur in thermally challenging climates. Many butterflies have bands or stripes that are much darker or lighter than the wing background, and this difference in brightness could enhance or reduce the solar radiation absorbed by the wing and thorax.

Anartia fatima (Fabricius, 1793) (Nymphalidae), the banded peacock butterfly, has a conspicuous yellowish band that fades to white with age and is set within a field of dark brown on both the dorsal and ventral wing surfaces of the fore and hind wings (Figure 1). The continuous band of 3 to 4 mm in width runs from the anterior to posterior wing edges about halfway between the wing base and distal wing margin. The continuous band is set approximately halfway on the wing, 1.25 cm from the thorax and 1.25 cm from the wing margin. *Anartia fatima* is found in disturbed habitats in Central and South America where the environment can be very hot and dry (Silberglied et al., 1979). Furthermore, *A. fatima* perches on vegetation in open areas and may not be able to seek shade for thermal relief (Silberglied et al., 1979). Hence, *A. fatima* may be under strong thermoregulatory pressures to reduce overheating in its habitat and the wing band may function to reduce the absorption of heat by the thorax. We tested the effect of the wing band in *A. fatima* on both heating rate and thermal equilibrium in a laboratory setting. We predicted that the band would have a minimal effect on the two thermal metrics due to the

band being further than .5 cm from the thorax, as is predicted from Wasserthal (1975).
These experiments are the first to explicitly test the thermoregulatory role of wing bands
in butterflies, which will further our understanding of evolutionary and ecological
significance of wing bands and coloration of butterflies.

Methods

Butterfly Collection

We collected 12 *Anartia fatima* males from gardens and fields in Gamboa,
Panama (9.1167N, 79.7000W) during May of 2015; all individuals had no evident wing
wear or scale loss (see Kemp 2006). All specimens were euthanized by freezing and then
stored individually in a freezer until use. For each trial, two butterflies were removed,
weighed, and mounted on a .33mm diameter type T thermocouple connected to a digital
recorder (Dual J-T-E-K Thermocouple Thermometer, Digi-sense, Vernon Hills, IL). The
thermocouple was inserted into the thorax, between the mesothoracic segments, to a
depth of ~2 mm at an angle perpendicular to the frontal plane. The wings were then
spread to an angle of approximately 180 degrees relative to each other to achieve
maximum light exposure.

Band Manipulation Experiment

We then measured reflectance at three locations on three parts of the wings: the
brown wing surface outside of the band, within the band (both unaltered and darkened),
and the brown inner wing surface (Fig. 1). We recorded reflectance perpendicular to the
wing surface and relative to a white spectralon standard (Ocean Optics, Dunedin, FL). All

reflectance spectra were recorded with SpectraSuite software (Ocean Optics, Dunedin, FL) from a bifurcated reflectance probe (Ocean Optics, Dunedin, FL) connected to a USB2000 spectrometer (Ocean Optics, Dunedin, FL). We recorded reflectance from 300 nm to 800 nm (UV to infrared). The spectra were then processed in R (cran) with the Pavo package (Maia et al., 2013). We extracted total reflectance (i.e. the total number of photons, calculated as the integral of the spectrum from 300 to 800nm) for each patch (i.e. distal brown, unaltered band, proximal brown, and darkened band) for each individual.

Mounted butterflies were positioned and allowed to thermally equilibrate to room conditions of illumination and temperature, which was maintained within a half degree of 27°C. Room illumination consisted of a single fluorescent light and had no detectable thermal effect, which we confirmed before each trial by measuring the surface temperature using a non-contact thermal gun (AR550, Smart Sensor, Intell Instruments) at four locations: external thorax, proximal side of band, band, and distal side of band. These temperatures were statistically no different from the air temperature and internal thoracic equilibrium temperature (repeated measures ANOVA; $F_{4, 50} = 1.29$, $p = 0.286$).

The trials consisted of heating each butterfly two times, once not-manipulated and then again with the darkened band, under a lamp (60 watt halogen, Satco S2248, Brentwood, NY) placed 28 cm above the butterflies. The light source was directly overhead of the two specimens, and light was passed through two band pass filters, one short wavelength transmissive and one long wavelength transmissive (L176R and L503R, Rosco Laboratories Inc, Stamford, CT), to achieve a spectrum closely resembling natural light (see Fig. 2 for irradiance spectra). Butterflies were heated for 15 minutes, which was sufficient for all samples to attain thermal equilibrium. Internal thoracic temperature was

recorded each minute during the trial. Before ending the trial, surface temperatures were again recorded with the non-contact thermal gun. After 15 minutes, the lamp was turned off and butterflies were allowed to reequilibrate to room temperature (27°C). We then painted each band black with a permanent marker (Sharpie, fine point, Downers Grove, IL). No obvious damage to wings occurred during the marking manipulation. Three spectral readings were taken of each artificially darkened band. The procedure was then repeated with the wings darkened: surface temperatures were recorded; butterflies were heated again for 15 minutes and internal thoracic temperatures were recorded at intervals of a minute: surface temperatures were recorded after 15 minutes. To control for body size, water loss, and order effects, post-trial weight and femur length were recorded at the end of each trial.

Wing Ablation Experiment

We selected four butterflies, in good condition, from the previous band manipulation experiment for the ablation experiment. They were weighed, then both wings were cut carefully along the proximal edge of the band, so each butterfly had only the portions of the wings proximal to the band intact. They were weighed, and then two butterflies were mounted on the thermocouples, on the same side of the lamp as they were in the previous experiment. After equilibration to room temperature (27°C), surface temperatures were recorded from the thorax and the proximal wing, and the lamp was turned on. Internal thoracic temperature was recorded every minute for 15 min, and after 15 minutes the surface temperatures were again recorded. Once all four had completed this trial, the butterflies were weighed, and their proximal wings were removed to leave

only the thorax. These were weighed and mounted again for a final heating trial. The heating procedure was repeated exactly as before, but surface temperature was recorded only from the thorax. A final individual mass, without wings, was collected.

Calculations and Statistics

The internal thoracic equilibrium temperature was calculated as the mean of the final three recorded values. Following Kammer and Bracchi (1973), we calculated the rate of heating as the slope of the internal thoracic temperature after the temperature had climbed higher than 3.2 °C above ambient. To ensure that this measurement correctly measured warming rate, we also calculated the rate of heating as the slope during the first two minutes of the trial and as a function of the time required to reach within one standard deviation of final equilibration temperature. As all metrics were statistically the same, we present only the first. The total reflectance was calculated as the mean of the proximal and distal integrals of each spectra. All data analyses were performed with the open software R (R Core Team 2014) and the lme4 package (Bates et al. 2015).

We used generalized linear mixed models to compare the effects of wing darkening on the final equilibrium temperature and the rate of heating. We included pre-trial mass, femur length, and total reflectance as independent variables. Two random variables were also included: individual to control for the repeated measures design, and position on the right or left side of the lamp to control for possible unequal heating. To compare external temperatures recorded at the thorax, proximal wing, band, and distal wing, we used the same model but included a comparison of pre- and post-trial temperatures. Significance of fixed factors were calculated using Satterthwaite's method

of denominator synthesis to generate the correct error terms. Likelihood ratios were used to test the significance of random factors (Satterthwaite, 1946). The effects of ablation on the equilibrium temperature and rate of heating of the internal thorax were analyzed using a repeated measures analysis of variance followed by a paired t-test with alpha error correction if significance was found.

Results

Wing Reflectance

The total reflectance was significantly different between the four patches ($F_{3,11}=116.2$, $p < 0.001$; Fig. 1), and post-hoc analyses revealed that the total reflectance of only the non-manipulated band was significantly different from other patches (i.e. distal, darkened band, and proximal). The non-manipulated band was much brighter than the darkened band, with mean total reflectance of 143.5 and standard deviation of 28.51, while the darkened band had a mean total reflectance of 32.47 and standard deviation of 21.15.

Band Manipulation Experiment

The initial masses of the butterflies were 0.74 ± 0.03 g. Femur length was 4.38 ± 0.31 mm. The internal thorax heating rate was faster in the darkened wing band treatment than the non-manipulated wing (Table 1; Fig. 4) and was significantly different across individuals (Table 1). Each butterfly reached an internal thoracic equilibrium temperature that was 12°C warmer than ambient temperature, Fig. 3B and Fig. 3C. However, internal thoracic equilibrium temperature did not differ between the non-manipulated band and

darkened wing manipulation (Table 1; Fig. 3A), although the position under the lamp did have a significant effect (Table 1). All parts of the wing increased in temperature during the trial (pre/post comparison in Table 1); however, there were no temperature differences between the two treatments for the wing surface, whether distal (Table 1; Fig. 3A), on the band (Table 1; Fig. 3A), or the proximal wing (Table 1; Fig. 3A). However, darkening the wing did increase the equilibrium temperature attained by the external thorax (Table 1; Fig. 3A). Femur length and pre-trial mass had no significant effect on any dependent variables. Average mass lost during heating was the same, approximately 6%, for both natural and darkened heating trials.

Wing Ablation Experiment

Full wing ablation significantly decreased the equilibrium temperature attained by the internal thorax ($F_{3,9} = 14.8$, $p < 0.001$, Fig. 5A & 6), but partial wing ablation had no significant effect on equilibrium temperature ($F_{3,8} = 0.18$, $p = 0.91$, Fig. 5A & 6). When the entire wing was removed, the internal equilibrium temperature of the thorax was about two and a half degrees less than the natural unaltered butterflies (a decrease of 6% from un-manipulated equilibrium temperature), while partial wing ablation was only half a degree less (a decrease of 1% from un-manipulated equilibrium temperature). Paired t-tests showed a significant difference between butterflies having darkened wings and completely ablated wings (Table 2; Fig. 5A & 6). There was no significant effect of the ablation experiment on the rate of heating (Fig. 5B & 6).

Discussion

We predicted that darkening the wing band would have minimal effects on the internal thoracic equilibrium temperature and the rate of heating in *Anartia fatima*. We did find that the band slows the rate of heating in *A. fatima*, but we did not find any differences between the two treatments for equilibrium temperatures. Our results support previous findings by Kammer and Bracchi (1973) and Wasserthal (1975) in that coloration further than 3mm away from the thorax has a minimal effect on internal thoracic temperatures. And like Wasserthal (1975) and Kammer and Bracchi (1973), we demonstrated that only the proximal portion of the wing is important for internal thoracic temperatures.

It must be noted that although we found an increased rate of heating after the darkened band manipulation, this result may be an order effect and caused by a decreased thermal inertia from evaporative water loss during the first heating trial. And again, this supports our initial hypothesis that the bands have minimal thermal effect. That is, when the plane of *Anartia fatima* dorsal wing surfaces are perpendicular to a light source mimicking the sun, with very little air moving across their bodies, and even accounting for a decreased thermal inertia, darkening their wing band does not allow them to reach a higher thoracic temperature, and it increases their rate of heating by less than 0.5°C/min.

Surprisingly, there was no significant effect of femur length, mass, and non-band wing total reflectance on equilibrium temperature and heating rate, although these metrics have been shown to have a strong effect on these variables in other studies (Berwaerts et al., 2001; Gilchrist, 1990; Kingsolver, 1987; Watt, 1968). In the Baltimore checkerspot (*Euphydryas phaeton*), females are better able to maintain body temperatures than males, which are half the size of females (Gilchrist, 1990). It is likely that we did not find a

254 difference in thermoregulatory ability dependent upon size in *A. fatima* due to the
255 similarity in size of the collected individuals. Although there is variation in size between
256 individuals, *A. fatima* does not have large size variation between individuals of the same
257 sex. However, females and males do differ in size and thus, there may be differences in
258 thermal properties of males and females. Lastly, Huey et al. (2003) provide a theoretical
259 construct on how thermoregulatory function may not be a leading driver of morphology
260 because organisms can mediate morphological constraints through behaviors.

261 The position of the butterflies, whether to the right or left of center in the paired
262 trials, had a significant effect on equilibrium temperature and distal wing temperature
263 (Table 1), indicating that the lamp provided uneven heating. We provided for this effect
264 experimentally by randomizing the assigned position of each butterfly and statistically by
265 including it as a random factor in our model to account for the variance. Similarly, to
266 account for the variance produced by taking temperatures before and after heating, we
267 included pre- and post-comparisons as a fixed factor in our model; its significance
268 verifying that each part of the butterflies increased in temperature (Table 1).

269 Our findings support both experimental and observational studies showing that
270 wings play a minimal role in thermoregulation of internal thoracic temperature (Kammer
271 and Bracchi, 1973; Wasserthal, 1975). The results of wing ablation in our experiment
272 showed that removing the distal portion of the wing, including the band, made no
273 difference compared to intact wings. We only found a thermal difference when we
274 removed the entire wing from an individual and this could be due to water loss from the
275 thorax, more than a result of radiation absorption of the wing. Furthermore, due to the
276 small sample size of the wing ablation experiment (n=4), we may have not detected

differences between the natural wing and the wing partially removed. Kammer and Bracchi (1973) showed that wing circulation does not play a role in heat conduction from the wings, and melanization had no effect on heating rates in the small lycaenid butterfly, *Polyommatus icarus* (De Keyser et al., 2015). Our results support the overall conclusion that absorption of energy by the thorax is more important than absorption by the whole wing, and that the thorax is the primary driver of temperature in lighter and darker butterflies under natural conditions (Heinrich 1972).

In a natural environment, the effects of convection are likely much stronger than any band heating effect. Convection can reduce heat load dramatically so that even in a species where there is a dark morph, for example in the speckled wood butterfly (*Pararge aegeria*), the dark morph engages on longer flights that make it consistently have a lower body temperature than the lighter morph, despite heating faster under controlled conditions (van Dyck 1998). We attempted to repeat our experiment in the meadow adjacent to the field station and were unable to get butterflies to reach an internal thoracic equilibrium temperature (i.e. the internal temperature reaches a horizontal asymptote) due to the effects of a mild breeze.

Two observations, that minimal convection removed the thermal effect of the band when we attempted to reproduce the experiment in a meadow, and that ablation up to and including the band did not affect equilibrium temperature (Table 1; Fig. 5A) or rate of heating (Table 1; Fig. 5B), warrant further research on the ecological role of the band in *A. fatima*. Although no published research has investigated thermoregulatory behavior in *A. fatima*, *A. fatima* individuals do perch with wings held at varying and ever changing angles and this may serve a thermoregulatory function as shown in other

butterflies (Barton et al., 2014; Berwaerts et al., 2001). Of course, the slower rate of heating due to the occurrence of a band, coupled with changes in wing position could be adaptive for *A. fatima*. *Anartia fatima* occupies open tropical habitats that are exposed to bright sunshine where the daily ambient temperatures can exceed 37°C and in captive settings, individuals are very susceptible to dying from heat stress and dehydration (Seymoure personal observation). Having a band that would slow the rate of overheating would allow *A. fatima* to spend more time searching for nectar sources, mates and host plants. Alternatively, the band may function in warming *A. fatima* individuals through reflective basking, in which solar radiation is reflected from light wing patches onto the thorax through behavioral adjustment of body position (Kingsolver, 1987). Now that there is evidence that a wing band has thermal effects, further research into the effects of a light band on thermoregulation in the field is needed.

Bands can serve ecological functions other than just thermoregulation. Emmel (1972) demonstrated that males use the wing band for selecting mates and Taylor (1973) showed that the total reflectance and color of the band can be informative of individual age as the wing band fades over time. Recently, Seymoure & Aiello (2015) supported the role of the band functioning as disruptive crypsis, as individual models with a natural band were less likely to be attacked than models that had the band moved to the wing margin. Therefore, *Anartia fatima* has many selective pressures acting upon its coloration and the three pressures mentioned above (thermoregulation, mate choice, and crypsis) may all contribute to advantages conferred by the wing band.

Conclusions

We tested the thermoregulatory role of the wing band in *Anartia fatima* and found that the band had an effect on the heating rate of the thorax but no effect on internal thoracic temperature equilibrium. We further supported that wings have a minimal effect on thermal properties by showing that when wings are removed, the thorax has a lower internal temperature. The equilibrium findings align with previous studies showing that wing coloration has a minimal effect on internal thoracic temperatures. However, the effect of band on heating rate is a novel find and urges further research into the thermoregulatory role of wing bands in butterflies, especially behavioral experiments in the field.

Acknowledgements

This work was supported by Arizona State University and the Smithsonian Tropical Research Institute (STRI). The School of Life Sciences Research Training Initiatives Office at ASU provided funding for the materials required for this study. We thank Autoridad Nacional del Ambiente and the Republic of Panamanian for the collecting permit (DAPVS-0478-2015). We are grateful for comments and suggestions from RL Rutowski, R. De Keyser, and an anonymous reviewer, which greatly improved this manuscript.

References

- Barton, M., Porter, W., Kearney, M., 2014. Behavioural thermoregulation and the relative roles of convection and radiation in a basking butterfly. *J. Therm. Biol.* 41, 65–71. doi:10.1016/j.jtherbio.2014.02.004
- Berwaerts, K., Van Dyck, H., Vints, E., Matthysen, E., 2001. Effect of manipulated wing characteristics and basking posture on thermal properties of the butterfly *Pararge aegeria* (L.). *J. Zool.* 255, 261–267. doi:10.1017/S0952836901001327
- Bosi, S.G., Hayes, J., Large, M.C.J., Poladian, L., 2008. Color, iridescence, and thermoregulation in

350 Lepidoptera. Appl. Opt. 47, 5235–5241.

351 De Keyser, R., Breuker, C.J., Hails, R.S., Dennis, R.L.H., Shreeve, T.G., 2015. Why Small Is Beautiful:
352 Wing Colour Is Free from Thermoregulatory Constraint in the Small Lycaenid Butterfly,
353 *Polyommatus icarus*. PLoS One 10, e0122623. doi:10.1371/journal.pone.0122623

354 Emmel, T.C., 1972. Mate Selection and Balanced Polymorphism in the Tropical Nymphalid Butterfly ,
355 *Anartia fatima*. Evolution (N. Y). 26, 96–107. doi:10.2307/2406986

356 Gilchrist, G.W., 1990. The consequences of sexual dimorphism in body size for butterfly flight and
357 thermoregulation. Funct. Ecol. 4, 475–487.

358 Heinrich, B., 1993. The hot-blooded insects, strategies and mechanisms of thermoregulation.
359 Springer-Verlag, Berlin.

360 Heinrich, B., 1972. Thoracic temperatures of butterflies in the field near the equator. Comp. Biochem.
361 Physiol. 43, 459–467.

362 Huey, R.B., Hertz, P.E., Sinervo, B., 2003. Behavioral drive versus behavioral inertia in evolution: a null
363 model approach. Am. Nat. 161, 357–366. doi:10.1086/346135

364 Kammer, A.E., Bracchi, J., 1973. Role of the wings in the absorption of radiant energy by a butterfly.
365 Comp. Biochem. Physiol. 45, 1057–1063.

366 Kemp, D.J., 2006. Heightened phenotypic variation and age-based fading of ultraviolet butterfly wing
367 coloration. Evol. Ecol. Res. 8, 515–527.

368 Kingsolver, J.G., 1987. Evolution and coadaptation of thermoregulatory behavior and wing pigmentation
369 pattern in pierid butterflies. Evolution (N. Y). 41, 472–490. doi:10.2307/2409250

370 Kingsolver, J.G., 1985. Butterfly thermoregulation: organismic mechanisms and population consequences.
371 J. Res. Lepid. 24, 1–20.

372 Kingsolver, J.G., Moffat, R.J., 1982. Thermoregulation and the determinants of heat transfer in *Colias*
373 butterflies. Oecologia 53, 27–33. doi:10.1007/BF00377132

374 Kingsolver, J.G., Wiernasz, D.C., 1991. Seasonal polyphenism in wing-melanin pattern and
375 thermoregulatory adaptation in *Pieris* butterflies. Am. Nat. 137, 816–830. doi:10.2307/2678832

376 Kleckova, I., Konvicka, M., Klecka, J., 2014. Thermoregulation and microhabitat use in mountain
377 butterflies of the genus *Erebia*: Importance of fine-scale habitat heterogeneity. J. Therm. Biol. 41, 50–
378 58. doi:10.1016/j.jtherbio.2014.02.002

379 Maia, R., Eliason, C.M., Bitton, P.P., Doucet, S.M., Shawkey, M.D., 2013. pavo: An R package for the
380 analysis, visualization and organization of spectral data. Methods Ecol. Evol. 4, 906–913.
381 doi:10.1111/2041-210X.12069

382 May, M.L., 1979. Insect thermoregulation. Annu. Rev. Entomol. 22, 313–349.

383 Nijhout, H.F., 1991. The Development and Evolution of Butterfly Wing Patterns. Smithsonian Institution
384 Scholarly Press.

385 Rawlins, J.E., 1980. Thermoregulation by the black swallowtail butterfly , *Papilio Polyxenes* (Lepidoptera:
386 Papilionidae). Ecology 61, 345–357. doi:10.2307/1935193

387 Rutowski, R.L., Demlong, M.J., Leffingwell, T., 1994. Behavioral Thermoregulation At Mate Encounter
388 Sites By Male Butterflies (*Asterocampa-Leilia*, Nymphalidae). Anim. Behav. 48, 833–841.
389 doi:10.1006/anbe.1994.1307

390 Satterthwaite, F.E., 1946. An Approximate Distribution of Estimates of Variance Components. Biometrics
391 Bull. 2, 110–114.

392 Schmitz, H., 1994. Thermal characterization of butterfly wings - 1. Absorption in relation to different color,
393 surface structure and basking type. J. Therm. Biol. 19, 403–412.

394 Seymoure, B.M., Aiello, A., 2015. Keeping the Band Together: Evidence for False Boundary Disruptive
395 Coloration in a Butterfly. J. Evol. Biol. n/a–n/a. doi:10.1111/jeb.12681

396 Silberglied, R.E., Aiello, A., Lamas, G., 1979. Neotropical Butterflies of the Genus *Anartia*: Systematics,
 397 Life Histories and General Biology (Lepidoptera: Nymphalidae). *Psyche A J. Entomol.* 86, 219–260.
 398 doi:10.1155/1979/50172
 399 Taylor, O.R., 1973. A non-genetic “polymorphism” in *Anartia fatima* (Lepidoptera: Nymphalidae).
 400 *Evolution* (N. Y). 27, 161–164.
 401 Team, R.C., 2014. R: A language and environment for statistical computing.
 402 Van Dyck, H., Matthysen, E., 1998. Thermoregulatory differences between phenotypes in the speckled
 403 wood butterfly□: hot perchers and cold patrollers□? *Oecologia* 114, 326–334.
 404 Wasserthal, L.T., 1975. The role of butterfly wings in regulation of body temperature. *J. Insect Physiol.* 21,
 405 1921–1930.
 406 Watt, W.B., 1968. Adaptive significance of pigment polymorphisms in *Colias* butterflies. I. Variation of
 407 melanin pigment in relation to thermoregulation. *Evolution* (N. Y). 22, 437–458.
 408 doi:10.1073/pnas.63.3.767
 409
 410

412

419 Figure 1. Experimental set up and reflectance spectra of the un-manipulated and
 420 manipulated wing patterns. Left, the reflectance spectra are mean (lines) and standard
 421 error (shading). Center and right, the photographs depict the natural coloration of *A.*
 422 *fatima* (left) and the sharpie darkened band of the manipulated wing (right). The colored
 423 shapes in each photograph match the reflectance spectra: green circle = proximal brown;
 424 red square = natural band; purple diamond = distal brown; blue triangle = sharpie
 425 darkened band.

Figure 2. Irradiance spectra of the experimental conditions and natural lighting conditions of *A. fatima*. A) Absolute irradiance in units of energy for the natural lighting of *A. fatima* habitat (green), unfiltered lamp (blue), and filtered lamp (red). B) Normalized irradiance for each condition. Normalized irradiance was calculated by dividing each energy value (binned by one nanometer) by the sum of all the energy values for each spectrum.

427

428

429

Figure 3. (A) Equilibrium temperature recordings for each location for the un-manipulated treatment (uncolored) and for the Sharpie darkened treatment (blue). The internal temperature recordings over the 15 minute trials for each individual for the un-manipulated treatment (B) and for the Sharpie darkened treatment (C). Blue line is air temperature. Asterisk indicates a significant difference ($p < 0.05$) between natural and manipulated conditions (Table 1).

Figure 4. Heating rates for the two treatment groups in experiment one. Asterisks indicate a significant difference ($p < 0.01$) between natural and manipulated conditions (Table 1).

440

441

444 Figure 5. Equilibrium temperatures (A) and heating rates (B) for the four treatment

445 groups in experiment two: wing ablation. Asterisk indicates a significant difference ($p <$

446 0.05) between butterflies with the band darkened and butterflies without wings (Table 2).

Figure 6. The four treatments of the wing ablation experiment for the four individuals.

Each graph shows the four treatments groups for one individual. The treatment groups

are: not manipulated (red squares), darkened band (blue triangles), band and distal brown

cut away (orange asterisks), wing cut away (cyan diamonds).

Table 1. Generalized linear mixed model analysis of the effects darkening the wing band on the internal equilibrium temperature, heating rate, and external temperatures taken at the thorax, proximal wing, band, and distal wing of *A. fatima* butterflies (n=12). Initial mass, femur length, and wing brightness were included as fixed factors, and the individual and position (right or left of center of lamp) were included as random factors. The significance of random factors was tested using likelihood ratio tests. Satterthwaite's (1946) method of denominator synthesis was used to calculate the appropriate error terms.

Equilibrium Temperature							
	Type of factor	df effect	MS effect	error df*	error MS*	F/X ²	p
band darkening	fixed	1	0.02	11.00	0.32	0.07	0.79
initial mass	fixed	1	0.19	7.01	0.32	0.62	0.45
femur length	fixed	1	0.02	7.00	0.32	0.08	0.77
brightness	fixed	1	0.16	7.00	0.32	0.50	0.49
individual	random	1	-	-	-	0.71	0.40
position	random	1	-	-	-	9.87	0.002

Heating Rate							
	Type of factor	df effect	MS effect	error df	error MS	F/X ²	p
band darkening	fixed	1	0.46	11.00	0.035	13.40	0.004
initial mass	fixed	1	0.05	7.07	0.033	1.51	0.25
femur length	fixed	1	0.01	7.00	0.033	0.03	0.87
brightness	fixed	1	0.001	7.00	0.005	0.20	0.66
individual	random	1	-	-	-	4.81	0.03
position	random	1	-	-	-	2.18	0.14

External Thorax							
	Type of factor	df effect	MS effect	error df	error MS	F/X ²	p
band darkening	fixed	1	2.13	34	0.39	5.4	0.02
initial mass	fixed	1	0.40	8	0.40	1.0	0.34
femur length	fixed	1	0.02	8	0.40	0.05	0.84
brightness	fixed	1	1.21	8	0.39	3.1	0.11
pre/post comp.	fixed	1	1524	1	0.39	3881	< 0.001
individual	random	1	-	-	-	1.89	0.2
position	random	1	-	-	-	< 0.001	1.0

Proximal Wing							
	Type of factor	df effect	MS effect	error df	error MS	F/X ²	p
band darkening	fixed	1	1.27	34.00	0.96	1.32	0.25
initial mass	fixed	1	0.12	7.08	1.00	0.12	0.73
femur length	fixed	1	0.04	7.00	1.00	0.04	0.84
brightness	fixed	1	2.43	7.00	0.96	2.53	0.15
pre/post comp.	fixed	1	2165	34.00	0.96	2249	< 0.001
individual	random	1	-	-	-	0.98	0.3

position	random	1	-	-	-	1.54	0.2
----------	--------	---	---	---	---	------	-----

465

Band							
	<i>Type of factor</i>	<i>df effect</i>	<i>MS effect</i>	<i>error df</i>	<i>error MS</i>	<i>F/X²</i>	<i>p</i>
band darkening	fixed	1	0.3	64	1.67	0.18	0.67
initial mass	fixed	1	1.1	8	1.48	0.74	0.41
femur length	fixed	1	0.1	8	1.11	0.09	0.77
brightness	fixed	1	1.7	8	1.53	1.11	0.32
pre/post comp.	fixed	1	3770	34	1.53	2459	< 0.001
individual	random	1	-	-	-	1.23	0.3
position	random	1	-	-	-	0.00	1.0

466

Distal Wing							
	<i>Type of factor</i>	<i>df effect</i>	<i>MS effect</i>	<i>error df</i>	<i>error MS</i>	<i>F/X²</i>	<i>p</i>
band darkening	fixed	1	0.3	41.00	2.98	0.1	0.74
initial mass	fixed	1	1.9	41.12	2.37	0.8	0.37
femur length	fixed	1	1.1	41.00	2.21	0.5	0.49
brightness	fixed	1	4.4	41.00	2.31	1.9	0.17
pre/post comp.	fixed	1	7397	41.00	2.32	3186	< 0.001
individual	random	1	-	-	-	0.00	1.00
position	random	1	-	-	-	8.89	0.003

467

468

469 Table 2. Results from post-hoc analysis, using holm alpha correction, of paired t test
 470 comparisons of internal equilibrium temperatures and heating rates when *A. fatima*
 471 butterflies (n=4) had their natural wings, the wing band darkened, the wing ablated just
 472 past the band, and the wing removed.

Ablation Equilibrium Temperature			
	<i>natural wing</i>	<i>band darkened</i>	<i>ablated to band</i>
natural wing	-	-	-
band darkened	0.23	-	-
ablated to band	0.67	0.23	-
wing removed	0.16	0.03	0.07

473

Ablation Heating Rate			
	<i>natural wing</i>	<i>band darkened</i>	<i>ablated to band</i>
natural wing	-	-	-
band darkened	0.94	-	-
ablated to band	0.19	0.40	-
wing removed	0.75	0.94	0.94

474