**Effects of some cations on the formation of
2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) in a model system**

Di Yu ^a, Shu-Juan Yu ^{a b *}

^a College of Light Industry and Food Sciences, South China University of
Technology, Guangzhou 510640, China

^b Guangdong Province Key Laboratory for Green Processing of Natural Products and
Product Safety, Guangzhou 510640, China

*Corresponding author: Shu-Juan Yu; e-mail: shujuanyu8@gmail.com; Postal address: #381,
Wushan road, Guangzhou, China; Tel.: +86 20 87113668, Fax: +86 20 87113668

1 **Abstract:** The present study aimed to investigate in detail the changes to
2 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) and its precursors in the
3 presence of some cations (i.e., K^+ , Na^+ , Ca^{2+} , Mg^{2+} , Fe^{2+} and Fe^{3+}) in a
4 creatinine/phenylacetaldehyde model system. Results showed that PhIP yields decreased
5 when Fe^{2+} and Fe^{3+} were added to a mixture of phenylacetaldehyde and creatinine. This
6 decrease may be attributed to the fact that Fe^{3+} can form complexes with various
7 properties with creatinine and accelerate creatinine degradation. This pathway can disturb
8 the reaction with phenylacetaldehyde, influence aldol condensation product formation,
9 and suppress PhIP formation. Furthermore, Ca^{2+} and Mg^{2+} enhanced PhIP content. Such
10 enhancement may be attributed to the fact that $CaCl_2$ and $MgCl_2$ promote aldol and aldol
11 condensation product reactions with ammonia and formaldehyde. A possible mechanism
12 for the action of cations during PhIP formation in a model system is also proposed.

13 **Keywords:** aldol condensation product, phenylacetaldehyde, creatinine, some cations,
14 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP)

- 15 Chemical compounds studied in this article
- 16 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) (PubChem CID: 1530);
- 17 Phenylacetaldehyde (PubChem CID: 998);
- 18 Creatinine (PubChem CID: 588);
- 19 *o*-Phenylenediamine (OPD) (PubChem CID: 7243)

1. Introduction

Heterocyclic aromatic amines (HAAs) produced in food during heating at high temperatures are a risk factor for certain human cancers. The formation of HAAs in food has been an ongoing concern among food chemists, nutritionists, and toxicologists because of the potential mutagenic/carcinogenic properties of these compounds (Damašius, Venskutonis, Ferracane, & Fogliano, 2011; Gibis & Weiss, 2012; Nakai & Nonomura, 2013; Turesky & Le Marchand, 2011; Yu, Chen, & Yu, 2016; Zamora, Alcón, & Hidalgo, 2013; Zur Hausen, 2012). Over 25 HAAs have been isolated and identified from food and model systems (Moon & Shin, 2013). Given their potent mutagenic activity and the fact that they can be formed even during ordinary household cooking (Warzecha, Janoszka, Błaszczuk, Stróżyk, Bodzek, & Dobosz, 2004), a great deal of research effort has focused on developing strategies that can effectively lower HAA contents in food and reduce human exposure to HAAs (Cheng, Chen, & Wang, 2006).

2-Amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) is one of the most abundant HAA produced in foods. Several researchers have studied the formation mechanism of PhIP, and a number of hypothetical formation pathways of PhIP in various reaction models have been proposed in recent years. PhIP is produced through the reaction of phenylacetaldehyde with creati(ni)ne (Zamora, Alcón, & Hidalgo, 2013, 2014; Zöchling & Murkovic, 2002). An aldol condensation product with a molecular mass of 215 was found to be an intermediate in this reaction, and the product was later identified as 2-amino-1-methyl-5-(2'-phenylethenyl)-imidazol-4-one (Zöchling & Murkovic, 2002). According to Zamora, Alcón, and Hidalgo (2014), PhIP originates from the reaction of this aldol condensation product, formaldehyde, and ammonia; hence, several factors may

influence the formation of PhIP.

Recent reports have indicated that metal cations can reduce acrylamide formation in thermally processed foods (Gökmen & Şenyuva, 2006, 2007). However, the effects of metal cations on PhIP formation have been ignored in recent literature. In the present paper, we investigated the effects of several cations (K^+ , Na^+ , Ca^{2+} , Mg^{2+} , Fe^{2+} and Fe^{3+}) at different concentrations on the formation of PhIP and its precursors in a creatinine/phenylacetaldehyde model system. A possible mechanism to explain the changes to PhIP and its precursors in the presence of these cations is proposed.

2. Materials and Methods

2.1 Materials

PhIP was purchased from Toronto Research Chemicals (North York, ON, Canada). HPLC-grade methanol and acetic acid were purchased from Tianjin Kermel Chemical Reagent Co., Ltd. (Tianjin, China). The NaCl, KI, $CaCl_2$, $MgCl_2 \cdot 6H_2O$, $FeCl_2 \cdot 4H_2O$, and $FeCl_3 \cdot 6H_2O$ used in this work were of reagent grade and purchased from Sigma-Aldrich (St. Louis, MO). All other chemicals were purchased from Sigma-Aldrich and of analytical-reagent grade. All solutions were prepared with double-distilled water.

2.2 Preparation of model reaction mixtures

A model system composed of creatinine and phenylacetaldehyde was used to study the formation of PhIP in the presence of certain cations (K^+ , Na^+ , Ca^{2+} , Mg^{2+} , Fe^{2+} , Fe^{3+}). Creatinine (0.6 mmol) and phenylacetaldehyde (0.6 mmol) were combined in 25-mL PTFE test tubes with a stainless steel exterior liner. Varying amounts of each cation were added to the reaction mixture. The total reaction volume was adjusted to 10 mL with water in each case. The final concentrations of each cation were 0, 0.5, 1, 2, 4, or 8 mM.

The pH of the reaction mixture was measured as 6.36 prior to addition of the cations. Adding up to 8 mM of cations to the reaction mixture did not change the pH significantly. The reactants were heated in a closed hood. After heating, the samples were taken and immediately cooled in iced water. Then, 1.0 mL of the reaction solution was mixed with 1 mL of (*o*-phenylenediamine) OPD solution (1 mg/mL in 1 M PBS of pH 7.4) and then reacted at 25 °C for 12 h in the dark. The remainder of the sample was stored at 4 °C for further analyses.

2.3 Analysis of PhIP, creatinine, and aldol condensation product

For HPLC-MS analysis of PhIP, creatinine, and the aldol condensation product, 10 µL of the reaction mixtures were injected into an HPLC-DAD-MS system, which consisted of a 5 µm Waters XBridge™ Shield RP18 250 × 4.6 mm column (Waters Corp., Milford, MA), a Waters 600 pump, a Waters 2707 autosampler, and a Waters 2998 diode array detector, connected to a LCQ-Fleet ion-trap mass spectrometer (Thermo Fisher Scientific, Waltham, MA) using an electrospray ionisation interface in positive ionisation mode (ESI+). Mobile phase **A** was composed of water/acetic acid (1000/1, v/v) at pH 3.6 (adjusted with 35% of ammonium hydroxide), and mobile phase **B** was composed of methanol. The gradient program was as follows: 5% **B**, 0 min; 5%–15% **B**, 0–15 min; 25%–35% **B**, 15–25 min; 35%–45% **B**, 25–30 min; 45%–100% **B**, 30–31 min; 100% **B**, 31–40 min; and 5% **B**, 41–50 min. The mobile phase was delivered at 0.5 mL/min in isocratic mode. The ESI-MS operating conditions included an electrospray voltage of 5.0 kV and nitrogen sheath and auxiliary gases set to 35 and 15 (arbitrary units), respectively. The temperature of the heated capillary was set to 275 °C. The 225.0→210.1 transition for PhIP was used for quantification purposes in this study. The

regression equation for PhIP standard was $y = 36.986x + 106.7$; $r^2 = 0.9994$; where x = PhIP concentration, $\mu\text{g/L}$; y = peak area of PhIP. The retention time of PhIP was about 26.5 min. The 114 $\rightarrow$ 86 transition for creatinine was used for quantification purposes in this study. The regression equation for creatinine standard was $y = 18.225x + 5984.4$; $r^2 = 0.9968$; where x = creatinine concentration, $\mu\text{g/L}$; y = peak area of creatinine. The retention time of creatinine was about 3.36 min.

According to Zöchling and Murkovic (2002), the first steps of the reaction involve formation of the aldol condensation product between creatinine and phenylacetaldehyde. The aldol condensation product is a substance with an expected molecular mass of 215 dominating the chromatogram. For data acquisition, the mass spectrometer was operated over the mass range of m/z 50–500 and for special identification the SIM (selected ion monitoring) mode was used (m/z 216) for the aldol condensation product of phenylacetaldehyde and creatinine in positive ionisation mode (ESI+).

2.4 HPLC analysis of benzimidazole formed from phenylacetaldehyde with *o*-phenylenediamine

Given that phenylacetaldehyde cannot be measured directly, it was measured using the corresponding benzimidazole 2-PB (Figure 1). After derivatisation with OPD, the samples were filtered through 0.45- μm Millex-HNnylon filters (Millipore, Billerica, MA). The percolates were then analysed by an HPLC-DAD system consisting of a 5 μm Waters XBridgeTM Shield RP18 250 $\times$ 4.6 mm column, a Waters 600 pump, and a Waters 2998 diode array detector. Approximately 1.0 mL of reaction solution was mixed with 1 mL of OPD solution (0.5 mg/mL) and reacted at 25 $^{\circ}\text{C}$ for 12 h in the dark. The injection volume was 10 μL (sterilised through 0.45- μm filters). The mobile phase was

5% methanol, 0–5 min, and the linear gradient program was as follows: 5%–30% methanol from 5 min to 10 min, 30%–40% methanol from 10 min to 45 min, and 100% methanol from 45 min to 65 min at a flow rate of 1 mL/min. The column temperature was set to 25 °C. Spectral data from all peaks were accumulated in the range 200–600 nm, and chromatograms were recorded at 280 nm. The retention time of 2-PB was about 42 min.

3. Results

3.1 Changes of phenylacetaldehyde in model reaction

In this study, the sample was derivatised using OPD before HPLC analysis. Table 1 shows the contents of phenylacetaldehyde in creatinine, phenylacetaldehyde, and some cation reaction mixtures. Addition of K^+ and Na^+ did not significantly influence the residual phenylacetaldehyde. By contrast, addition of Ca^{2+} and Mg^{2+} decreased the residual phenylacetaldehyde content. Interestingly, addition of Fe^{2+} and Fe^{3+} increased the residual phenylacetaldehyde content, and the effect of Fe^{3+} was more pronounced than that of Fe^{2+} .

3.2 Changes of creatinine in model reaction

Table 2 shows residual creatinine contents when creatinine, phenylacetaldehyde, and some cations were heated at 200 °C for 2 h. Addition of K^+ and Na^+ did not significantly influence the residual creatinine. By contrast, addition of Ca^{2+} , Mg^{2+} , Fe^{2+} , and Fe^{3+} decreased residual creatinine contents, and the effect of Fe^{3+} was more pronounced than that of Fe^{2+} .

3.3 Identification of the aldol condensation product

Identification of the aldol condensation product in a model system was conducted by comparing molecular masses and UV spectra with the values reported in a previous paper (Zöchling & Murkovic, 2002). The MS (SIM) and DAD chromatograms of the reaction product of creatinine and phenylacetaldehyde in the model system are shown in Figure 2. The results corresponded to the previously reported data (Zöchling & Murkovic, 2002).

3.4 Changes in the aldol condensation product in the reaction model

Certain amounts of the aldol condensation product were formed after heating the creatinine and phenylacetaldehyde mixture without including any cation in the reaction mixture. Addition of K^+ and Na^+ did not significantly influence the aldol condensation product. By contrast, the amounts of aldol condensation product formed during the reaction significantly decreased with increasing amounts of Ca^{2+} and Mg^{2+} in the reaction mixture. Increasing the amounts of Fe^{2+} and Fe^{3+} also decreased the amount of the aldol condensation product formed, and the inhibition of Fe^{3+} was stronger than that of Fe^{2+} in the reaction model (Table 3).

3.5 Changes of PhIP in model reaction

Table 3 shows the results of PhIP analysis in the model reaction when a solution mixture of creatinine, phenylacetaldehyde, and metal cations was heated at 200 °C for 2 h, and then added with 0, 5, 10, 20, 40, or 80 μ mol of metal cations. The amount of PhIP formation did not change significantly with increasing K^+ and Na^+ amounts. The amounts of PhIP formed during the reaction also increased with increasing amounts of Ca^{2+} and Mg^{2+} in the reaction mixture. PhIP content was significantly reduced with increasing Fe^{2+} and Fe^{3+} concentrations, and the effect of Fe^{3+} was more pronounced than that of Fe^{2+} .

4. Discussion

The present literature indicates that Fe^{3+} can induce oxidative decarboxylation of amino acids and formation of Strecker aldehydes because of its high oxidation potential (Bodiga, Eda, & Bodiga, 2014; Nashalian & Yaylayan, 2014; Zhuang, Pang, Zhang, Wu, Zhao, Yang, et al., 2012). Fe^{3+} can form complexes with various properties with carbonyl and promote further oxidative degradation of these complexes (Nashalian & Yaylayan, 2014). The amount of residual creatinine decreased with increasing Fe^{3+} , as shown in Table 2. By contrast, phenylacetaldehyde contents increased, as shown in Table 1. The amount of the aldol condensation product and PhIP formed also decreased. This phenomenon may be explained as follows: Fe^{3+} can form complexes of various properties with creatinine and accelerate creatinine degradation. This pathway can disturb the reaction with phenylacetaldehyde, influence aldol condensation product formation, and finally suppress PhIP formation. The results obtained in the present study show that the effects of Fe^{3+} and Fe^{2+} are very similar in inhibiting PhIP. Given that Fe^{2+} is oxidised to Fe^{3+} in the reaction, the effect of Fe^{3+} was stronger than that of Fe^{2+} in inhibiting PhIP in the reaction model.

Several studies have shown that CaCl_2 and MgCl_2 are very good catalysts for promoting formation of the aldol product and the aldol condensation product reactions with ammonia and formaldehyde (Asadullah, Kitamura, & Fujiwara, 2000; Gangadasu, Narender, ChinaRaju, & JayathirthaRao, 2006; Kossev, Koseva, & Troev, 2003; Miura, Nakagawa, & Hosomi, 2002). Thus, the amount of residual phenylacetaldehyde and creatinine decreased with increasing Ca^{2+} and Mg^{2+} contents in Tables 1 and 2, respectively. Meanwhile, because CaCl_2 and MgCl_2 promote PhIP formation of the aldol condensation product reactions with ammonia and formaldehyde, the amounts of the

aldol condensation product formed decreased with increasing amounts of Ca^{2+} and Mg^{2+} in the reaction mixture (Table 3). As a result, the amounts of PhIP formed during the reaction increased with increasing amounts of Ca^{2+} and Mg^{2+} in the reaction mixture (Table 3). Based on previous studies and the present results, the mechanism of some cations on the formation of PhIP in a model system is proposed in Figure 3. The mechanism contributes significantly to our understanding of the PhIP-inhibitory activity of metal cations, providing useful information for further studies in the field of food chemistry and toxicology.

5. Conclusion

Minimisation of the PhIP formed during thermal processing of food is important from the viewpoint of food safety. Metal ions play an important role in the human diet. This study found that Ca^{2+} and Mg^{2+} addition to the model system increased the formation of PhIP, whereas Fe^{3+} and Fe^{2+} addition decreased the formation of PhIP. This study also explained the mechanism of metal cations during PhIP formation in a model system. Drinking water contains various metal ions, and the hardness of water in different places varies despite the assurance offered by numerous regulatory agencies that the domestic drinking water in China is safe. The national standard hardness of drinking water in China is below $4.5 \mu\text{mol/mL}$ of CaCO_3 . The concentration range of cations is $0\text{--}8 \mu\text{mol/mL}$ in this study. So this study provides a theoretical basis on whether the use of water with different hardness levels in cooking food can reduce the formation of PhIP. It also provides answers on whether to use the iron wok in cooking.

Acknowledgements

All authors acknowledge National Science Foundation of China (GrantNo. 31271889),

science and Technology Planning Project of Guangdong Province of China (No.2014B020205001), the ministry of science and technology in agriculture science and technology Achievements Transformation Fund Project (No.2013GB23600669), science and Technology Planning Project of Guang Zhou of China (No.2013J4500036).

References

Asadullah, M., Kitamura, T., & Fujiwara, Y. (2000). Transformation of propane and CO to carboxylic acid and ester by highly active CaCl_2 catalyst. *Catalysis letters*, 69(1-2), 37-41.

Bodiga, V. L., Eda, S. R., & Bodiga, S. (2014). Advanced glycation end products: role in pathology of diabetic cardiomyopathy. *Heart Fail Rev*, 19(1), 49-63.

Cheng, K. W., Chen, F., & Wang, M. (2006). Heterocyclic amines: chemistry and health. *Mol Nutr Food Res*, 50(12), 1150-1170.

Damašius, J., Venskutonis, P. R., Ferracane, R., & Fogliano, V. (2011). Assessment of the influence of some spice extracts on the formation of heterocyclic amines in meat. *Food Chem*, 126(1), 149-156.

Gangadasu, B., Narender, P., ChinaRaju, B., & JayathirthaRao, V. (2006). Calcium chloride catalyzed three component, one-pot condensation reaction: An efficient synthesis of 3, 4-dihydropyrimidin-2 (1H)-ones. *Indian journal of chemistry*.

Gibis, M., & Weiss, J. (2012). Antioxidant capacity and inhibitory effect of grape seed and rosemary extract in marinades on the formation of heterocyclic amines in fried beef patties. *Food Chem*, 134(2), 766-774.

Gökmen, V., & Şenyuva, H. Z. (2006). Effects of some cations on the formation of acrylamide and furfurals in glucose–asparagine model system. *European Food*

226 *Research and Technology*, 225(5-6), 815-820.

227 Gökmen, V., & Şenyuva, H. Z. (2007). Acrylamide formation is prevented by divalent
 228 cations during the Maillard reaction. *Food Chem*, 103(1), 196-203.

229 Kossev, K., Koseva, N., & Troev, K. (2003). Calcium chloride as co-catalyst of onium
 230 halides in the cycloaddition of carbon dioxide to oxiranes. *Journal of Molecular*
 231 *Catalysis A: Chemical*, 194(1-2), 29-37.

232 Miura, K., Nakagawa, T., & Hosomi, A. (2002). Lewis base-promoted aldol reaction of
 233 dimethylsilyl enolates in aqueous dimethylformamide: Use of calcium chloride as
 234 a Lewis base catalyst. *Journal of the American Chemical Society*, 124(4),
 235 536-537.

236 Moon, S.-E., & Shin, H.-S. (2013). Inhibition of mutagenic
 237 2-amino-1-methyl-6-phenylimidazo [4, 5-b] pyridine (PhIP) formation using
 238 various food ingredients in a model systems. *Food Science and Biotechnology*,
 239 22(2), 323-329.

240 Nakai, Y., & Nonomura, N. (2013). Inflammation and prostate carcinogenesis.
 241 *International Journal of Urology*, 20(2), 150-160.

242 Nashalian, O., & Yaylayan, V. A. (2014). Thermally induced oxidative decarboxylation
 243 of copper complexes of amino acids and formation of strecker aldehyde. *J Agric*
 244 *Food Chem*, 62(33), 8518-8523.

245 Turesky, R. J., & Le Marchand, L. (2011). Metabolism and biomarkers of heterocyclic
 246 aromatic amines in molecular epidemiology studies: lessons learned from
 247 aromatic amines. *Chem Res Toxicol*, 24(8), 1169-1214.

248 Warzecha, L., Janoszka, B., Błaszczuk, U., Stróżyk, M., Bodzek, D., & Dobosz, C.

249 (2004). Determination of heterocyclic aromatic amines (HAs) content in samples
 250 of household-prepared meat dishes. *Journal of chromatography B*, 802(1),
 251 95-106.

252 Yu, D., Chen, M.-S., & Yu, S.-J. (2016). Effect of sugarcane molasses extract on the
 253 formation of 2-amino-1-methyl-6-phenylimidazo [4, 5-b] pyridine (PhIP) in a
 254 model system. *Food Chem*, 197, 924-929.

255 Zamora, R., Alcón, E., & Hidalgo, F. J. (2013). Effect of amino acids on the formation of
 256 2-amino-1-methyl-6-phenylimidazo [4, 5-b] pyridine (PhIP) in
 257 creatinine/phenylalanine and creatinine/phenylalanine/4-oxo-2-nonenal reaction
 258 mixtures. *Food Chem*, 141(4), 4240-4245.

259 Zamora, R., Alcón, E., & Hidalgo, F. J. (2014). Ammonia and formaldehyde participate
 260 in the formation of 2-amino-1-methyl-6-phenylimidazo[4,5-b]pyridine (PhIP) in
 261 addition to creati(ni)ne and phenylacetaldehyde. *Food Chem*, 155(0), 74-80.

262 Zhuang, X., Pang, X., Zhang, W., Wu, W., Zhao, J., Yang, H., & Qu, W. (2012). Effects
 263 of zinc and manganese on advanced glycation end products (AGEs) formation and
 264 AGEs-mediated endothelial cell dysfunction. *Life Sci*, 90(3-4), 131-139.

265 Zöchling, S., & Murkovic, M. (2002). Formation of the heterocyclic aromatic amine PhIP:
 266 identification of precursors and intermediates. *Food Chem*, 79(1), 125-134.

267 Zur Hausen, H. (2012). Red meat consumption and cancer: reasons to suspect
 268 involvement of bovine infectious factors in colorectal cancer. *International*
 269 *Journal of Cancer*, 130(11), 2475-2483.

270

270 Figure Legends:

271 **Figure 1.** Chemical structure of the corresponding benzimidazole of
272 phenylacetaldehyde obtained by derivatisation with OPD.

273 **Figure 2.** A, UV spectrum of the aldol condensation product in the model system; B,
274 DAD chromatogram of the reaction product of creatinine and phenylacetaldehyde; C,
275 MS chromatograms (SIM) showing the relative abundance of the selected analytes
276 (m/z 216) in the model system. Samples were heated at 200 °C for 120 min.

277 **Figure 3.** Proposed mechanism of some cations during PhIP formation in a model
278 system.

Table 1. Contents of phenylacetaldehyde in a model system containing phenylacetaldehyde, creatinine, and some cations. Samples were heated at 200 °C for 120 min.

cation amount (μmol)	phenylacetaldehyde mean $\pm$ SD (relative abundance)					
	K^+	Na^+	Mg^{2+}	Ca^{2+}	Fe^{2+}	Fe^{3+}
0	1094230 $\pm$ 56169 ^a	1094230 $\pm$ 56169 ^a	1094230 $\pm$ 56169 ^a	1094230 $\pm$ 56169 ^a	1094230 $\pm$ 56169 ^a	1094230 $\pm$ 56169 ^a
5	1084725 $\pm$ 386816 ^a	1074419 $\pm$ 125378 ^a	874607 $\pm$ 105269 ^b	435380 $\pm$ 307373 ^b	1402617 $\pm$ 252236 ^b	3015514 $\pm$ 114382 ^b
10	1031311 $\pm$ 53197 ^a	1084061 $\pm$ 211973 ^a	490066 $\pm$ 133403 ^c	170871 $\pm$ 119759 ^{b,c}	1861692 $\pm$ 99237 ^c	6981448 $\pm$ 196410 ^c
20	1059455 $\pm$ 169790 ^a	1175432 $\pm$ 104001 ^a	143171 $\pm$ 71132 ^d	60603 $\pm$ 41672 ^c	2662325 $\pm$ 131947 ^d	7946186 $\pm$ 234042 ^d
40	1090811 $\pm$ 100417 ^a	1161456 $\pm$ 215071 ^a	59839 $\pm$ 14551 ^e	7580 $\pm$ 1009 ^d	3343453 $\pm$ 96496 ^e	10400516 $\pm$ 874144 ^e
80	1081783 $\pm$ 337045 ^a	1075814 $\pm$ 202019 ^a	16747 $\pm$ 7529 ^f	6256 $\pm$ 742 ^e	4803449 $\pm$ 160756 ^f	12576443 $\pm$ 1433905 ^f

Means with different letters are significantly different $p < 0.05$.

Table 2. Contents of creatinine in a model system containing phenylacetaldehyde, creatinine, and some cations. Samples were heated at 200 °C for 120 min.

cation amount (μmol)	creatinine mean $\pm$ SD ($\mu\text{g/L}$)					
	K^+	Na^+	Mg^{2+}	Ca^{2+}	Fe^{2+}	Fe^{3+}
0	51542 $\pm$ 1308 ^a	51542 $\pm$ 1308 ^a	51542 $\pm$ 1308 ^a	51542 $\pm$ 1308 ^a	51542 $\pm$ 1308 ^a	51542 $\pm$ 1308 ^a
5	60350 $\pm$ 2129 ^b	55777 $\pm$ 628 ^b	32553 $\pm$ 776 ^b	46351 $\pm$ 1147 ^b	43737 $\pm$ 829 ^b	36636 $\pm$ 736 ^b
10	48766 $\pm$ 1234 ^c	54149 $\pm$ 1798 ^{a,b}	31483 $\pm$ 785 ^b	40310 $\pm$ 1838 ^c	37126 $\pm$ 385 ^c	21940 $\pm$ 790 ^c
20	51708 $\pm$ 630 ^a	56459 $\pm$ 885 ^b	15064 $\pm$ 274 ^c	29131 $\pm$ 1141 ^d	27950 $\pm$ 1371 ^d	21480 $\pm$ 1334 ^c
40	59197 $\pm$ 274 ^b	60568 $\pm$ 491 ^c	8874 $\pm$ 327 ^d	10105 $\pm$ 685 ^c	19169 $\pm$ 775 ^e	8152 $\pm$ 242 ^d
80	53704 $\pm$ 2355 ^a	50209 $\pm$ 759 ^a	6705 $\pm$ 228 ^e	8281 $\pm$ 480 ^f	11490 $\pm$ 987 ^f	4593 $\pm$ 91 ^e

Means with different letters are significantly different $p < 0.05$.

Table 3. Effect of some cations on the formation of the aldol condensation product and PhIP in a model system containing phenylacetaldehyde and creatinine. Samples were heated at 200 °C for 120 min.

cation amount (μmol)	aldol condensation product mean $\pm$ SD (relative abundance)					
	K^+	Na^+	Mg^{2+}	Ca^{2+}	Fe^{2+}	Fe^{3+}
0	1383817 $\pm$ 83929a	1383817 $\pm$ 83929a	1383817 $\pm$ 83929a	1383817 $\pm$ 83929a	1383817 $\pm$ 83929a	1383817 $\pm$ 83929a
5	1541864 $\pm$ 76748a,b	1315529 $\pm$ 97364a,b	1232183 $\pm$ 67839a,b	531596 $\pm$ 33784b	1006154 $\pm$ 87466b	566621 $\pm$ 17278b
10	1472886 $\pm$ 64778a,b	1212028 $\pm$ 87177a,b	1117545 $\pm$ 110938b,c	405443 $\pm$ 44029c	824141 $\pm$ 54747c	379666 $\pm$ 26698c
20	1517897 $\pm$ 54748a,b	1443396 $\pm$ 99827a,b	422209 $\pm$ 50297d	108895 $\pm$ 13409d	600029 $\pm$ 36828d	201576 $\pm$ 9029d
40	1246310 $\pm$ 89773a,c	1340741 $\pm$ 89871a,b,c	75597 $\pm$ 6932e	35806 $\pm$ 6847e	464263 $\pm$ 33647e	134263 $\pm$ 8889e
80	1441812 $\pm$ 98377a,b,c	1206419 $\pm$ 38177b,c	68559 $\pm$ 5874e	24800 $\pm$ 5847e	214244 $\pm$ 23788f	61092 $\pm$ 6289f

Means with different letters are significantly different $p < 0.05$.

Cation amount (μmol)	PhIP Mean $\pm$ SD ($\mu\text{g/L}$)					
	K^+	Na^+	Mg^{2+}	Ca^{2+}	Fe^{2+}	Fe^{3+}
0	634 $\pm$ 92.1 ^a	633 $\pm$ 92.1 ^a	633 $\pm$ 92.1 ^a	633 $\pm$ 92.1 ^a	633 $\pm$ 92.1 ^a	633 $\pm$ 92.1 ^a
5	607 $\pm$ 129 ^a	673 $\pm$ 181 ^a	1031 $\pm$ 98.3 ^b	1481 $\pm$ 104 ^b	484 $\pm$ 30.5 ^b	256 $\pm$ 44.0 ^b
10	659 $\pm$ 91.3 ^a	621 $\pm$ 121 ^a	1896 $\pm$ 104 ^c	2294 $\pm$ 61.7 ^c	413 $\pm$ 32.2 ^c	129 $\pm$ 12.4 ^c
20	569 $\pm$ 61.6 ^a	613 $\pm$ 84.0 ^a	2211 $\pm$ 76.4 ^d	2759 $\pm$ 102 ^d	317 $\pm$ 26.8 ^d	81 $\pm$ 10.0 ^d
40	571 $\pm$ 50.8 ^a	661 $\pm$ 129 ^a	2884 $\pm$ 36.4 ^e	3963 $\pm$ 51.3 ^e	232 $\pm$ 24.4 ^e	69 $\pm$ 8.9 ^d
80	613 $\pm$ 132 ^a	618 $\pm$ 159 ^a	2975 $\pm$ 62.9 ^e	4043 $\pm$ 159 ^e	204 $\pm$ 24.0 ^e	33 $\pm$ 3.0 ^e

Means with different letters are significantly different $p < 0.05$.

Figure 2

Figure 3