Genesis and evolution of regoliths: evidence from trace and major elements and Sr-Nd-Pb-U isotopes

C. Moragues-Quiroga^{1*,2}, J. Juilleret¹, L. Gourdol¹, E. Pelt², T. Perrone², A. Aubert², G. Morvan², F. Chabaux², A. Legout³, P. Stille², C. Hissler¹

¹Luxembourg Institute of Science and Technology (LIST), Catchment and Ecohydrology Research Group (CAT), 41 Rue du Brill L-4422 Belvaux-Luxembourg. cristina.moragues@list.lu, christophe.hissler@list.lu; ²Université de Strasbourg, LHyGeS (UMR 7517 CNRS/EOST/UdS), 1 Rue Blessig F-67084 Strasbourg Cedex-France. pstille@unistra.fr; ³INRA, UR 1138 Biogéochimie des Ecosystèmes Forestiers, Route d'Amance, 54280 Champenoux, Nancy-France. legout@nancy.inra.fr

Abstract

Regoliths encompass different materials from the fresh bedrock to the top of the organic horizons. The occurrence and evolution of these materials are determined by deposition, erosion and weathering processes that are specific for each region. The origin and interaction of the regolith layers are still important issues in the study of critical zone functioning. Studies that attempt to understand the processes responsible for the structure and the evolution of regoliths often focus either on the soil or on the bedrock compartment, and often from a narrowly focused approach using just a single tool such as major element patterns or mineralogy. This limits the understanding of the complete regolith composition and evolution. The present study proposes the combination of mineralogy, major and trace element concentrations, and Sr-Nd-Pb-U isotopes as an extremely useful procedure for tracking the

spatial and temporal origin of regolith materials and understanding their stratification. This multi-tracer approach was applied on a regolith profile from the south-western edge of the Rhenish Massif, which is characterised by the occurrence of slate bedrocks and overlying Pleistocene Periglacial Slope Deposits (PPSD). The data allowed the differentiation of three compartments within the regolith: 1) the organic compartment, where low $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{87}\text{Sr}/^{86}\text{Sr}$ isotopic ratios reflected the impact of atmosphere-derived components; 2) the PPSD compartment, whose upper horizons were characterised by the presence of atmosphere-derived particles rich in Cd, Sn, Sb, Hg and Pb and by relatively low $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{87}\text{Sr}/^{86}\text{Sr}$ ratios, whereas the lower horizons showed Nb enrichments in their matrix and a mineralogical composition pointing to ancient volcanic events; and 3) the weathered slate compartment, whose weathering progression and genetic relationship with the PPSD were particularly well tracked with $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{143}\text{Nd}/^{144}\text{Nd}$ isotope ratios, ($^{234}\text{U}/^{238}\text{U}$) activity ratios and positive Ce anomalies. The results prove the efficiency of radiogenic isotopes measurements combined with the analysis of major and trace elements to complete regolith studies. Moreover, these isotopes and elements are shown to be potential geochemical tracers of exchange processes at the water-mineral interface. This approach allows for the first time the description of an evolution scheme of a complete slate regolith, which covers a large part of the Rhenish Massif.

Keywords

Regolith; Sr-Nd-Pb-U Isotopes; Trace elements; Mineralogy; Critical zone; Periglacial coverbeds.

1. Introduction

Regolith represents the unconsolidated mantle of weathered rock and soil material on the Earth's surface (SSSA, 2008). In a broader sense, it encompasses all material from fresh rock to the atmosphere (Eggleton, 2001; Scott and Pain, 2008; NRC, 2001, Field et al., 2015). Regolith is a major compartment of the critical zone where fluxes of water, energy, solutes and matter occur. The production of regolith from the original bedrock influences the chemistry of surface waters and buffers the atmospheric CO₂ concentration (Banwart et al., 2011; Berner and Maasch, 1996; West et al., 2013). The regolith is the terrestrial environmental compartment where most of the water exchanges occur. Its bio-physico-chemical properties drastically impact the water that percolates and/or stores in its different parts (organic and mineral soil horizons, weathered bedrock, etc).

On a large scale from a space and time perspective, most of the *in situ* regolith systems are polygenetic. As an example, Felix-Henningsen (1994) showed that in the Rhenish Massif a weathering mantle representing the *in situ* regolith with a thickness up to 150 m was formed under warm and humid climates over a long period of time, from the Upper Mesozoic to the Tertiary. Additionally, loose materials produced in the *in situ* regolith move during erosion processes and contribute to form a *transported* regolith after redeposition. This means that actual *in situ* regoliths recorded successive weathering and erosion stages (Barbier, 2002). Therefore *in situ* regoliths can also be considered as polygenetic (Taylor and Eggleton, 2001), because they are the result of the accumulation of material from different origins. In the Rhenish Massif, the soil representing the upper part of the regolith is essentially developed from Pleistocene Periglacial Slope Deposits (PPSD - Kleber, 1997; Semmel and Terhorst, 2010) - also called periglacial coverbeds - that cover the weathered *in situ* regolith to constitute a polygenetic regolith system. These PPSD originate from the combination of

76 atmospheric deposition, solifluction and/or cryoturbation of the former active rock layer
77 during the last glacial period (Kleber, 1997; Semmel and Terhorst, 2010). The distances over
78 which these materials were transported range from a local to regional scale. Hence, the
79 alternation of such contrasting materials creates regolith components which may have a
80 proximal, but not direct, genetic link to the underlying bedrock. Their differentiation in the
81 regolith induced many lithic discontinuities that directly impact the evolution of the regolith
82 and control pedogenesis, water infiltration, interflow and root penetration (Lorz and Phillips,
83 2006; Lorz et al., 2011; Vökel et al., 2011).

84 Atmospheric deposition can significantly contribute to the polygenetic evolution of regoliths
85 and mask the autochthonous contribution coming from the bedrock. On the one hand,
86 atmosphere-derived anthropogenic depositions of trace metals originating from agricultural
87 tillage and fertilization, mining and other industrial activities tend to accumulate in the upper
88 soil layers due to their adsorption by organic matter (Aubert et al., 2002; Hissler and Probst,
89 2006; Steinmann and Stille, 1997). Stückrad et al. (2010) and Stille et al. (2011) suggested,
90 based on a combined trace element and Pb isotope study, a contribution of regional ore-vein
91 derived elements to a regolith from the south eastern edge of the Rhenish Massif and from the
92 Vosges Mountains in France, respectively. On the other hand, loess that was deposited during
93 the Pleistocene is widespread throughout Europe. These deposits form a more or less
94 continuous belt along a 2000 km east-west transect from Great Britain and Brittany in
95 northern France to the Dnieper Valley in Ukraine (Catt et al., 1986 ; Paepe and Sommé, 1970;
96 Rousseau et al., 2013, 2014). The origin of the European loess is still a matter of discussion.
97 Nevertheless, Sr and Pb isotopic data clearly demonstrate that the sources of the loess
98 deposits are proximal and different for each region (Rousseau et al., 2014). Some of these
99 aeolian deposits present a typical volcanic mineralogical contribution (Kleber and Terhorst,
100 2013; Pissart, 1995 Semmel and Terhorst, 2010). Impacts of different Pleistocene volcanic

eruptions were identified in the upper layers of the western European regoliths and can serve as efficient chronostratigraphic markers of these systems (Wörner and Schmincke, 1984; Pouclet and Juvigne, 2009; Pouclet et al., 2008). The studies of Gallet et al. (1998) and Chauvel et al. (2014) confirm an earlier conclusion reached by Taylor et al. (1983), that most of these aeolian deposits reflect the chemical composition of the upper continental crust. The matter and energy exchanges in the critical zone are partly controlled by the structure and evolution of the regoliths; therefore it is especially important to look at the entirety of the regolith when more than one formation occurs in the same profile. Relevant tools are required to improve the understanding of complex critical zone processes and to investigate this hidden part of the ecosystems. Over the last decades, radiogenic isotope, trace element and mineralogical analyses have become state-of-the-art tools for the characterisation of deposit and regolith formations (Debajyoti et al., 1967; Faure, 1986; Michard et al., 1985; Taylor and McLennan, 1981). Sr, Nd, Pb and U isotopes together with trace and rare earth element (REE) distribution patterns have been shown to be very suitable tools to answer open questions about regoliths formation (Aubert et al., 2001; Dequincey et al., 2002; Hissler et al., 2015; Stille et al., 2011, 2009). However, these techniques are rarely applied concurrently. In the present study, we combine mineralogical, major and trace element and Sr-Nd-Pb-U isotope analyses in order to reach a more comprehensive characterisation of a regolith profile. Our objective is to distinguish the different regolith strata by assessing their origin and evolution, and by evaluating the chemical and isotopical impact between the different strata. Hereby, we also address the question, how far dust from the late-Pleistocene volcanic eruption reached the south-western edge of the Rhenish Massif and mixed with the local loess deposits. To our knowledge, this is one of the few existing studies on a whole regolith system using such a multi-tracing approach.

2. Geomorphology and geological settings

The study site is a 4 ha beech (*Fagus sylvatica* L.) and oak (*Quercus petraea* (Matt.) Liebl.) dominated plateau located at an average altitude of 512 m asl in the Luxembourg Ardennes Massif (Oesling), close to the Belgian border (latitude: 49° 50' 05.5"N; longitude: 05° 47' 47.6"E) (Figure 1). The studied plateau is assumed to be representative of the regolith landform unit called “haute surface de l’Oesling” in Luxembourg, which developed at 500 m asl (Désiré-Marchand, 1985). A regolith landform unit represents an area characterised by similar landform and regolith attributes (Eggerton, 2001). Indeed, landforms are used as surrogates for mapping regoliths since both are usually spatially and genetically related (Craig et al., 1999). This “haute surface de l’Oesling” is the Luxembourg part of an extensive Danian-Montian surface of the eastern part of central Ardennes and Eifel at altitudes above 500 m (Demoulin, 2003) and correlates in Germany with the mapped “S2” surface, developed by pedimentation under semi-arid climate during the Upper Eocene/Lower Oligocene (Huguet, 1996, 1998). Désiré-Marchand (1985) explained that this “haute surface” was generated by gradual redevelopment of the original flat surface (S1 surface) and, consequently, the above *in situ* regolith recorded many weathering stages during the geological times and can be considered polygenetic.

The geological substratum of the studied plateau is composed of Devonian metamorphic schists, phyllites and slates (Juilleret et al., 2011) covered by Pleistocene Periglacial Slope Deposits (PPSD), which have been recognized to be the parent material of the soil in the Rhenish Massif (Müller, 1954; Deckers, 1996; Kwaad & Mùcher, 1977; Sauer, 2002; Sauer and Felix-Henningsen, 2006). PPSD generally consist of up to 3 different layers, with great variability in the occurrence and thickness of the layers depending on the geographical and topographical location (AG Boden, 2005; Dietze and Kleber, 2010; Semmel and Terhorst, 2010), and can be described as follows:

– an upper layer (UL) found in every topographic position. This layer always contains admixed loess and presents a homogenous thickness of 30 to 70 cm. This layer is characterised by higher silt content in comparison to the underlying soil.

– an intermediate layer (IL) which contains rock fragments originating from the bedrock below and also loess, as its presence is strongly controlled by topography and bound to sites especially prone to loess deposition.

– a basal layer (BL) composed almost exclusively of bedrock fragments present in the surrounding slopes. During the formation of the basal layer, bedrock and periglacial debris formed the surface before the onset of loess accumulation. The BL is therefore almost free of allochthonous material like loess and varies greatly in thickness. Rock fragments are usually oriented parallel to the slope and may appear in multiple layers.

According to Dietze and Kleber (2010), the above described layers form a continuous drape in Central European subdued mountains like the Rhenish Massif and may occur as a complete series of 3 layers (UL-IL-BL) or may be reduced to only upper and basal layers (UL-BL). The latter sequence (UL-BL) is the most widespread (Sauer and Henningsen, 2006).

3. Methodological approach

3.1. Regolith stratigraphy and classification

Prior to the sampling, the studied regolith profile was observed and described (i) from a soil pit for the upper part (0 to 140 cm depth); and (ii) by core drilling for the deeper part (140 to 735 cm depth). The following description was made according to the Guidelines for Soil Description (FAO, 2006) and completed with the description of the *subsolum* material according to Juilleret et al (2016). As regolith encompasses in its upper parts the *solum* (where pedogenic processes and biota are dominant) and in its lower parts the *subsolum* (where the original rock structure or fabric of the Bedrock is preserved), we described and

subdivided the profile according to *solum* and *subsolum* part following the methodology of Juilleret et al (2016) (Table 1).

The *solum* (from -1 to 45 cm) showed 3 horizons described as follows (Table 1): a very thin layer of fresh to highly decomposed organic material (O horizon) on top of a thin very dark greyish brown (10YR 3/2) silty clay Ah horizon and a brownish yellow (10 YR 6/6) silty clay loam B Cambic horizon. Both organo-mineral horizon were not sticky, non-plastic and showed a soft dry consistency. We estimated fine and very fine root density on vertical plane at 8 to 16 roots cm⁻² for Ah and B horizons. No earthworm channels were observed. Rock fragments of slate nature and of fine-to-medium gravel size (2 to 20 mm) were common in the Ah and B horizons with respective volume contributions of 13% and 27%.

The upper *Subsolum* (45-140 cm) observed from the soil pit (Table 1) is represented by: an olive yellow (2.5 Y 6/8) loam 2Cg1 horizon, and an olive yellow (2.5 Y 6/6) sandy loam 2Cg2 horizon. The estimated root density on the vertical plane was below 8 roots cm⁻² in both C layers. The stone content increases significantly in these two horizons and ranges from 40% to 50% of the volume for the respective 2Cg1 and 2Cg2 horizons. Common faint mottling of medium size and a few soft iron-manganese-oxide traces in the earth material around rock fragments were observed. This upper *subsolum* is defined as *Regolithic* layer (Juilleret et al., 2016).

The colours of the deeper *subsolum* material (140-735 cm) ranged from light grey (2.5 Y 7/2) to brownish yellow (10YR 6/6). *Gleyic* properties were observed between 340 and 380 cm depth (WRB, IUSS Working Group 2015). As the material was mechanically weakened by water, we classified it as *saprolithic* (Juilleret et al., 2016). From 450 to 735 cm depth, the material consisted of longer intact slate rock cores of several centimetres length (5 to 7 cm) mainly of grey colour (GLEY 1 5/N). Some of the longer cores showed cracks covered by sesquioxides interpreted as a mechanical weakened part inherited from the cleavage planes

where waters flow. This deeper material is classified as *paralithic* according to Juilleret et al. (2016).

The abrupt increase and the change in size of rock fragments between B and 2Cg1 horizons allowed us to diagnose a lithic discontinuity. The irregular orientation of the longer axes of slate rock fragments inherited from the underlying geological substratum in the 2Cg1 and 2Cg2 layers were interpreted as indications of cryoturbation. Below the 2Cg2 layer, the observation of the cores from the deeper *subsolum* showed that rock fragments have a different orientation, mainly vertical. The orientation of the slate rock fragments is inherited from the almost vertical cleavage planes of the geological substratum (Juilleret et al., 2011). This observation allows us to diagnose a lithic discontinuity between the 2Cg2 horizon and the geological substratum 3CR layer below.

According to the PPSD classification, we concluded that Ah and B horizons developed in the UL. Indeed, Ah and B horizons present a higher silt content with a cumulate thickness of 45 cm, while the underlying 2Cg1 and 2Cg2 are characterised by a dominant amount of rock fragments originated from cryoturbation. Consequently, the studied profile presents a *transported* regolith part made of PPSD on top of the *in situ* slate weathered substratum and is organized as follows (Table 1):

- 1) An organic topsoil, which encompasses the O horizon and constitutes the upper part of the *solum*.
- 2) An organo-mineral compartment developed in the PPSD which can be subdivided in *solum* and upper *subsolum*. The *solum* part is composed of Ah and B horizons. The upper *subsolum* can be divided into 2Cg1 and 2Cg2 horizons and contains *regolithic* material according to Juilleret et al. (2016).

3) A lower mineral *subsolum*, which contains *saprolithic* and *paralithic* materials according to Juilleret et al. (2016). This third compartment represents the *in situ* regolith profile made of weathered slate substratum and referred to as 3CR and 3R layers. These layers are separated from the above PPSD compartment by a lithic discontinuity.

The studied regolith profile can be classified as Haplic HEMIMODER (Jabiol et al., 2013) developing on a Dystric Cambisol (Ruptic, Endoskeletal, Siltic, Protosodic) (IUSS Working Group, 2015) overlying a Regolithic Saprolite (Gleyic, Ruptic, Rootic, Siltic, Skeletic) [Slatic] (Juilleret et al., 2016). In order to simplify the presentation of the results and the discussions, the name of the samples of the regolith layers (Table 1 – OH, PPSD, SP) will be used in the following document as reference of the regolith compartments.

We fulfilled the same pedo-stratigraphic report for four other profiles in the same plateau area. We searched for a profile that minimized the effect of erosion of the plateau unit (“haute surface de l’Oesling”) and permitted thicker layers to be investigated. Given the homogeneity we observed *in situ* and in later chemical analyses, this extensive sampling allowed us to choose the studied profile as a representative example of the geological system, on which we therefore focus this work.

The profile description is considered as a general framework that, although very informative, does not allow by itself the distinction of the origin of the various contributions to the regolith composition, for which geochemical analyses are needed. Indeed, the pedo-stratigraphic study informs about the physical properties of the material and its position within the regolith profile. But, this does not yield any precise information about in how far the *in situ* soil has been modified by external contributions such as aeolian depositions (loess or volcanic dust).

In addition, only geochemical and isotopic investigations allow u to understand the chemical evolution of the different parts of the regolith.

3.2. Samples collection and preparation

The sampling strategy was based on the description of the entire regolith presented above. We collected samples from the regolith profile including organic, organo-mineral and mineral material in three different steps from the surface to the deeper layers. First, we manually collected the highly decomposed organic material of the OH topsoil horizon on a 4 m² area around the opened soil pit in order to obtain a sufficient volume for further analyses. Second, we performed a soil sampling from the soil pit according to the ISO 10381 guidelines for soil sampling. Special attention was given to collect a representative amount of material for each horizon. According to the stone content, we collected 5 to 10 kg of material for each layer and homogenized it prior to taking representative aliquots for the laboratory analyses. Third, we sampled 5 layers from the core drilling from 140 cm depth (weathered slate) to 735 cm depth (fresh slate bedrock). We encountered some difficulties in obtaining continuous cores as the technique used requires water flow to remove cuttings (Gabrielli and McDonnell, 2012). The material extracted from 140 to 450 cm was made of pieces of rock fragments with muddy matrix. All the samples were stored in plastic bags and air-dried in the laboratory before being prepared for analyses.

We sieved the organic topsoil sample to 1 mm in order to remove the coarse vegetal debris that could remain after the sampling of the OH horizon. Then we fully ground it to 63 µm in a Fritsch centrifugal ball mill Pulverisette 6 in order to homogenise potential semi-decomposed organic particles. We sieved the *solum* and upper *subsolum* first at 2 mm in order to separate the coarse elements (>2 mm fraction), composed of slate rock fragments, and the fine earth (<2 mm fraction), called PPSD matrix in the manuscript. A representative aliquot of coarse

elements (PPSDce) was separated and ground to 63 μm . Then, the fine earth was sieved at 63 μm for the mineralogical and chemical characterization of the PPSD matrix. The material collected from the core drillings (lower *subsolum* samples - SP samples) was ground to 63 μm in a Fritsch centrifugal ball mill Pulverisette 6.

3.3 Analytical methods

We analysed aliquots of all regolith powdered samples for mineralogical contents by X-Ray Diffractometry (XRD, diffractometer RX Bruker, D5000) and Scanning Electron Microscopy (SEM, TESCAN Vega II fitted with EDAX X-ray energy dispersive spectrometer). We used Inductively Coupled Plasma – Optical Emission Spectrometry (ICP-OES, Thermo Fischer ICap 6500) and Inductively Coupled Plasma – Mass Spectrometry (ICP-MS, Thermo Elemental X7/Perkin-Elmer[®] DRC-e) for determining major and trace element concentrations respectively for all samples after digestion by LiBO_2 alkaline fusion. The analytical errors were < 5% for both instruments.

Regolith sample aliquots used for isotopic composition analyses were preliminary digested using $\text{HNO}_3\text{:HF:HClO}_4$ concentrated acid mixture in Savillex[®] Teflon vessels. We separated Sr, Pb and Nd from other elements for radiogenic isotope analyses by extraction chromatography on Eichrom Sr Spec, TRU Spec and Ln Spec resins, respectively, following the procedures of Geagea et al. (2007) and Guéguen et al. (2012) adapted from Pin and Zalduegui (1997) and Deniel and Pin (2001). U was separated by chromatography on Biorad AG1x8 anionic resin following procedures developed at the LHyGeS (Strasbourg, France) (Chabaux et al., 1997; Dequincey et al., 2002; Granet et al., 2010; Pelt et al. 2008). We measured Sr and Nd isotopic ratios on a Thermo-Scientific Triton (TIMS) and Pb and U isotopic ratios on a Neptune Thermo-Scientific multicollector (MC-ICP-MS). Mass fractionation of Pb isotopic ratios was corrected online using a SRM 997 Tl isotopic standard following the procedure of the laboratory (Stille et al., 2011). Mass fractionation and the

Faraday/SEM drift during U isotopic measurements by MC-ICPMS were corrected by bracketing against the IRMM-184 natural U standard. The 2014 mean for the Sr SRM 987 standard yields a $^{87}\text{Sr}/^{86}\text{Sr}$ ratio of 0.71025 ± 0.00002 (2SD, $n=21$) and for the La Jolla standard a $^{143}\text{Nd}/^{144}\text{Nd}$ ratio of 0.511842 ± 0.000006 (2SD, $N=7$). The Pb SRM 981 standard measurement during the session gave the following ratios: $^{208}\text{Pb}/^{206}\text{Pb} = 2.16604 \pm 0.00002$, $^{207}\text{Pb}/^{206}\text{Pb} = 0.914651 \pm 0.000007$, $^{206}\text{Pb}/^{204}\text{Pb} = 16.9267 \pm 0.0004$ (2SE), which are within the 2014 0.2‰ dispersion observed in the lab and consistent within less than 1‰ with the reference values of Doucelance and Manhès (2001). The HU1 uraninite was measured 5 times during the uranium MC-ICPMS session and yields a ($^{234}\text{U}/^{238}\text{U}$) activity ratio of 0.9983 ± 0.0004 (2SD) consistent with secular equilibrium within 2‰. The analysis of the ID-blanks suggests a contamination of less than 30 pg for Sr and Nd, ~400 pg for Pb and 23 pg for U which is negligible compared to the amount processed (>200 ng of Sr, Nd, Pb and U for all samples except for the humus sample with 20-30 ng U).

3.4. Statistical methods

In order to determine groups of major and trace elements that follow similar variation patterns in the regolith with depth, we performed a hierarchical cluster analysis (e.g. Hartigan, 1975; Kaufman and Rousseeuw, 1990, 2005). Such statistical technique has been widely used for drawing meaningful information from geochemical data (e.g. Bini et al., 2011; Levitan et al., 2015; Schot and van der Wal, 1992). For the linkage rule, we chose the Ward's method (Ward, 1963), which has been successfully used in many previous studies (Gourdol et al., 2013; Lin et al., 2014). For the linkage distance, the Pearson correlation distance ($1 - \text{Pearson correlation coefficient}$) was retained, which is suitable for clustering variables (Reimann et al., 2008). Prior to the analysis, the most universal z-transformation was applied to each parameter (mean subtraction and division by standard deviation) to ensure that each major

and trace element is weighted equally (Templ et al., 2008). The resulting dendrogram, illustrating the similarity of the parameters, was cut using two phenon lines to define groups and subgroups of parameters. Defining the number of groups by selecting the position of the phenon line up or down the dendrogram is a subjective evaluation step (Güler et al., 2002). The heights of these two phenon lines were retained by visual inspection of the dendrogram and the associated ordered z-transformed concentrations matrix and gave us the most satisfactory geochemical parameters groups and subgroups in terms of variations with depth.

4. Results

4.1. Mineralogical composition of the regolith components

The XRD analyses (Figure 2) indicate that PPSD3 and PPSD4 are enriched in phyllosilicates and clay minerals, mainly chamosite, a Fe-rich polytype of chlorite, and kaolinite. At these depths, chamosite and kaolinite show relative abundances ranging from 17 to 20%. Other chlorite polytypes are much less abundant (rel. abund. < 3%). The illite/mica group shows increasing abundances with depth in the matrix of the PPSD compartment between 7 and 28%. Similarly, one observes enrichments in orthoclase (rel. abund. 8%) in PPSD3 and PPSD4 and in albite (3%) at the PPSD2 and PPSD3 layers. Conversely, anorthite is depleted between PPSD2 and PPSD4 (rel. abund. < 3%), and quartz is depleted at the PPSD3 and PPSD4 layers (rel. abund. 16 to 24% respectively), both compared to the above *solum* horizons (PPSD1 and PPSD2) and the *subsolum* saprolithic material below (SP), where the relative abundances scatter between 8 and 11% for anorthite and 36 and 57% for quartz. Rock fragments from the PPSD (PPSDce) generally present a mineralogical composition rather similar to the SP compartment, being slightly more enriched in kaolinite (12%) and notably in orthoclase (14%).

SEM analyses on all PPSD samples indicate that in this compartment there are REE-bearing minerals such as monazite, xenotime, zircon and florencites. Other Ti-bearing and iron oxide trace minerals observed at these depths are rutile, ilmenite and Ti-magnetite.

4.2. Chemical composition of the regolith components

The chemical composition of the entire regolith is shown in Table 2. Na₂O and TiO₂ are enriched in the PPSD matrix between PPSD2 and PPSD4 compared to SP. K₂O, Fe₂O₃, MgO and Al₂O₃ tend to increase with depth within the matrix of the PPSD compartment and reach values similar to those of the underlying saprolite (SP samples). CaO concentrations are high at the OH horizon and decrease with depth in PPSD, showing an important depletion on top of the SP regolith compartment, where it slightly increases again towards the fresh bedrock. On the contrary, P₂O₅ is highly enriched at the top (PPSD1) and bottom (PPSD4) of the PPSD and in SP3 similarly to MnO and Fe₂O₃. PPSDce is, compared to the matrix between PPSD2 and PPSD4, depleted in TiO₂, K₂O and Al₂O₃. PPSDce is enriched in Fe₂O₃ with respect to the rest of the profile, with the exception of SP3. SiO₂, CaO, MgO and MnO show in PPSDce similar concentrations as in the SP1 compartment, whereas Na₂O and P₂O₅ PPSDce concentrations are rather close to the ones of the soil matrix at the PPSD bottom horizons.

The trace elements can be classified in three groups according to the vertical evolution of their concentrations within the regolith profile (Figure 3). Group I contains Cd, Sn, Sb, Hg and Pb. Their concentrations are higher in the top layer of the PPSD compartment and are low and almost constant throughout the rest of the regolith profile (SP). Group II is composed of Co, Ni, Cu, Zn and As, which present high concentrations in PPSD4 and are enriched in SP3. Group III includes trace elements having increasing concentrations with depth in the PPSD and a notable decrease in the concentrations between PPSD and SP. These elements can be divided in three subgroups according to their behaviour in the SP compartment: Group IIIa

375 encloses those presenting a concentration increase at 380 cm depth, similar to Mg and K (U,
 376 Pr, Nd, Gd, Sm, Eu); Group IIIb those that, similar to Al, Si and Ti, remain stable below 200
 377 cm depth (Cr, Rb, Sr, Y, Nb, Th, La Ce, Tb, Dy, Ho, Er, Tm, Yb, Lu) and Group IIIc those
 378 that continue decreasing after 200 cm depth, like Na (Zr, Hf).

379 Major and trace element concentrations of PPSD and SP samples are normalized to the deeper
 380 SP lithic material (SP5), which can be considered, at this study site, as the fresh slate (Figure
 381 4). Compared to this local reference, OH and PPSD1 are depleted in most of the elements,
 382 except the trace elements of group I. The other PPSD and SP samples are slightly enriched
 383 and show almost identical distribution patterns with ratios close to 1 for most of the analysed
 384 elements. However, the enrichments of the elements of Group II in SP3 and SP4, especially
 385 Mn, Co, As and U are different. Compared to the slate reference, Nb is generally slightly
 386 enriched in the PPSD compartment (ratio = 1.4), whereas the underlying SP samples present a
 387 ratio of 1.0. This is also not the case for the rock fragments of PPSD (PPSDce), whose Nb
 388 ratio is 0.77.

389 Post Archean Australian Shale (PAAS) normalized Rare Earth Element (REE) concentrations
 390 of PPSD and SP samples show similar distribution patterns (not shown). They display middle
 391 REE (MREE) enrichments, which are more notable in coarse materials (PPSDce), in the
 392 matrix of PPSD4 and in SP, especially at SP4, as indicated by the $(Eu/Yb)_N$ ratios (Figure 5).
 393 Similarly, though to a lesser extent, $(La/Yb)_N$ ratios point to a strong enrichment of light-
 394 REE (LREE) in PPSD4 and SP4, whereas a positive Ce anomaly ($Ce_N/Ce_N^*=1.05$) is only
 395 observable in SP3. Ce is an especially interesting element as it is one of the most reactive
 396 REE. Conversely to most of the other REE, it can pass from state 3+ to 4+ and precipitate as
 397 cerianite (CeO_2) in oxidizing conditions (Braun et al., 1990). Here, Ce anomalies are
 398 calculated as the enrichment of Ce with respect to other LREE (namely La and Pr) normalised

to Post Archean Australian Shales (PAAS) with the equation $[Ce_N/Ce_N^* = Ce_N/(0.5La_N + 0.5Pr_N)]$.

4.3. Pb, Sr, Nd and U isotopic compositions of the regolith components

All isotope data are given in Table 3. $^{87}Sr/^{86}Sr$ and $^{206}Pb/^{207}Pb$ ratios are low in OH and PPSD1 (0.7218 and 1.153 respectively) and increase with depth through PPSD2 and PPSD3 up to 0.7387 and 1.220, respectively (Figure 6a and b). $^{87}Sr/^{86}Sr$ ratio decreases in PPSD4 (0.7366), increases again in the saprolithic material of the SP and has the highest isotopic ratio in the lithic material at SP5 (0.7416). $^{206}Pb/^{207}Pb$ ratio still increases beyond PPSD4 and, after a remarkable decrease at the lithic discontinuity between PPSD and SP, increases again until the deeper part of the regolith (1.214), similar to $^{87}Sr/^{86}Sr$. The coarser materials of the PPSD (PPSDce) display $^{87}Sr/^{86}Sr$ ratios close to the values of the lower PPSD (0.73736) and high $^{206}Pb/^{207}Pb$ ratios compared to both PPSD and SP compartments. The $^{143}Nd/^{144}Nd$ ratios show rather small variations between 0.51192 and 0.51198 throughout the profile (Figure 6c). The ratio is slightly higher for the OH horizon (0.51195) compared to the other soil horizons (0.51192).

The $(^{234}U/^{238}U)$ activity ratios scatter between 0.947 and 0.960 along the profile with the exception of the 270 and 320 cm depth horizons. At these depths, SP2 and SP3 show $(^{234}U/^{238}U)$ activity ratios higher than 1 (1.05 and 1.04 respectively - Figure 5d).

5. Discussion

5.1. Major and trace element behaviour within the studied regolith system

The cluster analysis on the PPSD matrix and SP samples indicates that there are three groups of elements that have different origins and contrasted behaviours during weathering and pedogenetic processes (Figure 3). Even though the coarse elements of the PPSD are not

included in this statistical assessment, and thus uncertainty increases, the results allow discerning the chemical zonations according to the soil fraction that is responsible for most of the processes described below.

Group I includes elements such as Cd, Sn, Sb, Hg and Pb, which are only enriched in the surface Ah horizon (PPSD1) and are probably atmosphere-derived and of anthropogenic origin. The Ah horizon is located directly under the organic horizon (OH) and is as a consequence rich in organic matter, which is known to trap heavy metals and subsequently release them by leaching into the soil (Hissler and Probst, 2006; Steinmann and Stille, 1997; Stille et al., 2011, 2009). This is especially observable in the Pb isotope data (Figure 6). According to the work of Redon et al (2013), depth patterns of Sn, Sb, Hg and Pb present a disconnection between upper soil horizons and the rest of the regolith as well as the PPSD coarse materials (PPSDce; Table 2), supporting the hypothesis that they do not derive from the underlying saprolite but from the atmosphere. In contrast is the slight Cd enrichment at 320 cm depth (SP3) compared to the horizons underneath, which resembles that of the elements of Group II and thus cannot be attributed to anthropogenic deposition at the top of the regolith.

Group II comprises Co, Ni, Cu, Zn and As, which show similar distribution patterns to P_2O_5 , MnO and Fe_2O_3 with a strong enrichment at 320 cm depth (SP3) in the saprolithic material. At the same time, Group IIIa comprises U and Pr, Nd, Gd, Sm, Eu, whose depth patterns resemble those of MgO and K_2O with a strong enrichment at SP4. All of them also show also high concentrations in the lowermost horizon of the PPSD. These trace elements are sensitive to being mobilised during changing redox conditions in the regolith profile, e.g. groundwater table fluctuations. The fractionation and mobilisation/accumulation can be either direct by valence alteration into soluble/stable forms or due to the dissolution-precipitation of Fe and Mn-(oxy)hydroxides in which these trace elements get preferentially sorbed. Of particular

449 interest here is the slight but meaningful positive Ce anomaly that can be observed in SP3
 450 ($Ce_N/Ce_N^*=1.05$ compared to ratios between 0.92 and 0.98 for the other samples). During
 451 water saturation periods (winter), the anoxic conditions in the saprolite result in a reducing
 452 environment which favours the leaching of REE. In the formed solution, Ce, as the majority
 453 of the lanthanides, presents a trivalent state ($3+$). When the water table flows downwards
 454 (summer), water is mainly retained only at the smallest pores of the soil aggregates and in the
 455 clay fraction of the saprolithic material. Oxygen can get into the interfaces, favouring Ce^{3+} to
 456 Ce^{4+} oxidation. In these oxic conditions, Ce can precipitate as cerianite e.g. on the hydrated
 457 surface of manganese oxides (Braun et al., 1990; Steinmann and Stille, 1997). This means, in
 458 this case Ce might be mobilised from the upper horizons and immobilised at 320 cm depth. In
 459 agreement is the evolution of the U geochemistry within the studied profile (Figure 3 and 6,
 460 and Table 2). U concentrations tend to increase from 3.09 ppm at the top of the SP
 461 compartment to 5.84 ppm at SP4. Under oxidising conditions, Uranium can oxidise to the
 462 uranyl ion ($U(VI)O_2^{+2}$) and form compounds that are soluble in water. As with Ce, the uranyl
 463 ion can get released into the solution at SP1, migrate downwards and re-deposit as it co-
 464 precipitates with Fe oxy-hydroxides at the “summer” redox interface (SP3-SP4) (Duff et al.,
 465 2002; Bruno et al., 1995). In figure 6, the ($^{234}U/^{238}U$) activity ratios lower than 1 above SP2
 466 (270 cm) and below SP3 (320 cm) (<0.96) but higher than 1 in between (>1.04) indicate,
 467 respectively, a ^{234}U depletion and enrichment of the regolith. During water-rock interaction,
 468 ^{234}U and ^{238}U have different mobilities, with a preferential leaching of ^{234}U compared to ^{238}U
 469 due to the so-called alpha recoil process (more details in e.g., Chabaux et al., 2003; 2008;
 470 DePaolo et al., 2006; 2012). For a material old enough to be at secular equilibrium (i.e.,
 471 $^{234}U/^{238}U = 1$), such a process leads to a preferential enrichment of ^{234}U in waters compared to
 472 its parent ^{238}U , and hence to ($^{234}U/^{238}U$) activity ratios higher than 1 in the waters and lower
 473 than 1 in the residual materials (see also Schaffhauser et al., 2014; Pierret et al., 2014; Prunier

et al., 2015; and references therein). Thus, during long periods of water saturation and intense weathering of the rock, ^{234}U is leached from the minerals, giving ($^{234}\text{U}/^{238}\text{U}$) activity ratios below 1 in the saprolithic materials. This seems to be the case in the studied saprolithic material above SP2 and below SP3. In the studied regolith, the water table fluctuates between 320 cm depth during low hydrological conditions and 108 cm depth during winter saturation periods (according to field monitoring hydrological data, not shown). Thus, we hypothesise that ^{234}U is mobilised above 270 cm depth and potentially accumulated around the permanent water table level (SP3-SP4), where the redox potential allows the above-mentioned co-precipitation. Conversely, the permanently saturated layer between SP3 and SP5 (735 cm) favours the downward flow of ^{234}U , whereas below SP5 depth we are on the impermeable and slightly weathered slate bedrock. This hypothesis is supported by occurrences of gleyic properties with rusty patches observed from drill cores of the saprolithic material between 340 and 380 cm depth, which correspond to strong reduction processes and iron segregation (WRB, IUSS Working Group 2015).

In summary, the depth patterns of the Ce anomaly and ($^{234}\text{U}/^{238}\text{U}$) activity ratios together with the other redox sensitive elements, definitely point to the effect of the seasonal water table fluctuation between 108 and 380 cm depth. The regolith above SP1 is preserved from the weathering process and conserves element concentrations untouched. Whereas between SP1 and SP4, the historical exposure of the material to alternating oxic and anoxic conditions (redox conditions) favoured the dissolution-precipitation processes that control the dynamics of the redox sensitive and allied elements. Below SP4, most labile elements may be flushed away from the saprolite into deeper groundwater.

On the other hand, the behaviour of all elements from Group III in the matrix of the PPSD compartment is also related to the stability of some specific residual trace mineral phases. Zr and Hf reside in zircons, Th in monazite, Y in xenotime and Nb in Ti-bearing minerals.

Similarly, PPSD4, representing the lowermost part of PPSD matrix, is enriched in LREE compared to the upper horizons (Figures 3 and 5) pointing to an important presence of phosphate and LREE bearing trace minerals such as monazite and florencite. At last, the Eu positive anomaly of PPSD's coarse materials (PPSDce), surely relates to their greater feldspars abundance (Vázquez Ortega et al., 2015) (Figures 2 and 5).

Finally, vegetation life cycles (nutrition, evapotranspiration) might also have an important impact in the overall chemical composition. In this regolith, beech and oak have the greatest root density between 20 to 40 cm depth (up to 16 roots cm⁻²). The roots uptake of macronutrients and other oligoelements is probably a major cause of element fractionation between the upper and lower PPSD layers. In this sense, Ca must be accumulated in the OH horizon due to biological cycling and litter decomposition (Stille et al., 2009). Phosphate concentration is high in the Ah horizon for the same reason. Then it decreases at the level of the maximum root uptake at 20 to 40 cm depth (PPSD2-3). At these depths, also Mg, K, Fe, Al and most of the REE are depleted compared to the lower PPSD4 horizon. Further studies on the vegetation are needed to better understand its geochemical impact on the regolith studied here.

5.2. Impact of atmosphere-derived anthropogenic depositions on the PPSD

The ⁸⁷Sr/⁸⁶Sr and ²⁰⁶Pb/²⁰⁷Pb isotope ratios decrease with decreasing depth within the organic (OH) and organo-mineral (PPSD) compartments (Figure 6a and b and Table 3), whereas the corresponding ¹⁴³Nd/¹⁴⁴Nd ratios are only very weakly scattered, ranging between 0.51192 and 0.51195 (Figure 6c and Table 3). The comparatively lower Pb and Sr isotopic compositions of the organic compartment (OH sample) can be related to anthropogenic impacts (Stille et al., 2011). Indeed, the current local atmospheric baseline isotopic

524 compositions determined on lichens show significantly lower $^{206}\text{Pb}/^{207}\text{Pb}$ (1.162) and $^{87}\text{Sr}/^{86}\text{Sr}$
 525 (0.7152) but higher $^{143}\text{Nd}/^{144}\text{Nd}$ (0.51205) ratios (Hissler et al., 2008). In a $^{87}\text{Sr}/^{86}\text{Sr}$ vs Rb/Sr
 526 diagram the PPSD data show a mixing trend with the OH horizon and the Ah organo-mineral
 527 soil horizon (PPSD1) with the lowest isotopic composition values and Rb/Sr ratios similar to
 528 the lichen (0.87 and 1.54 respectively) (Figure 7a). Similar relationships are observable in the
 529 $^{206}\text{Pb}/^{207}\text{Pb}$ vs U/Pb diagram (Figure 7b). Again, OH and PPSD1 samples have Pb isotopic
 530 composition and U/Pb ratios similar to lichen (U/Pb = 0.01 for both samples). Therefore, the
 531 Rb-Sr and U-Pb isotope systems of the uppermost regolith samples are strongly influenced by
 532 recent atmosphere-derived inputs. This is not the case for the Sm-Nd isotope system, which is
 533 not to be affected by atmospheric deposition (not shown). If mixing occurred between
 534 anthropogenic, atmosphere-derived and soil particles then the regolith isotope data should
 535 describe a mixing hyperbola in the $^{87}\text{Sr}/^{86}\text{Sr}$ vs. $^{143}\text{Nd}/^{144}\text{Nd}$ diagram (Faure, 1986). This is
 536 indeed the case (Figure 8). The shape of the curve is controlled by the end-members Sr/Nd
 537 ratios. The atmospheric end member is the lichen (Sr : 1.17ppm, $^{87}\text{Sr}/^{86}\text{Sr}$: 0.7152 ; Nd :
 538 0.17ppm, $^{143}\text{Nd}/^{144}\text{Nd}$: 0.51205) (Hissler et al., 2008) and the geogenic end member
 539 corresponding to the lithic material representing fresh slate bedrock (SP5) (Sr : 98.2ppm,
 540 $^{87}\text{Sr}/^{86}\text{Sr}$: 0.7416 ; Nd : 37.9ppm, $^{143}\text{Nd}/^{144}\text{Nd}$: 0.51193). The resulting Sr/Nd ratios are 6.9
 541 for the lichen and 2.6 for the slate. One calculates that the uppermost OH and Ah horizons
 542 (PPSD1) contain up to 90% and 50% of recent atmosphere-derived Sr and 95% and 50% of
 543 recent atmosphere-derived Nd, respectively. Such enrichments are rather surprising,
 544 suggesting that these elements have been transported over distances greater than 20 km. The
 545 closest active steel industrial parks are situated around 25 km southeast (Bissen, Luxembourg)
 546 and 120 km north-west (Liège, Belgium) of the studied site, and a historic mine of mainly Pb
 547 and Zn was located around 20 km north of the study site. Alternatively, these enrichments

might be attributed to the heavy fighting nearby during the “Battle of the Bulge” in the vicinity of Bastogne during the Second World War (Cole, 1965).

5.3. Impact of volcanic events on the PPSD

In central Europe, the upper layer of PPSD (UL) often presents a typical volcanic mineralogical composition related to the Laacher See Eruption in the late Pleistocene (12900 years BP) (Kleber and Terhorst, 2013; Semmel and Terhorst, 2010; Pissart, 1995; Wörner and Schmincke, 1984). Wörner and Schmincke (1984) and Wörner et al. (1985) also highlighted the very specific trace element and isotopical composition of the Laacher See tephra, which permits its identification and differentiation from other material contributions. These mineralogical and isotopical fingerprints allow us to use the Laacher See deposits as a stratigraphic marker (Schmincke et al., 1999). Nevertheless, relatively older volcanic events took place in the same region, such as the Rocourt (74-90.3 ka) and Eltville (16-30 ka) eruptions, whose tephras spread in the south east of Belgium, central Germany and north of Luxembourg (Poucllet and Juvigne, 2009; Poucllet et al., 2008). Poucllet et al. (2008) pointed out the potential of the Rocourt tephra as chronostratigraphic marker for the Upper Pleistocene loess deposits in these areas thanks to its glass fragment shape, chemical and mineral composition, which differs from the Laacher See tephra.

One of the principal questions raised in this study concerns the stratigraphic and pedogenetic evolution of the periglacial coverbeds (PPSD compartment) and, more specifically, the origin of their constituting mineral phases. Are these phases only derived from the underlying slate or do some of them have other origins; e.g. originate from volcanic dust and ashes from the late-Pleistocene Laacher See eruption, and reached also the south-western edge of the Rhenish Massif. Indeed during solifluction processes the Laacher See tephra, which was already deposited, was admixed with other materials to become part of the upper coverbed

(Kleber, 1997; Schmincke et al., 1999; Semmel and Terhorst, 2010; Stückrad et al., 2010; Terhorst, 2007). Previous studies have shown that Laacher See particles have been incorporated to the upper PPSD layer in central Europe (Kleber & Terhorst 2013; Pissart, 1995; Semmel and Terhorst, 2010). At the same time, several tephra layers have been recognized in the Upper Pleistocene Loess deposits in Belgium, as the ones of the above mentioned Laacher See, Eltville and Rocourt (Pouclet & Juvigné, 2009). Among those tephra sequences, Wörner and Wright (1984) noticed that the Laacher See tephra has a phonolite-like mineral composition dominated by alkali feldspar, especially sanidine phenocrysts, followed by plagioclase (mainly albite), hyne, amphibole, clinopyroxene, sphene, Ti-magnetite, apatite, phlogopite and traces of zircon phenocrysts. The abundance of the different observed mineral phases of the studied regolith vary along the profile but manifest specific mineral enrichments in the matrix of the basal layer of PPSD (PPSD3 and PPSD4), which might be related to a Pleistocene eruption (Figure 2).

In that regard, of great importance is the finding of chamosite as a main mineral phase (up to 20 vol.%) in the matrix of the basal layer of the PPSD compartment. It is significantly less present in the saprolithic material (SP, 10-13 vol.%). This mineral phase is the Fe-rich end-member of the chlorite group. Its presence explains the rather high Fe₂O₃ contents (7-11 wt.%) in the basal layer of PPSD. Chamosite is typically a replacement mineral phase and alteration product of ferromagnesian minerals such as pyroxenes and amphiboles (Deer et al., 1974). However, also other mineral phases confirm the presence of tephra in the basal layer of PPSD, such as kaolinite, which results from feldspar alteration and reaches 20 vol % which represents a strong enrichment compared to the underlying saprolite (SP, 3-5 vol %). Primary feldspar minerals such as albite and orthoclase are also observable in smaller quantities in the PPSD. Orthoclase, respectively sanidine (a not distinguishable polymorph of orthoclase),

occurs only in PPSD (up to 8 vol.% in the matrix and 14 vol.% in the coarse materials) and is not at all observable in SP.

In a similar way, one might suggest that the titanium enrichments observed for the same PPSD samples (Figure 3) result from sphene and Ti-magnetite. Indeed, the important presence of Ti-bearing minerals such as ilmenite, rutile and Ti-magnetite observed under the scanning electron microscope (SEM) confirm the presence of tephra-derived mineral constituents in the PPSD.

The deposited tephra dust particles caused elevated Nb concentrations in the samples (Schmincke et al., 1999; Haase et al., 2007) because the erupted tephra (Lower Laacher See Tephra deposits, LLST) is enriched in Nb with concentrations of more than 200 ppm (Wörner and Schmincke, 1984). The Nb concentrations of the studied covered matrix are well correlated with TiO_2 (Table 2) indicating, as suggested (Bonjour and Dabard, 1991), that titaniferous mineral phases are the principal Nb carrying phases. Nb concentrations vary according to depth, being lowest close to the surface (10.28 ppm) (Figure 3 and Table 2). Between PPSD2 and PPSD4, the Nb concentrations range between 19.4 and 19.9 ppm and are higher than those of the underlying bedrock showing concentrations between 14 and 16 ppm. The Nb concentrations of coverbeds from two other sites of the studied plateau range, at same depths as PPSD2 to PPSD4, between 18.5 and 22.5 ppm (not shown). In the coarser materials (PPSDce) of these horizons and in the underlying bedrock, Nb concentrations decrease again to values below 11 and 16.6 ppm respectively. According to Taylor and McLennan (1985) the UCC Nb concentration is higher (25 ppm) and, therefore, the PPSD samples show Nb depletions similar to European loess studied by Gallet et al. (1998). These authors provided plausible arguments that the depletions are artefacts resulting from an over-estimation of the UCC Nb concentration. Similarly Condie (1993) suggest that the Nb concentrations of clastic sediments are seldom higher than 15 ppm. This value is in agreement with the Nb

concentrations found in the bedrock below the covered. Thus, the fact that the PPSD matrix shows slightly higher Nb concentrations than the PPSD coarse materials and the bedrock is suggested to be the result of deposition of Nb-enriched mineral phases from the late-Pleistocene Laacher See eruption.

The Nb contribution from this eruption to the loess component of the PPSD sediments can be estimated by using a two component mixing equation, a Nb concentration of the Laacher See tephra (100 ppm) (Wörner and Schmincke, 1984) for the volcanic mixing end-member and a Nb concentration of the underlying bedrock (15 ppm) for the PPSD end-member. This rough estimation indicates that the volcanic Nb contribution is less than 6% in the PPSD matrix. Mixing calculations can also be performed by using Nd concentrations and $^{143}\text{Nd}/^{144}\text{Nd}$ isotope ratios (see Faure 1986). However, the Sr-Nd isotopic and concentration similarities between slate bedrock and PPSD samples and, thus, the resulting mixing relationships (Figure 8) suggest that the PPSD below 45 cm depth do not contain visible amounts of tephra-derived Sr and Nd. Nevertheless, Nb enriched PPSD samples plot close to the crust-mantle mixing hyperbola (Figure 8), pointing to a certain impact of the volcanic materials in their signature. Thus, in the case of our study only Nb enrichments and mineralogical composition allow the identification of tephra contributions in the lowermost PPSD horizons. Given the fractionation of some minerals and trace elements within the PPSD, we propose that the discontinuities observed from PPSD2 and PPSD3 respond both to the impact of volcanic depositions and pedogenetic processes such as diagenesis and neoformation of minerals.

5.4. Impact of the saprolite (SP) on PPSD

With the exception of the two uppermost soil horizons (OH and Ah), no impact of anthropogenic Sr and Nd is observable for the rest of the regolith compartments. Similarly we have seen that neither Sr nor Nd isotope ratios are impacted by tephra depositions during

Pleistocene volcanic eruption events. PPSD and SP material are isotopically very similar pointing to similar origins. This is confirmed by the major element concentrations of the slate saprolithic material (SP) and the coverbeds (PPSD) below the OH horizon, which scatter in nearly the same range (Figure 3) and show similar depletions and enrichments compared to the reference slate layer of SP (Figure 4). Only very few trace elements such as Nb and some redox-sensitive elements manifest, as discussed above, differences between PPSD and underlying saprolithic material. Consequently, if the PPSD profile contains, as suggested for other neighboured coverbeds (Stückrad et al., 2010), pre-industrial atmosphere-derived particles, then they have not travelled long distances but originate from the slate of the region. In Figure 9, the ϵNd (T) values of the Devonian slates of our study site (“initial” epsilon Nd values, calculated back to the stratigraphic age) are plotted *versus* their stratigraphic age and compared with other European phanerozoic shales. This type of diagram was first used by Michard et al. (1985) in order to show that during major orogenic events sediments have been impacted by mantle-derived material such as volcanogenic detrital material. One observes that the slates in this study show initial ϵNd (T) values very similar to those of other Devonian shales collected in Brittany and Wales. These sediments have been deposited in the same oceanic environment in a period between the Hercynian and Caledonian orogenesis and are not visibly impacted by volcanogenic detritus during their deposition. Volcanogenic deposits containing significant quantities of mantle-derived material are characterised by high $^{143}\text{Nd}/^{144}\text{Nd}$ isotope ratios and low $^{87}\text{Sr}/^{86}\text{Sr}$ isotope ratios, whereas crustal materials devoid of mantle-derived material have low $^{143}\text{Nd}/^{144}\text{Nd}$ isotope ratios and high $^{87}\text{Sr}/^{86}\text{Sr}$ isotope ratios (e.g. SP samples). Continental crust plots in function of the proportion of integrated mantle-derived Sr and Nd on a mixing hyperbola defined by the end-member compositions of continental crust and mantle (see hypothetical mixing hyperbola in Figure 8). Thus, when comparing the regolith data of this study with European loess and tephra data from the

literature, one observes that each data set has a very particular isotopic signature. Each of these coordinates plot inside or on a mixing curve ranging from mantle or anthropogenic (low $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{143}\text{Nd}/^{144}\text{Nd}$ ratios) materials to crustal materials (Figure 8). The tephra signature from Wörner et al. (1985) being the most mantle rich material. This indicates that the different loess deposits originate from very distinct rock units containing different proportions of mantle and crust material. In this context, our PPSD show Sr and Nd isotope signatures rather close to the studied slate regolith compartment than to other loesses (Figure 8), which we hypothesise reflects the genetic link of the materials of both compartments.

6. Conclusion

The study we performed on a typical polygenetic regolith profile from a remnant plateau from the western part of the Rhenish Massif allowed three distinct compartments to be distinguished. Their geochemical and mineralogical characteristics present contrasting evolutions, which can be related to different atmospheric deposition events and to the seasonal water saturation dynamics. The estimations based on $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{143}\text{Nd}/^{144}\text{Nd}$ ratios indicate that the recent atmospheric deposition contributes to about 50% of the Sr and Nd content in the organo-mineral part of the regolith (0-45 cm depth). Nb enrichments and specific mineralogical compositions in the upper subsoil (45-140 cm depth) point to tephra contributions in the loess of the PPSD. However, we cannot refer to a specific Upper Pleistocene volcanic event. Finally, we identified a close genetic link of both upper and lower *subsolum* materials. The fresh slate and the central European loess from the literature defines a mixing curve between crustal and mantle materials, in which our PPSD layers show signatures close to the studied slate compartment and continental crust rather than to other European loess with lower Sr and higher Nd isotopic compositions.

The combination of mineralogical, major and trace element pattern and Sr-Nd-Pb-U isotope ratio analyses appears to be an extremely powerful approach for understanding the evolution of the polygenetic regolith systems. This procedure offers the possibility of i) characterising and tracking the origin of the components that constitute the different compartments of the regolith; ii) identifying the genetic links between the different regolith compartments; and iii) doing further research on its great potential for the identification of important hydrological tracers, which are able to shed light on water pathways and water-rock interactions within the complex regolith system of the critical zone.

Acknowledgement

This study is part of a project financed by the National Research Fund (FNR) of the Grand Duchy of Luxembourg (AFR Grant 9188481). We want to thank Jean François Iffly, François Barnich and Johanna Ziebel for their help in the field and laboratories and Francis Huguet for his scientific advice. We also thank Richard Coupe from the USGS for a preliminary review of the manuscript, as well as Markus Egli, editor of the journal, and the reviewers for their helpful revision.

References

- AG (Ad-hoc-Arbeitsgruppe) Boden, 2005. Bodenkundliche Kartieranleitung. E. Schweizerbart, Hannover.
- Aubert, D., Stille, P., Probst, A., 2001. REE fractionation during granite weathering and removal by waters and suspended loads: Sr and Nd isotopic evidence. *Geochim. Cosmochim. Acta* 65, 387–406. doi:10.1016/S0016-7037(00)00546-9

719 Aubert, D., Stille, P., Probst, A., Gauthier-lafaye, F., Pourcelot, L., Del nero, M., 2002.
 720 Characterization and migration of atmospheric REE in soils and surface waters.
 721 Geochim. Cosmochim. Acta 66, 3339–3350. doi:10.1016/S0016-7037(02)00913-4

722 Banwart S., Bernasconi S.M., Bloem J., Blum W., Brandao M., Brantley S., Chabaux F.,
 723 Duffy C., Kram P., Lair G., Lundin L., Nikolaidis N., Novak M., Panagos P.,
 724 Ragnarsdottir K.V., Reynolds B., Rousseva S., de Ruiter P., van Gaans P., van
 725 Riemsdijk W., White T., Zhang B., 2011. Soil Processes and Functions in Critical Zone
 726 Observatories: Hypotheses and Experimental Design, Vadose zone Journal, 10, 974-987.

727 Berner R.A., Maasch K.A., 1996, Chemical weathering and controls on atmospheric CO₂ and
 728 CO₂: fundamental principles were enunciated by J.J. Ebelmen in 1845. Geochim.
 729 Cosmochim. Acta, 60, 1633-1637.

730 Bini, C., Sartori, G., Wahsha, M., Fontana, S., 2011. Background levels of trace elements and
 731 soil geochemistry at regional level in NE Italy. J. Geochemical Explor. 109, 125–133.
 732 doi:10.1016/j.gexplo.2010.07.008

733 Barbier, F. 2012. Etude de paléoaaltération météorique (Crétacé Inférieur à Néogène) sur socle
 734 silicoclastique hétérogène : caractérisation et essai de cartographie d'altérites en Région
 735 Wallonne (Belgique). Ph D Thesis. Prom. : Yans, Johan

736 Bojko, O., Kabala, C., 2016. Transformation of physicochemical soil properties along a
 737 mountain slope due to land management and climate changes — A case study from the
 738 Karkonosze Mountains, SW Poland, Catena, Volume 140, 43-54.

739 Bonjour, J.-L., Dabard, M.-P., 1991. Ti/Nb ratios of clastic terrigenous sediments used as an
 740 indicator of provenance. Chem. Geol. 91, 257–267. doi:10.1016/0009-2541(91)90003-A

741 Brantley, S.L., Kubicki, J.D., White, A.F. (Eds.), 2008. Kinetics of Water-Rock Interaction.
 742 Springer New York, New York, NY. doi:10.1007/978-0-387-73563-4

743 Braun, J.-J., Pagel, M., Muller, J.-P., Bilong, P., Michard, A., Guillet, B., 1990. Cerium
 744 anomalies in lateritic profiles. *Geochim. Cosmochim. Acta* 54, 781–795.
 745 doi:10.1016/0016-7037(90)90373-S

746 Braun, J.-J., Viers, J., Dupré, B., Polve, M., Ndam, J., Muller, J.-P., 1998. Solid/Liquid REE
 747 Fractionation in the Lateritic System of Goyoum, East Cameroon: The Implication for
 748 the Present Dynamics of the Soil Covers of the Humid Tropical Regions. *Geochim.*
 749 *Cosmochim. Acta* 62, 273–299. doi:10.1016/S0016-7037(97)00344-X

750 Bruno, J., De Pablo, J., Duro, L., Figuerola, E., 1995. Experimental study and modeling of the
 751 U(VI)-Fe(OH)₃ surface precipitation/coprecipitation equilibria. *Geochim. Cosmochim.*
 752 *Acta* 59, 4113–4123. doi:10.1016/0016-7037(95)00243-S

753 Catt J.A. (1986) Soils and Quaternary geology. A Handbook for Field Scientist, Oxford
 754 Sciences Publications, Clarendon Press, Oxford, 1986.

755 Chabaux, F., O'Nions, R.K., Cohen, A.S., Hein, J.R., 1997. ²³⁸U-²³⁴U-²³⁰Th disequilibrium
 756 in Fe-Mn crusts: Palaeoceanographic record or diagenetic alteration? *Geochim.*
 757 *Cosmoch. Acta* 61, 3619-3632.

758 Chabaux, F., Riotte, J., Dequincey, O., 2003. U–Th–Ra fractionation during weathering and
 759 river transport. *Rev. Mineral. Geochem.* 52, 533–576.

760 Chabaux, F., Bourdon, B., Riotte, J., 2008. Chapter 3 U-series geochemistry in weathering
 761 profiles, river waters and lakes. In: Krishnaswami, S., Cochran, J.K. (Eds.), *Radioactiv-*
 762 *ity in the Environment*. Elsevier, pp. 49–104.

763 Chauvel, C., Garçon, M., Bureau, S., Besnault, A., Jahn, B., Ding, Z., 2014. Constraints from
 764 loess on the Hf–Nd isotopic composition of the upper continental crust. *Earth Planet.*
 765 *Sci. Lett.* 388, 48–58. doi:10.1016/j.epsl.2013.11.045

766 Cole, H. M. (1965). *The Ardennes: Battle of the Bulge* (Vol. 8). Government Printing Office.

767 Condie, K.C., 1993. Chemical composition and evolution of the upper continental crust:
 768 Contrasting results from surface samples and shales. *Chem. Geol.* 104, 1–37.
 769 doi:10.1016/0009-2541(93)90140-E

770 Craig, M.A., Wilford, J.R., Tapley, I.J. 1999. Regolith-landform mapping in the Gawler craton.
 771 *MESA journal* 12: 17-21.

772 Debajyoti, P., White, W. M., Turcotte, D.L., 1967. Modelling isotopic evolution of Earth. *Br.*
 773 *Med. J.* 4, 438.

774 Deckers J., 1966. Contribution à l'étude de la composition et de la capacité de production des
 775 sols de l'Ardenne centrale et de la Famenne orientale. Mémoire n°3. Gand, Belgique:
 776 Société belge de Pédologie

777 Deer, W.A., Howie, R. A., Zussmann, J., 1974. *An Introduction to the Rock-forming*
 778 *Minerals*. London: Longman.

779 Demoulin, A., 2003. Paleosurfaces and residual deposits in Ardenne-Eifel: historical
 780 overview and perspectives. *Géol. Fr.* 2003 (1), 17–21.

781 Deniel, C., Pin, C., 2001. Single-stage method for the simultaneous isolation of lead and
 782 strontium from silicate samples for isotopic measurements. *Anal. Chim. Acta* 426, 95–
 783 103. doi:10.1016/S0003-2670(00)01185-5

784 DePaolo, D.J., Maher, K., Christensen, J.N., McManus, J., 2006. Sediment transport time
785 measured with U-series isotopes: results from ODP North Atlantic drift site 984. *Earth*
786 *Planet. Sci. Lett.* 248, 394–410.

787 DePaolo, D.J., Lee, V.E., Christensen, J.N., Maher, K., 2012. Uranium comminution ages:
788 sediment transport and deposition time scales. *C. R. Geosci.* 344, 678–687.

789 Dequincey, O., Chabaux, F., Clauer, N., Sigmarsson, O., Liewig, N., Leprun, J.C., 2002.
790 Chemical remobilizations in laterites: Evidence from trace elements and ^{238}U - ^{234}U -
791 ^{230}Th disequilibria *Geochim. Cosmoch. Acta* , 66, 1197-1210

792 Désiré-Marchand, J., 1985. Notice de la carte géomorphologique du Grand-Duché de
793 Luxembourg. Service Géologique de Luxembourg, Bulletin 13, 47 p.
794

795 Dietze, M., Kleber, A., 2010. Characterisation and prediction of thickness and material
796 properties of periglacial cover beds, Tharandter Wald, Germany, *Geoderma*, Volume
797 156, Issues 3–4, 346-356.

798 Doucelance, R., Manhès, G., 2001. Reevaluation of precise lead isotope measurements by
799 thermal ionization mass spectrometry. *Chem. Geol.* 176, 361-377.

800 Duff, M.C., Coughlin, J.U., Hunter, D.B., 2002. Uranium co-precipitation with iron oxide
801 minerals. *Geochim. Cosmochim. Acta* 66, 3533–3547. doi:10.1016/S0016-
802 7037(02)00953-5

803 Durand, S., Chabaux, F., Rihs, S., Düringer, P., Elsass, P., 2005. U isotope ratios as tracers of
804 groundwater inputs into surface waters: Example of the Upper Rhine hydrosystem.
805 *Chem. Geol.* 220, 1–19. doi:10.1016/j.chemgeo.2005.02.016

806 Eggleton, R. A., 2001. The regolith glossary: surficial geology, soils and landscapes.
807 Canberra: Cooperative Research Centre for Landscape Evolution and Mineral
808 Exploration, Floreat Park, WA. 144p.

809 FAO, 2006. Guidelines for Soil Description. fourth ed. FAO, Rome, p. 97.

810 Faure, G., 1986. Principles of isotope geology. John Wiley and Sons, Inc., New York, NY

811 Felix-Henningsen, P. , 1994. Mesozoic Tertiary weathering and soil formation on slates of the
812 Rhenish Massif (Germany). Catena, 21, 229-242.

813 Field, J.P., Breshears, D.D., Law, D.J., Villegas, J.C., Lopez-Hoffman, L., Brooks,
814 P.D., Chorover, J., Barron-Gafford, G.A., Gallery, R.E., Litvak, M.E., Lybrand, R.A.,
815 McIn-tosh, J.C., Meixner, T., Niu, G.Y., Papuga, S.A., Pelletier, J.D., Rasmussen, C.R.,
816 Troch, P.A., 2015. Critical zone services: expanding context, constraints, and currency
817 beyond ecosystem services. Vadose Zone J. 14. Doi:10.2136/vzj2014.10.0142

818 Gabrielli, C.P., McDonnell, J.J., 2012. An inexpensive and portable drill rig for bedrock
819 groundwater studies in headwater catchments. Hydrol. Process. 26, 622–632.
820 doi:10.1002/hyp.8212

821 Gallet, S., Jahn, B., Van Vliet Lanoë, B., Dia, A., Rossello, E., 1998. Loess geochemistry and
822 its implications for particle origin and composition of the upper continental crust. Earth
823 Planet. Sci. Lett. 156, 157–172. doi:10.1016/S0012-821X(97)00218-5

824 Geagea, M.L., Stille, P., Millet, M., Perrone, T., 2007. REE characteristics and Pb, Sr and Nd
825 isotopic compositions of steel plant emissions. Sci. Total Environ. 373, 404–19.
826 doi:10.1016/j.scitotenv.2006.11.011

827 Gourdol, L., Hissler, C., Hoffmann, L., Pfister, L., 2013. On the potential for the Partial
828 Triadic Analysis to grasp the spatio-temporal variability of groundwater hydrochemistry.
829 *Appl. Geochemistry* 39, 93–107. doi:10.1016/j.apgeochem.2013.10.002

830 Granet, M., Chabaux, F., Stille, P., Dosseto, A., France-Lanord, C., Blaes, E. (2010) U-series
831 disequilibria in suspended river sediments and implication for sediment transfer time in
832 alluvial plains: the case of the Himalayan rivers *Geochim. Cosmochim. Acta*, 74, 2851-
833 2865

834 Guéguen, F., Stille, P., Dietze, V., Gieré, R., 2012. Chemical and isotopic properties and
835 origin of coarse airborne particles collected by passive samplers in industrial, urban, and
836 rural environments. *Atmos. Environ.* 62, 631–645. doi:10.1016/j.atmosenv.2012.08.044

837 Güler, C., Thyne, G.D., McCray, J.E., Turner, K. a., 2002. Evaluation of graphical and
838 multivariate statistical methods for classification of water chemistry data. *Hydrogeol. J.*
839 10, 455–474. doi:10.1007/s10040-002-0196-6

840 Haase, D., Fink, J., Haase, G., Ruske, R., Pécsi, M., Richter, H., Altermann, M., Jäger, K.D.,
841 2007. Loess in Europe-its spatial distribution based on a European Loess Map, scale
842 1:2,500,000. *Quat. Sci. Rev.* 26, 1301–1312. doi:10.1016/j.quascirev.2007.02.003

843 Hartigan, J., 1975. *Clustering Algorithms*. John Wiley and Sons, NY.

844 Hissler, C., Probst, J.-L., 2006. Impact of mercury atmospheric deposition on soils and
845 streams in a mountainous catchment (Vosges, France) polluted by chlor-alkali industrial
846 activity: the important trapping role of the organic matter. *Sci. Total Environ.* 361, 163–
847 78. doi:10.1016/j.scitotenv.2005.05.023

848 Hissler, C., Stille, P., Juilleret, J., Iffly, J.F., Perrone, T., Morvan, G., 2015. Elucidating the
849 formation of terra fuscas using Sr–Nd–Pb isotopes and rare earth elements. Appl.
850 Geochemistry 54, 85–99. doi:10.1016/j.apgeochem.2015.01.011

851 Hissler, C., Stille, P., Krein, A., Geagea, M.L., Perrone, T., Probst, J.L., Hoffmann, L., 2008.
852 Identifying the origins of local atmospheric deposition in the steel industry basin of
853 Luxembourg using the chemical and isotopic composition of the lichen *Xanthoria*
854 *parietina*. Sci. Total Environ. 405, 338–344. doi:10.1016/j.scitotenv.2008.05.029

855 Huguet, F., 1998. Le Hunsrück: un jalon dans l'évolution géomorphologique de la chaîne
856 varisque/Between Hohes Venn and Schwarzwald, geomorphological investigations in
857 Hunsrück (Rhenish Massif). In: Géomorphologie : relief, processus, environnement,
858 vol. 4, n°3. pp. 233-250.

859 IUSS Working Group WRB. 2015. World Reference Base for Soil Resources 2014, update
860 2015. International soil classification system for naming soils and creating legends for
861 soil maps. World Soil Resources Reports No. 106. FAO, Rome.Jabiol, B., Zanella, A.,
862 Ponge, J.F., Sartori, G., Englisch, M., van Delft, B., de Waal, R., Le Bayon, R.C., 2013.
863 A proposal for including humus forms in the World Reference Base for Soil Resources
864 (WRB-FAO). Geoderma 192, 286–294. doi:10.1016/j.geoderma.2012.08.002

865 Juilleret, J., Dondeyne, S., Vancampenhout, K., Deckers, J., Hissler, C., 2016. Mind the gap: A
866 classification system for integrating the subsolum into soil surveys, Geoderma,
867 <http://dx.doi.org/10.1016/j.geoderma.2015.08.031>

868 Juilleret J., Iffly J.-F., Pfister L. and Hissler C., 2011. Remarkable Pleistocene periglacial
869 slope deposits in Luxembourg (Oesling): pedological implication and geosite potential.
870 Bull. Soc. Nat. Luxemb. 112, 125-130.

871 Pouclet, A., Juvigne, E., 2009. The Eltville tephra, a late Pleistocene widespread tephra layer
872 in Germany, Belgium and The Netherlands; symptomatic compositions of the minerals.
873 Geol. Belgica. 12 (1/2), 93-103.

874 Kasprzak, M., Migoń, P., 2015. DEM-based analysis of geomorphology of a stepped
875 sandstone plateau, Stołowe Mountains (SW Poland) Stołowe Mountains (SW Poland).
876 Zeitschrift für Geomorphol. Suppl. Issues 59, 245–268. doi:10.1127/zfg_suppl/2015/S-
877 00183.

878 Kaufman, L., Rouseeuw, P.J., 1990. Finding Groups in Data. John Wiley and Sons Inc., NY.

879 Kaufman, L., Rouseeuw, P.J., 2005. Finding Groups in Data. John Wiley and Sons Inc., NY ,
880 368pp.

881 Kleber, A., 1997. Cover-beds as soil parent materials in midlatitude regions. CATENA 30,
882 197–213. doi:10.1016/S0341-8162(97)00018-0.

883 Kleber, A., Terhorst, B., 2013. Mid-Latitude Slope Deposits (Cover Beds), Developments in
884 Sedimentology, Developments in Sedimentology. Elsevier. doi:10.1016/B978-0-444-
885 53118-6.00001-5.

886 Kwaad, F.J.P.M., Múcher, H.J., 1977. The evolution of soils and slope deposits in the
887 Luxembourg Ardennes near Wiltz. Geoderma 17, 1–37. doi:10.1016/0016-
888 7061(77)90002-7

889 Laveuf, C., Cornu, S., Guilherme, L.R.G., Guerin, A., Juillot, F., 2012. The impact of redox
890 conditions on the rare earth element signature of redoximorphic features in a soil
891 sequence developed from limestone. Geoderma 170, 25–38.
892 doi:10.1016/j.geoderma.2011.10.014

893 Levitan, D.M., Zipper, C.E., Donovan, P., Schreiber, M.E., Seal, R.R., Engle, M. a.,
 894 Chermak, J. a., Bodnar, R.J., Johnson, D.K., Aylor, J.G., 2015. Statistical analysis of soil
 895 geochemical data to identify pathfinders associated with mineral deposits: An example
 896 from the Coles Hill uranium deposit, Virginia, USA. *J. Geochemical Explor.* 154, 238–
 897 251. doi:10.1016/j.gexplo.2014.12.012

898 Lin, X., Wang, X., Zhang, B., Yao, W., 2014. Multivariate analysis of regolith sediment
 899 geochemical data from the Jinwozi gold field, north-western China. *J. Geochemical*
 900 *Explor.* 137, 48–54. doi:10.1016/j.gexplo.2013.11.006

901 Lorz C. and Philips J.D., 2006. Pedo-ecological consequences of lithological discontinuities
 902 in soils-examples from Central Europe. *J. of Plant Nutrition and Soil Science* 169, 573-
 903 581.

904 Michard, A., Gurriet, P., Soudant, M., Albarede, F., 1985. Nd isotopes in French Phanerozoic
 905 shales: external vs. internal aspects of crustal evolution. *Geochim. Cosmochim. Acta*
 906 49, 601–610. doi:10.1016/0016-7037(85)90051-1

907 Müller, E.H., 1954. Die Bedeutung des eiszeitlichen Bodenfließens (Solifluktion) für die
 908 Bodenbildung im nördlichen Teil des Rheinischen Schiefergebirges. *Zeitschrift für*
 909 *Pflanzenernährung, Düngung, Bodenkd.* 65, 52–61. doi:10.1002/jpln.19540650107

910 National Research Council (NRC). 2001. Basic research opportunities in the earth sciences.
 911 National Academies Press, Washington, DC.

912 Paepe R. and Sommé J., 1970. Les loess et la stratigraphie du Pléistocene récent dans le nord
 913 de la France et en Belgique. *Ann. Soc. Géol. Nord* 90, 191-201.

914

915 Pelt, E., Chabaux, F., Innocent, C., Navarre-Sitchler, A.K., Sak, P.B., Brantley, S.L., 2008.
 916 Uranium–thorium chronometry of weathering rinds: Rock alteration rate and paleo-
 917 isotopic record of weathering fluids. *Earth Planet. Sci. Lett.* 276, 98–105.
 918 doi:10.1016/j.epsl.2008.09.010
 919
 920 Pierret, M.C., Stille, P., Prunier, J., Viville, D., Chabaux, F., 2014. Chemical and U-Sr
 921 isotopic variations of stream and source waters at a small catchment scale (the
 922 Strengbach case; Vosges mountains; France), *Hydrology and Earth System Sciences*
 923 Discussions. doi:10.5194/hessd-11-3541-2014
 924
 925 Pin, C., Zalduegui, J.S., 1997. Sequential separation of light rare-earth elements, thorium and
 926 uranium by miniaturized extraction chromatography: Application to isotopic analyses of
 silicate rocks. *Anal. Chim. Acta* 339, 79–89. doi:10.1016/S0003-2670(96)00499-0
 927
 928 Pissart A. (1995) L'Ardenne sous le joug du froid. Le modelé périglaciaire du massif
 929 ardennais. In : L'Ardenne, essai de Géographie physique (A. Demoulin, éd.),
 Département de Géographie physique, Uni. De Liège, pp. 136-154
 930
 931 Pouclet, A., Juvigné, E., Pirson, S., 2008. The Rocourt Tephra, a widespread 90–74 ka
 932 stratigraphic marker in Belgium. *Quat. Res.* 70, 105–120.
 doi:10.1016/j.yqres.2008.03.010
 933
 934 Prunier, J., Chabaux, F., Stille, P., Gangloff, S., Pierret, M.C., Viville, D., Aubert, A., 2015.
 935 Geochemical and isotopic (Sr, U) monitoring of soil solutions from the Strengbach
 936 catchment (Vosges mountains, France): Evidence for recent weathering evolution.
Chem. Geol. 417, 289–305. doi:10.1016/j.chemgeo.2015.10.012

- 937 Redon, P.-O., Bur, T., Guiesse, M., Probst, J.-L., Toiser, A., Revel, J.-C., Jolivet, C., Probst,
938 A., 2013. Modelling trace metal background to evaluate anthropogenic contamination in
939 arable soils of south-western France. *Geoderma* 206, 112–122.
940 doi:10.1016/j.geoderma.2013.04.023
- 941 Reimann, C., Filzmoser, P., Garrett, R.G., Dutter, R., 2008. *Statistical Data Analysis*
942 *Explained: Applied Environmental Statistics with R*. John Wiley and Sons Ltd., NY,
943 343pp.
- 944 Rousseau, D.-D., Chauvel, C., Sima, A., Hatté, C., Lagroix, F., Antoine, P., Balkanski, Y.,
945 Fuchs, M., Mellett, C., Kageyama, M., Ramstein, G., Lang, A., 2014. European glacial
946 dust deposits: Geochemical constraints on atmospheric dust cycle modeling. *Geophys.*
947 *Res. Lett.* 41, 7666–7674. doi:10.1002/2014GL061382
- 948 Rousseau, D.D., Ghil, M., Kukla, G., Sima, a., Antoine, P., Fuchs, M., Hatté, C., Lagroix, F.,
949 Debret, M., Moine, O., 2013. Major dust events in Europe during marine isotope stage 5
950 (130–74 ka): A climatic interpretation of the “markers.” *Clim. Past* 9, 2213–2230.
951 doi:10.5194/cp-9-2213-2013
- 952 Sauer, D., 2002. Genese, Verbreitung und Eigenschaften periglaziärer Lagen im Rheinischen
953 Schiefergebirge- anhand von Beispielen aus Westerwald, Hunsrück und Eifel. *Boden u.*
954 *Landschaft* 36, 1-294.
- 955 Sauer, D., Scholten, T., Spies, E.-D., Felix-Henningsen, P., 2002a. Pleistocene periglacial
956 slope deposits and colluviums in the Rhenish Massif—a result of climatic changes and
957 human landuse. In: Rubio, J.L., Morgan, R.P.C., Asins, S., Andreu, V. (Eds.), *Man and*
958 *Soil at the Third Millennium*, Vol. 1. Geoforma Ediciones/Centro de Investigaciones
959 *sobre Desertificacio´ n*, Logroñ o pp. 763–776.

- 960 Sauer, D., Felix-Henningsen, P., 2006. Saprolite, soils, and sediments in the Rhenish Massif
961 as records of climate and landscape history. *Quat. Int.* 156-157, 4–12.
962 doi:10.1016/j.quaint.2006.05.001
- 963 Schaffhauser, T., Chabaux, F., Ambroise, B., Lucas, Y., Stille, P., Reuschlé, T., Perrone, T.,
964 Fritz, B., 2014. Geochemical and isotopic (U, Sr) tracing of water pathways in the
965 granitic Ringelbach catchment (Vosges Mountains, France). *Chem. Geol.* 374-375, 117–
966 127.
- 967 Schmincke, H.-U., Park, C., Harms, E., 1999. Evolution and environmental impacts of the
968 eruption of Laacher See Volcano (Germany) 12,900 a BP. *Quat. Int.* 61, 61–72.
969 doi:10.1016/S1040-6182(99)00017-8
- 970 Schot, P.P., van der Wal, J., 1992. Human impact on regional groundwater composition
971 through intervention in natural flow patterns and changes in land use. *J. Hydrol.* 134,
972 297–313. doi:10.1016/0022-1694(92)90040-3
- 973 Scott, K.M., Pain, C.F. (Eds), 2008. *Regolith Science*. Springer Science and CSIRO,
974 Dordrecht, The Netherlands, and Collingwood, Australia.
- 975 Semmel, A., Terhorst, B., 2010. The concept of the Pleistocene periglacial cover beds in
976 central Europe: A review. *Quat. Int.* 222, 120–128. doi:10.1016/j.quaint.2010.03.010
- 977 Shields, G., Stille, P., 2001. Diagenetic constraints on the use of cerium anomalies as
978 palaeoseawater redox proxies: an isotopic and REE study of Cambrian phosphorites.
979 *Chem. Geol.* 175, 29–48. SSSA 2008
- 980 Soil Science Glossary Terms Committee (SSGT). 2008. Internet glossary of soil science
981 terms. <https://www.soils.org/publications/soils-glossary> (accessed 17.08.2015.)

982 Steinmann, M., Stille, P., 1997. Rare earth element behavior and Pb, Sr, Nd isotope
983 systematics in a heavy metal contaminated soil. *Appl. Geochem.* 12, 607–623.

984 Stille, P., Pierret, M.C., Steinmann, M., Chabaux, F., Boutin, R., Aubert, D., Pourcelot, L.,
985 Morvan, G., 2009. Impact of atmospheric deposition, biogeochemical cycling and water-
986 mineral interaction on REE fractionation in acidic surface soils and soil water (the
987 Strengbach case). *Chem. Geol.* 264, 173–186. doi:10.1016/j.chemgeo.2009.03.005

988 Stille, P., Pourcelot, L., Granet, M., Pierret, M.C., Guéguen, F., Perrone, T., Morvan, G.,
989 Chabaux, F., 2011. Deposition and migration of atmospheric Pb in soils from a forested
990 silicate catchment today and in the past (Strengbach case): Evidence from ²¹⁰Pb
991 activities and Pb isotope ratios. *Chem. Geol.* 289, 140–153.
992 doi:10.1016/j.chemgeo.2011.07.021

993 Stückerad, S., Sabel, K.J., Wilcke, W., 2010. Contributions of different parent materials in
994 soils developed from periglacial cover-beds. *Eur. J. Soil Sci.* 61, 844–853.
995 doi:10.1111/j.1365-2389.2010.01288.x

996 Taylor, G., Eggleton, R. A., 2001. *Regolith Geology and Geomorphology*. 375 pp.
997 Chichester, New York, Weinheim, Brisbane, Singapore, Toronto: John Wiley & Sons
998 Ltd.

999 Taylor, S.R., McLennan, S.M., 1981. The composition and evolution of the continental crust:
1000 rare earth element evidence from sedimentary rocks. *Phil. Trans. R. Soc. Lond. A* 301,
1001 381–399.

1002 Taylor S.R., McLennan S.M. and McCulloch M.T. (1983) Geochemistry of loess, continental
1003 crust composition and crustal model ages. *Geochim. Cosmochim. Acta* 47, 1897-1905.

- 1004 Taylor, S.R., McLennan, S.M., 1985. The Continental Crust: Its Composition and Evolution.
1005 Blackwell, Oxford, 312p.
- 1006 Templ, M., Filzmoser, P., Reimann, C., 2008. Cluster analysis applied to regional
1007 geochemical data: Problems and possibilities. *Appl. Geochemistry* 23, 2198–2213.
1008 doi:10.1016/j.apgeochem.2008.03.004
- 1009 Terhorst, B., 2007. Periglacial cover beds and soils in landslide areas of SW-Germany.
1010 *Catena*, 71, 467–476. doi:10.1016/j.catena.2007.03.021
- 1011 Thiry, M., 2006. Continental France and Belgium during the early Cretaceous:
1012 paleoweatherings and paleolandforms. *Bull. la Société Géologique Fr.* 177, 155–175.
1013 doi:10.2113/gssgfbull.177.3.155
- 1014 Vázquez-Ortega, A., Perdrial, J., Harpold, A., Zapata-Ríos, X., Rasmussen, C., McIntosh, J.,
1015 Schaap, M., Pelletier, J.D., Brooks, P.D., Amistadi, M.K., Chorover, J., 2015. Rare earth
1016 elements as reactive tracers of biogeochemical weathering in forested rhyolitic terrain.
1017 *Chem. Geol.* 391, 19–32. doi:10.1016/j.chemgeo.2014.10.016
- 1018 Völkel, J., Huber, J., Leopold., M., 2011. Significance of slope sediments layering on
1019 physical characteristics and interflow within the Critical Zone - Examples from the
1020 Colorado Front Range, USA, *Applied Geochemistry*, Volume 26, Supplement, June
1021 2011, Pages S143-S145, ISSN 0883-2927,
1022 <http://dx.doi.org/10.1016/j.apgeochem.2011.03.052>.
- 1023 Ward, J.H., 1963. Hierarchical grouping to optimize an objective function. *J. Am. Stat. Assoc.*
1024 doi:10.1080/01621459.1963.10500845

1025 West, N., Kirby, E., Bierman, P., Slingerland, R., Ma, L., Rood, D., Brantley, S., 2013.
 1026 Regolith production and transport at the Susquehanna Shale Hills Critical Zone
 1027 Observatory, Part 2: Insights from meteoric ¹⁰Be. *J. Geophys. Res. Earth Surf.* 118,
 1028 1877–1896. doi:10.1002/jgrf.20121

1029 Wörner, G., Staudigel, H., Zindler, A., 1985. Isotopic constraints on open system evolution of
 1030 the Laacher See magma chamber (Eifel, West Germany). *Earth Planet. Sci. Lett.* 75, 37–
 1031 49. doi:10.1016/0012-821X(85)90048-2

1032 Wörner, G., Schmincke, H.-U., 1984. Mineralogical and Chemical Zonation of the Laacher
 1033 See Tephra Sequence (East Eifel, W. Germany). *J. Petrol.* 25, 805–835.
 1034 doi:10.1093/petrology/25.4.805

1035 Wörner, G., Wright, T.L., 1984. Evidence for magma mixing within the Laacher See magma
 1036 chamber (East Eifel, Germany). *J. Volcanol. Geotherm. Res.* 22, 301–327.
 1037 doi:10.1016/0377-0273(84)90007-6

1038
 1039

Table 1. Regolith structure and stratigraphic characterisation, including location of the samples (LD: Lithic discontinuities; UL: Upper Layer; BL: Basal Layer).

Stratigraphy	Regolith profile	Horizons layers	Diagnostic material	Sample depth (cm)	Sample name	
PPSD (LU)		O		-1 – 0	OH	
		Ah		0 – 4	PPSD1	
		B	Cambic	4 – 45	PPSD2	L.D
PPSD (BL)		2Cg1	Regolithic	45 – 80	PPSD3	
		2Cg2	Regolithic	110	PPSD4	L.D
Slate substrate		3CR	Saprolithic	190	SP1	
		3CR	Saprolithic	270	SP2	
		3CR	Saprolithic	320	SP3	
		3CR	Saprolithic	380	SP4	
		3R	Paralithic	735	SP5	

Table 2. Major (% oxide) and trace element (ppm) composition of the regolith (D.L.: Detection Limit; NA: not analysed).

	D.L.	OH	PPSD1	PPSD2	PPSD3	PPSD4	PPSDce	SP1	SP2	SP3	SP4	SP5
Major elements (% oxide)												
Na ₂ O	0.01	0.05	0.24	0.38	0.38	0.40	0.44	0.30	0.21	0.24	0.27	0.26
MgO	0.015	0.10	0.30	1.02	1.69	1.92	1.57	1.90	1.85	1.73	2.18	1.83
Al ₂ O ₃	0.02	1.31	7.70	16.49	19.12	22.86	13.11	19.13	16.69	18.76	20.01	18.29
SiO ₂	0.02	<0.02	37.82	62.82	60.27	52.92	66.35	62.51	64.16	58.53	59.41	64.35
P ₂ O ₅	0.04	<0.04	0.27	0.16	0.12	0.22	0.21	0.07	0.17	0.29	0.15	0.15
K ₂ O	0.01	0.28	1.31	2.63	3.41	3.92	1.96	3.62	3.07	3.72	4.19	3.77
CaO	0.03	<0.03	0.10	0.08	0.05	0.00	<0.03	0.00	0.00	0.04	0.06	0.15
TiO ₂	0.02	<0.02	0.55	1.01	1.05	1.12	0.64	0.96	0.86	0.93	0.95	0.87
Fe ₂ O ₃	0.02	0.66	3.48	5.77	6.66	7.99	10.90	5.26	7.15	9.15	6.31	5.56
MnO	<0.001	0.05	0.03	0.06	0.05	0.09	0.05	0.02	0.03	0.94	0.05	0.05
Trace elements (ppm)												
Cr	4	15.4	72.6	127.3	151.6	173.8	120.8	166.9	144.0	144.4	149.8	133.6
Co	0.4	1.7	5.0	15.3	17.5	20.9	20.5	16.0	13.8	200.8	61.9	15.3
Ni	5	8	20	60	71	78	83	74	74	131	79	68
Cu	5	12	20	20	23	39	30	17	41	68	36	44
Zn	11	46	84	145	107	147	110	96	115	175	116	101
As	1.5	2.5	14.9	8.2	9.5	13.8	10.1	8.3	14.9	18.2	12.5	7.6
Rb	0.4	12.2	74.1	159.3	175.6	193.3	91.0	166.6	146.1	167.0	187.7	170.4
Sr	2	14	48	94	110	126	61	96	81	102	113	98
Y	0.2	<0.2	15.8	31.5	38.7	36.6	22.2	34.8	31.8	32.6	33.7	28.0
Zr	1	<1	154	283	305	293	168	260	190	177	174	149
Nb	0.09	<0.09	10.28	19.87	19.44	19.66	10.77	15.96	14.61	14.94	15.28	13.93
Cd	0.12	0.25	0.32	0.25	0.27	0.25	0.2	0.25	0.24	0.28	0.19	0.16
Sn	0.45	0.12	9.63	3.76	4.13	4.63	2.44	4.13	3.72	4.30	4.45	4.04
Sb	0.2	0.4	2.9	0.6	0.5	0.5	0.4	0.3	0.7	0.9	0.7	0.4
La	0.09	2.23	22.03	45.30	51.45	61.33	30.62	48.00	41.00	48.29	57.40	44.04
Ce	0.14	4.26	43.15	91.90	102.90	128.00	61.53	96.75	82.07	109.30	124.10	89.11
Pr	0.015	0.502	4.876	10.340	11.640	14.550	7.052	11.580	9.511	11.880	15.150	10.180
Nd	0.06	1.87	17.93	37.48	42.35	54.10	25.80	43.11	35.83	45.39	59.72	37.89
Sm	0.015	0.313	3.250	6.923	7.778	9.712	4.968	8.233	7.068	9.039	11.680	7.216
Eu	0.005	0.058	0.659	1.408	1.562	1.875	1.030	1.602	1.458	1.871	2.348	1.471
Gd	0.013	0.176	2.682	5.668	6.465	7.161	4.099	6.494	6.111	7.474	8.833	5.970
Tb	0.003	0.029	0.446	0.916	1.074	1.104	0.670	1.001	0.973	1.097	1.187	0.892
Dy	0.01	0.17	2.81	5.80	6.89	6.81	4.11	6.21	5.99	6.50	6.76	5.41
Ho	0.002	0.036	0.601	1.201	1.451	1.395	0.847	1.286	1.250	1.326	1.340	1.112
Er	0.01	0.10	1.63	3.26	3.93	3.79	2.28	3.50	3.32	3.53	3.46	2.94
Tm	0.001	0.016	0.241	0.470	0.571	0.563	0.336	0.512	0.484	0.519	0.506	0.429
Yb	0.007	0.110	1.647	3.274	3.904	3.829	2.296	3.447	3.315	3.531	3.409	2.996
Lu	0.003	0.016	0.252	0.508	0.592	0.586	0.357	0.520	0.500	0.543	0.516	0.451
Hf	0.03	<0.03	3.98	7.53	8.02	7.93	4.55	6.99	5.49	5.16	5.18	4.40
Hg	<0.001	<0.001	0.342	0.091	0.050	0.038	<0.001	<0.001	0.040	0.049	0.043	0.031
Pb	0.7	41.6	128.9	19.4	14.6	14.6	8.9	30.3	24.3	33.7	24.5	17.8
Th	0.06	0.66	7.07	13.21	15.07	17.26	10.12	14.53	13.22	14.43	15.21	13.71
U	0.03	0.33	1.93	3.14	3.14	3.51	2.45	3.09	3.82	5.22	5.84	3.83

Table 3. Sr, Nd, Pb and U radiogenic isotope ratios of all analysed regolith samples(NA: not analysed).

Sample	$^{87}\text{Sr}/^{86}\text{Sr}$	$^{143}\text{Nd}/^{144}\text{Nd}$	$^{206}\text{Pb}/^{207}\text{Pb}$	$^{234}\text{U}/^{238}\text{U}$
OH	0.72181	0.511951	1.153	0.933
PPSD1	0.73023	0.511919	1.162	NA
PPSD2	0.73399	0.511922	1.204	NA
PPSD3	0.73866	0.511923	1.215	0.961
PPSD4	0.73657	0.511931	1.220	NA
PPSDce	0.73736	NA	1.226	NA
SP1	0.74141	0.511961	1.200	0.949
SP2	0.74205	0.511927	1.202	1.053
SP3	0.73912	0.511959	1.199	1.038
SP4	0.73915	0.511983	1.206	0.953
SP5	0.74160	0.511933	1.214	0.947

FIGURE CAPTIONS

Figure 1. Situation and extension of the Rhenish Massif in Europe and location of the study site in the Luxembourg Ardennes region.

Figure 2. Depth-dependent patterns of mineralogical composition from XRD analyses expressed in relative abundances (PPSDce: PPSD coarse elements; dashed lines: lithic discontinuities; shaded area: redox sensitive zone).

Figure 3. Hierarchical cluster analysis performed on major and traces elements: (a) dendrogram defined using Pearson correlation distance as distance measure and Ward's method for the linkage rule; (b) coloured z-transformed concentrations matrix (row names correspond to regolith layers, parameter columns are ordered as the dendrogram); and (c) z-transformed concentrations as a function of depth for the 3 parameters groups defined (the thick black lines correspond to the median of each group, the thin coloured lines correspond to individual parameters of each subgroup as defined by the colour of the parameter labels in the dendrogram (a)).

Figure 4. Major and trace element concentrations of (a) OH and PPSD samples and (b) SP samples, normalized to the SP5 sample concentrations, representing the fresh slate bedrock.

Figure 5. Depth dependent patterns of PAAS-normalised $\text{La}_\text{N}/\text{Yb}_\text{N}$ and $\text{La}_\text{N}/\text{Eu}_\text{N}$ ratios and Ce anomaly ($\text{Ce}_\text{N}/\text{Ce}_\text{N}^*$). (dashed lines: lithic discontinuities; shaded area: redox sensitive horizons).

Figure 6. Depth-dependent variation of (a) $^{206}\text{Pb}/^{207}\text{Pb}$, (b) $^{87}\text{Sr}/^{86}\text{Sr}$, (c) $^{143}\text{Nd}/^{144}\text{Nd}$ and (d) $^{234}\text{U}/^{238}\text{U}$ of all analysed samples (shaded area: redox sensitive horizons).

Figure 7. Relationships between (a) $^{87}\text{Sr}/^{86}\text{Sr}$ vs. Rb/Sr ratio and (b) $^{206}\text{Pb}/^{207}\text{Pb}$ vs. U/Pb ratio of all analysed samples and a lichen sample from Hissler et al. (2008) representing the anthropogenic end member.

Figure 8. $^{87}\text{Sr}/^{86}\text{Sr}$ vs. $^{143}\text{Nd}/^{144}\text{Nd}$ diagram describing mixing hyperbolas between the unweathered slate sample of this study (SP5) as the continental crust end member (Sr: 98.2ppm, $^{87}\text{Sr}/^{86}\text{Sr}$: 0.7416 ; Nd: 37.9ppm, $^{143}\text{Nd}/^{144}\text{Nd}$: 0.51193) and a mantle representing end member according to the study of

26 Wörner et al. (1985) (Sr: 500ppm; $^{87}\text{Sr}/^{86}\text{Sr}$: 0.70477; Nd: 49ppm; $^{143}\text{Nd}/^{144}\text{Nd}$: 0.51262); and between
27 SP5 and lichen representing the atmospheric derived anthropogenic end member (Sr: 1.17ppm,
28 $^{87}\text{Sr}/^{86}\text{Sr}$: 0.7152 ; Nd: 0.17ppm, $^{143}\text{Nd}/^{144}\text{Nd}$: 0.51205, from Hissler et al., 2008). Included are the
29 loess signatures from UK, France and Belgium from Gallet et al. (1998) and Kaiserstuhl (Germany)
30 from Taylor et al. (1983) (grey circles: Nb enriched PPSD samples).

31 Figure 9. $\epsilon\text{Nd(T)}$ values vs. stratigraphic ages of the PPSD and SP samples of this study and of rock
32 materials which originated from similar palaeographic environments reported by Michard et al. (1995)
33 and Ohr et al. (1994). Modified after Michard et al. (1995).

Figure 1

Figure 2

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9