1 TITLE

2 Disentangling canid howls across multiple species and subspecies: structure in a complex
3 communication channel

4

5 AUTHORS

6 Arik Kershenbaum¹, Holly Root-Gutteridge², Bilal Habib³, Jan Koler-Matznick⁴, Brian Mitchell⁵,
7 Vicente Palacios⁶, Sara Waller⁷

8 ¹Department of Zoology, University of Cambridge, UK

9 ²Department of Biology, Syracuse University, USA

10 ³ Department of Animal Ecology and Conservation Biology, Wildlife Institute of India, Dehradun,
11 India

12 ⁴ The New Guinea Singing Dog Conservation Society, Central Point, OR, USA

13 ⁵ The Rubenstein School of Environment and Natural Resources, University of Vermont, USA

14 ⁶Instituto Cavanilles de Biodiversidad y Biología Evolutiva. University of Valencia, Spain

15 ⁷Montana State University, Bozeman, MT, USA

16

17 CORRESPONDING AUTHOR: Arik Kershenbaum, arik.kershenbaum@gmail.com

18

19 WORD COUNT: 5376

20

21

22

23 ABSTRACT

24

25 Wolves, coyotes, and other canids are members of a diverse genus of top predators of considerable
26 conservation and management interest. Canid howls are long-range communication signals, used both
27 for territorial defence and group cohesion. Previous studies have shown that howls can encode
28 individual and group identity. However, no comprehensive study has investigated the nature of
29 variation in canid howls across the wide range of species. We analysed a database of over 2,000 howls
30 recorded from 13 different canid species and subspecies. We applied a quantitative similarity measure
31 to compare the modulation pattern in howls from different populations, and then applied an
32 unsupervised clustering algorithm to group the howls into natural units of distinct howl types. We
33 found that different species and subspecies showed markedly different use of howl types, indicating
34 that howl modulation is not arbitrary, but can be used to distinguish one population from another. We
35 give an example of the conservation importance of these findings by comparing the howls of the
36 critically endangered red wolves to those of sympatric coyotes *C. latrans*, with whom red wolves may
37 hybridise, potentially compromising reintroduced red wolf populations. We believe that quantitative
38 cross-species comparisons such as these can provide important understanding of the nature and use of
39 communication in socially cooperative species, as well as support conservation and management of
40 wolf populations.

41

42 Keywords: Bioacoustics, Coyote, Dog, Howling, Jackal, Social communication, Wolf

1. INTRODUCTION

The genus *Canis* comprises several species and subspecies that share many ecological and behavioural similarities (Bekoff et al. 1981). Most are apex predators, and although some hunt in packs and others alone, all species are strongly social, living in groups ranging in size from a handful of close family members, e.g. coyotes *Canis latrans* (Bekoff 1977), to large groups of 20 or more animals, e.g. Ethiopian wolves *C. simensis* (Sillero-Zubiri & Gottelli 1994). For humans, one of the most familiar canid behaviours is the howl, a long-range communication channel (i.e. a mode through which communication can occur) thought to play a role both in territorial advertising and in group cohesion (Theberge & Falls 1967; Harrington & Mech 1979; Harrington 1987). Howling is most familiar in grey wolves *C. lupus* (Harrington et al. 2003), but all species in the genus produce howl-like vocalisations in addition to other, shorter range communication, such as barks, yips, and growls (Cohen & Fox 1976). These diverse short-range vocalisations are thought to mediate much of canid social behaviour (Yin & McCowan 2004; Siniscalchi et al. 2008), such as maintaining dominance relationships, but it has been speculated that howling too plays a role in inter- and intra-group dynamics (Harrington & Mech 1979; Jaeger et al. 1996; Gese 2001). Support for this hypothesis includes recent studies showing that wolves recognise the howl vocalisations of familiar individuals, and that howls show affective changes in response to the removal of individuals from the group (Mazzini et al. 2013; Palacios et al. 2015). In addition, individual differences in howls have been found in other canid species, e.g. coyotes (Hallberg 2007), and dingoes (Déaux & Clarke 2013). Canids of all species pose a number of management and conservation challenges. As apex predators, canids have a major influence on prey populations, and changes in canid numbers can result in trophic cascades (Elmhagen & Rushton 2007; Beschta & Ripple 2009). Some species, such as the Ethiopian wolf (Sillero-Zubiri & Gottelli 1994) and the red wolf *C. rufus* (Paradiso & Nowak 1972) are critically endangered, whereas the grey wolf *C. lupus* is frequently in conflict with human populations due to livestock depredation (Sillero-Zubiri & Laurenson 2001), and golden jackals *C. aureus* and domestic dogs *C. familiaris* are considered to be significant reservoirs of rabies (Seimenis 2008; Davlin & VonVille 2012). Management of these issues requires an in-depth understanding of the

behavioural ecology of these species and subspecies, which would appear to be incomplete without an understanding of the role of long-range vocal communication. In addition, phylogenetic relationships in the genus as a whole are unclear (Bardeleben et al. 2005; Koepfli et al. 2015), with most component species being capable of producing fertile hybrids, and there is considerable lack of agreement over the status of several grey wolf subspecies and populations (Chambers et al. 2012). As a result, the possible role of vocal behaviour as an isolating factor (or otherwise) between populations is important for the conservation of genetic diversity in subspecies that, while genetically compatible, maintain considerable phenotypic adaptation to their local habitats (Chambers et al. 2012). Partly because of the lack of agreement on the taxonomic status of many canid species and subspecies, and partly for reasons of simplicity, in this paper we will use the term "species" as a shorthand for "species and subspecies".

Early studies of canid howling behaviour emphasised qualitative descriptions of howl types (McCarley 1975; Cohen & Fox 1976; Tembrock 1976; Lehner 1978) and overall acoustic characteristics, such as mean fundamental frequency and frequency range, as well as modulation shape measures (Theberge & Falls 1967; Tooze et al. 1990). Multiple variables describing changes in the frequency and amplitude of the howl over time can be used for individual discrimination, among which important discriminative variables are the mean, maximum, and coefficient of variation of the fundamental frequency, and the amplitudes of the various harmonics (Root-Gutteridge et al. 2014a; Root-Gutteridge et al. 2014b). However, there are reasons to consider that information exists in the precise frequency modulation of wolf howls, as well as in simpler acoustic characteristics. Firstly, howls are predominantly narrow-band vocalisations, meaning that most of the acoustic energy is concentrated at a small range of frequencies at any one time. Further, this well-defined frequency varies throughout the course of the howl (Figure 1). This "frequency modulation" is known to be used to encode information in other species with similar vocalisations; particularly bottlenose dolphins *Tursiops truncatus* (Janik & Slater 1998; Quick & Janik 2012), in which individual identity can be reliably extracted from the frequency modulation patterns of whistles (Kershenbaum et al. 2013). Frequency modulation is an effective encoding technique in terrestrial communication (Wiley & Richards 1978), and in addition, considerations of signal transmission indicate that long-range

communication in an absorptive environment (e.g. forest) would tend to favour narrow-band frequency modulation over other encoding modalities (Henry & Lucas 2010). Therefore, we consider it appropriate to analyse the frequency modulation of canid howls in a similar way to that of dolphin whistles, to test for characteristic differences between species and populations.

Figure 1. Example spectrogram of multiple wolves howling. The x-axis represents time (seconds) and the y-axis frequency (Hertz).

A few studies have examined frequency modulation in canid howls, e.g. in coyotes (Hallberg 2007) and Iberian wolves (Palacios et al. 2007), by defining stereotyped modulation patterns such as, "rising", "step down", and "warble to flat". However, these arbitrary categories may not be perceived as distinct units by the focal animal (Kershenbaum et al. 2014), and are potentially subject to selective bias by researchers focusing on "interesting" spectral patterns. Therefore, a thorough analysis of frequency modulation must include (a) a quantitative measure of howl similarity (Deecke & Janik 2006), and (b) a quantitative and objective method for grouping howls into distinct howl types, without relying on subjective interpretation. The latter requirement is particularly acute, as a quantitative comparison between the vocal behaviours of different populations is problematic if both repertoires include vocalisations that are qualitatively of a different nature. For example, comparing the howls of one population to the barks of another would be an unproductive effort. Therefore, an alternative paradigm is required that takes into account the partitioning of a vocal repertoire into distinct types, whether arising from functionally different mechanisms (such as howls and barks), or

whether being discrete variations of the same functional mechanism (such as different notes in a bird song). We propose that, where multiple distinct vocalisation types are used with overlapping repertoires between populations, the only meaningful way to compare behaviour is to compare the vocalisation type histograms, rather than compare the individual vocalisations. This approach has also been carried out in previous studies of birdsong syntax (Jin & Kozhevnikov 2011). In essence, we interpret the howl type usage histograms as a "fingerprint" of vocal behaviour. In this work, we define and implement a howl similarity metric, as well as an automated clustering technique, and analyse a large database of over 2,000 howls from 21 different species of canids. We classify these howls into distinct types, and compare the relative use of this global repertoire by different populations, thereby testing for objective differences that distinguish between different species. Our results show a diversity of different howl types between species and, although we do not explicitly test for contextual reference in canid howling, we cannot exclude the possibility that specific howl types may be more common in some behavioural contexts than others.

2. METHODS

We collected a database of canid howling recordings from a wide range of sources. Altogether, we collected 6,009 howls from 21 distinct species, from 207 sources. Recordings were made both of captive and wild animals. The number of sources for each species varied from one (dingo *C. lupus dingo* or *C. familiaris dingo*, status unclear, Tibetan wolf *C. l. chanco*, and others) to 20 (eastern timber wolf *C. l. lycaon*). However, we excluded all species with only a single source to avoid confounding individual distinctiveness with species distinctiveness, providing a dataset with 13 distinct species from 131 sources. Of these, 2,005 howls were considered to be of sufficient quality for further analysis (no overlapping howls, sufficient signal strength). A breakdown of the recordings is given in Table 1. For each howl, we traced the frequency modulation using a combination of manual and automatic extraction tools, using an image-processing ridge tracker (Kershenbaum &

Roch 2013), or by fitting the harmonic peaks to a Lorentzian function (Root-Gutteridge et al. 2014b).

Each analysis was reviewed by both AK and HRG for validation.

Table 1. Number of howls, and number of recording sources (packs) for each of the species in the database.

Canid species	Common name	Reference	Number of howls	Number of sources
<i>C. aureus</i>	Golden jackal		28	3
<i>C. latrans</i>	Coyote		187	4
	Red wolf	(Chambers et al. 2012)		4
<i>C. rufus</i>			79	
<i>C. lycaon</i> or <i>C. l. lycaon</i>	Eastern wolf	(Chambers et al. 2012)	510	20
<i>C. lupus</i>	Grey wolf			
	Mackenzie Valley wolf	(Chambers et al. 2012)	127	8
<i>C. l. occidentalis</i>	Mexican wolf	(Chambers et al. 2012)	31	2
<i>C. l. baileyi</i>				
	Arctic wolf	(Chambers et al. 2012)	26	7
<i>C. l. arctos</i>				
<i>C. l. lupus</i>	European wolf	(Nowak 1995)	65	13
<i>C. l. signatus</i>	Iberian wolf	(Vilà et al. 1999)	25	3
<i>C. l. pallipes</i>	Indian wolf	(Nowak 1995)	175	7
	North African wolf	(Rueness et al. 2011)	33	5
<i>C. l. lupaster</i>				
	Domestic dog			53
<i>C. familiaris</i> or <i>C. l. familiaris</i>	(as companion animal)		375	
<i>C. hallstromi</i> or <i>C. l. hallstromi</i>	New Guinea singing dog	(Koler-Matznick et al. 2003)	344	2

Once the frequency modulation of the howls had been recorded, we compared every howl pairwise to generate a 2,005 x 2,005 matrix of howl similarity/dissimilarity. We used dynamic time warping (DTW) (Kruskal 1983) to deliver a quantitative metric of this distance (or dissimilarity) between every pair of howls. Dynamic time warping has been widely used for comparing frequency data such as these, particularly in the analysis of dolphin vocalisations (Buck & Tyack 1993; Deecke & Janik 2006; Sayigh et al. 2007). The DTW algorithm allows the time points of each sampled frequency measurement to vary freely, until an optimum match between the two curves is achieved. The amount

of time-axis distortion necessary to achieve this match is then taken as a measurement of the quantitative difference between the curves.

Using the dissimilarity matrix obtained by DTW, we applied the k-means unsupervised clustering algorithm to group the howls into natural clusters based on their similarity. We chose the appropriate number of clusters using a combination of cluster purity, measured as the mean cluster silhouette value (Rousseeuw 1987), and stability using a bootstrap technique; repeatedly applying the clustering to a random subset of 90% of the howls, and measuring similarity of the clustering results using normalised mutual information (Zhong & Ghosh 2005).

We then examined the usage of each of the different howl types by the 13 different species. As multiple recordings were obtained from the same individual, or from individuals within the same pack, potential issues of pseudoreplication arise if howls are directly compared to each other; howls from the same individual are likely to be more similar to each other than howls from separate individuals or from different packs. Therefore, rather than analysing howl difference distributions directly, we examined only differences in the use of different howl types, by calculating the proportion of howls $P(t)$ that belong to each howl type t , for each species: $P(t)=n(t)/N$, where $n(t)$ is the number of howls of type t for a particular species, and N is the total number of howls from that species. This provides a "fingerprint" of howl type usage, which can then be compared between species. We calculated the sum of squared differences Δ between the howl type histograms of different sources (packs) within each of the 13 species:

$$\Delta(a, b) = \sum_t^T (P_a(t) - P_b(t))^2$$

where $\Delta(a, b)$ is the sum of square differences between sources a and b , $P_a(t)$ and $P_b(t)$ are the incidences of howl type t in sources a and b respectively, and T is the total number of distinct howl types for this species.

To test the ability of the howl type usage fingerprint to identify canid species, we measured the sum of squared differences Δ between each source (pack) and the mean histograms of each of the 13 species (with the target source excluded), and recorded which species was most similar to the target source as

indicated by the lowest value of Δ . From this we constructed a confusion matrix showing the classification of each of the sources, whether to the correct species type or to an incorrect species. We then used an exact test (Fisher 1925) to estimate the significance of the similarity within a species. We randomised the howl type distributions 10^5 times within each species by reordering the incidences $P(t=1...T)$ randomly, and recalculated $\Delta'(a,b)$ to generate a null distribution of sum of squared differences. We then calculated the proportion of randomised differences Δ' that were less than the measured intra-species difference Δ . We also identified the most common howl type in each species and examined various exemplar howls of this type, as an illustration of what may be a typical howl type for this species.

Finally, we examined more closely the similarity in the howling behaviour of three sympatric species, the red wolf, eastern timber wolf, and coyote. Red wolves and coyotes hybridise in the wild, which poses a threat to reintroduction programs for the critically endangered red wolf (Hinton et al. 2013; Gese et al. 2015). The eastern timber wolf *C. l. lycaon*, is considered a subspecies of grey wolf, but whose taxonomic status is unclear, and is considered to be very closely related to *C. rufus*, if not conspecific (Wilson et al. 2000; Koblmuller et al. 2009; Chambers et al. 2012). We tested for significant differences between the howls of these three species, to determine whether howling behaviour may potentially provide a form of behavioural isolation, or alternatively encourage admixing and introgression. We reclustered the DTW data, using only howls from the red wolf, coyote, and eastern wolf. We then repeated the sum of square difference analysis, comparing the red wolf-coyote-eastern difference to a null distribution generated by randomising the order of the histogram of howl types, as well as comparing the histogram fingerprints between sources, as with the full data set.

3. RESULTS

Applying multidimensional scaling (Cox & Cox 2000) to the full 2,005 x 2,005 matrix of howl distances found 37 significant dimensions, which were then passed to the k-means clustering algorithm. Analysis of silhouette values in k-means led to 21 distinct clusters (howl types). Figure 2 shows the howl distance matrix reduced to two dimensions (for visualisation), with cluster assignment indicated. The clustering appeared robust; 99.3% of all howls were classified with posterior probability > 0.5. Bootstrapping and re-clustering with 90% of the data produced a normalised mutual information in comparison to the full data set of 0.760 ± 0.033 , i.e. 76% of the cluster assignment information was retained even when applying the algorithm to a reduced data set.

Figure 2. Multidimensional scaling of the 2,005 x 2,005 howl distance matrix into two dimensions. Each point is a howl, and points closer together are more similar than those further apart. Colours indicate k-means clustering assignment. The size of each point is for ease of visualisation only.

Within-species comparisons show that for the eastern timber wolf, the domestic dog, the coyote, the red wolf, the North African wolf *C.l. lupaster*, and the Arctic wolf *C. l. arctos*, howl type usage was more similar among sources of that species than would be expected by chance (Table 2). This indicates that in these species, the different sites from which recordings were taken showed a species-specific pattern of howl type usage.

232

233 Table 2. Exact test of similarity of howl type use within each species. The p-value represents the
 234 proportion of randomised trials where the mean difference between sources within a particular species
 235 was less than the actual mean difference within the species. Starred values are significant at 5%.

Species	p	Number of sources
Golden jackal	0.718	3
Coyote	0.019 *	4
Red wolf	0.007 *	4
Eastern Timber wolf	0.014 *	20
Mackenzie Valley wolf	0.955	8
Mexican wolf	0.891	2
Arctic wolf	0.006 *	7
European wolf	0.237	13
Iberian wolf	0.935	3
Indian wolf	0.144	7
North African wolf	<0.001 *	5
Domestic dog	0.003 *	53
New Guinea singing dog	0.899	2

236

237

238 The use of each howl type, adjusted for overall howl use frequency, for each of the species show
 239 species-specific fingerprints (Figure 3). The red wolf and coyote share howl type 3 as the most
 240 common; the European *C. l. lupus* and Iberian *C. l. signatus* wolves share type 5; and the Mackenzie
 241 Valley *C. l. occidentalis*, Indian *C. l. pallipes*, and Mexican *C. l. baileyi* wolves share type 7. Each
 242 other species has a distinct call type that is most commonly used, relative to its overall usage in the
 243 sample database. Apart from these distinctive howl types, the different species have different
 244 repertoire diversities, with for instance the North African wolf making use of many fewer howl types
 245 than the golden jackal, despite being represented by a similar overall number of sources and howls
 246 (Figure 3). One qualitative trend noticeable from the exemplar howls (chosen as those nearest to the
 247 cluster centroid) is that the smaller species (red wolf, coyote, New Guinea singing dog, domestic dog,
 248 golden jackal) favoured howls that ended with a sharp drop in frequency, whereas larger species
 249 (arctic wolf, eastern timber wolf, European wolf, Mackenzie Valley wolf) used howls with much less
 250 frequency modulation, particularly at the end of the howl (Figure 4), although this may be an artefact
 251 of the lower fundamental frequency used by larger species.

253

254 Figure 3. Howl use histograms for each of the 13 species, showing the relative use of each of the 21
 255 howl types, adjusted for overall howl type frequency. Red bars show the most commonly distinctive
 256 howl type for each species, with the index number of that type appearing above each histogram. N
 257 indicates the number of howls, and S indicates the number of sources.

258

259

Figure 4. Three examples of the howls of the particular howl types identified as characteristic of each species, and represented in Figure 2 by the red bars. Note that the howls within a type are similar in a dynamic time warping sense, although they may vary somewhat in length.

The confusion matrix for the identification of species by source, and the results of the species identification assessment (Table 3) shows that the coyote, Arctic wolf, and North African wolf all were well identified by howl usage fingerprint comparison, with identification of the red wolf and Mackenzie Valley wolf also higher than expected. The New Guinea singing dog *C. l. hallstromi*, domestic dog, golden jackal, and North African wolf appeared to form a cluster of similar howl usage types, and the coyote and red wolf seem to form a separate cluster, with heavy use of type 15 howls (which only seem to be used by 3 other species, and at very low frequency).

Table 3a. Classification success by comparing howl type usage histograms as fingerprints. The % correct column indicates how many recording sources (animal packs) were correctly identified as their particular species when compared to all other sources in the database. The Best guess column indicates which species were most frequently identified as the most similar species to the target source.

Species	% correct	Best guess
Golden jackal	33.3	Domestic dog
Coyote	50	Coyote
Red wolf	25	Red wolf, Coyote, Domestic dog, Arctic
Eastern Timber	5	Arctic
Mackenzie Valley	25	Mackenzie Valley, Indian
Mexican	0	Red wolf
Arctic	57.1	Arctic
European	0	Mackenzie Valley
Iberian	0	Eastern Timber , European , Mackenzie Valley
Indian	0	Mackenzie Valley
North African	40	North African
Domestic dog	13.2	North African
New Guinea Singing Dog	0	Domestic dog, North African

Table 3b. Confusion matrix showing the number of sources identified as each species type.

	Predicted species												
	RW	COY	NGSD	DD	GJ	ARC	ETIM	EUR	IBER	MV	MEX	NAFR	IND
Actual species	RW	1	1	1		1							
	COY	1	2								1		
	NGSD			1								1	
	DD	3	3	1	7	4	3	8	3	2	2	9	8
	GJ			2	1								
	ARC		1	1		4							1
	ETIM	3				5	1	1	3	4			3
	EUR		2		2	2			2	3	1		1
	IBER						1	1		1			
	MV	1		1			1	1		2			2
	MEX	1							1				
	NAFR	1		1	1							2	
	IND		1		1			1		4			

281

282 In the reduced analysis of just red wolf, coyote, and eastern timber wolf, there were a total of 776
 283 howls, 510 eastern timber wolf, 187 coyote, and 79 red wolf. Applying multidimensional scaling to
 284 the full 776 x 776 matrix led to 42 significant dimensions, and 11 k-means clusters. All howls (100%)
 285 were classified with posterior probability > 0.5 , and bootstrapping followed by reclustering led to a
 286 normalised mutual information of 0.706 ± 0.059 . With these data, the red wolf and coyote also
 287 showed significant similarity between the different sources (packs) of the same species ($p=0.006$ and
 288 $p=0.009$ respectively), whereas the eastern timber wolf was only marginally significant ($p=0.052$).
 289 Comparison of the howl type fingerprints showed that the coyote was well identified from most
 290 sources (3 out of 4 sources correctly identified), whereas the red wolf and eastern timber wolf were
 291 often misclassified one as the other, with the red wolf identified as eastern timber wolf in 2 out of 4
 292 sources, and eastern timber wolf as red wolf in 6 out of 20 sources. Red wolves and coyotes share
 293 their most common howl type – type 3 – which is rarely used by timber wolves. Red wolves will often
 294 use howl type 6, which coyotes and timber wolves rarely use, and may be intermediate in
 295 characteristics between coyote (type 3) and timber wolf (type 11) howls, by being lower in frequency
 296 and less frequency-modulated (Figure 5).

297

298

299 Figure 5. Examples of coyote howls of type 3 (left), red wolf howls of type 6 (middle) and eastern
 300 timber wolf howls of type 11 (right). Type 6 howls are rarely used by coyotes and eastern timber
 301 wolves, but commonly used by red wolves, and may represent an intermediate form.

302

303

4. DISCUSSION

In this study we analysed over 2,000 howls from 13 different species and subspecies belonging to the genus *Canis* around the world. Using dynamic time warping as a quantitative measure of howl dissimilarity, we applied an objective unsupervised clustering algorithm to group the howls into distinct howl types. The k-means algorithm produced 21 clusters that were stable to bootstrapping, and that probably represent genuine howl type categories, which we define without the need for subjective description of howl characteristics.

Each population recorded made different use of these 21 howl types, with many species/subspecies having a particular howl type that was characteristic of that species/subspecies. Within six of the species - the eastern timber wolf, the domestic dog, the coyote, the red wolf, the North African wolf, and the Arctic wolf - a statistically significant similarity existed in their howl type usage. Further, we found that individual populations of five of the species - coyote, red wolf, Arctic wolf, North African wolf, and Mackenzie Valley wolf - could be identified using the howl type histograms of the remaining populations in the data set.

In general, we conclude that canid howling is not an arbitrary signal, but possesses species-specific information, which may reflect adaptive and/or neutral processes of isolation.

We also performed a more detailed analysis of the howls of three American canids - the red wolf, the coyote, and the eastern timber wolf - because of the conservation importance of hybridisation between the critically endangered red wolf and the coyote, as well as continuing disagreement over the phylogenetic relationship between the red wolf and both the coyote and eastern timber wolf. We found that coyote and red wolf howl type usage differs significantly, which could be a useful tool for managing red wolf conservation in the face of competition from sympatric coyotes. Red wolf howling was similar to that of eastern timber wolves, further complicating the challenge of red wolf introduction both at the southern end of its range (coyotes) and at the northern end (eastern). In contrast, red wolves and coyotes share their most common howl type, whereas red wolves will often use howl type 6, which coyotes and timber wolves almost never use. The intermediate nature of howl type 6 may provide potential evidence of hybridization between these sepecies.

We note in passing that the smaller species - the red wolf, domestic dog, New Guinea singing dog, golden jackal - show a greater diversity of howl types than the larger species, and are similar to each other in their howl type usage. We lack sufficient data to examine this further; however, this phenomenon could be due to peculiarity of the habitat or niche of these smaller species, or could be due to a different emphasis on long and short range communication between larger and smaller species, differences in the vocal production mechanism in different sized species, or a different emphasis on the social role of howling.

Given the diverse and non-arbitrary nature of howl differences, it is natural to ask whether variations in howl structure reflect referential or context-specific information. Early studies of wolf communication pointed out that different vocalisation types (e.g. howl vs. growl, yelp, etc.) were associated with different behavioural contexts (Cohen & Fox 1976; Tembrock 1976), but stopped short of suggesting that particular features within howls themselves represented certain arousal states or environmental contexts (Theberge & Falls 1967; Lehner 1978). More recent studies have begun to address this question in dingoes (Déaux & Clarke 2013), as well as dogs (Faragó et al. 2014), and there is some evidence that vocal communication may be used in *Canis* to coordinate hunting activity (Muntz & Patterson 2004). Experimentally, it has been shown that howl modulation patterns convey individual identity, and that animals attend to this information (Palacios et al. 2015). Thus, individual identity in howl structure is more than just an epiphenomenon, and may be of relevance to conservation and management programs (Llaneza et al. 2005; Terry et al. 2005; Brennan et al. 2013; Hansen et al. 2015). Depredation of livestock by coyotes (Knowlton et al. 1999) and wolves (Sillero-Zubiri & Laurenson 2001), in particular, is a cause for concern, but attempts to use vocalisation playbacks as active deterrents have largely been unsuccessful (Gable 2010).

Our results have shown clear differences in howl structure between populations. Whether populations in geographical proximity represent separate species, subspecies, or otherwise, it seems clear that distinct ecotypes exist. The presence of discrete differences in vocal behaviour suggests that consideration should be given to conservation of populations such as *C. rufus* and *C. lupus lycaon*, even if genetic isolation does not exist. Recent studies have shown multiple examples of dialects not just in birdsong (Kroodsma 2004), but also in multiple mammalian taxa including rodents

(Slobodchikoff & Coast 1980; Gannon & Lawlor 1989), primates (de la Torre & Snowdon 2009; Thinh et al. 2011; Meyer et al. 2012), and hyraxes (Kershenbaum et al. 2012). Our study adds to recent work showing dialectic differences between the howls of wolves in Europe and North America (Palacios et al. 2007), and fits into an increasingly important trend of understanding the proximal causes and ultimate significance of dialectic variation (Lameira et al. 2010).

In the case of the critically endangered red wolf, hybridisation with coyotes represents the largest threat to reintroduced populations (Hinton et al. 2013; Gese et al. 2015). Although howling behaviour has long been identified in *Canis* as a mechanism for separating competing populations (Harrington & Mech 1979; Jaeger et al. 1996; Gese 2001), and vocal behaviour as a mechanism for genetic isolation in other mammalian taxa (Braune et al. 2008), to our knowledge no studies have addressed the question whether vocal differences can act to reduce interspecific hybridisation in *Canis*, or may in fact be the result of past hybridisation. Coyotes fail to respond to stimuli of wolf howling (Petroelje et al. 2013); detailed analysis of *C. rufus* recordings have uncovered non-howl vocalisations that have not been reported in *C. latrans* (Schneider & Anderson 2011); and the behavioural responses of individual wolves vary according to the familiarity of playback howls (Palacios et al. 2015). All these findings raise the possibility that vocal differences between *C. rufus* and *C. latrans* may have conservation significance. Our work adds to this body of evidence, and should encourage further investigation of the possibility of behavioural isolation between these populations.

Our study made use of data sources of widely varying size and quality - something inevitable when integrating recordings from around the world and from species of greatly varying abundance. We have endeavoured to minimise the statistical artefacts arising from this imbalance, and have been careful to use the recording source (essentially, a single pack) as the unit of comparison. Some pseudo-replication may remain, as we cannot ensure that the proportion of howls in each type is constant for a species. However, in most cases there are insufficient howls from specific individuals to look at how the pattern varies by individual within species. In addition, it is possible that differences between populations reflect differences in contextual stimuli. Despite these statistical limitations, we believe that such broad comparative studies have great value in understanding behaviour across a

wider taxonomic basis than just the species, and we hope that this utility compensates somewhat for the patchy nature of the data sources.

Automatic clustering using unsupervised algorithms is potentially problematic, as the presence of computer-identified clusters does not guarantee that these elements have cognitive significance for the animals involved. Indeed, we have no mechanistic indication that canids perceive and compare howls in a way similar to our dynamic time warping. To date, what we know is that wolves detect changes in the fundamental frequency of howls outside their natural range of variability, and changes in the frequency modulation pattern of howls (Palacios et al. 2015). However we feel confident that DTW provides a useful comparative tool, because consideration of acoustic propagation would indicate that frequency modulation of howls is likely an important encoding technique in long-range communication. Also, we took care to evaluate our clustering results using multiple metrics, and assessing their stability in the face of bootstrapping, to maximise confidence that the howl type partitions did, in fact, represent a division of howls into realistic howl types.

This study has involved only correlative analyses, but we believe that this kind of quantitative categorisation of vocalisation types is necessary before carrying out manipulative and playback experiments. Being armed with an objective set of howl types, or a methodology for arriving at such a definition, allows researchers to test the cognitive significance of different howl compositions, and look for potential behavioural correlates, such as territorial advertising and group cohesion. Any experimental work with critically endangered species such as the red wolf can be problematic, but we hope that with a firmer understanding of the vocal behaviour of these animals, it will be possible to design experiments that will benefit the conservation and management of this and other species.

5. CONCLUSIONS

Howling is a social communication process that is likely of major importance in the overall behaviour of all canid species. A deeper understanding of their social behaviour is not possible without a framework within which to understand their vocal behaviour. In particular, quantitative and objective

assessment of howling is highly preferable to subjective interpretation by humans, who lack the auditory and cognitive instruments of the focal animals. Techniques such as ours providing quantitative comparisons are important for any future experiments to investigate functionally referential elements to the canid howl repertoire, which would be a highly significant finding for two reasons. Firstly, the role of vocal communication in mediating social behaviour in canids may contribute to understanding the evolution of human language (Seyfarth & Cheney 2014). To our knowledge, no animal species other than humans possess any form of true language, not even any form of "proto-language". Therefore, it has been problematic to explain the evolution of human language as a continual progression from "non-language" to "language", through increasing adaptive advantage at each step (Tomasello 2008). The presence of complex referential communication in species that must cooperate to survive was probably a crucial step in the evolution of language (Jackendoff 1999). Understanding the communication systems of extant social species is essential to understanding the potential evolutionary trajectories to more complex communication that have occurred in the past, eventually leading to human linguistic abilities. Although taxonomically distant from early hominids, wolves and other canids show remarkable parallels with humans in, for example, social behaviour, intelligence, and vocal communication (Miklósi 2014). Further understanding of canid social communication could serve as a model system for the evolution of more complex vocal communication and language. Secondly, canid conservation and management can benefit from acoustic methods for surveying and assessing population size and health/genetic purity (Llaneza et al. 2005; Brennan et al. 2013), which can be difficult using traditional methods, particularly when snow is absent (Blanco & Cortés 2011). Active acoustic deterrence has also been suggested as a tool in the control of animal movements for mitigating wolf conflict with farmers (Gable 2010), but such techniques cannot be successfully implemented without understanding the message being transmitted. Therefore, clearer and quantitative techniques for the description of the long-range communication of canids could have multiple benefits for the conservation and management of these species: through understanding the processes controlling behavioural isolation, identifying populations by remote surveying, and active techniques to reduce human-animal conflict.

443

444 6. ACKNOWLEDGEMENTS

445 We thank Barbara Smuts for her helpful comments in the preparation of this manuscript and for the
446 provision of domestic dog recordings. AK thanks Sheela Hira and Knoxville Zoo, Laura Pearson and
447 Asheville Zoo, and Chris Lasher and North Carolina Zoo for access to record their red wolves, and
448 Arthur Povey for assistance coding the data. Recording work was approved by the Institutional
449 Animal Care and Use Committee of the University of Tennessee. AK is supported by a Herchel Smith
450 postdoctoral fellowship at the University of Cambridge. Part of this work was carried out while AK
451 was a Postdoctoral Fellow at the National Institute for Mathematical and Biological Synthesis, an
452 Institute sponsored by the National Science Foundation through NSF Award #DBI-1300426, with
453 additional support from The University of Tennessee, Knoxville. BH is thankful to the State Forest
454 Departments of Himachal Pradesh, J&K, and Maharashtra, and to various zoos in India for permitting
455 us to record howls. HRG is grateful to all who helped with the project: the staff at Colchester Zoo; the
456 Wildwood Trust, the Bior Laboratory of Bioacoustics; the British Library; Lupus Laetus; Polish
457 Mammal Research Institute; Tigress Productions; the BBC Natural History Unit; Longleat Safari
458 Park; Tierstimmen Archiv; Wild Sweden; Wolf Park; the Macaulay Sound Library and the UK Wolf
459 Conservation Trust; and Mike Collins, Teresa Palmer, Monty Sloan, Karl-Heinz Frommolt, Yorgos
460 Iliopoulos, Christine Anhalt, Louise Gentle, Richard Yarnell, Victoria Allison Hughes and Susan
461 Parks. BRM thanks the USDA/APHIS/WS/National Wildlife Research Center for supporting his
462 doctoral research and providing access to captive coyotes; recording work was approved by the
463 NWRC IACUC. SW thanks Mariana Olsen for assistance with data collection, and Yellowstone
464 National Park for permission to record.

465

466

- 468 **Bardeleben, C., Moore, R. L. & Wayne, R. K.** 2005. A molecular phylogeny of the Canidae based
469 on six nuclear loci. *Molecular Phylogenetics and Evolution*, **37**, 815-831.
- 470 **Bekoff, M.** 1977. *Canis latrans*. *Mammalian species*, **79**, 1-9.
- 471 **Bekoff, M., Diamond, J. & Mitton, J. B.** 1981. Life-history patterns and sociality in canids: body
472 size, reproduction, and behavior. *Oecologia*, **50**, 386-390.
- 473 **Beschta, R. L. & Ripple, W. J.** 2009. Large predators and trophic cascades in terrestrial ecosystems
474 of the western United States. *Biological Conservation*, **142**, 2401-2414.
- 475 **Blanco, J. C. & Cortés, Y.** 2011. Surveying wolves without snow: a critical review of the methods
476 used in Spain. *Hystrix, the Italian Journal of Mammalogy*, **23**, 35-48.
- 477 **Braune, P., Schmidt, S. & Zimmermann, E.** 2008. Acoustic divergence in the communication of
478 cryptic species of nocturnal primates (*Microcebus ssp.*). *BMC Biology*, **6**, 19-7007-6-19. doi:
479 10.1186/1741-7007-6-19.
- 480 **Brennan, A., Cross, P. C., Ausband, D. E., Barbknecht, A. & Creel, S.** 2013. Testing automated
481 howling devices in a wintertime wolf survey. *Wildlife Society Bulletin*, **37**, 389-393.
- 482 **Buck, J. R. & Tyack, P. L.** 1993. A quantitative measure of similarity for *Tursiops truncatus*
483 signature whistles. *The Journal of the Acoustical Society of America*, **94**, 2497-2506.
- 484 **Chambers, S. M., Fain, S. R., Fazio, B. & Amaral, M.** 2012. An account of the taxonomy of North
485 American wolves from morphological and genetic analyses. *North American Fauna*, **77**, 1-67.
- 486 **Cohen, J. & Fox, M.** 1976. Vocalizations in wild canids and possible effects of domestication.
487 *Behavioural Processes*, **1**, 77-92.
- 488 **Cox, T. F. & Cox, M. A. A.** 2000. *Multidimensional Scaling*. Berlin: Springer.
- 489 **Davlin, S. L. & VonVille, H. M.** 2012. Canine rabies vaccination and domestic dog population
490 characteristics in the developing world: a systematic review. *Vaccine*, **30**, 3492-3502.
- 491 **de la Torre, S. & Snowdon, C. T.** 2009. Dialects in pygmy marmosets? Population variation in call
492 structure. *American Journal of Primatology*, **71**, 333-342. doi: 10.1002/ajp.20657.
- 493 **Déaux, É C. & Clarke, J. A.** 2013. Dingo (*Canis lupus dingo*) acoustic repertoire: form and contexts.
494 *Behaviour*, **150**, 75-101.
- 495 **Deecke, V. B. & Janik, V. M.** 2006. Automated categorization of bioacoustic signals: avoiding
496 perceptual pitfalls. *The Journal of the Acoustical Society of America*, **119**, 645-653.
- 497 **Elmhagen, B. & Rushton, S. P.** 2007. Trophic control of mesopredators in terrestrial ecosystems:
498 top-down or bottom-up? *Ecology Letters*, **10**, 197-206.
- 499 **Faragó, T., Townsend, S. & Range, F.** 2014. The information content of wolf (and dog) social
500 communication. In: *Biocommunication of Animals* (Ed. by G. Witzany), pp. 41-62. Springer.

501 **Fisher, R. A.** 1925. *Statistical Methods for Research Workers*. Edinburgh: Genesis Publishing Pvt
502 Ltd.

503 **Gable, E.** 2010. Wolves: 'Howl boxes' could provide new tool to reduce conflicts.
504 <http://www.eenews.net/stories/93834>.

505 **Gannon, W. L. & Lawlor, T. E.** 1989. Variation of the chip vocalization of three species of
506 Townsend chipmunks (genus *Eutamias*). *Journal of Mammalogy*, **70**, 740-753.

507 **Gese, E. M.** 2001. Territorial defense by coyotes (*Canis latrans*) in Yellowstone National Park,
508 Wyoming: who, how, where, when, and why. *Canadian Journal of Zoology*, **79**, 980-987.

509 **Gese, E. M., Knowlton, F. F., Adams, J. R., Beck, K., Fuller, T. K., Murray, D. L., Steury, T. D.,**
510 **Stoskopf, M. K., Waddell, W. T. & Waits, L. P.** 2015. Managing hybridization of a recovering
511 endangered species: The red wolf *Canis rufus* as a case study. *Current Zoology*, **61**, 191-205.

512 **Hallberg, K. I.** 2007. Information in a long-distance vocal signal: Chorus howling in the coyote
513 (*Canis latrans*). PhD thesis, The Ohio State University.

514 **Hansen, S. J., Frair, J. L., Underwood, H. B. & Gibbs, J. P.** 2015. Pairing call-response surveys
515 and distance sampling for a mammalian carnivore. *The Journal of Wildlife Management*, **79**, 662-671.

516 **Harrington, F. H., Asa, C. S., Mech, L. & Boitani, L.** 2003. Wolf communication. In: *Wolves:*
517 *Behavior, Ecology, and Conservation* (Ed. by L. D. Mech & L. Boitani), pp. 66-103.
518 Chicago:University of Chicago Press.

519 **Harrington, F. H.** 1987. Aggressive howling in wolves. *Animal Behaviour*, **35**, 7-12.

520 **Harrington, F. H. & Mech, L. D.** 1979. Wolf howling and its role in territory maintenance.
521 *Behaviour*, **68**, 207-249. doi: 10.1163/156853979X00322.

522 **Henry, K. S. & Lucas, J. R.** 2010. Habitat-related differences in the frequency selectivity of auditory
523 filters in songbirds. *Functional Ecology*, **24**, 614-624.

524 **Hinton, J. W., Chamberlain, M. J. & Rabon, D. R.** 2013. Red Wolf (*Canis rufus*) Recovery: A
525 Review with Suggestions for Future Research. *Animals*, **3**, 722-744.

526 **Jackendoff, R.** 1999. Possible stages in the evolution of the language capacity. *Trends in Cognitive*
527 *Sciences*, **3**, 272-279.

528 **Jaeger, M. M., Pandit, R. K. & Haque, E.** 1996. Seasonal differences in territorial behavior by
529 golden jackals in Bangladesh: howling versus confrontation. *Journal of Mammalogy*, **77**, 768-775.

530 **Janik, V. M. & Slater, P. J. B.** 1998. Context-specific use suggests that bottlenose dolphin signature
531 whistles are cohesion calls. *Animal Behaviour*, **56**, 829-838.

532 **Jin, D. Z. & Kozhevnikov, A. A.** 2011. A compact statistical model of the song syntax in Bengalese
533 finch. *PLoS Computational Biology*, **7**, e1001108.

534 **Kershenbaum, A. & Roch, M. A.** 2013. An image processing based paradigm for the extraction of
535 tonal sounds in cetacean communications. *Journal of the Acoustical Society of America*, **134**, 4435-
536 4445.

- 537 **Kershenbaum, A., Sayigh, L. S. & Janik, V. M.** 2013. The encoding of individual identity in
538 dolphin signature whistles: how much information is needed? PLoS One, **8**, e77671. doi:
539 10.1371/journal.pone.0077671.
- 540 **Kershenbaum, A., Ilany, A., Blaustein, L. & Geffen, E.** 2012. Syntactic structure and geographical
541 dialects in the songs of male rock hyraxes. Proceedings of the Royal Society B: Biological Sciences,
542 **279**, 2974-2981.
- 543 **Kershenbaum, A., Blumstein, D. T., Roch, M. A., Akçay, Ç., Backus, G., Bee, M. A., Bohn, K.,**
544 **Cao, Y., Carter, G. & Căsar, C.** 2014. Acoustic sequences in non-human animals: a tutorial review
545 and prospectus. Biological Reviews, (**online before print**).
- 546 **Knowlton, F. F., Gese, E. M. & Jaeger, M. M.** 1999. Coyote depredation control: an interface
547 between biology and management. Journal of Range Management, **52**, 398-412.
- 548 **Koblmueller, S., Nord, M., Wayne, R. K. & Leonard, J. A.** 2009. Origin and status of the Great
549 Lakes wolf. Molecular Ecology, **18**, 2313-2326.
- 550 **Koepfli, K., Pollinger, J., Godinho, R., Robinson, J., Lea, A., Hendricks, S., Schweizer, R. M.,**
551 **Thalmann, O., Silva, P. & Fan, Z.** 2015. Genome-wide Evidence Reveals that African and Eurasian
552 Golden Jackals Are Distinct Species. Current Biology, (**online before print**).
- 553 **Koler-Matznick, J., Brisbin, I. L., Feinstein, M. & Bulmer, S.** 2003. An updated description of the
554 New Guinea singing dog (*Canis hallstromi*, Troughton 1957). Journal of Zoology, **261**, 109-118.
- 555 **Kroodtsma, D.** 2004. The diversity and plasticity of birdsong. In: *Nature's Music: The Science of*
556 *Birdsong* (Ed. by P. R. Marler & H. Slabbekoorn), San Diego, CA:Elsevier Academic Press.
- 557 **Kruskal, J. B.** 1983. An overview of sequence comparison: time warps, string edits, and
558 macromolecules. SIAM Review, **25**, 201-237.
- 559 **Lameira, A., Delgado, R. & Wich, S.** 2010. Review of geographic variation in terrestrial mammalian
560 acoustic signals: Human speech variation in a comparative perspective. Journal of Evolutionary
561 Psychology, **8**, 309-332.
- 562 **Lehner, P. N.** 1978. Coyote vocalizations: a lexicon and comparisons with other canids. Animal
563 Behaviour, **26**, 712-722.
- 564 **Llaneza, L., Ordiz, A., Palacios, V. & Uzal, A.** 2005. Monitoring wolf populations using howling
565 points combined with sign survey transects. Wildlife Biology in Practice, **1**, 108-117.
- 566 **Mazzini, F., Townsend, S. W., Virányi, Z. & Range, F.** 2013. Wolf howling is mediated by
567 relationship quality rather than underlying emotional stress. Current Biology, **23**, 1677-1680.
- 568 **McCarley, H.** 1975. Long-distance vocalizations of coyotes (*Canis latrans*). Journal of Mammalogy,
569 **56**, 847-856.
- 570 **Meyer, D., Hodges, J. K., Rinaldi, D., Wijaya, A., Roos, C. & Hammerschmidt, K.** 2012.
571 Acoustic structure of male loud-calls support molecular phylogeny of Sumatran and Javanese leaf
572 monkeys (genus *Presbytis*). BMC Evolutionary Biology, **12**, 16-2148-12-16. doi: 10.1186/1471-2148-
573 12-16.
- 574 **Miklósi, Á.** 2014. *Dog Behaviour, Evolution, and Cognition*. Oxford: Oxford University Press.

575 **Muntz, E. M. & Patterson, B. R.** 2004. Evidence for the use of vocalization to coordinate the killing
576 of a white-tailed deer, *Odocoileus virginianus*, by coyotes, *Canis latrans*. The Canadian Field-
577 Naturalist, **118**, 278-280.

578 **Nowak, R.** 1995. Another look at wolf taxonomy. Ecology and conservation of wolves in a changing
579 world, **375**, 357-389.

580 **Palacios, V., Font, E., Márquez, R. & Carazo, P.** 2015. Recognition of familiarity on the basis of
581 howls: a playback experiment in a captive group of wolves. Behaviour, **152**, 593-614.

582 **Palacios, V., Font, E. & Márquez, R.** 2007. Iberian wolf howls: acoustic structure, individual
583 variation, and a comparison with North American populations. Journal of Mammalogy, **88**, 606-613.
584 doi: 10.1644/06-MAMM-A-151R1.1.

585 **Paradiso, J. L. & Nowak, R. M.** 1972. *Canis rufus*. Mammalian species, **22**, 1-4.

586 **Petroelje, T. R., Belant, J. L. & Beyer Jr, D. E.** 2013. Factors affecting the elicitation of vocal
587 responses from coyotes *Canis latrans*. Wildlife Biology, **19**, 41-47.

588 **Quick, N. J. & Janik, V. M.** 2012. Bottlenose dolphins exchange signature whistles when meeting at
589 sea. Proceedings of the Royal Society B: Biological Sciences, **279**, 2539-2545.

590 **Root-Gutteridge, H., Bencsik, M., Chebli, M., Gentle, L. K., Terrell-Nield, C., Bourit, A. &**
591 **Yarnell, R. W.** 2014a. Identifying individual wild Eastern grey wolves (*Canis lupus lycaon*) using
592 fundamental frequency and amplitude of howls. Bioacoustics, **23**, 55-66.

593 **Root-Gutteridge, H., Bencsik, M., Chebli, M., Gentle, L. K., Terrell-Nield, C., Bourit, A. &**
594 **Yarnell, R. W.** 2014b. Improving individual identification in captive Eastern grey wolves (*Canis*
595 *lupus lycaon*) using the time course of howl amplitudes. Bioacoustics, **23**, 39-53.

596 **Rousseeuw, P. J.** 1987. Silhouettes: a graphical aid to the interpretation and validation of cluster
597 analysis. Journal of Computational and Applied Mathematics, **20**, 53-65.

598 **Rueness, E. K., Asmyhr, M. G., Sillero-Zubiri, C., Macdonald, D. W., Bekele, A., Atickem, A. &**
599 **Stenseth, N. C.** 2011. The cryptic African wolf: *Canis aureus lupaster* is not a golden jackal and is
600 not endemic to Egypt. PloS one, **6**, e16385. doi: 10.1371/journal.pone.0016385.

601 **Sayigh, L. S., Esch, H. C., Wells, R. S. & Janik, V. M.** 2007. Facts about signature whistles of
602 bottlenose dolphins, *Tursiops truncatus*. Animal Behaviour, **74**, 1631-1642.

603 **Schneider, J. N. & Anderson, R. E.** 2011. Tonal vocalizations in the red wolf (*Canis rufus*):
604 potential functions of nonlinear sound production. The Journal of the Acoustical Society of America,
605 **130**, 2275-2284.

606 **Seimenis, A.** 2008. The rabies situation in the Middle East. Developments in Biologicals, **131**, 43-53.

607 **Seyfarth, R. M. & Cheney, D. L.** 2014. The evolution of language from social cognition. Current
608 Opinion in Neurobiology, **28**, 5-9.

609 **Sillero-Zubiri, C. & Laurenson, M.** 2001. Interactions between carnivores and local communities:
610 Conflict or co-existence? In: *Carnivore Conservation* (Ed. by J. Gittleman, K. Funk, D. Macdonald &
611 R. Wayne), pp. 282-312. Press Syndicate of the University of Cambridge.

- 612 **Sillero-Zubiri, C. & Gottelli, D.** 1994. *Canis simensis*. Mammalian Species, **485**, 1-6.
- 613 **Siniscalchi, M., Quaranta, A. & Rogers, L. J.** 2008. Hemispheric specialization in dogs for
614 processing different acoustic stimuli.
- 615 **Slobodchikoff, C. & Coast, R.** 1980. Dialects in the alarm calls of prairie dogs. Behavioral Ecology
616 and Sociobiology, **7**, 49-53. doi: 10.1007/BF00302518.
- 617 **Tembrock, G.** 1976. Canid vocalizations. Behavioural Processes, **1**, 57-75.
- 618 **Terry, A. M., Peake, T. M. & McGregor, P. K.** 2005. The role of vocal individuality in
619 conservation. Frontiers in Zoology, **2**, 10.
- 620 **Theberge, J. B. & Falls, J. B.** 1967. Howling as a means of communication in timber wolves.
621 American Zoologist, **7**, 331-338.
- 622 **Thinh, V. N., Hallam, C., Roos, C. & Hammerschmidt, K.** 2011. Concordance between vocal and
623 genetic diversity in crested gibbons. BMC Evolutionary Biology, **11**, 36.
- 624 **Tomasello, M.** 2008. *Origins of Human Communication*. Cambridge, MA: MIT press.
- 625 **Tooze, Z., Harrington, F. & Fentress, J.** 1990. Individually distinct vocalizations in timber wolves,
626 *Canis lupus*. Animal Behaviour, **40**, 723-730.
- 627 **Vilà, C., Amorim, I. R., Leonard, J. A., Posada, D., Castroviejo, J., Petrucci-Fonseca, F.,**
628 **Crandall, K. A., Ellegren, H. & Wayne, R. K.** 1999. Mitochondrial DNA phylogeography and
629 population history of the grey wolf *Canis lupus*. Molecular Ecology, **8**, 2089-2103.
- 630 **Wiley, R. H. & Richards, D. G.** 1978. Physical constraints on acoustic communication in the
631 atmosphere: implications for the evolution of animal vocalizations. Behavioral Ecology and
632 Sociobiology, **3**, 69-94.
- 633 **Wilson, P. J., Grewal, S., Lawford, I. D., Heal, J. N., Granacki, A. G., Pennock, D., Theberge, J.**
634 **B., Theberge, M. T., Voigt, D. R. & Waddell, W.** 2000. DNA profiles of the eastern Canadian wolf
635 and the red wolf provide evidence for a common evolutionary history independent of the gray wolf.
636 Canadian Journal of Zoology, **78**, 2156-2166.
- 637 **Yin, S. & McCowan, B.** 2004. Barking in domestic dogs: context specificity and individual
638 identification. Animal Behaviour, **68**, 343-355.
- 639 **Zhong, S. & Ghosh, J.** 2005. Generative model-based document clustering: a comparative study.
640 Knowledge and Information Systems, **8**, 374-384.

641

642

643 FIGURE CAPTIONS

644

645 Figure 1. Example spectrogram of multiple wolves howling.

646

647 Figure 2. Multidimensional scaling of the 2,005 x 2,005 howl distance matrix into two dimensions.

648 Each point is a howl, and points closer together are more similar than those further apart. Colours

649 indicate k-means clustering assignment. The size of each point is for ease of visualisation only.

650

651 Figure 3. Howl use histograms for each of the 13 species, showing the relative use of each of the 21

652 howl types, adjusted for overall howl type frequency. Red bars show the most commonly distinctive

653 howl type for each species, with the index number of that type appearing above each histogram. N

654 indicates the number of howls, and S indicates the number of sources.

655

656 Figure 4. Three examples of the howls of the particular howl types identified as characteristic of each

657 species, and represented in Figure 2 by the red bars. Note that the howls within a type are similar in a

658 dynamic time warping sense, although they may vary somewhat in length.

659

660 Figure 5. Examples of coyote howls of type 3 (left), red wolf howls of type 6 (middle) and eastern

661 timber wolf howls of type 11 (right). Type 6 howls are rarely used by coyotes and eastern timber

662 wolves, but commonly used by red wolves, and may represent an intermediate form.

663

664

Graphical Abstract

(1) Identify multiple wild and captive canid populations from around the world

(2) Record howls and convert them frequency patterns

(4) Similarity in the use of howl types can reveal population processes

(3) Cluster the howls into similar modulation shapes, i.e. howl types