Design, implementation and interpretation of *in vitro* batch culture experiments to
assess enteric methane mitigation in ruminants – a review

Yáñez-Ruiz D.R.^{a*}, Bannink A.^b, Dijkstra J.^c, Kebreab E.^d, Morgavi D.P.^e, O’Kiely P.^f,
Reynolds C.K.^g, Schwarm A.^h, Shingfield K.J.^{ij}, Yu Z.^k, Hristov A.N.^l

^aEstación Experimental del Zaidín (CSIC), Profesor Albareda, 1, 18008, Granada, Spain

^bAnimal Nutrition, Wageningen UR Livestock Research, PO Box 65, Lelystad, The Netherlands

^cAnimal Nutrition Group, Wageningen University, PO Box 338, 6700 AH Wageningen, The
Netherlands

^dDepartment of Animal Science, University of California, Davis 95616, USA

^eINRA, UMR1213 Herbivores, F-63122 Saint-Genès-Champanelle, and Clermont Université,
VetAgro Sup, UMR Herbivores, BP 10448, F-63000 Clermont-Ferrand, France

^fTeagasc, Animal & Grassland Research and Innovation Centre, Grange, Dunsany, Co. Meath,
Ireland

^gThe University of Reading, Earley Gate, P.O. Box 237, Reading RG6 6AR, UK

^hETH Zurich, Institute of Agricultural Sciences, Animal Nutrition, 8092 Zurich, Switzerland

ⁱNatural Resources Institute Finland (Luke), FI-31600 Jokioinen, Finland

^jInstitute of Biological, Environmental and Rural Sciences, Aberystwyth University,
Aberystwyth, SY23 3EB, UK

^kDepartment of Animal Sciences, The Ohio State University, Columbus 43210, USA

^lDepartment of Animal Science, The Pennsylvania State University, University Park 16802,
USA

Abstract

In vitro fermentation techniques (IVFT) have been widely used to evaluate the nutritive value of feeds for ruminants and in the last decade to assess the effect of different nutritional strategies on methane production. However, many technical factors may influence the results obtained. The present review has been prepared by the ‘Global Network’ FACCE-JPI international research consortium to provide a critical evaluation of the main factors that need to be considered when designing, conducting and interpreting IVFT experiments investigate nutritional strategies to mitigate methane (CH₄) emission from ruminants. Given the increasing and wide-scale use of IVFT techniques, there is a need to critically review reports in the literature and establish what criteria are essential to the establishment and implementation of *in vitro* techniques. Key aspects considered include: i) donor animal species and number of animal used, ii) diet fed to donor animals, iii) collection and processing of rumen fluid as inoculum, iv) choice of substrate and incubation buffer, v) incubation procedures and CH₄ measurements, vi) headspace gas composition and vii) comparability of *in vitro* and *in vivo* measurements. Based on an evaluation of experimental evidence, a set of technical recommendations are presented that allowing the harmonization of IVFT laboratory methods employed with IVFT, of and procedures for feed evaluation, assessment of rumen function and CH₄ production.

Keywords: feed evaluation, *in vitro* gas production, methane, rumen, mitigation, microbial inoculum

Abbreviations: DM, dry matter; DMI, dry matter intake; IVFT, *in vitro* fermentation technique; NDF, neutral detergent fibre; OM, organic matter; VFA, volatile fatty acids.

1. Introduction

In vitro fermentation techniques (IVFT) that involve incubations of substrates with rumen fluid have been used extensively to evaluate the nutritive value of ruminant feeds. Measurements based on IVFT complement standard laboratory analysis of chemical composition and therefore offer a rapid and less expensive alternative to the determination of nutrient digestibility *in vivo* (Rymer et al., 2005). Also, application of the IVFT to reduce the use of experimental animals represents an advantage when large number of treatments needs to be tested. More recently, IVFT techniques have been used to assess the potential of diet, dietary ingredients and modifiers of rumen fermentation to decrease methane (CH₄) emissions from ruminant livestock (Bodas et al., 2008; Durmic et al. 2010). For many research groups with limited resources, the use of *in vitro* tools is often the only option available for investigating potential agents for CH₄ mitigation. Depending on the research question, *in vitro* studies can be valuable for screening and informing on the suitability for further evaluation *in vivo*. However, a positive outcome *in vitro* does not guarantee that the same treatment will have a similar effect *in vivo*. In some cases, IVFT results for feed evaluation and CH₄ mitigation can be misleading when the inherent characteristics of a batch culture system are not carefully considered (reviewed by Dijkstra et al., 2005). Furthermore, the goals, experimental design, results and conclusions of *in vitro* experiments require cautious and considered interpretation. Often the findings from *in vitro* studies have little relevance to commercial conditions, simply because in the amounts tested, an additive would be too expensive for use on-farm, efficacy cannot be confirmed *in vivo* or have detrimental effect on animal health and function.

The FACCE-JPI ‘Global Network’ project is an international initiative that intends, among other goals, to develop reliable and robust guidelines for generating and evaluating data from *in vitro* and *in vivo* experiments examining the potential to mitigate greenhouse gas (GHG) emissions from ruminant livestock systems. The present review is one output from the project that provides a critical evaluation of the key points for consideration when planning and performing *in vitro* studies and guidance for end users in the interpretation of experimental data from IVFT experiments. Major emphasis is placed on factors that influence microbial activity within *in vitro* systems typically used to assess CH₄ production, and how these can be balanced or accounted for, rather than simply providing a contrast and comparison of reports in the scientific literature. It is not intended to provide an in-depth and comprehensive description of different *in vitro* systems available, but rather provide an appraisal of key aspects that are central to undertaking robust, representative and reproducible *in vitro* experiments.

2. History and use of *in vitro* batch culture

Early *in vitro* studies focused on endpoint measurements such as the extent of substrate degradation (Tilley and Terry, 1963). In the 1970’s, researchers recognized that measurement of fermentation gases in combination with dietary chemical composition could be used to estimate both feed metabolisable energy content and ruminal organic matter degradability. Czerkawski and Breckenridge (1975) developed a system that involved recording the direct displacement of a piston by gases produced during the fermentation of feeds by rumen fluid in a glass syringe. This was the basis of the ‘Hohenheim Gas Test’ developed by Menke et al. (1979, Table 1). The ‘syringe technique’ was originally developed to determine end-point fermentation of feeds after

24 h of incubation. Blümmel and Ørskov (1993) modified the technique by incubating syringes in a water bath rather than a rotating incubator. By recording gas production at more frequent intervals, the kinetics of fermentation could also be determined.

Wilkins (1974) described a different approach to measure fermentation kinetics *in vitro*, whereby fermentation took place in a sealed vessel containing rumen fluid, buffer and substrate, and a pressure transducer was used to measure gas accumulation in the vessel headspace. In the simplest setup of this system, headspace pressure is measured manually, as described by Theodorou et al. (1994, Table 1), while gas samples are collected for the analysis of CO₂, CH₄ and/or H₂ concentrations when gas pressure is released (Tekippe et al., 2013). In the 1990's the first automated pressure based systems were developed (Pell and Schofield, 1993, Table 1), providing real-time measurements of gas accumulation allowing for a better understanding of the kinetics of fermentation for a range of substrates (Groot et al., 1996). During the development of these systems, it became increasingly clear that increased pressure within the fermentation container could affect fermentation end-products (Jouany and Lassalas, 2002) and the rate and extent of fermentation (Tagliapietra et al., 2010). More advanced systems periodically release and collect the gas via a solenoid valve (Cone et al., 1996; Davies et al., 2000, Table 1) thereby avoiding the build-up of pressure. Even in automated systems, analysis of gas composition (e.g. CH₄) typically requires manual injection of sample gases into a gas analyzer (Martínez et al., 2010; Pellikaan et al., 2011).

Cornou et al. (2013) described the results of a ring-test evaluating the use of a wireless system for automated gas release developed by Ankom (Ankom Technology, Macedon, NY, USA). This system is being used in various laboratories, but still relies on manual gas sampling and analysis.

More recently Muetzel et al. (2014) developed an automated gas measurement system by which gas production is monitored in real-time via pressure sensors and the proportion of CH₄ and H₂ in the vented fermentation gases is measured automatically by gas chromatography. The main difference with previous automated systems (Cornou et al., 2013) is that fermentation gases are collected and analysed by a computer-controlled gas chromatograph, rather than being released into the air once a threshold pressure is reached.

In vitro gas production systems have been used extensively for rapid screening of chemical substances, plant species, plant extracts and dietary ingredients on CH₄ emissions from rumen fermentation. Such experimental approaches have allowed the mode of action of a range of chemicals (Busquet et al., 2005; Bodas et al., 2008; Garcia-Gonzalez et al., 2008; Durmic et al., 2010) and dietary substrates (Patra and Yu, 2013; Hatew et al., 2015) to be investigated. Use of IVFT offers the opportunity to evaluate a broad spectrum of chemical agents alone or in a number of combinations over a wide range of concentrations (e.g., Busquet et al., 2005; Garcia-Gonzalez et al., 2008). However, this technique does not generate reliable information for agents that are only effective for decreasing CH₄ emissions over an extended period (Castro-Montoya et al., 2015). Furthermore, results from screening studies (Bodas et al., 2008; Durmic et al., 2010) are often inconclusive and may be conflicting due to variation in dosage, chemical structure of the test substance or compound, diet, combination of treatments applied, adaptation of rumen microbes or the form in which an agent is introduced into the system (Cardozo et al., 2004, 2005). Substantial decreases of CH₄ production *in vitro* (Tan et al., 2011) have been reported, but in several cases these have been accompanied by adverse effects on feed degradation, with the implication that a similar effect may occur *in vivo* which would compromise diet digestibility and animal

performance.

Please insert Table 1 around here

3. Aspects to consider

3.1. Donor animal species and animal numbers

Different animal species (sheep, goats, cattle and buffaloes) may vary in their response to the same CH₄ mitigation strategy. The obvious recommendation is to use the same species as donors of rumen fluid for *in vitro* incubations as the intended target species. However, this is not always possible, and due to cost small ruminants are often used as donors of rumen contents, even when cattle are the target species. One key question is, therefore, whether sheep or goats can be used as suitable surrogates for cattle for the study of CH₄ production *in vitro*.

Bueno et al. (1999) compared the microbial biomass in bovine and ovine rumen fluid and, although no intake data were reported, the inocula were adjusted to provide the same microbial biomass. It was concluded that rumen fluid from sheep could replace that from cattle or vice versa as rumen inoculum and that the two sources were comparable under tropical feeding conditions. Cone et al. (2002) reported a comparison of rumen samples collected in the same way from sheep and cattle maintained under similar conditions. Incubations of 22 different feeds were performed. A close association was observed ($r = 0.98$) for gas production at 24 (ranging from 100 to 325 ml/g OM incubated) and 48 (ranging from 150 to 350 ml/g OM) h for incubations with rumen fluid from cows and sheep. However, the relationship based on the rate of gas production (ml/g OM/h) was weaker ($r = 0.79$). Calabro et al (2005) compared rumen fluid from buffalo and sheep as a source of inoculum and observed higher fermentation rates and extent of degradation during incubations with rumen fluid from sheep.

Differences in fermentation kinetics were greater when fibre-rich substrates were tested, such as straw and hay, but negligible for barley grain.

Muetzel et al. (2014) compared rumen fluid from cattle (Holstein $\times$ Jersey cows) and sheep using a newly developed automated *in vitro* system and reported that total gas production is unaffected by donor animal species. Prior to rumen sampling, cattle and sheep had been adapted to a medium quality hay diet fed to meet maintenance energy requirements for 14 days. However, proportions of CH₄ in vented fermentation gases were lower during incubations with rumen fluid from sheep than cattle (150 vs. 158 ml/L, P=0.003) associated with a lower proportion of acetate and a higher proportion of propionate. There is no evidence of differences in methanogen communities in sheep and cattle (Jeyanathan et al., 2011), which suggests that the differences are driven by H₂ production from the bacterial or protozoal communities. No interaction between animal species (sheep vs. cattle) and the type of substrate incubated (chicory, lucerne, ryegrass, straw and white clover) was observed (Muetzel et al., 2014). Bueno et al. (2015) compared *in vitro* CH₄ production using rumen fluid from taurine dairy cattle (*Bos taurus taurus*), zebu beef cattle (*Bos taurus indicus*), water buffaloes (*Bubalus bubalis*), sheep (*Ovis aries*) and goats (*Capra hircus*) fed similar diets while testing the effect of condensed tannins from an Acacia extract. Rumen fluid from cattle resulted in higher CH₄ per unit of degraded organic matter (OM) formation than rumen fluid from small ruminants.

Although microbiota of ruminant species housed in close contact and fed a similar diet may be of a similar composition, the microbial ecology of rumen samples between sheep and goats, for example, may vary due to differences in dentition, eating and ruminating behaviour, digestive tract physiology and ruminal retention time (Ammar et al., 2004). For this reason, collection of inoculum from animals of the same target

species fed a diet containing the same feedstuffs would be recommended. Differences due to feeding behaviour and diet composition can to some extent be overcome by the collection of rumen samples before morning feeding, when the effect of diet composition on rumen metabolites or microbiota are likely to be minimized (Martinez et al., 2010) (see section 3.3).

There remains some uncertainty on the number of animals that need to be sampled to provide a representative sample of rumen inocula. Several studies using different ruminant species (e.g., Pinares-Patiño et al., 2003; Waghorn et al., 2006; Yan et al., 2006) have reported that CH₄ emissions per unit dry matter intake vary between individual animals. Such variation has been associated, among other factors, with differences in the rumen microbiome associated with between-animal variation in passage rates, rumen volume and morphology, eating behaviour, etc. (Kittelman et al., 2014). Martínez et al. (2010) observed consistent differences in CH₄ production and H₂ recovery *in vitro* during incubations with rumen liquor collected from 6 different sheep fed the same diet. Such differences can only be explained by differences in microbial populations or activities in the starting inocula or variation in the survival or activity of microbes over the incubation *period*. To reduce the effects of unusual rumen inocula, it has been suggested that, normally, at least 3 animal sources should be used to provide a representative source of rumen inoculum (Editorial, Animal Feed Science and Technology, 2012). However, this might be difficult when large ruminants are used. Another issue to consider is the number of independent incubation runs to be conducted in different days, which should be at least 3. Unfortunately, there is no work available assessing in statistical terms the minimum number of donor animals needed to provide a representative source of rumen inoculum.

RECOMMENDATION: Where possible the target animal species should be used as the donor of rumen fluid. Sampling before feeding is advantageous for minimizing diet by animal interactions. We recommend using 3 or more (optimum) or 2 (minimum) animals as donors of rumen inoculum and at least 3 independent incubation runs.

3.2. Donor animal diet

Diet composition and nutrient intake are major factors affecting both microbial populations in the rumen and microbial activity of rumen inoculum (Mould et al., 2005). Compared with ruminants fed high-concentrate diets, a greater proportion of fibrolytic bacteria and methanogenic archaea can be expected in rumen fluid collected from animals on high-forage diets (Demeyer and Fievez, 2000). However, the extent to which concentrate feeds affect rumen digestion and microbial populations may depend on the source and proportion of concentrate ingredients in the diet, as well as forage quality (Dijkstra, 1994).

Martínez et al. (2010) assessed the effect of feeding sheep diets differing in forage:concentrate ratios (F:C; 70:30 vs 30:70) and forage source (alfalfa hay vs grass hay) on rumen fermentation and CH₄ production. *In vitro* CH₄ production (per g of incubated DM) was increased by decreases in the F:C ratio of diets fed to donor animals or when Lucerne hay was replaced by grass hay. Differences in the F:C ratio altered pH and the activity of certain glycoside hydrolases (carboxymethylcellulase, xylanase and amylase) in rumen fluid. Forage type also influenced NH₃ content and carboxymethylcellulase activity in rumen fluid. These results suggest that feeding donor animals a diet similar to the substrate to be incubated *in vitro* may be advantageous.

Of particular importance is the observation that some mitigation strategies have been evaluated using rumen fluid from animals fed poor quality feeds. This raises the

question of whether outcomes on the efficacy of mitigation agents under these circumstances can be considered reliable when donor animals have not been fed a diet of similar characteristics as that offered to target animals. The effects of differences in diet composition fed to donor animals may be minimized by obtaining rumen fluid immediately before feeding. Huntington and Givens (1998) obtained fluid from cows fed either a silage:barley diet (80:20) or a barley straw diet. Although microbial activity of rumen fluid collected on the straw diet was lower than the silage:barley diet, this did not alter the gas production profile, which was attributed to sampling prior to morning feeding.

The diet fed to donor animals also needs to be considered when testing additives as CH₄ mitigation agents. Mateos et al. (2013) reported that the effect of garlic oil and cinnamaldehyde on *in vitro* fermentation and CH₄ production varied depending on whether the donor animals were fed a typical dairy diet (alfalfa hay:concentrate 50:50) or a fattening diet (barley straw: concentrate 15:85). Observations from several *in vitro* studies suggest that the effects of essential oils on rumen function are pH-dependent, and this also appears to be true for garlic oil and some of its components (Cardozo et al. 2005; Kamel et al. 2008). Cardozo et al. (2005) found that garlic oil had a more pronounced impact on rumen volatile fatty acids (VFA) profile at low compared with high rumen pH (5.5 versus 7.0), an effect explained by differences in the status of the active molecules (i.e. dissociated or un-dissociated) possibly mediated by changes in rumen pH. However, batch cultures are usually highly buffered systems allowing ruminal microorganisms to grow for a prolonged period despite the accumulation of fermentation end-products. In such cases, factors other than pH (e.g. microbial composition) may explain differences in the efficacy of additives. Hatew et al. (2015) provided further evidence on the importance of diet fed to the donor animal.

Experiments involved the incubation of the same substrate (grass silage or beet pulp) with rumen inoculum obtained from donor cows fed on diets that differed in starch source (native vs. gelatinized maize grain) and starch level (270 vs 530 g/kg concentrate DM). A higher level of starch and gelatinized rather than native maize were found to lower gas and CH₄ production after 24-h incubations.

Level of feed intake is also an important consideration, given that a higher DM intake (DMI) lowers retention time in the rumen, decreasing the amount of time available for feed degradation of feeds and hence ruminal digestibility (Clauss et al., 2007). Rumen pH, proteolytic and cellulolytic activities are thought to be influenced by the level of DMI which can in turn influence growth rates and the metabolic activity of inoculum used in *in vitro* systems. Increasing feeding frequency will generally lower diurnal variation in rumen fermentation parameters. For example, in lactating cows increasing feeding frequency from two to six times-daily was found to decrease post-feeding variation in rumen pH, osmolality, VFA and NH₃ concentrations (Le Liboux and Peyraud, 1999).

The period of adaptation to a given diet by the donor animal probably needs to be revisited. It is common to collect rumen fluid from animals fed a diet for 2 weeks. However, there are indications that the methanogenic archaeal population requires an adaptation period of around 30 days after a change in diet (Williams et al., 2009). Also Monteils et al., (2012) reported that protozoa counts in the rumen needed 25 days to stabilize after different dietary changes. Given the lower generation time that protozoa and archaea have as compared to bacteria (Dehority, 2003), further research is required to assess the effect of dietary treatments on the adaptation time needed for methanogenesis tests.

RECOMMENDATION: The diet fed to donor animals should be similar in composition to the substrate incubated in vitro. Care should be taken to ensure sufficient buffering capacity when investigating diets or dietary ingredients or additives that promote differences in rumen pH (see section 3.4), in particular when samples are not taken immediately before feeding. It is recommended that donor animals are fed a diet at a restricted level of feeding as frequent meals to ensure constancy of diet composition and digestion, minimize variation in feed intake and avoid diurnal variation in rumen fermentation.

3.3. Rumen fluid sampling: time, location and processing

In all *in vitro* fermentation systems it is essential to create an environment, which, for any set of parameters, mimics the fermentation in a specific section of the gastrointestinal tract *in vivo* (e.g. reticulo-rumen or caecum). Therefore, the inoculum should be representative of that environment with respect to both the composition and abundance of the microbial population. For *in vitro* systems to be robust (i.e., reproducible over time and representative of conditions *in vivo*), the inoculum must meet certain criteria. Making a valid assessment of whether a given study has met these criteria may be problematic, as often-essential information is not reported. Given the precision of gas release kinetic techniques relative to degradability at a set end-point over extended periods, variations in inoculum characteristics due to host animal effects, nutrition and sampling time, as well as sample preparation and inoculation, can have substantial cumulative effects on *in vitro* fermentation. It seems pertinent, not only to permit comparison between studies, but also to limit potential errors, to have a set of accepted guidelines and standard procedures for preparing inoculum for measuring CH₄ *in vitro*, as proposed for animal studies *in vivo*. Such guidelines should include host

animal management, sampling techniques (time, location, alternatives) and inoculum preparation.

Sampling time:

Diurnal changes of the rumen microbiome, both in terms of abundance and metabolic activity have been documented. Concentrations of viable microbial populations in the rumen typically decrease 4 h post-feeding, due to dilution with feed, water and saliva, and peaks at 6-12 h post feeding (depending on diet and level of feed intake) (Leedle et al., 1982; Dehority, 2003). Furthermore, microbial abundances were found to decline during the degradation of available nutrients. Cone et al. (1996) observed that the rate of fermentation was highest when rumen fluid was collected after morning feeding, although rumen sampling time had no effect on the total gas production. Menke and Steingass (1988) stated that sampling rumen contents just before feeding lowered variation in composition and activity of the inoculum and minimized the influence of diet fed to donor animals. However, Payne et al. (2002) observed that the total gas production from both starch and ground straw were less variable between replicate bottles and between weeks of collection when rumen fluid inoculum was collected either 4 or 8 h post feeding, compared with samples collected just before, or 2 h after feeding. Presumably, the activity of the inoculum is determined just as much by time of rumen sampling relative to feeding as by feeding pattern and eating time. Furthermore, microbial diversity and activity is at its lowest before feeding and while this reduces the variability, it may not reflect the ‘true’ effect that the feed/additive may have on much diverse population and their activities present after feeding. This deserves further research in the future.

RECOMMENDATION: There is no general recommendation on the ideal rumen sampling time as it depends on the objectives of a specific experiment. Given the difficulty of minimizing diurnal variation, rumen fluid samples are best collected immediately before feeding, based on a consistent protocol for dietary access by donor animals across experiments. When a series of studies are conducted over time, feeding and sampling procedures should be kept as identical as possible.

Sampling procedure: Rumen digesta is comprised of different fractions (large and small particulate matter and liquid). Ample evidence exists on the different abundance and diversity of bacteria associated with the liquid and solid ruminal contents (Pei et al., 2010). The pH and VFA concentration varies between different sections of the reticulo-rumen in cattle (Bryant 1964). De Visser et al. (1993) indicated that rumen VFA concentrations were about 20% higher for the entire evacuated rumen content compared with calculations based on samples of rumen fluid collected in a standardized manner. Storm and Kristensen (2010) indicated differences of 0.4 to 0.6 pH units and of 40 to 50 mM VFA between the central and ventral regions of the rumen, with the lowest pH and highest VFA concentrations in the medial rumen. Rumen fermentation parameters have also been found to differ between samples collected at different locations in the rumen (Shen et al., 2012), implying possible differences in microbial abundance and activity within the rumen.

Rumen cannulation is considered the reference method allowing the collection of representative samples of rumen digesta from donor animals (Komarek, 1981; Kristensen et al., 2010). Access to surgically-modified animals is not universal, and therefore less invasive techniques, such as oral stomach probing, have been used as an alternative. In the relatively few studies that have compared sampling through the

rumen cannula or by stomach probing, differences in fermentation profile and microbiota have been reported in some (e.g., Geishauser and Gitzel, 1996; Duffield et al., 2004), but not all cases (e.g., Lodge-Ivey et al., 2009; Shen et al., 2012; Terré et al., 2013). Part of the discrepancy between studies may reflect differences in the procedures used to avoid salivary dilution and contamination, the type of samples collected and rumen sampling site. Stomach probing results in the collection of samples containing a high proportion of liquid, whereas sampling via a rumen cannula allows both solid and liquid digesta fractions to be obtained. Differences in the methods used to collect rumen samples are of greater relevance when treatments are not expected to have the same effect on microbial populations attached to solids or inhabiting the liquid phase (Martínez et al., 2010). The study of Shen et al. (2012), attributed the differences between samples collected via cannula or stomach tube to rumen sampling site, as a consequence of the probe not being inserted to a depth sufficient to reach the ventral sac. Accurate probe insertion to a desired location within the rumen is extremely challenging in small ruminants. In a recent study, Ramos-Morales et al. (2014) found that stomach-tubing in sheep and goats detected the same differences in rumen fermentation due to species, diet or sampling time as sampling via the cannula. However, certain differences were more readily detected in rumen cannula samples, while substantial differences in the bacterial community structure were detected between the sampling methods.

Faeces has also been used as an alternative source of inoculum to rumen fluid (El Shaer et al., 1987). Cultures of ruminal or faecal microorganisms appear to result in similar fermentation processes (El-Meadaway et al., 1998). However, fewer microorganisms in faecal inocula may result in lower degradation capacity and decreased gas production (Cone et al., 2002; Váradyová, et al., 2005), a longer lag

phase and a slower rate of degradation at the outset (Mauricio et al., 2001). Furthermore, for poor-quality forages, there is only a weak relationship between gas production during incubations with inocula sourced from faeces and rumen fluid (El-Meadaway et al., 1998; Varadyova et al., 2005). Dhanoa et al. (2004) proposed a method allowing mathematical adjustments to convert or translate the degradation profiles produced by faecal inoculum to correspond with ruminal fluid, but this requires application of different prediction equations for each group of feeds incubated.

Rumen contents collected post-mortem at abattoir can also be used as an alternative to rumen fluid (Mould et al., 2005). Several IVFT have been performed using rumen fluid collected from slaughtered cattle, sheep, buffalo and dromedary (Haddi et al., 2003; Salem, 2005). To date, there are no reports directly comparing the use of rumen fluid from slaughtered animals with oral or rumen sampling in the same animal. Such an approach requires sampling of rumen contents soon after slaughter, as well as the same criteria for other sources of inocula being met. While the intake and diet composition of slaughtered animals are not known, access to entire rumen contents allows the collection of representative samples that can also be used to inform on nutrient supply. If sampling of rumen cannulated animals is not possible, collection of rumen content from slaughtered animals may prove a viable alternative.

RECOMMENDATION: To be as representative of the rumen environment as possible, samples of ruminal contents for the preparation of in vitro inoculum need to be collected from several locations. This is more feasible in large ruminants and requires the collection of rumen contents from animals fitted with rumen cannula following a clearly defined and standardized sampling protocol. Stomach tubing or faecal inoculae may serve as an alternative for ranking purposes, but quantitative data using these

427 *alternatives may differ from sampling of rumen contents in cannulated animals.*
428 *Stomach tubing should be performed by well-trained persons to minimize salivary*
429 *dilution.*

430

431 Preservation of inoculum: Anaerobiosis is essential to culture rumen microorganisms,
432 methanogenic archaea, in particular (Joblin, 2005). Certain experiments may require
433 rumen inoculum to be stored until culturing *in vitro*. Short-term storage (<1 h), for
434 example during transport from donor animals to the IVFT laboratory, should exclude
435 exposure to air. It is equally important that any increase in headspace pressure does not
436 cause CO₂ to go into solution, thus lowering pH. Excessive fermentation due to
437 extended storage at 39°C should be restricted to prevent any microbial group from
438 becoming dominant and modifying the composition of the original inoculum.

439 The influence of storage time and temperature on neutral detergent fibre (NDF)
440 degradation by rumen microorganisms has been investigated (Robinson et al., 1999).
441 Studies involved the use of the ANKOM end-point system to examine the effect of
442 delaying inoculum storage at 39°C by up to 6.5 h, or up to 48 h after storage over a
443 range of temperatures from – 22 and 39 °C. No apparent effect on 48 h end-point
444 degradation of medium-term (6.5 h) storage compared with long-term storage (48 h)
445 was identified. Authors concluded that no storage method, irrespective of temperature,
446 would maintain rumen inoculum activity for up to 48 h that support normal
447 fermentation *in vitro*. Subsequent works (Cone et al., 2000; Hervás et al., 2005; Prates
448 et al., 2010) presented data on the effect of using rumen fluid directly or stored
449 anaerobically at 39°C for increments of up to 24 h or at –24°C for 1, 3, 10, 40 or 76
450 days before use. In general, gas production in terms of kinetics and cumulative yields,
451 decreased as storage period increased. Final gas volumes were similar when rumen fluid

was stored up to 4-6 h compared with no prior storage. Gas production rates were lower for inocula stored for 8 or 24 h, while gas production was considerably decreased by extended storage at -20°C of more than 10 days. Microbial activity was lowered by freezing, with the decrease being substrate-dependent with the degradation of pure starch and cellulose being less affected compared with lucerne hay and barley straw. Freezing in liquid nitrogen is preferred over storage at -20°C (Prates et al., 2010) and thawing of small volumes (approximately 20 ml) at 39°C for 2 min. The time taken for freezing and thawing appears to be as equally important as storage temperature. Overall, studies suggest that preservation of rumen fluid at 0°C for up to 6 h offered a practical alternative, where necessary, to freshly collected inocula. Protozoa are lost after freezing which could have an impact on the fermentation. There are, however, no reports documenting the effect of different preservation methods on methanogenic activity.

RECOMMENDATION: Fresh rumen fluid maintained under anaerobic conditions at 39°C should be inoculated into in vitro vessels as soon as possible, ideally within 1 h post collection. When this is not possible, rumen fluid can be preserved at 0-4 °C for up to 6 h or frozen in liquid nitrogen following addition of a cryo-protectant (i.e. 15% glycerol) for longer periods for use as inoculum. In either case, implementation of standardized procedures to avoid undesirable variation in microbial activity is highly recommended.

Preparation of inoculum prior to incubation: Method of preparation also influences the microbial activity of rumen inoculum. The rumen microbiome consists of three sub-populations of microbes: those in the fluid phase, adherent to the particulate phase

(further divided into loosely- and firmly-associated with the feed particles) (Cheng et al., 1993) or attached to the epithelium (Sadet et al., 2007). The latter tends to be primarily involved in the release of ammonia from urea absorbed across the rumen epithelium and, as such, has only a minor role in feed degradation, and for this reason does not need to be sampled (Mueller et al., 1984). Fluid and particulate associated bacterial populations (Kim et al., 2004) and methanogenic archaea (Shin et al., 2004) differ in growth characteristics and in the activities of most enzymes (Moharrery and Das, 2001). Microorganisms in rumen liquid (20 to 30% of total microbes) including free-living bacteria and bacteria detached from solid substrate, have little direct involvement in structural carbohydrate digestion (Miron et al., 2001). Microbes attached to feed particles predominate (70 to 80% of microbial matter and microbial ATP production) and play a key role in feed particle digestion in the rumen (Miron et al., 2001; Tralbalza-Marinucci et al., 2006). Consequently, once samples of whole rumen contents have been collected, the problem arises of how to effectively detach the microorganisms associated with feed particles. Failure to do so will result in a high proportion remaining to be attached after filtration, while the use of multiple layers of cheesecloth, muslin or surgical gauze leads to different microbial fractions being retained and inoculated *in vitro*. For example, use of a cloth with 50 µm pores would substantially lower the number of large protozoa (which range from 15–250 µm in size, Dehority, 2003) in inocula. Williams and Coleman (1991) reported that most protozoa are retained using a 100 µm pore size filter. Therefore, if total protozoa are to be included in the microbial culture, a 250 µm pore size cloth should be used. Furthermore, some of the physical methods used to detach particle-associated microbes (e.g., a stomacher or maceration of rumen contents in a food processor) may also cause cell damage. Rymer et al. (1999) examined four methods of inoculum preparation:

strained, blended fluid, “stomacher” or strained plus blended residues. No consistent treatment effect was observed, apart from blending which tended to decrease substrate degradation. Authors concluded that there was little advantage from blending inoculum, particularly in light of the risk of exposing microorganisms to oxygen. Even though homogenizing rumen fluid may increase numbers of particle-associated bacteria in the inoculum, Pell and Schofield (1993) excluded this step on the grounds that it i) introduced an extra procedure into the laboratory protocol, ii) increased the risk of exposing rumen microorganisms to oxygen, iii) increased the quantity of gas released from the blanks, and iv) had no obvious effect on the results of IVFT.

A number of techniques that have been proposed for detaching microorganisms from rumen feed particles that involve various combinations of chemical and physical treatments and h yield removal rates of between 20 and 80% (Hristov et al., 1999; Ranilla et al., 2001). While the use of these techniques is essential when collecting microbial biomass to determine their composition to accurately assess passage of nutrients of microbial origin to the intestine, it is not clear whether a standard protocol for microbial detachment should be applied for *in vitro* methods. In recent work, Soto et al. (2013) studied the development of the microbiota in different *in vitro* rumen simulation systems inoculated with intact or filtered rumen fluid from goats. Incubation of filtered rumen fluid fraction in batch culture resulted in lower microbial diversity compared with non-filtered rumen fluid inoculum. Substantial growth of fibrolytic bacteria and methanogenic archaea over the first 24 h partially compensated for low numbers in the inoculum due to filtering.

RECOMMENDATION: Filtration of rumen fluid using the same pore size across incubation runs is a straightforward method for preparing inoculum suitable for in vitro

experiments. A larger pore size results in greater numbers of small particulate associated bacteria and protozoa. A pore size of 250 μm is recommended. Use of multiple layers of cheesecloth is not recommended due to inconsistencies in pore size.

3.4. Substrate and incubation buffer

Substrate:

Provided donor animals are fed the same or a similar diet as that tested *in vitro*, there remains an uncertainty of the choice of substrate to be used in the evaluation of additives on CH_4 production *in vitro*. The composition of diets fermented *in vitro* determines the production of dissolved H_2 that serves as a substrate for methanogens. Most *in vitro* studies have tested additives in incubations with a single substrate, but there are reports on the effects of additives using different fermentation substrates in a single experiment. Machmüller et al. (2001) investigated effects of medium chain fatty acids in incubations containing high or low amounts of fiber. The efficacy of monensin has been examined using corn meal or timothy hay (Russell and Strobel, 1988) and corn meal or soyabean hulls (Pellikaan et al., 2011) as substrates. In certain instances a substrate by additive interaction has been observed. Pellikaan et al. (2011) reported a complete inhibition of CH_4 production during the first 30 h of incubation, irrespective of substrate composition (soyhulls or maize), with bromoethanesulphonate (BES) and cinnamaldehyde. However, CH_4 production from soybean hulls with BES or cinnamaldehyde was 65% lower compared with soyhulls without additive after 72 h, suggesting an influence on adaptation, whereas no CH_4 was produced after 72 h production when maize was incubated with these additives. Subsequent studies covering a range of plant extracts and fatty acids (Castro-Montoya et al., 2012; Klevenhusen et al., 2012; O'Brien et al., 2013) have provided further evidence that responses of CH_4 to

a given additive differs depending on the feed substrate incubated. For example, addition of fatty acids were found more effective in lowering CH₄ during incubations containing higher proportions of concentrate ingredients, an effect attributed to greater protonation of fatty acids at a lower pH (e.g. Zhou et al. 2015). While specific experiments have provided examples of substrate by additive interactions, drawing firm conclusions on the magnitude of these effects remains challenging. Nevertheless, there is a need for end users to recognize that specific characteristics of incubated substrates impact on the outcomes of IVFT and be aware of a possible mismatch between the diet of the donor animal and incubated substrate.

Implementing a standardized protocol for preparing substrates to be incubated is also critical in allowing for between IVFT comparisons (Rymer et al., 2005). The most critical issue appears the methods used to dry fresh material, such as grass. Comparisons of freeze-drying with oven drying at 60 or 105°C are often contradictory. There are reports on the effect of feed processing on CH₄ production. Nevertheless, *a priori* freeze-drying is the method of choice for minimizing cell damage that potentially alters the dynamics of microbial attachment, substrate degradation and altering bioactive compounds (Rymer et al., 2005).

With regards to particle size, using cereals as substrates resulted in particle size having little effect on *in vitro* fermentation, provided that the grain kernel had been cut (Lowman et al., 2002), although some evidence to contradict this argument exists with maize (Rymer et al., 2005). With fibrous and more slowly degraded feeds, fermentation rate increases as particle size decreases (Lowman et al., 2002) and it seems likely that this is a consequence of increased surface area as a result of grinding, thereby allowing better microbial access. The most common procedure is to mill the substrate through a 1 mm screen and store under dry, cool and dark conditions in sealed containers prior to

use. Adoption of a standardized approach to sample preparation may enable comparison between independently produced *in vitro* fermentation data of different feeds.

RECOMMENDATION: It is recommended that a range of substrates that reflect the types of feeds used in commercial production systems are used in the initial screening of new additives, unless the objective of an in vitro experiment requires a predefined substrate. Freeze-drying is preferred to oven drying for the drying of high moisture substrates.

Incubation medium and rumen fluid:medium ratio (RF:M)

There is considerable variation in the composition of the medium used for *in vitro* studies reported in the literature. It is important to make a distinction between the term ‘medium’ (i.e., a solution containing a number of components including buffering agents, trace elements, true protein and reducing agents) and ‘buffer’ (Williams, 1998). The types of buffers used in IVFT and the implications on fermentation has been comprehensively reviewed (Rymer et al., 2005). A medium with a high buffering capacity, when used in IVFT, may be disadvantageous because it creates conditions that are not representative of the rumen *in vivo*. This is particularly important when assessing the effectiveness of CH₄ mitigations strategies that may rely on a decrease in rumen pH that include an inhibition of fibrolytic bacteria and/or methanogens (Argyle and Baldwin, 1988; Van Kessel and Russell, 1996; Navarro-Villa et al., 2011). Furthermore, through the prevention of a sharp decline in pH, a highly buffered medium may increase acetic:propionic acid ratios more than would otherwise occur *in vivo*, which impacts on the availability of H₂ for CH₄ production (Lana et al., 1998).

The effect of the incubation medium and rumen fluid to medium ratio (RF:M) on IVFT has been investigated (Rymer et al., 1999; Pell and Schofield, 1993; Cone et al., 2000). Such studies have demonstrated that increases in the proportion of rumen fluid is associated with a decrease in lag phase and a higher rate of gas production, while the effect on total gas production varies. Navarro-Villa et al. (2011) investigated the effect of variable RF:M ratios (1:2, 1:4, and 1:6) in IVFT involving the incubation of different amounts (0.3, 0.5, and 0.7 g) of three contrasting feeds (barley grain, grass silage and barley straw). The results indicated that CH₄ per unit of DM degraded was a more appropriate unit for expressing *in vitro* CH₄ output than CH₄ per unit of DM incubated. Incubation of 0.3 g dried milled feed in 50 ml of *in vitro* culture containing 1:2 RF:M was an acceptable combination for 24 h incubations allowing for a decline in pH declined and maximizing the difference in CH₄ output between substrates.

For a given substrate, pH in the incubation vessel should ideally mimic that in the rumen, i.e. between 6.0-7.0 for forage based diets and 5.5- 6.0 for concentrate based diets. Patra and Yu (2013) evaluated IVFT containing different bicarbonate concentrations (80, 100, and 120 mM) in buffer. Results indicated that bicarbonate concentrations above 80 mM should be avoided to minimize non-microbial CO₂ production associated with changes in pH. A recent comparison of two buffers commonly used in IVFT (McDoughall's (McDoughal, 1948) and Mould's (Mould et al., 2005) buffer indicated that buffer composition had no effect on total gas production (Muetzel et al., 2014). These findings are in direct contrast to earlier reports that a higher phosphate concentration decreased gas production over a 9 h fermentation period (Mould et al., 2005). The differences the latter authors reported, however, were about 4% and decreasing over incubation time. A trend towards higher gas production for McDoughall buffer compared with Mould's buffer may be related to a higher carbonate

concentration (Muetzel et al., 2014). The buffer composition had no effect on CH₄ production or on the percentage of CH₄ released or on VFA production. However, the proportions of major VFA were altered; molar proportion of acetate was higher and that of propionate was lower for incubations with Mould's buffer compared with McDougall's buffer. It is possible that differences in the phosphate to carbonate ratio and associated changes in pH may be responsible (Broudiscou et al., 1999). Based on the work of Kohn and Dunlap (2008) a recent report demonstrated that by adjusting the concentration of buffer bicarbonate, pH can be reasonably well controlled at specific target pH (6.50, 6.25, 6.00, 5.75 and 5.50) during 12 h incubations (Amanzougarene et al., 2015). What is less clear is whether different antimethanogenic additives perform similarly in IVFT over a range of pH.

RECOMMENDATION: Available data do not allow for recommendations on an ideal RF:M. A ratio of 1:2 appears to generate the most reliable results for 24 h incubations. However, this ratio and the amount of substrate incubated needs to be considered on the basis of the frequency of gas sampling and the duration of the incubation depending on the research objectives. Bicarbonate concentration in buffer may influence methanogenesis. To minimize non-metabolic CO₂ production use of buffers containing bicarbonate concentrations above 80 mM should be avoided. Furthermore, adjusting the concentration of bicarbonate in the buffer or the RF:M ratio offers the possibility of setting a target pH according to the substrate incubated.

3.5. Incubation procedure and CH₄ measurements

In vitro gas production systems are typically conducted over intervals of between 16 to 72 h. As such, IVFT do not mimic important physiological processes *in vivo* such as

ruminal digesta turnover. Removal of soluble particles in the liquid medium may have adverse effects on microbial fermentation, by decreasing the amount of soluble substrate available for microbial growth or may conversely stimulate activity (Roger et al., 1990). Soto et al. (2013) reported that the numbers of all quantified microorganisms (total bacteria, protozoa, methanogens, fungi, *Fibrobacter succinogenes* and *Ruminococcus flavefaciens*) declined sharply during 24 h to 72 h of incubation. This is likely due to the exhaustion of fermentable substrate and the accumulation of fermentation end products. Different substrates (soluble carbohydrates, starch, pectins, cellulose, hemicellulose and protein) that are fermented can be degraded at different rates also have a variable contribution to CH₄ production (Bannink et al., 2006). In cows, mean retention time of NDF components in the reticulorumen is around 28 h (using external markers, Schwarm et al., 2015) or 28 to 60 h (using intrinsic stable isotope labelling techniques, Warner et al., 2014), being longer than the standard *in vitro* incubation time of 24 h. Retention time in the reticulorumen is related to particle size (Schwarm et al. 2008). It can be argued that NDF fermentation during *in vitro* incubations probably approaches a plateau after 24 h given that relatively small feed particles (ca. 1 mm) are usually incubated.

Some IVFT have tested the effects of different substrates on CH₄ production based on the collection of a single gas sample after 24 h (García-González et al., 2008). Such an approach may result in gas pressure in the headspace exceeding a given threshold (48 kPa) and consequently an impairment of microbial activity (Rymer et al., 2005; Taglapietra et al., 2010). Several protocols, such as those described by Theodorou et al., 1994, Cone et al., 1996 and Davies et al., 2000, stipulate that headspace gases being released at pre-determined intervals or when a pre-set threshold of pressure is reached (Muetzel et al., 2014). Venting the gas produced requires that CH₄ concentration is

measured simultaneously, given that different CH₄ concentrations can be expected depending on substrate. Although the most common method to measure CH₄ concentration on gas samples collected from incubation vessels is by using gas chromatography, as for *in vivo* measurements, other techniques are available and have been used such as infrared methane analysis (Goel et al., 2008, Cobellis et al., 2015) and absorption of CO₂ (Fievez et al., 2005).

For successful IVFT, it is essential that experimental treatments be randomly allocated to bottle positions. The order in which bottles are inoculated also needs to be randomized across bottle positions and treatments. Randomization serves to minimize possible confounding effects of bottle position (or water baths used), treatments and timing of inoculation. Such an approach is analogous to the random allocation of treatments to animals for *in vivo* experiments. It is also important to consider establishing incubations with a single, identical source of medium, and a single, identical source of substrate. Finally, independently of the number of independent incubation runs performed, different replicates (bottles) of each treatment need to be included. Three bottles minimum is sensible as it allows possible outliers to be identified.

RECOMMENDATION: The duration of the incubation should be adjusted based on the composition and physical properties of substrate incubated that determines the frequency of gas sampling required for measuring CH₄ production. Gas composition should be determined at the same time gas pressure is vented. It is highly desirable that the timing of inoculation and the allocation of treatments with respect to bottle position are randomized as much as possible. It is recommended that 3 bottles be used per unique treatment in each incubation run.

3.6. Headspace gas composition

It is well established that H_2 concentration can affect the thermodynamics of fermentation and the growth rate of hydrogenotrophic methanogens in the rumen (Janssen, 2010). Hydrogen produced in the rumen is present in two forms, as dissolved H_2 and as H_2 gas, but methanogens only utilize dissolved H_2 (Wang et al., 2014). Reports on the influence of headspace gas composition on *in vitro* gas production and rumen fermentation are scarce. In one study, Patra and Yu (2013) noted that initial CO_2 headspace, but not N_2 headspace, was positively correlated with CH_4 production after fermentation. This prompted the hypothesize that headspace gas composition, CO_2 in particular, which is in exchange with H_2CO_3/HCO_3^- in the medium, depending on concentration, acid-base balance and gas pressure, may affect fermentation characteristics and gas production in ruminal *in vitro* cultures. A range of initial headspace composition of *in vitro* cultures has been reported, including 100% CO_2 (Anderson et al., 2003; Weimer et al., 2005), 100% N_2 (Hoover et al., 1976) and a mixture of gases typical of an anaerobic chamber (85% N_2 , 10% H_2 , and 5% CO_2 ; Zhou et al., 2011; Patra et al., 2012). Patra and Yu (2013) investigated the effects of three different headspace gases ($N_2 + CO_2 + H_2$ in the ratio of 90:5:5, 100 % CO_2 , and 100 % N_2) and the interaction with type of substrate (alfalfa hay or alfalfa hay and concentrate) and media bicarbonate concentration on gas and CH_4 production. Methane production was much higher when CO_2 was present in the headspace. It is conceivable that equilibrium is established between CO_2 dissolved in the inoculum and CO_2 in headspace gas (Alford, 1976), such that a higher concentration of CO_2 in the headspace would result in a greater concentration of dissolved CO_2 in the media. Higher CH_4 production corresponding to CO_2 in the headspace may be explained by an immediate

and greater availability of CO₂ in the inoculum that serves as the electron acceptor for the primary hydrogenotrophic methanogenesis pathway. An increase in dissolved CO₂ may also promote growth and activity of methanogens. Even though CH₄ production in the study of Patra and Yu (2013) was greater when the headspace contained CO₂ rather than N₂, total or net gas production was lower for the former compared with the latter. Further investigations are required to understand the impact of a mixture of CO₂ and N₂ that best mimics rumen gas composition *in vivo*.

RECOMMENDATION: Owing to the effect of headspace gas composition on gas production, including CH₄ production, it is recommended that all future studies should both consider this as an influencing factor and report headspace gas composition. Rumen fluid should be flushed continuously, and once added into the medium, continued to be flushed for at least 10 min before incubation is initiated. Following the addition of inoculum into the bottle, the headspace should be flushed continuously until the bottle is sealed.

Please insert Table 2 around here

3.7. *In vitro* versus *in vivo*

Numerous studies have examined the influence of antimethanogenic compounds on CH₄ production *in vitro*, but few have undertaken a simultaneous evaluation *in vivo* and *in vitro*. Direct comparison of effects *in vitro* and *in vivo* would allow a better interpretation of IVFT data and inform on the treatments suitable for further evaluation

in vivo. When addressing inconsistencies between results from *in vitro* and *in vivo* studies it is worthwhile considering:

- i) The accuracy of *in vitro* systems to predict the CH₄ production of a given diet per unit feed intake or digested matter
- ii) The ability to simulate the direction of changes (not absolute values) in CH₄ production when anti-methanogenic agents are tested relative to an appropriate control.

i) Flachowsky and Lebzien (2012), using data from Moss and Givens (1997) reported a poor relationship ($r^2=0.264$) in CH₄ production obtained *in vivo* and *in vitro* methods. Bhatta et al. (2007) compared IVFT measurements of CH₄ production with the SF₆-technique across a range of diets. Methane production (ml/g DM) estimated from 48 h *in vitro* gas production was higher than measurements *in vivo* for all diets. Of particular note is that the average of CH₄ production at 24 h and 48 h was closely correlated with values based on SF₆ ($R^2=0.78$, 5 diets and 4 animals used) (Table 2).

Blümmel et al. (2005) conducted a study to compare feed intake, digestibility and CH₄ production by open-circuit respiration measurements in sheep fed 15 untreated, sodium hydroxide (NaOH) treated and anhydrous ammonia (NH₃) treated wheat, barley and oat straws also evaluated using IVFT. Total daily CH₄ production, calculated from *in vitro* fermentation characteristics (i.e., true degradability, SCFA ratio and efficiency of microbial production) and OM intake were found to be closely correlated with CH₄ emissions (L/d) measured in respiration chambers ($y = 2.5 + 0.86x$, $R^2 = 0.89$, $P < 0.001$, 15 diets and 4 animals used; $y=CH_4$ production *in vivo*; $x=CH_4$ predicted *in vitro*). Intake of OM measured *in vivo* OM intake was also used to calculate CH₄ production *in vitro*. As such OM intake was common to both the independent and the dependent variable, which could explain the close association between *in vitro* and *in*

vivo measurements. It is important to point that Blümmel et al. (2005) and Bhatta et al. (2007) did not use the same animals from the *in vivo* trials as donor of rumen fluid for the *in vitro* incubations.

More recently, Hatew et al. (2015) reported a study comparing measurements of CH₄ production *in vitro* and *in vivo*, and those were conducted simultaneously (animals adapted to the same substrate as incubated in bottles were used as a source of rumen inocula for 72 h *in vitro* incubations). Measurements of CH₄ production for 24 h *in vitro* (expressed per unit of OM incubated) were found to be moderately correlated ($R^2 = 0.54$; $P = 0.04$, 4 diets and 16 animals used) with *in vivo* CH₄ production (when expressed per unit of estimated rumen-fermentable OM) across a range of diets differing in source and amount of starch in dietary concentrates. However, no association was found when *in vivo* CH₄ production was expressed per unit of ingested OM ($R^2 = 0.04$; $P = 0.88$). More research is needed using a wider range of diets representing different production systems.

ii) Few direct comparisons of antimethanogenic compounds *in vitro* and *in vivo* are available (Table 2). Martínez-Fernández et al. (2013) compared the effectiveness of bromochloromethane and propyl propane thiosulfinate to inhibit CH₄ production *in vitro* and *in vivo*. Even though both compounds were found to decrease CH₄ production *in vitro* by as much as 90% per unit of DM intake responses in goats were much lower (-33%), although measurements *in vivo* and *in vitro* were not made simultaneously. Two newly developed molecules (ethyl-3-nitrooxy propionate, ENP, and 3-nitrooxypropanol, 3NOP) have also been evaluated *in vitro* and *in vivo* (Martinez-Fernandez et al., 2014). Both compounds were given to sheep at two different doses of 50 and 500 mg/animal per day, corresponding to around 10 and 100 mg/L of rumen content or concentrations of 68 and 681 μ M, respectively. Administration of 500 mg/d of h 3NOP at decreased

CH₄ production by -29% on d 14, which was much lower than a value of -95%, determined *in vitro* (Martínez-Fernández et al., 2014). Similar differences in responses to Japanese horseradish oil have been reported between *in vitro* and *in vivo* studies (Mohammed et al., 2004). Much larger decreases in CH₄ production were observed *in vitro* (-89%) than *in vivo* (-18.7%, Table 2), findings that are in agreement a recent meta-analysis (Hristov et al. 2012). Such discrepancies in the effectiveness of test compounds when given in similar doses may be explained by a combination of several factors: (1) test compounds used are typically administered in 1-2 pulses via the ruminal cannula that often coincide with feeding times, and as a consequence not be rapidly and well mixed with rumen contents; (2) differences in the degradation rate of the active compounds *in vitro* and *in vivo*; (3) decrease in microbial density and changes in bacterial community structure of rumen contents during processing as inoculum for *in vitro* studies associated with the exposure of microorganisms to oxygen and the removal of solids during filtration (Soto et al., 2012); (4) potential washout of these compounds from the rumen or absorption through the rumen wall and (5) adaptation of the rumen microbial ecosystem to the tested compound *in vivo* that is not emulated by inoculated microbiota *in vitro*.

A different scenario as described above has been recently reported by Castro-Montoya et al. (2015): a blend of essential oil was effective reducing daily emissions of methane in dairy cattle and emissions relative to body weight in beef cattle, interestingly, these effects were not observed *in vitro* regardless of the technique used to replicate *in vivo* results (IVFT or consecutive batch culture). This might be due to differences in the mode of action of the essential oils *in vitro* and *in vivo*, which merits attention for future research.

826 **RECOMMENDATIONS:**

827 *In vitro* CH₄ production is more closely correlated with *in vivo* CH₄ production across a
828 range of feeding regimes when *in vivo* CH₄ emissions are expressed per unit of
829 degraded material rather than per unit of material ingested, although the ratio of
830 substrate (feed) to volume of rumen fluid needs to be considered. The ability of IVFT to
831 reliably predict effects *in vivo* is affected by adaptation of the rumen inoculum to the
832 substrate tested *in vitro*. IVFT offers a valuable tool for the study and screening of anti-
833 methanogenic additives before testing *in vivo*. It is recommended that *in vitro* data are
834 confirmed *in vivo* before conclusions on the effectiveness of feed ingredients or
835 additives for lowering CH₄ production are drawn given that inhibition potential is often
836 overestimated *in vitro*.

837

838 **Please insert Table 3 around here**

839

840 **3.8. Units to express CH₄ production**

841 *In vitro* CH₄ production is usually expressed as ml (or g or mmol) of CH₄/g of
842 substrate incubated (Bodas et al., 2008), ml (or g or mmol) of CH₄/g of substrate
843 degraded (García-González et al., 2008), or ml of CH₄/ml of gas produced (Goel et al.,
844 2008). As rumen fermentation modifiers may affect substrate degradation in a dose
845 dependent manner (Russell and Strobel, 1988), effects on CH₄ production are better
846 expressed per unit of substrate degraded, rather than per unit of substrate incubated. For
847 example, Navarro-Villa et al. (2011) highlighted the importance of measuring substrate
848 degradability over the incubation period based on the observation that the ranking of
849 CH₄ production potential of incubated feeds (barley grain, grass silage, barley straw)
850 differed depending on whether CH₄ was expressed per unit DM incubated or DM

degraded. Methane output/g DM incubated (CH_4/DMi) was in the order barley grain > grass silage > barley straw, whereas when CH_4 output was expressed per g DM disappeared (CH_4/DMd) the ranking was barley straw > grass silage > barley grain. This last observation is consistent among all quantities of feed, and ratios of rumen fluid to buffer combinations, and concurs with findings from *in vivo* studies, which compared low and high concentrate diets (Yan et al., 2000).

To calculate the volume of CH_4 produced per unit of substrate degraded, net values of gas and CH_4 production, as well as net amount of substrate degraded, are needed. These are obtained by blanks (i.e., flasks without substrate which contain only inoculum and medium), which are necessary to correct for gas, CH_4 and fermentation of residual OM in the inoculum (Rymer et al., 2005; Navarro-Villa et al., 2011). The CH_4/DMd values assume that all DM which disappeared is digested, and that no undigested soluble or particulate DM passed through the sintered glass crucible when the post-incubation DM residue was being isolated. Total gas or VFA produced is not as precise as substrate degraded but a good, simple proxy when degradation of substrate can not be determined.

RECOMMENDATIONS:

Regardless of the duration of the incubation in vitro, the amount of CH_4 produced should ideally be expressed relative to the amount of substrate degraded, rather than the amount of substrate incubated. If not possible, it is recommended to be expressed relative to total gas or VFA produced.

4. Conclusion

There is no standard protocol for assessing enteric methane mitigation in ruminants using *in vitro* gas production technique, as conditions need to be adjusted according to the research question. However, numerous technical issues relating to donor animals, microbial inoculum and general procedures need to be considered (Table 3) to ensure the objectives of experiments can be properly fulfilled. This would allow harmonization of laboratory methods, better interpretation of results and facilitate inter-studies comparisons. Results from *in vitro* gas production technique studies need to be carefully interpreted before assessing mitigation strategies *in vivo*.

Acknowledgements

This review is part of the FACCE-JPI ‘Global Network’ project and the ‘Feeding and Nutrition Network’ of Livestock Research Group, in the Global Research Alliance for Agricultural Greenhouse Gases (www.globalresearchalliance.org). Authors acknowledge national funding from INIA (Spain, project MIT01-GLOBALNET-EEZ), the Ministry of Economic Affairs (The Netherlands; project Global Research Alliance on Agricultural Greenhouses Gases, BO-20-007-006), USDA-NIFA (USA), French National Research Agency through the program FACCE-JPI program, Agricultural GHG Research Initiative for Ireland (AGRI-I), Department for Environment Food & Rural Affairs (UK), BLW (Switzerland) and Academy of Finland, Helsinki, Finland (Project 281337).

References

Alford, J. S., Jr., 1976. Measurement of dissolved carbon dioxide. Can. J. Microbiol. 22, 52–56.

900 Amanzougarene, Z., Schauf, S., Fondevila, M., 2015. Control of pH in vitro for the
 901 simulation of fermentation in high concentrate feeding conditions. AIDA, XVI
 902 Jornadas sobre producción Animal, I, 215-217 (ISBN 978-84-606-7969-1).
 903 Ammar, H., S. López, S. Andrés, M.J. Ranilla, R. Bodas, J.S. González., 2008. In vitro
 904 digestibility and fermentation kinetics of some browse plants using sheep or goat
 905 ruminal fluid as the source of inoculum. Anim. Feed Sci. Technol. 147, 90-104.
 906 Anderson, R. C., Callaway, T.R., Van Kessel, J.A.S., Jung, Y.S., , Edrington, T. S.,
 907 Nisbet, D. J., 2003. Effect of select nitrocompounds on ruminal fermentation; an
 908 initial look at their potential to reduce economic and environmental costs
 909 associated with ruminal methanogenesis. Bioresour. Technol. 90, 59–63.
 910 Argyle, J.L., Baldwin, R.L., 1988., Modeling of rumen water kinetics and effects of
 911 rumen pH changes. J. Dairy Sci. 71, 1178-1188.
 912 Bannink, A., Kogut, J., Dijkstra, J., Kebreab, E., France, J., Tamminga, A., Van
 913 Vuuren, A.M., 2006. Estimation of the stoichiometry of volatile fatty acid
 914 production in the rumen of lactating cows. J. Theor. Biol. 238, 36–51.
 915 Blümmel, M., Givens, D.I., Mossb, A.R., 2005. Comparison of methane produced by
 916 straw fed sheep in open-circuit respiration with methane predicted by
 917 fermentation characteristics measured by an in vitro gas procedure. Anim. Feed
 918 Sci. Technol. 123–124, 379–390.
 919 Blümmel, M., Ørskov, E.R., 1993. Comparison of in vitro gas production and nylon bag
 920 degradability of roughages in predicting feed intake in cattle. Anim. Feed Sci.
 921 Technol. 40, 109–119.
 922 Bodas, R., Lopez, S., Fernandez, M., García-González, R., Rodríguez, A.B., Wallace,
 923 R.J., Gonzalez, J.S., 2008. In vitro screening of the potential of numerous plant

924 species as antimethanogenic feed additives for ruminants. *Anim. Feed Sci.*
 925 *Technol.* 145, 245–258.

926 Broudiscou, L.P., Papon, Y., Broudiscou, A.F., 1999. Optimal mineral composition of
 927 artificial saliva for fermentation and methanogenesis in continuous culture of
 928 rumen microorganisms. *Anim. Feed Sci. Technol.* 79, 43–55.

929 Bryant, A. M., 1964. *N. Z. J. Agric. Res.* 7, 695

930 Bueno, I.C.H., Brandi, R.B., Franzolin, Benetel, G., Adibe G., Abdalla, L., Muir, H.L.
 931 2015. In vitro methane production and tolerance to condensed tannins in five
 932 ruminant species. *Anim. Feed Sci. Technol.* 205, 1-19.

933 Busquet, M., Calsamiglia, S., Ferret, A., Cardozo, P., Kamel, C., 2005. Effects of
 934 cinnamaldehyde and garlic oil on rumen microbial fermentation in a dual flow
 935 continuous culture. *J. Dairy Sci.* 88, 2508–2516.

936 Calabrò, S., Lopez, S., Piccolo, V., Dijkstra, J., Dhanoa, M.S., France, J., 2005.
 937 Comparative analysis of gas production profiles obtained with buffalo and sheep
 938 ruminal fluid as the source of inoculum. *Anim. Feed Sci. Technol.*, 123, 51-65.

939 Cardozo, P., Calsamiglia, S., Ferret, A., Kamel, C., 2004. Effects of natural plant
 940 extracts on ruminal protein degradation and fermentation profiles in continuous
 941 culture. *J. Anim. Sci.* 82, 3230–3236.

942 Cardozo, P., Calsamiglia, S., Ferret, A., Kamel, C., 2005. Screening for the effects of
 943 natural plant extracts at different pH on in vitro rumen microbial fermentation of a
 944 high-concentrate diet for beef cattle. *J. Anim. Sci.* 83, 2572–2579.

945 Castro-Montoya, J., De Campeneere, S., Van Ransta, G., Fievez, V., 2012. Interactions
 946 between methane mitigation additives and basal substrates on in vitro methane
 947 and VFA production. *Anim. Feed Sci. Technol.* 176, 47– 60.

948 Castro-Montoya, J., Peiren, N., Cone, J.W., Zweifel, B., Fievez, V., De Campeneere, S.,
949 2015. In vivo and in vitro effects of a blend of essential oils on rumen methane
950 mitigation. *Livest. Sci.* 180, 134–142.

951 Cheng, K.J., McAllister, T.A., Mathiesen, S.D., Blix, A.S., Orpin, C.G., Costerton,
952 J.W., 1993. Seasonal changes in the adherent microflora of the rumen in high-
953 arctic Svalbard reindeer. *Can. J. Microbiol.* 39, 101-8.

954 Clauss, M., Schwarm, A., Ortmannb, S., Streichb, W.J., Hummel, J., 2007. A case of
955 non-scaling in mammalian physiology? Body size, digestive capacity, food intake,
956 and ingesta passage in mammalian herbivores. *Comp. Bioch. Physiol.* 148, 249–
957 265.

958 Cobellis, G., Petrozzi, A., Forte, C., Acuti, G., Orrù, M., Marcotullio, M.C., Aquino, A.,
959 Nicolini, A., Mazza, V., Trabalza-Marinucci, M., 2015. Evaluation of the Effects
960 of Mitigation on Methane and Ammonia Production by Using *Origanum vulgare*
961 *L.* and *Rosmarinus officinalis L.* Essential Oils on in Vitro Rumen Fermentation
962 Systems. *Sustainability.* 7, 12856-12869.

963 Cone, J.W., Van Gelder, A.H., Bachmann, H., 2000. Influence of inoculum source,
964 dilution and storage of rumen fluid on gas production profiles. In: *Gas Production:*
965 *Fermentation Kinetics for Feed Evaluation and to Assess Microbial Activity.*
966 *Proceedings of the EAAP Satellite Symposium on Gas Production, Wageningen,*
967 *The Netherlands. Proc. Br. Soc. Anim. Sci.,* 15–16.

968 Cone, J.W., Van Gelder, A.H., Bachmann, H., 2002. Influence of inoculum source on
969 gas production profiles. *Anim. Feed Sci. Technol.* 99, 221–231.

970 Cone, J.W., Van Gelder, A.H., Visscher, G.J.W., Oudshoorn, L., 1996. Influence of
971 rumen fluid and substrate concentration on fermentation kinetics measured with a

972 fully automated time related gas production apparatus. Anim. Feed Sci. Technol.
 973 61, 113–128.

974 Cornou, C., Storm, I.D., Hindrichsen, I., Worgan, H., Bakewell, E., Yáñez-Ruiz, D.,
 975 Abecia, L., 2013. A ring test of a wireless in vitro gas production system. Anim.
 976 Prod. Sci. 53, 585–592.

977 Czerkawski, J.W., Breckenridge, G., 1975. New inhibitors of methane production by
 978 rumen micro-organisms.

979 Davies, Z.S., Mason, D., Brooks, A.E., Griffith, G.W., Merry, R.J., Theodorou, M.K.,
 980 2000. An automated system for measuring gas production from forages inoculated
 981 with rumen fluid and its use in determining the effect of enzymes on grass silage.
 982 Anim. Feed Sci. Technol. 83, 205–221.

983 De Visser, H., Huisert, H., Klop, A., Ketelaar, R.S. 1993. Autumn-cut silage as
 984 roughage component in dairy cow rations. 2. Rumen degradation, fermentation
 985 and kinetics. Neth. J. Agric. Sci. 41, 221–234

986 Dehority, B.A., 2003. Rumen Microbiology. Nottingham: Nottingham University Press.

987 Demeyer, D., Fievez, V. 2000. Ruminants and environment: methanogenesis. Ann.
 988 Zootech. 49, 95–112.

989 Dhanoa, M.S., France, J., Crompton, L.A., Mauricio, R.M., Kebreab, E., Mills, J.A.N.,
 990 Sanderson, R., Dijkstra, J., López, S., 2005. Technical note: A proposed method
 991 to determine the extent of degradation of a feed in the rumen from the degradation
 992 profile obtained with the in vitro gas production technique using feces as the
 993 inoculum. J. Anim. Sci. 82, 733-746.

994 Dijkstra, J., 1994. Simulation of the dynamics of protozoa in the rumen. Brit. J. Nut. 72,
 995 679-699.

996 Dijkstra, J., Kebreab, E., Bannink, A., France, J., Lopez, S., 2005. Application of the
 997 gas production technique to feed evaluation systems for ruminants. *Anim. Feed*
 998 *Sci. Technol.* 123, 561–578.

999 Duffield, T., Plaizier, J.C., Fairfield, A., Bagg, R., Vessie, G., Dick, P., Wilson, J.,
 1000 Aramini, J., McBride, B., 2004. Comparison of techniques for measurement
 1001 of rumen pH in lactating dairy cows. *J. Dairy Sci.* 87, 59–66.

1002 Durmic, Z., Hutton, P., Revell, D.K., Emms, J., Hughes, S., Vercoe, P.E., 2010. In vitro
 1003 fermentative traits of Australian woody perennial plant species that may be
 1004 considered as potential sources of feed for grazing ruminants. *Anim. Feed Sci.*
 1005 *Technol.* 160, 98–109.

1006 Editorial, 2012. Use of replicates in statistical analyses in papers submitted for
 1007 publication in *Animal Feed Science and Technology*. *Anim. Feed Sci. Technol.*
 1008 171, 1-5.

1009 El Shaera, H.M., Omeda, H.M., Chamberlain, A.G., Axford, R.F.E., 1987. Use of
 1010 faecal organisms from sheep for the in vitro determination of digestibility. *J.*
 1011 *Agric. Sci.* 109, 257-259

1012 El-Meadaway, A., Z. Mir, P. S. Mir, M. S. Zaman, L. J. Yanke., 1998. Relative efficacy
 1013 of inocula from rumen liquor or fecal solution for determining in vitro
 1014 digestibility and gas production. *Can. J. Anim. Sci.* 78, 673–679.

1015 Fievez, V., O.J. Babayemi and D. Demeyer, 2005. Estimation of direct and indirect gas
 1016 production in syringes: A tool to estimate short chain fatty acid production
 1017 requiring minimal laboratory facilities. *Anim. Feed Sci. Technol.* 123-124, 197-
 1018 210.

1019 Flachowsky, G., Lebzien, P., 2012. Effects of phytogenic substances on rumen
1020 fermentation and methane emissions: A proposal for a research process. Anim.
1021 Feed Sci. Technol. 176, 70-77.

1022 García-González, R., Lopez, S., Fernandez, M., Bodas, R., Gonzalez, J.S., 2008.
1023 Screening the activity of plants and spices for decreasing ruminal methane
1024 production in vitro. Anim. Feed Sci. Technol. 147, 36–52.

1025 Geishauser, T., Gitzel, A., 1996. A comparison of rumen fluid sampled by oro-ruminal
1026 probe versus rumen fistula. Small Rumin. Res. 21, 63–69.

1027 Goel, G., Makkar, H.P.S., Becker, K., 2008. Effect of Sesbania sesban and Carduus
1028 pycnocephalus leaves and fenugreek seeds (*Trigonella foenum-graecum* L.) and
1029 their extracts on partitioning of nutrient from roughage and concentrate based
1030 feeds to methane. Anim Feed Sci Technol. 147, 72–89.

1031 Groot, J.C.J., Cone, J.W., Williams, B.A., Debersaques, F.M.A., Lantinga, E.A., 1996.
1032 Multiphasic analysis of gas production kinetics for in vitro fermentation of
1033 ruminant feeds. Anim. Feed Sci. Technol. 64, 77–89.

1034 Haddi, M.L., Filacorda, S., Meniai, K., Rollin, F., Susmel, P., 2003. In vitro
1035 fermentation kinetics of some halophyte shrubs sampled at three stages of
1036 maturity. Anim. Feed Sci. Technol. 104, 215-225.

1037 Hatew, B., Conea, J.W., Pellikaana, W.F., Podesta, S.C., Bannink, A., Hendrik, W.H.,
1038 Dijkstra, J., 2015. Relationship between in vitro and in vivo methane production
1039 measured simultaneously with different dietary starch sources and starch levels in
1040 dairy cattle. Anim. Feed Sci. Technol. 202, 20–31

1041 Hervás, G., Frutos, P., Giráldez, F.J., Mora, M.J., Fernández, B., Mantecón, A.R., 2005.
1042 Effect of preservation on fermentative activity of rumen fluid inoculum for in
1043 vitro gas production techniques. Anim. Feed Sci. Technol., 123–124, 107–118.

1044 Hoover, W. H., B. A. Crooker, and C. J. Sniffen., 1976. Effects of differential solid-
 1045 liquid removal rates on protozoa numbers in continuous cultures of rumen
 1046 contents. *J. Anim. Sci.* 43, 528–534.

1047 Hristov A.N., Lee, C., Hristov, R., Firkins, J.L. 2012. A meta-analysis of variability in
 1048 continuous-culture ruminal fermentation and digestibility data. *Journal of Dairy*
 1049 *Science* 95, 5299-307 .

1050 Hristov, A. N., T. A. McAllister, F. H. Van Herk, K. -J. Cheng, C. J. Newbold and P. R.
 1051 Cheeke. 1999. Effect of *Yucca schidigera* on ruminal fermentation and nutrient
 1052 digestion in heifers. *J. Anim. Sci.* 77, 2554-2563.

1053 Huntington, J.A., Givens, D.I. 1998. The effects of host diet on the cumulative gas
 1054 production profile of grass hay and high temperature dried grass. In *vitro*
 1055 Techniques for Measuring Nutrient Supply to Ruminants. In: E.R., Deaville, E.,
 1056 Owen, A.T., Adesogan, C., Rymer, J.A., Huntington, T.L.J., Lawrence (Eds.),
 1057 Occ. Publication no. 22. *Br. Soc. Anim. Sci.* 224–226.

1058 Janssen, P.H. 2010. Influence of hydrogen on rumen methane formation and
 1059 fermentation balances through microbial growth kinetics and fermentation
 1060 thermodynamics. *Anim. Feed Sci. Technol.* 160, 1–22

1061 Jeyanathan, J., Kirs, M., Ronimus, R.S., Hoskin, S.O., Janssen, P.H., 2011. Methanogen
 1062 community structure in the rumens of farmed sheep, cattle and reddeer fed
 1063 different diets. *FEMS Microbiol. Ecol.* 76, 311–326.

1064 Joblin, K.N., 2005. Classical methods for isolation, enumeration, cultivation and
 1065 functional assays of rumen microbes. Methanogenic archaea. In *Methods in Gut*
 1066 *Microbial Ecology for Ruminants*. Springer, Edited by H.P.S. Makkar and C.S.
 1067 McSweeney. ISBN-10 1-4020-3790-2 (HB).

1068 Jouany, J.P., Lassalas, B., 2002. Gas pressure inside a rumen in vitro systems stimulates
 1069 the use of hydrogen. Presented at 3. Joint RRI-INRA Gastrointestinal Tract
 1070 Microbiology Symposium. Beyond Antimicrobials. The Future of Gut
 1071 Microbiology, Aberdeen, GBR (2002-06-12 - 2002-06-15).

1072 Kamel, C., Greathead, H.M.R., Ranilla, M.J., Tejido, M.L., Carro, M.D., 2008. Effects
 1073 of allicin and diallyl disulfide on in vitro fermentation of a mixed diet. *Anim.*
 1074 *Feed Sci. Technol.* 145, 351–363.

1075 Kim, C. H., J. N. Kim, J. K. Ha, S. G. Yun, S. S. Lee., 2004. Effects of dietary addition
 1076 of surfactant Tween 80 on ruminal fermentation and nutrient digestibility of
 1077 Hanwoo steers. *Asian-Aust. J. Anim. Sci.* 17:337-342.

1078 Kittelmann, S., Pinares-Patiño, C.S., Seedorf, H., Kirk, M.R., Ganesh, S.V., McEwan,
 1079 J.C., Janssen, P.H., 2014. Two Different Bacterial Community Types Are Linked
 1080 with the Low-Methane Emission Trait in Sheep. *PLoS ONE* 9(7): e103171.

1081 Klevenhusen, F., A. Muro-Reyes, R. Khiaosa-ard, B. U. Metzler-Zebeli, and Q.
 1082 Zebeli. 2012. A meta-analysis of effects of chemical composition of incubated
 1083 diet and bioactive compounds on in vitro ruminal fermentation. *Anim. Feed Sci.*
 1084 *Technol.* 176:61–69.

1085 Kohn, RA., Dunlap, TF., 2008. Calculation of the buffering capacity of bicarbonate in
 1086 the rumen and in vitro. *J. Anim. Sci.* 76, 1702-1709.

1087 Komarek, R.J., 1981. Rumen and abomasal cannulation of sheep with specially
 1088 designed cannulas and a cannula insertion instrument. *J. Anim. Sci.* 53,790–795.

1089 Kristensen, N.B., Engbæk, M., Vestergaard, M., Harmon, D.L., 2010. Technical note:
 1090 ruminal cannulation technique in young Holstein calves: effects ofcannulation on
 1091 feed intake, body weight gain, and ruminal development at six weeks of age. *J.*
 1092 *Dairy Sci.* 93, 737–742.

1093 Lana, R.P., Russell, J.B., Van Amburgh, M.E., 1998. The role of pH in regulating
1094 ruminal methane and ammonia production. *J. Anim. Sci.* 76, 2190-2196.

1095 Le Liboux, S., Peyraud, J.L., 1999. Effect of forage particle size and feeding frequency
1096 on fermentation patterns and sites and extent of digestion in dairy cows fed mixed
1097 diets. *Anim. Feed Sci. Technol.* 76, 297-319.

1098 Leedle, M.P. Bryant, R.B. Hespell., 1982. Diurnal variations in bacterial numbers and
1099 fluid parameters in ruminal contents of animals fed low- or high-forage diets.
1100 *Appl. Environ. Microbiol.* 44, 402–412

1101 Lodge-Ivey, S.L., Browne-Silva, J., Horvath, M.B., 2009. Technical note: bacterial
1102 diversity and fermentation end products in rumen fluid samples collected via oral
1103 lavage or rumen cannula. *J. Anim. Sci.* 87, 2333–2337.

1104 Lowman, R.S., Theodorou, M.K., Cuddeford, D., 2002. The effect of sample processing
1105 on gas production profiles obtained using the pressure transducer technique.
1106 *Anim. Feed Sci. Technol.* 97, 221–237.

1107 Machmüller, A., Dohme, F., Soliva, C.R., Wanner, M., Kreuzer, M., 2001. Diet
1108 composition affects the level of ruminal methane suppression by medium-chain
1109 fatty acids. *Aust. J. Agric. Res.* 52, 713–722.

1110 Martinez-Fernandez, G., Abecia, L., Martin-Garcia, A.I., Ramos-Morales, E., Hervás,
1111 G., Molina-Alcaide, E., Yáñez-Ruiz, D.R., 2013. In vitro–in vivo study on the
1112 effects of plant compounds on rumen fermentation, microbial abundances and
1113 methane emissions in goats. *Animal* 7, 1925–1934.

1114 Martínez-Fernández, G., L. Abecia , A. Arco , G. Cantalapiedra-Hijar , A. I. Martín-
1115 García, E. Molina-Alcaide ,M. Kindermann , S. Duval , Yáñez-Ruiz, D.R., 2014.
1116 Effects of ethyl-3-nitrooxy propionate and 3-nitrooxypropanol on ruminal

1117 fermentation, microbial abundance, and methane emissions in sheep. *J. Dairy Sci.*
1118 97, 3790–3799.

1119 Martínez, M.E., Ranilla, M.J., Tejido, M.L., Ramos, S., Carro, M.D., 2010. The effect
1120 of the diet fed to donor sheep on in vitro methane production and ruminal
1121 fermentation of diets of variable composition. *Anim. Feed Sci. Technol.* 158,
1122 126–135.

1123 Mateos, I., Ranilla, M.J., Tejido, M.L., Saro, C., Kamel, C. and Carro, M.D., 2012. The
1124 influence of diet type (dairy versus intensive fattening) on the effectiveness of
1125 garlic oil and cinnamaldehyde to manipulate in vitro ruminal fermentation and
1126 methane production. *Anim. Prod. Sci.* 53, 299–307

1127 Mauricio, R.M., Owen, E., Mould, F.L., Givens, I., Theodorou, M.K., France, J.,
1128 Davies, D.R. and Dhanoa, M.S., 2001. Comparison of bovine rumen liquor and
1129 bovine faeces as inoculum for an in vitro gas production technique for evaluating
1130 forages. *Anim. Feed Sci. Technol.* 89, 33-48.

1131 McDougall, E.J., 1948. Studies on ruminant saliva, 1. The composition and output of
1132 sheep's saliva. *Biochem. J.* 43, 99–109.

1133 Menke, K.H., Raab, L., Salewski, A., Steingass, H., Fritz, D., Schneider, W., 1979. The
1134 estimation of the digestibility and metabolisable energy content of ruminant
1135 feeding stuffs from the gas production when they are incubated with rumen liquor.
1136 *J. Agric. Sci.*, 93, 217–222

1137 Menke, K.H., Steingass, H., 1988. Estimation of the energetic feed value obtained from
1138 chemical analysis and in vitro gas production using rumen fluid. *Anim. Res. Dev.*
1139 28, 7–55.

1140 Miron, J., Ben Ghedalia, D., Morrison, M.. 2001. Invited Review: Adhesion
1141 mechanisms of rumen cellulolytic bacteria. *J. Dairy Sci.* 84, 1294-1309.

1142 Mohammed, N., N. Ajisaka, Z. Lila, K. Hara, K. Mikuni, K. Hara, S. Kanda, and H.
1143 Itabashi., 2004. Effect of Japanese horseradish oil on methane production and
1144 ruminal fermentation in vitro and in steers. *J. Anim. Sci.* 82, 1839–1846.

1145 Moharrery, A., Das, M., 2001. Correlation between microbial enzyme activities in the
1146 rumen fluid of sheep under different treatments. *Reprod. Nutr. Dev.* 41, 513-29.

1147 Monteils, V., Rey, M., Silberberg, M., Cauquil, L., Combes, S., 2012. Modification of
1148 activities of the ruminal ecosystem and its bacterial and protozoan composition
1149 during repeated dietary changes in cows. *J. Anim. Sci.* 90, 4431–4440.

1150 Mould, F.L, Kliem, K.E., Morgan, R., Mauricio, R.M. 2005. In vitro microbial
1151 inoculum: A review of its function and properties. *Anim. Feed Sci. Technol.* 123–
1152 124, 31–50.

1153 Moss, A.R., Givens, D.J., 1997. Methane production from a range of feedstuffs as
1154 determined in vitro using the cumulative gas production technique and compared
1155 with that measured in vivo. *Proc. Brit. Soc. Anim. Sci.* 194.

1156 Mould, F.L., Morgan, R., Kliem, K.E., Krystallidou, E., 2005. A review and
1157 simplification of the in vitro incubation medium. *Anim. Feed Sci. Technol.*
1158 123,155–172.

1159 Mueller, R.E., Asplund, J.M., Lannotti, E.L., 1984. Successive changes in the epimural
1160 bacterial community of young lambs as revealed by scanning electron
1161 microscopy. *Appl. Environ. Microbiol.* 47, 715–723.

1162 Muetzel, S., Hunt, C., Tavendale, M.H., 2014. A fully automated incubation system for
1163 the measurement of gas production and gas composition. *Anim. Feed Scie*
1164 *Technol.* 196, 1-11.

1165 Navarro-Villa, A., O'Brien, M., López, S., Boland, T.M., O'Kiely, P., 2011.
 1166 Modifications of a gas production technique for assessing in vitro rumen methane
 1167 production from feedstuffs. *Anim. Feed Sci. Technol.* 166, 163–174.

1168 O'Brien, M., Navarro-Villa, A., Purcell, P. J., Boland, T. M., O'Kiely P., 2013.
 1169 Reducing in vitro rumen methanogenesis for two contrasting diets using a series
 1170 of inclusion rates of different additives. *Anim. Prod. Sci.* 54, 141–157.

1171 Patra, A. K., J. Stiverson, Z. Yu., 2012. Effects of Quillaja and yucca saponins on
 1172 communities and select populations of rumen bacteria and archaea, and
 1173 fermentation in vitro. *J. Appl. Microbiol.* 113, 1329–1340.

1174 Patra, A.K., Yu, Z., 2013. Effects of gas composition in headspace and bicarbonate
 1175 concentrations in media on gas and methane production, degradability, and rumen
 1176 fermentation using in vitro gas production techniques. *J. Dairy Sci.* 96, 4592–
 1177 4600.

1178 Payne, J.S., Hamersley, A.R., Milligan, J.C., Huntington, J.A., 2002. The affect of
 1179 rumen fluid collection time on its fermentative capacity and the stability of rumen
 1180 conditions in sheep fed a constant diet. *Proc. Br. Soc. Anim. Sci.* 165.

1181 Pei, C.X., Mao, S.Y., Chenga, Y.F., Zhua, W.Y., 2010. Diversity, abundance and novel
 1182 16S rRNA gene sequences of methanogens in rumen liquid, solid and epithelium
 1183 fractions of Jinnan cattle. *Animal.* 4, 20-29.

1184 Pell, A.N., Schofield, P., 1993. Computerized Monitoring of Gas Production to Measure
 1185 Forage Digestion In Vitro. *J. Dairy Sci.* 76, 1063-1073.

1186 Pellikaan, W., Hendriks, W., Uwimana, G., Bongers, L., Becker, P., Cone, J., 2011. A
 1187 novel method to determine simultaneously methane production during in vitro gas
 1188 production using fully automated equipment. *Anim. Feed Sci. Technol.* 168, 196–
 1189 205.

1190 Pinares-Patiño, C.S., Ulyatt, M.J., Lassey, K.R., Barry, T.N., Holmes, C.W. 2003.
1191 Rumen function and digestion parameters associated with differences between
1192 sheep in methane emissions when fed chaffed lucerne hay. J. Agric. Sci. 140,
1193 205–214.

1194 Prates, A., de Oliveira, J.A., Abecia, L., Fondevila, M., 2010. Effects of preservation
1195 procedures of rumen inoculum on in vitro microbial diversity and fermentation.
1196 Anim. Feed Sci. Technol. 155, 186–193.

1197 Ramos-Morales, E., Arco-Pérez, A., Martín-García, A.I., Yáñez-Ruiz, D.R., Frutos, P.,
1198 Hervás, G. 2014. Use of stomach tubing as an alternative to rumen cannulation to
1199 study ruminal fermentation and microbiota in sheep and goats. Anim. Feed Sci.
1200 Technol. 198, 57–66

1201 Ranilla M.J., Tejido, M.L., Carro M.D., 2001. Comparación de diversos métodos de
1202 desligamiento de bacterias ruminales asociadas a la fase sólida de la digesta en un
1203 sistema in vitro (Rusitec). ITEA (Vol. extra), 22, pp. 382-384.

1204 Robinson, P., Campbell, M., Fadel, J., 1999. Influence of storage time and temperature
1205 on in vitro digestion on neutral detergent fibre at 48h, and comparison to 48h in
1206 sacco neutral detergent fiber digestion. Anim. Feed Sci. Technol. 80, 257-266.

1207 Roger, V.R., G. Fonty, S. Komisarczuk-Bondy, Gouet, P., 1990. Effects of
1208 physicochemical factors on the adhesion to cellulose avicel of the rumen bacteria
1209 *Ruminococcus flavefaciens* and *Fibrobacter succinogenes* subsp. *succinogenes*.
1210 Appl. Environ. Microbiol. 56:3081-3087.

1211 Russell, J.B., Strobel, H.J., 1988. Effects of additives on in vitro ruminal fermentation: a
1212 comparison of monensin and bacitracin. Another Gram-positive antibiotic. J.
1213 Anim. Sci. 66, 552–558.

1214 Rymer, C., Huntington, J.A., Givens, D.I., 1999. Effects of inoculum preparation
1215 method and concentration, method of inoculation and pre-soaking the substrate on
1216 the gas production profile of high temperature dried grass. *Anim. Feed Sci.*
1217 *Technol.* 78, 199–213.

1218 Rymer, C., Huntington, J.A., Williams B.A., Givens, D.I., 2005. In vitro cumulative gas
1219 production techniques: History, methodological considerations and challenges.
1220 *Anim. Feed Sci. Technol.* 123–124, 9–30

1221 Sadet, S., Martin, C., Meunier, B., Morgavi, D.P., 2007. PCR-DGGE analysis reveals a
1222 distinct diversity in the bacterial population attached to the rumen epithelium.
1223 *Animal.* 1, 939-944.

1224 Salem, A.F.Z.M., 2005. Impact of season of harvest on in vitro gas production and dry
1225 matter degradability of *Acacia saligna* leaves with inoculum from three ruminant
1226 species. *Anim. Feed Sci. Technol.* 123, 67-79.

1227 Schwarm, A., Ortmann, S., Wolf, C., Streich, W.J., Clauss, M., 2008. Excretion patterns
1228 of fluid and different sized particle passage markers in banteng (*Bos javanicus*)
1229 and pygmy hippopotamus (*Hexaprotodon liberiensis*): two functionally different
1230 foregut fermenters. *Comp. Biochem. Physiol. A*, 150: 32-39

1231 Schwarm, A., Schweigel-Röntgen, M., Kreuzer, M., Ortmann, S., Gill, F., Kuhla, B.,
1232 Meyer, U., Lohölter, M., Derno, M., 2015. Methane emission, digestive
1233 characteristics and faecal archaeol in heifers fed diets based on silage from brown
1234 midrib maize as compared to conventional maize. *Arch. Anim. Nutr.* 69, 159-176.

1235 Shen, J.S., Chai, Z., Song, L.J., Liu, J.X., Wu, Y.M., 2012. Insertion depth of oral
1236 stomach tubes may affect the fermentation parameters of ruminal fluid collected
1237 in dairy cows. *J. Dairy Sci.* 95, 5978–5984.

1238 Shin, E.C., Choi, B.R., Lim, W.J., Hong, S.Y., An, C.L., Cho, K.M., Kim, Y.K., An,
1239 J.M., Kang, J.M., Lee, S.S., Kim, H., Yun, H.D., 2004. Phylogenetic analysis of
1240 archaea in three fractions of cow rumen based on the 16S rDNA sequence.
1241 *Anaerobe*. 10, 313-319.

1242 Soto, E.C., Molina-Alcaide, E., Khelil, H., Yáñez-Ruiz, D.R., 2013 Ruminant microbiota
1243 developing in different in vitro simulation systems inoculated with goats' rumen
1244 liquor. *Anim. Feed Sci. Technol.* 185, 9-18.

1245 Soto, E.C., Yáñez-Ruiz, D.R., Cantalapiedra-Hijar, G., Vivas, A., Molina-Alcaide, E.,
1246 2012. Changes in ruminal microbiota due to rumen content processing and
1247 incubation in single-flow continuous culture fermenters. *Anim. Prod. Sci.* 52, 813–
1248 822.

1249 Storm, A.C., Kristensen, N.B., 2010. Effects of particle size and dry matter content of a
1250 total mixed ration on intraruminal equilibration and net portal flux of volatile fatty
1251 acids in lactating dairy cows. *J. Dairy Sci.* 93, 4223–4238

1252 Tagliapietra, F., Cattani, M., Bailoni, L., Schiavon, S. 2010. In vitro rumen
1253 fermentation: Effect of headspace pressure on the gas production kinetics of corn
1254 meal and meadow hay. *Anim. Feed Sci. Technol.* 158, 197–201.

1255 Tan, H.Y., Sieo, C.C., Abdullah, N., Liang, J.B., Huang, X.D., Ho, Y.W., 2011. Effects
1256 of condensed tannins from *Leucaena* on methane production, rumen fermentation
1257 and populations of methanogens and protozoa in vitro. *Anim. Feed Sci. Technol.*
1258 169, 185–193.

1259 Tekippe, J.A., Tacoma, R., Hristov, A.N., Lee, C., Oh, J., Heyler, K.S., Cassidy, T.W.,
1260 Varga, G.A., Bravo, D., 2013. Effect of essential oils on ruminal fermentation and
1261 lactation performance of dairy cows. *J. Dairy Sci.* 96, 7892-7903.

1262 Terré, M., Castells, L., Fàbregas, F., Bach, A., 2013. Comparison of pH, volatile fatty
1263 acids, and microbiome of rumen samples from preweaned calves obtained via
1264 cannula or stomach tube. *J. Dairy Sci.* 96, 5290–5294.

1265 Theodorou, M.K., Williams, B.A., Dhanoa, M.S., McAllan, A.B., France, J., 1994. A
1266 simple gas production method using a pressure transducer to determine the
1267 fermentation kinetics of ruminant feeds. *Anim. Feed Sci. Technol.* 48, 185–197.

1268 Tilley, J.M.A., Terry, R.A., 1963. A two-stage technique for the in vitro digestion of
1269 forage crops. *J. Br. Grass Soc.* 18, 104–111.

1270 Trbalza-Marinucci, M., Poncet, C., Delval, E., Fonty, G., 2006. Evaluation of
1271 techniques to detach particle-associated microorganisms from rumen contents.
1272 *Anim. Feed Sci. Technol.* 125, 1–16.

1273 Van Kessel J.A.S., Russell, J.B., 1996. The effect of pH on ruminal methanogenesis.
1274 *FEMS Microbiol. Ecol.* 20, 205–210.

1275 Váradyová, Z., M. Baran and I. Zelenák., 2005. Comparison of two in vitro
1276 fermentation gas production methods using both rumen fluid and faecal inoculum
1277 from sheep. *Anim. Feed Sci. Technol.* 123-124:81-94.

1278 Waghorn, G.C., Woodward, S.L., Tavendale, M., Clark, D.A., 2006. Inconsistencies in
1279 rumen methane production—effects of forage composition and animal genotype.
1280 *International Congress Series* 1293, 115–118.

1281 Wang, X.Z., Sunb, P.H. Janssenb, S.X. Tanga, Z.L. Tana., 2014. Responses of methane
1282 production and fermentation pathways to the increased dissolved hydrogen
1283 concentration generated by eight substrates in in vitro ruminal cultures. 2014.
1284 *Anim. Feed Sci. Technol.* 194, 1–11.

1285 Warner, D., Dijkstra, J., Hendriks, W.H. & Pellikaan, W.F., 2014. Stable isotope-
 1286 labelled feed nutrients to assess nutrient-specific feed passage kinetics in
 1287 ruminants. *J. Sci. Food Agric.* 94, 819-824.

1288 Weimer, P. J., B. S. Dien, T. L. Springer, and K. P. Vogel., 2005. In vitro gas
 1289 production as a surrogate measure of the fermentability of cellulosic biomass to
 1290 ethanol. *Appl. Microbiol. Biotechnol.* 67, 52–58.

1291 Wilkins, J., 1974. Pressure transducer method for measuring gas production by
 1292 microorganisms. *Appl. Microbiol.* 27, 135–140.

1293 Williams, A.G., Coleman, G.S., 1992. *The Rumen Protozoa*. New York, NY: Springer-
 1294 Verlag.

1295 Williams, B.A., 1998. Poster discussion session report. In: Deaville, E.R., Owen, E.,
 1296 Adesogen, A.T., Rymer, C.,

1297 Williams, Y.J., Popovski, S., Rea, S.M., Skillman, L.C., Toovey, A.F., Northwood,
 1298 K.S., Wright, A.D.G., 2009. A vaccine against rumen methanogens can alter the
 1299 composition of archaeal populations. *Appl. Environm. Microbiol.* 75, 1860–1866.

1300 Yan, T., Agnew, R.E., Gordon, F.J., Porter, M.G., 2000. Prediction of methane energy
 1301 output in dairy and beef cattle offered grass silage-based diets. *Livest. Prod. Sci.*
 1302 64, 253–263.

1303 Zhou X, Zeitz JO, Meile L, Kreuzer M, Schwarm A (2015) Influence of pH and the
 1304 degree of protonation on the inhibitory effect of fatty acids in the ruminal
 1305 methanogen *Methanobrevibacter ruminantium* strain M1, *J Appl Microbiol*, in
 1306 press, doi: 10.1111/jam.12955.

1307 Zhou, Z., Q. Meng, Z. Yu., 2011. Effects of methanogenic inhibitors on methane
 1308 production and abundance of methanogen and cellulolytic bacteria in in vitro
 1309 ruminal cultures. *Appl. Environ. Microbiol.* 77, 2634–2639.

1 Table 1.

2 Typical *in vitro* fermentation techniques (IVFT) used to quantify CH₄ (and CO₂ and H₂) production (adapted from Rymer et al., 2005).

In vitro system	Device and volume	Inoculum	Inoculum collection time	Incubation volume, ml	Inoculum: medium ratio	Buffer reference	Duration of incubation, h	Dietary substrate, mg	Gas release/collected	Pressure control	Gas sampling and analysis (GC)
Menke et al. (1979)	Syringes kept in rotor	Liquid phase	Before feeding	30	1:3	Menke et al. (1979)	24	200-300	Manual/End-point sampling	Yes, moveable glass piston	Manual
Theodorou et al. (1994)	Bottle in incubator	Liquid and solid phases	Before feeding	60	1:5	Theodorou (1993)	24-72	500-1000	Manual/End-point sampling	No, pressure increases	Manual
Mauricio et al. (1999)	Bottle in incubator	Liquid phase	Before feeding	100	1:10	Theodorou (1993)	n.s.	1000	Manual/End-point sampling	No, pressure increases	Manual
Pell and Schofield (1993)	Bottle and stirrer kept in incubator	Liquid phase	2 h after feeding	10	1:5	Goering and Van Soest (1970)	n.s.	100	Manual/End-point sampling	No, pressure increases	Manual, CH ₄ estimated by difference to CO ₂
Cone et al. (1996)	Bottle in shaking water bath	Liquid phase	2 h after feeding	100	1:3	Steingass (1983)	48	400-500	Automated/Fixed pressure	Yes	Manual
Davies et al. (1998)	Bottle in incubator	Liquid and solid phases	Before feeding	100	1:10	Theodorou (1993)	n.s.	1000	Automated/Fixed pressure	Yes	n.s.
Cornou et al. (2012)	Bottle in incubator	Liquid phase	n.s.	60	1:3	Menke and Steigass (1988)	72	500	Automated/Fixed pressure	Yes	Manual
Muetzel et al. (2014)	Bottle in incubator	Liquid phase	n.s.	60	1:5	Mould et al. (2005)	48	600	Automated/Automated	Yes	Automated GC

3

4 GC, gas chromatograph; n.s., not specified

5

6

7

8

9 Table 2.

10 Comparison of results based on *in vitro* gas production and studies *in vivo* evaluating the effect of a) diets or b) feed additives on methane production

Diets	Animal species /Intake level	Adaptation period	IVFT used	CH ₄ <i>in vivo</i>	R ² , P value	Reference	
Straw ± NH ₃ /NaOH	Sheep / Maintenance	10 days	Menke et al. (1979) ³	Open circuit respiration chamber, g/kg OM	0.89, <i>P</i> <0.0001	Blümmel et al. (2005)	
AH, CS+SBM, RG+SBM, RS+SBM, SG+SBM ¹	Cattle / Maintenance	12 days	Menke and Steingass (1988)	SF6 ml/kg DMI	0.98, <i>P</i> <0.0001	Bhatta et al. (2007)	
GS:CO 60:40 ²	Cattle / 95 % voluntary DMI	12 days	Cone et al. (1996)	Open circuit respiration chamber g/kg OMI (a) or OMFR (b)	0.04, <i>P</i> = 0.878 (a) 0.54, <i>P</i> = 0.040 (b)	Hatew et al. (2015)	
Additive ⁴	Dose <i>in vitro</i> // Inclusion rate <i>in vivo</i>	Animal species	Adaptation period	IVFT used/ incubation time	CH ₄ <i>in vivo</i>	Change in CH ₄ (% , L/kg DMI) <i>in vitro</i> // <i>In vivo</i>	Reference
Horseradish oil	0.17, 0.85 and 1.7 g/L // 20 g, kg DM	Cattle	14 days	Russel and Martin 1984/6 h	Headhood collection chamber	-18, -89 // -18.6	Mohamed et al. (2004)
PTS	100, 320 µl/L // 50, 100, 200 mg/L rumen content	Goats	7 days	Theodorou et al. (1994)/24 h	Open circuit respiration chamber	-28, -96 // -13, -18	Martínez-Fernández et al. (2013)
BCM	100, 320 µl/L // 50, 100, 160 mg/L rumen content	Goats	7 days	Theodorou et al. (1994)/24 h	Open circuit respiration chamber	-94, -96 // -34, - 45	Martínez-Fernández et al. (2013)
E3NP	25, 50 µM // 100 mg/d	Sheep	30 days	Theodorou et al. (1994)/24 h	Open circuit respiration chamber	-90, - 96 // -21	Martínez-Fernández et al. (2014)

3NOP	33, 66 µm // 100 mg/d	Sheep	30 days	Theodorou et al. (1994), 24 h	Open circuit respiration chamber	-99, -99 // -24	Martínez-Fernández et al. (2014)
Blend essential oils	0.25, 0.5, 1, 2, 5, 15, 30 mg/L // 0.2 g/day	Cattle	42 days	Theodorou et al. (1994), 24 h, 72 h	Open circuit respiration chamber	0 // - 15 (dairy cattle), 0 (beef cattle)	Castro-Montoya et al. (2015)

¹AH=alfalfa hay, CS= corn silage, SBM=soyabean meal, RS=rice straw, SG=sudan grass; ²GS= grass silage + 4 types of concentrate containing low or high amounts of slowly or rapidly degraded sources of starch; ³CH₄ estimated from fermentation end products. ⁴PTS: propyl thiosulfanite, BCM: brochloromethane, E3NP: ethyl-3-nitrooxy propionate, 3NPOP: and 3-nitrooxypropanol, 3NOP

14 Table 3.

15 Summary of technical recommendations on the use of *in vitro* gas production methods for measuring methane production

Aspect		Recommendation
Donor animals		
Species		Donor animals should be the same as target species
Numbers		3 or more animals where possible, minimum 2 animals.
Replications		Minimum of 3 independent incubation runs
Diet		Donor animals should be fed the same diet as incubated or of similar nutrient composition
Adaptation		Collect inoculum from animals that have been adapted to treatment for at least 2 weeks
Microbial inoculum		
Collection		Collection before feeding unless the experimental design requires a different sampling regimen.
Preservation		Ideally maintain inocula under anaerobic conditions at 39°C and use within 1 h. Preservation at 0°C up to 6 h or after freezing in liquid N are viable alternatives.
Processing		Filter through 250 µm pore size cloth.
Procedure		
Substrate incubated		Use a range of feeds for additives screening. Include a blank substrates to control for intra and inter variability and as a reference for comparisons between laboratories.
Buffer composition		No single buffer is recommended. For all incubations adjust bicarbonate concentration to maintain a target culture pH culture. Concentrations should not exceed 80 mM.
Inoculum:medium ratio		Adjust depending on the duration of incubation and frequency of gas sampling.
Duration of incubation		Adjusted according to the composition and particle size of substrate based on retention time in the rumen <i>in vivo</i> .

Gas measurements	Determine CH ₄ and CO ₂ when gas pressure is vented.
Units	Preferably express data relative to substrate (DM, OM, NDF) degradation. If this is not possible, data should be expressed relative to total gas, CO ₂ or VFA production.
Mandatory technical details to be reported in peer-reviewed publications	Numbers of donor animals used to obtain inocula. Composition of diet fed to donor animal and the time on diet. Inoculum collection time relative to feeding. Headspace gas composition. Pore size of filter used to strain rumen fluid. Buffer and media composition. Rumen inoculum storage conditions, both temperature and the time from collection to inoculation. Ratio of medium to rumen inoculum. Substrate processing procedures, frequency of gas sampling and CH ₄ measurements

16

17

18