Coarse woody debris and the carbon balance of a moderately disturbed forest

Amy V. Schmid^a, Christoph S. Vogel^b, Eli Liebmann^c, Peter S. Curtis^d, Christopher M. Gough^{a*}

^a Virginia Commonwealth University, Department of Biology, Richmond, VA 23284-2012, USA

^b University of Michigan, Biological Station & Department of Ecology and Evolutionary
Biology, Pellston, MI 49769, USA

^c Macalester College; 1600 Grand Ave., Saint Paul, MN 55105, USA

^d Ohio State University, Department of Evolution, Ecology and Organismal Biology, Columbus,
OH 43210, USA

*Author for correspondence:

Christopher M. Gough

Virginia Commonwealth University

Department of Biology

Box 842012

1000 W. Cary Street

Richmond, VA 23284-2012

Phone: 804-827-8596

Email: cmgough@vcu.edu

Abstract

Forested landscapes are shaped by disturbances varying in severity and source. Moderate disturbance from weather, pathogens, insects, and age-related senescence that kills only a subset of canopy trees may increase standing woody debris and alter the contribution of coarse woody debris (CWD) to total ecosystem respiration (R_E). However, woody debris carbon (C) dynamics are rarely examined following moderate disturbances that increase standing dead wood pools. We used an experimental manipulation of moderate disturbance in an upper Great Lakes forest to: 1) examine multi-year changes in CWD mass through a moderate disturbance; 2) quantify *in situ* CWD respiration during different stages of decay for downed and standing woody debris and; 3) estimate the annual contribution of CWD respiration to the ecosystem C balance through comparison with R_E and net ecosystem production (NEP). Six years following disturbance, we found that the standing dead wood mass of $24.5 \text{ Mg C ha}^{-1}$ was an order of magnitude greater than downed woody debris mass and a large source of ecosystem C flux. Instantaneous *in situ* respiration rates from standing and minimally decayed downed woody debris were not significantly different from one another. Separate estimates of ecosystem CWD respiration of 1.1 to $2.1 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$ six years following disturbance were comparable in magnitude to NEP and 12.5 % to 23.8 % of R_E , representing a substantial increase relative to pre-disturbance levels. Ecosystem respiration and NEP were stable following moderate disturbance even though ecosystem CWD respiration increased substantially, suggesting a reduction in the respiratory C contribution from other sources. We conclude that standing and downed CWD can be essential components of the ecosystem C balance following moderate severity disturbance.

43

44 *Keywords:* Coarse woody debris; Disturbance; Carbon cycling; Respiration; Wood

45 decomposition; Standing woody debris

46

47

48

49

50

51

52

53

54

55

56

1. Introduction

Forest disturbances alter the balance between ecosystem carbon (C) uptake and loss, and are primary determinants of the terrestrial C balance (Bond-Lamberty *et al.*, 2007; Gough *et al.*, 2007; Amiro *et al.*, 2010; Pan *et al.*, 2011). Tree mortality from disturbance may affect ecosystem C balance in two ways: by reducing the amount of C fixed via canopy photosynthesis (or gross primary production, GPP), and by altering the quantity of C lost through ecosystem respiration (R_E), particularly as detritus-fueled microbial respiration increases (Liu *et al.*, 2006; Harmon *et al.*, 2011). The difference between these two large opposing C fluxes determines ecosystem C balance, or net ecosystem production (NEP), with a small post-disturbance change in either GPP or R_E potentially causing large alterations in NEP. Numerous studies show that tree mortality can considerably reduce NEP, with some forests becoming C sources immediately after severe, stand-replacing disturbance (Janisch and Harmon, 2002; Amiro *et al.*, 2010; Harmon *et al.*, 2011; Hicke *et al.*, 2012). The effects of moderate disturbances that kill only a subset of canopy trees – such as those caused by insect pests, pathogens, and severe weather – on NEP are less certain even as lower severity disturbances increase in frequency and extent in many forested regions (Amiro *et al.*, 2010). Coarse woody debris (CWD), defined here as standing and downed dead wood > 10 cm diameter, is an important source of detritus following disturbance, but is often unaccounted in estimates of R_E and, as a result, the extent to which CWD C cycling dynamics following disturbance alter ecosystem C balance is poorly understood (Amiro *et al.*, 2010; Harmon *et al.*, 2011; Renninger *et al.*, 2014).

Coarse woody debris C cycling dynamics are most thoroughly characterized for the extreme end-members on the disturbance severity continuum, with most prior work conducted severely and minimally disturbed forests (Gough *et al.*, 2007; Harmon *et al.*, 2011); considerably

less is known about CWD C cycling dynamics following moderate disturbances that kill a substantial fraction of, but not all, canopy trees within an ecosystem (Renninger *et al.*, 2014). Studies conducted in recently undisturbed forests show that CWD is a relatively small C pool that contributes little to R_E and thus minimally affects NEP (Harmon *et al.*, 2004; Janisch *et al.*, 2005; Liu *et al.*, 2006; Gough *et al.*, 2007; Luyssaert *et al.*, 2008; Tang *et al.*, 2008). Following severe, stand-replacing disturbance, CWD is a primary substrate for microbial decomposition and, therefore, a large component of R_E and a source of C losses that reduce NEP (Amiro *et al.*, 2010; Harmon *et al.*, 2011; Russell *et al.*, 2014; Woodall *et al.*, 2015). How CWD progresses through stages of decay and contributes to R_E following more moderate disturbance is less clear, though one recent study suggests that accounting for changes in CWD mass and respiration is essential to interpreting how and why NEP may shift following lower intensity disturbances (Renninger *et al.*, 2014).

The motivation for studying forest C, including CWD, dynamics across a range of disturbance severities stems from observation and theory that suggests moderate disturbances modify ecosystem structure and function differently than severe disturbances (Nave *et al.*, 2011), and these differences extend to CWD structure, mass and decomposition (Harmon *et al.*, 2011). Some moderate disturbances initially leave more CWD standing than downed, with implications for woody debris microclimate and microbial colonization, and therefore microbial activity and CWD respiration (Pedlar *et al.*, 2002; Forrester *et al.*, 2012; Forrester *et al.*, 2013; Brazee *et al.*, 2014; Lewandowski *et al.*, 2015; White *et al.*, 2015). For example, partial defoliation of forest canopies from insects increased standing dead wood mass, while severe disturbances such as clear-cut harvesting and fire felled, removed, or reduced standing wood (Pedlar *et al.*, 2002; Renninger *et al.*, 2014). Moreover, standing, but not downed, CWD has increased broadly in

eastern US forests owing to patchy disturbance and natural tree senescence, with uncertain consequences for the C cycle (Woodall *et al.*, 2015). Yet, C cycling studies tend to focus on downed rather than increasingly prevalent standing CWD, concluding the latter contributes nominally to detritus pools and decays slowly, therefore contributing little to R_E (Harmon *et al.*, 1986; Yatskov *et al.*, 2003; Liu *et al.*, 2006; Jomura *et al.*, 2008; Tang *et al.*, 2008). However, a recent study reported similar *in situ* respiration rates for standing and downed CWD, with standing woody debris contributing substantially to R_E (Renninger *et al.*, 2014).

We evaluated CWD dynamics following a moderate forest disturbance at the University of Michigan Biological Station (UMBS) in which all mature aspen (*Populus*) and birch (*Betula*) were killed via experimental stem girdling but not immediately felled. The treatment produced a forest structure and composition similar to that which is broadly emerging in the upper Great Lakes region as secondary forests reach the century mark in age (Gough *et al.*, 2010). The experimental disturbance, which temporarily reduced live tree basal area by 39 % and leaf area index by 44 %, is similar in severity to increasingly prevalent naturally occurring disturbances in the region (Gough *et al.*, 2013). Prior C cycling studies resulting from this experiment emphasized NEP, GPP and net primary production (NPP) following disturbance, demonstrating sustained production in the moderately disturbed forest despite the experimental disturbance-related transfer of 35 Mg ha⁻¹ of wood from live to dead pools (Nave *et al.*, 2011; Gough *et al.*, 2013; Stuart-Haentjens *et al.*, 2015). Our prior emphasis on the unexpected resistance of forest production to moderate disturbance revealed mechanisms underlying sustained C uptake, notably improved canopy resource-use efficiency and the rapid reallocation of limiting nitrogen and light resources in support of leaf area recovery. Here, we provide a first focus on C losses, asking whether a large pulse of CWD that followed moderate disturbance was a prominent contributor

to R_E and, by extension, how this C lost by the ecosystem from CWD decomposition affects NEP. Specific objectives were to: 1) examine multi-year changes in CWD pools through moderate disturbance; 2) quantify *in situ* CWD respiration during different stages of decay for downed and standing woody debris and; 3) calculate the annual contribution of CWD respiration to the ecosystem C balance through comparison with R_E and NEP. We hypothesized that the large influx of CWD mass would constitute a substantial fraction of R_E and, because the disturbance did not cause the immediate felling of dead trees, standing woody debris would be the principle contributor to the respiratory flux from CWD.

2. Methods

2.1 Site Description

The study was conducted at the University of Michigan Biological Station (UMBS) in northern, lower Michigan, USA (45° 35N 84° 43W). The site has a mean annual temperature of 5.5° C and a mean annual precipitation of 817 mm (1942-2003) (Gough *et al.*, 2013). The UMBS forest is a representative secondary broadleaf deciduous forest in the transition zone between temperate and boreal forests. The forest developed following a clear-cut and wildfire regime in the early 20th century and has since undergone only low severity, patchy disturbances. Early successional species such as bigtooth aspen (*Populus grandidentata*), trembling aspen (*Populus tremuloides*), and paper birch (*Betula papyrifera*) are nearing or past maturity and senescing naturally, or in response to experimental disturbance described below. Canopy dominance is shifting toward longer-lived, later successional species, including red oak (*Quercus rubra*), red maple (*Acer rubrum*), and to a lesser extent, sugar maple (*Acer saccharum*) and eastern white pine (*Pinus strobus*). The average overstory tree age is 100 years old (Gough *et al.*,

2007). Downed woody debris represented a pool of 2.2 Mg C ha⁻¹ or 1.2 % of total ecosystem C prior to the decline of mature aspen and birch (Gough *et al.*, 2007).

The Forest Accelerated Succession Experiment (FASET) is an ecosystem-level manipulation that was initiated in May 2008 to quantify how forest C pools and fluxes are affected by moderate disturbance, ecological succession, and climate change. All early successional aspen and birch trees (~6,700) were killed via stem girdling within a 39 ha area. Carbon dioxide exchange between the atmosphere and the forest is continuously measured using a meteorological tower established in 2007, with estimates of annual NEP and R_E through 2013 reported by Gough *et al.* (2013) and Bond-Lamberty *et al.* (2015). Coarse woody debris mass, respiration, and microclimate measurements, detailed below, occurred in the experimentally disturbed area positioned within the meteorological flux tower footprint and no more than 300 m away from the tower base.

2.2 Coarse woody debris carbon mass

In 2009, 2011, 2013, and 2014, downed coarse woody debris mass was inventoried by decay class in eight, 0.1 ha subplots within the experimentally manipulated area. In the first sampling year, all standing and downed CWD samples were tagged for repeated measures in subsequent years of volume and decay class; newly produced CWD identified for the first time in following sampling years was tagged and included in future inventories. Each inventory year, all downed woody debris (> 10 cm diameter, < 45° from the forest floor) samples within a plot were individually classified into one of the following decay classes (Marra and Edmonds, 1994): (1) recently downed with bark and tissue fully intact, (2) sapwood is still present but beginning to show signs of decay and bark may be beginning to peel or crack, (3) heartwood is intact,

sapwood is present but softening, (4) heartwood is beginning to decay, sapwood and bark are mostly gone, (5) heartwood shows signs of substantial decay, sapwood and bark are completely missing. Standing woody debris (> 45° from the forest floor), which was identified as an additional category of woody debris, was inventoried in the same plots as downed wood in 2011 and 2014.

We measured the length and diameter, at the ends and middle, of each piece of downed CWD and measured total height, from sample base to top, and diameter at breast height of each piece of standing woody debris. The surface area and volume of each piece of CWD were calculated using the equation for the frustum of a cone (Harmon and Sexton, 1996). Site- and species-specific wood densities and C concentrations for each decay class were used to calculate C mass from CWD volume (Gough *et al.*, 2007). In the case of standing CWD, which is difficult to representatively sample without felling, we used the wood density and C fraction of downed CWD, decay class 1 since values for these two categories are comparable (e.g., Koster *et al.*, 2015).

2.3 Coarse woody debris respiration and microclimate

Instantaneous *in situ* respiration from the surface of aspen CWD (R_{CWD} ; $\mu\text{mol m}^{-2} \text{s}^{-1}$) was repeatedly measured on downed and standing aspen CWD in the field during July and August 2014. Though aspen and birch were both targeted in the disturbance experiment, aspen comprised over 80 % of the biomass stem girdled (Gough *et al.*, 2010) and has similar decay characteristics as birch (Russell *et al.*, 2014); therefore, our analysis focused on aspen respiration. Five PVC respiration collars (2 cm high, 10 cm diameter) were permanently affixed with duct seal putty to each of three replicate CWD samples (> 1.5 m long, > 10 cm center

diameter) per downed decay class and three standing CWD samples (> 10 cm diameter at breast height). The five subsample collars were evenly spaced and positioned at random angles that did not interfere with measurements along each CWD replicate, including the ends of downed woody debris and the tops of standing woody debris when possible. For standing dead wood, collar heights above the ground ranged from 1.2 to 12.5 m. We measured R_{CWD} weekly beginning early July for downed CWD and mid-July for standing CWD through mid-August using a Li-Cor LI-6400 Portable Photosynthesis System (Li-Cor, Lincoln, NE, USA). Wood temperature (1 cm depth; T_{CWD}) and wood moisture (12 cm integrated depth; ϕ_{CWD}) were measured concurrently with R_{CWD} using a type-E thermocouple and a Campbell Scientific HydroSense II soil moisture sensor (Model HS2-12, Campbell Scientific, Logan, UT, USA), using a drill when necessary to produce holes for sensor placement in standing and downed CWD. Wood water content was converted from volumetric to gravimetric values using site-specific CWD densities (Gough *et al.*, 2007).

2.4 Annual carbon flux estimates

We estimated the annual respiratory C flux from downed and standing CWD ($Mg\ C\ ha^{-1}\ yr^{-1}$), hereafter termed ecosystem CWD respiration, for 2014 using three different methods. We employed three different approaches because of the high uncertainty associated with estimates of heterotrophic respiration (Harmon *et al.*, 2011), focusing on 2014 because this was the year in which we measured *in situ* CWD respiration. A mass balance estimate of ecosystem CWD respiration was calculated as the change in total woody debris mass across all decay classes over one year, from 2013 to 2014. Standing dead wood was inventoried in 2011 and 2014 only, with mass values for 2013 estimated from linear interpolation between measurement years (Figure 1).

218 In this approach, ecosystem CWD respiration is the summed or aggregate change in CWD mass,
 219 including those that transitioned to a more advanced decay class. One limitation of this approach
 220 is that, similar to other cohort-based mass balance approaches (e.g., Johnson *et al.*, 2001), it does
 221 not account for CWD production and losses occurring between inventories. The total annual loss
 222 of C mass from CWD was assumed to be equal to the flux to the atmosphere, as CWD-C
 223 leaching and export is considered negligible (Harmon *et al.*, 1986; Laiho and Prescott, 1999). A
 224 second approach integrated measured R_{CWD} over time, with fluxes interpolated between
 225 measurement dates. To estimate R_{CWD} outside of the measurement period, we linearly
 226 extrapolated fluxes to zero on the dates in the Spring and Autumn when temperatures $< 0^{\circ} \text{C}$;
 227 winter R_E (all sources) is $< 10\%$ of the annual total at our site (Curtis *et al.*, 2005). The
 228 integrated area under the R_{CWD} time series curve, shown in Figure 2a (illustrated without
 229 extrapolated flux values), was calculated using SigmaPlot (SYSTAT Inc., San Jose, CA, USA).
 230 We scaled fluxes to the ecosystem by multiplying integrated R_{CWD} by the surface area of downed
 231 and standing dead wood. Lastly, we used a published site- and species-specific model to infer
 232 half-hourly R_{CWD} from T_{CWD} and ϕ_{CWD} ($R_{CWD} = -1.28 * e^{-0.034 * T_{CWD}} + 0.25 * \ln(f_{CWD})$; $r^2 = 0.64$, $P <$
 233 0.001 ; Gough *et al.*, 2007). Because continuous ϕ_{CWD} and T_{CWD} were required to model R_{CWD}
 234 but not available from direct measurements, regressions between (3 to 4 per sample) point
 235 measurements of ϕ_{CWD} and T_{CWD} and continuously (half-hourly) logged soil moisture and soil
 236 temperature, respectively, were developed following the approach of Gough *et al.*, 2007 to derive
 237 continuous half-hourly estimates of ϕ_{CWD} and T_{CWD} for 2014. Respiration was assumed to be
 238 zero when T_{CWD} was $< 0^{\circ} \text{C}$ (Gough *et al.*, 2007). Half-hourly estimates of R_{CWD} were summed
 239 and multiplied by decay class specific CWD mass to estimate ecosystem CWD respiration.
 240 Ecosystem CWD respiration was compared with published mean annual R_E and NEP (2009-

2013) obtained from meteorological tower measurements of the exchange of CO₂ between the forest and atmosphere (Gough *et al.*, 2013; Bond-Lamberty *et al.*, 2015); we report mean, rather than single-year, R_E because this derived C flux, made with relatively high uncertainty, did not change significantly over time following disturbance (Gough *et al.*, 2013), and therefore is a higher confidence estimate with which to compare ecosystem CWD respiration.

2.5 Statistical analysis

Statistical differences over time and among decay classes in CWD pools, R_{CWD}, ϕ_{CWD} , and T_{CWD} were assessed using a repeated measures analysis of variance (ANOVA), with values ± 1 S.E. presented in the text. Multiple comparisons were made using post hoc Fisher's LSD analysis, selected because our hypotheses were based on *a priori* knowledge of differences among decay classes in CWD pools, fluxes, and microclimate from an earlier, more tightly controlled field and laboratory manipulation conducted at our site prior to disturbance (Gough *et al.*, 2007). Linear and non-linear (2-parameter exponential function) regression analysis was used to examine decay class specific relationships between R_{CWD}, and both ϕ_{CWD} and T_{CWD} (Curtis *et al.*, 2005; Gough *et al.*, 2007) and standing woody debris measurement height, and ϕ_{CWD} and R_{CWD}. The uncertainty in each ecosystem CWD respiration estimate was expressed as a standard error (S.E.) derived from the cumulative error in model parameters (model approach only) and accounting for spatial variation in R_{CWD} (integration approach only) and CWD mass distribution (all approaches) (Gough *et al.*, 2007). All statistical analyses used a significance level of $\alpha = 0.05$ and were performed in SAS (ANOVA; SAS 9.1, SAS Institute, Cary, NC, USA) or SPSS (regression; IBM SPSS Statistics, version 22, IBM Corp., Armonk, NY, USA).

3. Results

3.1 Coarse woody debris carbon pools

The allocation of CWD among decay classes and the total mass of woody debris changed over time following moderate disturbance. One year after the stem-girdling treatment, in 2009, and prior to peak tree mortality, downed CWD mass was evenly distributed among decay classes, totaling $0.3 \pm 0.09 \text{ Mg C ha}^{-1}$ for all decay classes combined (Figure 1). By 2011, 97 % of stem-girdled aspen and birch were defoliated and considered dead, resulting in a 35 Mg C ha^{-1} pulse of CWD from experimental disturbance (Gough *et al.*, 2013) plus an additional 2.1 Mg C ha^{-1} of naturally senesced CWD in early stages of decomposition, including standing dead wood. In 2013 and 2014, following multiple years of decomposition, the mass of downed wood trended upward as standing CWD began to fall and advance through stages of decay (Figure 1); this increase over time was not statistically significant, (time main effect, $p = 0.27$; time x decay class interaction, $p = 0.39$), however, because of the very high spatial variation among replicate plots in CWD mass, a signature of disturbance-related increases in the spatial heterogeneity of C pools and fluxes (Gough *et al.*, 2013). Six years following disturbance in 2014, standing dead wood remained the largest pool of CWD ecosystem-wide at $24.5 \pm 7.6 \text{ Mg C ha}^{-1}$; downed dead wood comprised $1.5 \pm 0.7 \text{ Mg C ha}^{-1}$.

3.2 Coarse woody debris respiration and microclimate

Instantaneous *in situ* respiration from the surface of CWD, R_{CWD} , was variable among decay classes and over time (Figure 2A). Mean R_{CWD} varied by an order of magnitude among decay classes and across time, from $1.2 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ to $11.4 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$, with individual point measurements as high as $64.3 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$. In general, R_{CWD} increased with

increasing extent of decay (decay class main effect, $p = 0.008$), but significant pairwise differences ($p < 0.05$) were only observed between decay class 1 and decay classes 2, 4, and 5, and decay class 5 and standing dead wood, decay class 1, and decay class 3. Notably, standing dead wood and downed decay class 1 CWD had comparable R_{CWD} (Figure 2A). Wood water content, ϕ_{CWD} , increased with increasing extent of decay, up to decay class 4 (decay class main effect, $p < 0.0001$) (Figure 2B). Wood temperature, T_{CWD} , increased with increasing extent of downed wood decay (decay class main effect, $p = 0.04$), and was highest in downed decay classes 4 and 5, and in standing dead wood (Figure 2C).

Although T_{CWD} and ϕ_{CWD} varied considerably over time and among decay classes, temperature and moisture were not significant predictors of R_{CWD} in downed wood; contrastingly, mean moisture, which varied significantly by standing CWD height, was correlated with R_{CWD} in standing wood (Figure 3A). There was no significant relationship between downed R_{CWD} , and either ϕ_{CWD} or T_{CWD} when each point measurement was treated as an independent observation or when averaged across measurement dates for any decay class individually or when all decay classes were combined ($p > 0.1$; data not shown). In contrast to downed CWD, standing dead wood exhibited a significant decline in mean ϕ_{CWD} , averaged across dates, with increasing measurement height ($p < 0.0001$, $r^2 = 0.77$) (Figure 3A) and a significant relationship between mean ϕ_{CWD} and mean R_{CWD} ($p = 0.015$, $r^2 = 0.402$) (Figure 3B).

3.3 Annual carbon flux estimates

We used three different approaches to estimate the contribution of standing and downed CWD respiration to ecosystem respiration, R_E , in 2014: a mass balance approach calculating ecosystem CWD respiration from changes in C pools, a simple under the curve (Figure 2A)

integration and scaling of mean *in situ* instantaneous CWD respiration from point measurements, R_{CWD} , and a previously developed site-specific model driven by continuous ϕ_{CWD} and T_{CWD} estimations. Estimates of ecosystem CWD respiration from downed and standing dead wood combined did not differ significantly among approaches, totaling $1.1 \pm 1.1 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$ for the mass balance approach, $2.1 \pm 1.3 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$ for the integration approach, and $1.2 \pm 1.6 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$ for the modeling approach. Standing dead wood was the largest contributor to ecosystem CWD respiration in all estimations (Figure 4), comprising 65 %, 57 %, and 74 % of the total annual CWD respiration from the mass balance, integration, and model estimates, respectively. This large contribution from standing dead wood was a function of its high surface area relative to downed wood (Figure 1) rather than high R_{CWD} . Decay class 2 downed woody debris was the second largest source of respired C from CWD in all estimations. Total C flux from downed woody debris classes 1, 3, 4, and 5 contributed < 20 % of the total ecosystem CWD respiration in all estimations. Given total CWD mass (26 Mg C ha^{-1}) and ecosystem CWD respiration values (1.1 to $2.1 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$) six years following disturbance, the CWD decomposition rate-constant (ecosystem CWD mass/ecosystem CWD respiration) was between 0.4 and 0.8 yr^{-1} .

Ecosystem CWD respiration (1.1 to $2.1 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$) following moderate disturbance was a substantial contributor to R_E and similar in magnitude, but opposite in sign, to NEP. Mean R_E following disturbance (2009-2013) was an $8.81 \pm 0.23 \text{ Mg C ha}^{-1} \text{ yr}^{-1}$ flux to the atmosphere (Bond-Lamberty *et al.*, 2015) and did not differ significantly over time or from a nearby undisturbed, control forest (Gough *et al.*, 2013). The different estimation approaches suggest that ecosystem CWD respiration comprised 12.0 – 24.3 % of the total respiratory C flux ecosystem-wide 6 years following disturbance (Figure 4). Despite the mortality of > one-third of all canopy trees, the ecosystem remained a modest C sink following disturbance, with mean NEP (= GPP -

R_E) comparable to that of the control forest (Gough *et al.*, 2013). Ecosystem CWD respiration was only slightly lower in magnitude than mean NEP of $2.59 \pm 0.18 \text{ Mg C ha}^{-1}$ following disturbance (2009-2013) (Bond-Lamberty *et al.*, 2015).

4. Discussion

We have shown that ecosystem CWD respiration was a substantial contributor to R_E , and therefore a large C loss from the ecosystem, following a moderate forest disturbance, approaching NEP in magnitude. Our results indicate that standing woody debris mass, though spatially variable, was a particularly important contributor to the ecosystem's total CWD pool and respiration following moderate disturbance, with *in situ* standing woody debris respiration rates comparable to those of downed woody debris in the earliest stage of decay. Despite an increase in ecosystem CWD respiration following tree mortality, comparable levels of R_E before and after disturbance, and relative to a nearby undisturbed control site suggest that the respiratory contribution from other sources may have declined as ecosystem CWD respiration increased, helping to stabilize NEP.

Our experimental disturbance was similar in severity to moderate disturbances caused by insect defoliation, pathogen outbreaks, and extreme weather, resulting in a comparable input of CWD. Our experiment gradually transferred 35 Mg C ha^{-1} from live to dead wood pools, with standing woody debris accounting for $> 90 \%$ of the total CWD mass 6 years following disturbance. This value is considerably higher than CWD pools of 2.2 Mg C ha^{-1} in a nearby undisturbed forest (Gough *et al.*, 2007) and of 0.3 Mg C ha^{-1} one year following the disturbance treatment. Sudden oak death similarly increased standing woody debris and total CWD mass from 1.4 Mg C ha^{-1} to $33.9 \text{ Mg C ha}^{-1}$ (Cobb *et al.*, 2012). Gypsy moth defoliation increased

CWD mass to a lesser extent, from 2.5 Mg C ha⁻¹ to 11 Mg C ha⁻¹ three years post-disturbance, with standing woody debris making up 75 % of dead wood mass (Renninger *et al.*, 2014).

Although measurements of *in situ* CWD respiration, R_{CWD} , were highly variable within and among decay classes, and across time, our measured values are similar to previously reported field measurements expressed on a surface area basis from our site and others (Gough *et al.*, 2007; Forrester *et al.*, 2012; Renninger *et al.*, 2014). Our observation of increasing R_{CWD} with increasing extent of decay was previously observed at our site (Gough *et al.*, 2007) and others (Forrester *et al.*, 2012; Renninger *et al.*, 2014). In contrast, less decayed CWD respired at a higher rate than more moderately decayed wood following pest-related mortality in an oak forest (Renninger *et al.*, 2014). Our inability to detect significant relationships between downed R_{CWD} and known microclimatic drivers (Zhou *et al.*, 2007; Vanderhoof *et al.*, 2013) was associated with high variance in our field measurements, which are considerably more variable than controlled laboratory based measurements of whole CWD pieces (Gough *et al.*, 2007). Poor correlations between R_{CWD} and microclimate in the field may occur because CO₂ emitted from the CWD surface may originate far from the location of measurement and, therefore, be decoupled from the local microclimate (Teskey *et al.*, 2008). Nevertheless, we did detect a relationship between R_{CWD} and respiration collar height in standing dead wood, which corresponded with differences in ϕ_{CWD} .

Moderate disturbance caused a considerable increase in ecosystem CWD respiration without a corresponding increase in total R_E or a decrease in NEP, suggesting that the quantity respired C from other sources may have declined and, in doing so, stabilized ecosystem C losses. Though ecosystem CWD respiration estimates varied nearly 2-fold among the three approaches, this level of variation among methods is typical of heterotrophic C fluxes, which are made with

high uncertainty (Gough *et al.*, 2008b; Harmon *et al.*, 2011). Our estimates of ecosystem CWD respiration before and after moderate disturbance suggest an increase from 0.2 Mg C ha⁻¹ yr⁻¹ (Gough *et al.*, 2007) to a conservative estimate of 1.1 Mg C ha⁻¹ yr⁻¹, or from 2 % to 12.5 % of R_E six years after disturbance chiefly because of the increase in total downed and standing CWD mass. An increase in ecosystem CWD respiration without a concomitant increase in R_E suggests that the contributions of other respiratory sources shifted by a similar amount but in the opposite direction, stabilizing the total ecosystem respiratory C flux. Interestingly, this shift in the relative contributions of different respiratory C fluxes parallels findings from our site demonstrating that, following disturbance, a rapid compensatory increase in leaf photosynthesis and new growth by undisturbed vegetation maintained ecosystem C uptake, thereby sustaining GPP and NPP despite the mortality of more than a third of all canopy trees (Gough *et al.*, 2013; Bond-Lamberty *et al.*, 2015; Stuart-Haentjens *et al.*, 2015). Though we did not quantify respiration from other sources in the current study, soil respiration, the primary contributor to R_E in most ecosystems, temporarily declined at our site and others following disturbance perhaps due to an initial decrease in rhizosphere activity (Hogberg *et al.*, 2001; Bhupinderpal *et al.*, 2003; Nave *et al.*, 2011; Levy-Varon *et al.*, 2012; Moore *et al.*, 2013). Similarly, total plant respiration decreased following pine-beetle disturbance because fewer live trees remained (Moore *et al.*, 2013). Placed within the broader context of prior results from our experimental disturbance, sustained NEP in the first several years following moderate disturbance may be attributed to the maintenance of C uptake and primary production (Gough *et al.*, 2013) and, given the present findings, stability in C losses owing to compensatory shifts in respiratory sources that maintained, rather than increased, R_E.

We found that standing woody debris, an often unmeasured source of respiratory C, was the largest component of ecosystem CWD respiration, with relatively high *in situ* CWD respiration rates. Respiration from standing dead wood is infrequently measured or accounted for in ecosystem C budgets because it is assumed to be a small fraction of total CWD in recently undisturbed ecosystems (Harmon *et al.*, 2011), and have very low *in situ* respiration rates (Harmon *et al.*, 1986; Yatskov *et al.*, 2003; Liu *et al.*, 2006; Jomura *et al.*, 2008; Tang *et al.*, 2008). However, recent region-wide forest inventories suggest standing dead wood pool size, in contrast to downed CWD, is broadly increasing in aging forests of the Eastern US (Woodall *et al.*, 2015). In our moderate experimental disturbance standing wood was a large fraction of the total CWD pool, with *in situ* CWD respiration rates comparable to those for downed wood in decay class 1. Similarly, respiratory C losses from standing woody debris were substantial following disturbances that initially left dead trees standing rather than felled and when *in situ* standing CWD respiration rates were comparable to those of downed wood (Jomura *et al.*, 2008; Renninger *et al.*, 2014).

The large remaining pool of CWD that originated following moderate disturbance demonstrates the lasting, but presently uncertain, effect of disturbance on the forest's future C cycle. Six years following disturbance, 26.0 Mg C ha⁻¹ of mostly legacy woody debris remained in the ecosystem, a large C pool approaching 15 % of the ecosystem's total carbon stores (Gough *et al.*, 2008a). At present, the CWD decomposition rate-constant of 0.4 to 0.8 yr⁻¹ is slightly lower than the 0.9 yr⁻¹ reported for our undisturbed, control forest (Gough *et al.*, 2007) because of the dominance of slowly decaying standing CWD in the disturbed forest. However, CWD decomposition rate-constants, as the term implies, assume unchanged environmental conditions and decomposer activity, an assumption that is violated by disturbance and which suggests the

rate-constants we report here are likely to change in the future (Harmon *et al.*, 2011; Russell *et al.*, 2014; Russell *et al.*, 2015; Woodall *et al.*, 2015). Nonetheless, aspen CWD can take > 70 years to fully decompose (Russell *et al.*, 2014), suggesting that dead wood could remain a source of C until the end of this century. Although respiration from decomposing CWD may continue for decades, the contribution of ecosystem CWD respiration to R_E is likely to change. For example, respiration from other sources returned to pre-disturbance levels within several years of an ecosystem perturbation (Levy-Varon *et al.*, 2014; Trahan *et al.*, 2015). However, uncertainty exists in precisely when ecosystem CWD respiration will peak following disturbance. A lag in time between disturbance and peak R_E is commonly observed following perturbations that do not immediately kill or fell trees (Harmon *et al.*, 2011; Renninger *et al.*, 2014). In our system, the abrupt felling of standing CWD from wind or ice, for example, could accelerate decomposition and consequently increase ecosystem CWD respiration. Rates of dead wood decomposition could also be affected by climate change, as the climate in northern lower Michigan becomes warmer and drier with more variable precipitation (Duveneck *et al.*, 2014; Peterson *et al.*, 2014).

We conclude that following a moderate severity forest disturbance the accounting of downed and standing CWD pools and fluxes was essential to accurately quantifying the ecosystem C budget, and interpreting the relative stability in R_E and NEP following tree mortality. Our findings are in agreement with those for a moderately disturbed oak forest (Renninger *et al.*, 2014), but contrast with many other studies demonstrating that CWD was minimally important to the overall C budget during periods of low disturbance (Liu *et al.*, 2006; Gough *et al.*, 2007; Tang *et al.*, 2008). Standing woody debris, which our study suggests may respire more than often assumed, may be an especially important C pool and flux to quantify

following disturbances that do not immediately fell trees, such as insect defoliation, pathogen outbreaks, and age-related senescence. Finally, our work here and that of others from our site show that stability in the ecosystem C balance, NEP, during the first several years following disturbance was tied not only to sustained C uptake and primary production (Gough *et al.*, 2013; Stuart-Haentjens *et al.*, 2015) but also to unexpectedly stable R_E , despite a ≥ 5 -fold increase in ecosystem CWD.

Acknowledgements

Funding for the UMBS AmeriFlux core site was provided by the U.S. Department of Energy's (DOE) Office of Science and by the DOE Climate and Environmental Sciences Division, Office of Science (Award Nos. DE-SC0006708, DE-SC0006708). EL was supported by a UMBS Research Experience for Undergraduates (REU) program through NSF Award No. AGS-0851421. We thank UMBS for facilities and logistical support, three anonymous reviewers for their comments, and Drs. Rima Franklin, Scott Neubauer, and Julie Zinnert for reviewing our manuscript.

References

- Amiro, B.D., Barr, A.G., Barr, J.G., Black, T.A., Bracho, R., Brown, M., Chen, J., Clark, K.L., Davis, K.J., Desai, A.R., Dore, S., Engel, V., Fuentes, J.D., Goldstein, A.H., Goulden, M.L., Kolb, T.E., Lavigne, M.B., Law, B.E., Margolis, H.A., Martin, T., McCaughey, J.H., Misson, L., Montes-Helu, M., Noormets, A., Randerson, J.T., Starr, G., Xiao, J., 2010. Ecosystem carbon dioxide fluxes after disturbance in forests of North America. *Journal of Geophysical Research-Biogeosciences* 115.
- Bhupinderpal, S., Nordgren, A., Lofvenius, M.O., Hogberg, M.N., Mellander, P.E., Hogberg, P., 2003. Tree root and soil heterotrophic respiration as revealed by girdling of boreal Scots pine forest: extending observations beyond the first year. *Plant Cell and Environment* 26, 1287-1296.
- Bond-Lamberty, B., Fisk, J.P., Holm, J.A., Bailey, V., Bohrer, G., Gough, C.M., 2015. Moderate forest disturbance as a stringent test for gap and big-leaf models. *Biogeosciences* 12, 513-526.
- Bond-Lamberty, B., Peckham, S.D., Ahl, D.E., Gower, S.T., 2007. Fire as the dominant driver of central Canadian boreal forest carbon balance. *Nature* 450, 89-+.
- Brazee, N.J., Lindner, D.L., D'Amato, A.W., Fraver, S., Forrester, J.A., Mladenoff, D.J., 2014. Disturbance and diversity of wood-inhabiting fungi: effects of canopy gaps and downed woody debris. *Biodiversity and Conservation* 23, 2155-2172.

479 Cobb, R.C., Chan, M.N., Meentemeyer, R.K., Rizzo, D.M., 2012. Common Factors Drive
 480 Disease and Coarse Woody Debris Dynamics in Forests Impacted by Sudden Oak Death.
 481 Ecosystems 15, 242-255.

482 Curtis, P.S., Vogel, C.S., Gough, C.M., Schmid, H.P., Su, H.B., Bovard, B.D., 2005. Respiratory
 483 carbon losses and the carbon-use efficiency of a northern hardwood forest, 1999-2003. New
 484 Phytologist 167, 437-455.

485 Duveneck, M.J., Scheller, R.M., White, M.A., Handler, S.D., Ravenscroft, C., 2014. Climate
 486 change effects on northern Great Lake (USA) forests: A case for preserving diversity. Ecosphere
 487 5.

488 Forrester, J.A., Mladenoff, D.J., Gower, S.T., 2013. Experimental Manipulation of Forest
 489 Structure: Near-Term Effects on Gap and Stand Scale C Dynamics. Ecosystems 16, 1455-1472.

490 Forrester, J.A., Mladenoff, D.J., Gower, S.T., Stoffel, J.L., 2012. Interactions of temperature and
 491 moisture with respiration from coarse woody debris in experimental forest canopy gaps. Forest
 492 Ecology and Management 265, 124-132.

493 Gough, C.M., Hardiman, B.S., Nave, L.E., Bohrer, G., Maurer, K.D., Vogel, C.S., Nadelhoffer,
 494 K.J., Curtis, P.S., 2013. Sustained carbon uptake and storage following moderate disturbance in a
 495 Great Lakes forest. Ecological Applications 23, 1202-1215.

496 Gough, C.M., Vogel, C.S., Hardiman, B., Curtis, P.S., 2010. Wood net primary production
 497 resilience in an unmanaged forest transitioning from early to middle succession. *Forest Ecology*
 498 *and Management* 260, 36-41.

499 Gough, C.M., Vogel, C.S., Kazanski, C., Nagel, L., Flower, C.E., Curtis, P.S., 2007. Coarse
 500 woody debris and the carbon balance of a north temperate forest. *Forest Ecology and*
 501 *Management* 244, 60-67.

502 Gough, C.M., Vogel, C.S., Schmid, H.P., Curtis, P.S., 2008a. Controls on annual forest carbon
 503 storage: Lessons from the past and predictions for the future. *Bioscience* 58, 609-622.

504 Gough, C.M., Vogel, C.S., Schmid, H.P., Su, H.B., Curtis, P.S., 2008b. Multi-year convergence
 505 of biometric and meteorological estimates of forest carbon storage. *Agricultural and Forest*
 506 *Meteorology* 148, 158-170.

507 Harmon, M.E., Bible, K., Ryan, M.G., Shaw, D.C., Chen, H., Klopatek, J., Li, X., 2004.
 508 Production, respiration, and overall carbon balance in an old-growth *Pseudotsuga-tsuga* forest
 509 ecosystem. *Ecosystems* 7, 498-512.

510 Harmon, M.E., Bond-Lamberty, B., Tang, J.W., Vargas, R., 2011. Heterotrophic respiration in
 511 disturbed forests: A review with examples from North America. *Journal of Geophysical*
 512 *Research-Biogeosciences* 116.

513 Harmon, M.E., Franklin, J.F., Swanson, F.J., Sollins, P., Gregory, S.V., Lattin, J.D., Anderson,
 514 N.H., Cline, S.P., Aumen, N.G., Sedell, J.R., Lienkaemper, G.W., Cromack, K., Cummins,
 515 K.W., 1986. Ecology of coarse woody debris in temperate ecosystems. *Advances in Ecological*
 516 *Research* 15, 133-302.

517 Harmon, M.E., Sexton, J., 1996. Guidelines for Measurements of Woody Detritus in Forest
 518 Ecosystems. (US LTER Publication No. 20) US LTER Network Office, University of
 519 Washington, Seattle, WA, USA.

520

521 Hicke, J.A., Allen, C.D., Desai, A.R., Dietze, M.C., Hall, R.J., Hogg, E.H., Kashian, D.M.,
 522 Moore, D., Raffa, K.F., Sturrock, R.N., Vogelmann, J., 2012. Effects of biotic disturbances on
 523 forest carbon cycling in the United States and Canada. *Global Change Biology* 18, 7-34.

524 Hogberg, P., Nordgren, A., Buchmann, N., Taylor, A.F.S., Ekblad, A., Hogberg, M.N., Nyberg,
 525 G., Ottosson-Lofvenius, M., Read, D.J., 2001. Large-scale forest girdling shows that current
 526 photosynthesis drives soil respiration. *Nature* 411, 789-792.

527 Janisch, J.E., Harmon, M.E., 2002. Successional changes in live and dead wood carbon stores:
 528 implications for net ecosystem productivity. *Tree Physiology* 22, 77-89.

529 Janisch, J.E., Harmon, M.E., Chen, H., Fasth, B., Sexton, J., 2005. Decomposition of coarse
 530 woody debris originating by clearcutting of an old-growth conifer forest. *Ecoscience* 12, 151-
 531 160.

532 Johnson, M.G., Tingey, D.T., Phillips, D.L., Storm, M.J., 2001. Advancing fine root research
 533 with minirhizotrons. *Environmental and Experimental Botany* 45, 263-289.

534 Jomura, M., Kominami, Y., Dannoura, M., Kanazawa, Y., 2008. Spatial variation in respiration
 535 from coarse woody debris in a temperate secondary broad-leaved forest in Japan. *Forest Ecology*
 536 *and Management* 255, 149-155.

537 Koster, K., Metslaid, M., Engelhart, J., Koster, E., 2015. Dead wood basic density, and the
 538 concentration of carbon and nitrogen for main tree species in managed hemiboreal forests. *Forest*
 539 *Ecology and Management* 354, 35-42.

540 Laiho, R., Prescott, C.E., 1999. The contribution of coarse woody debris to carbon, nitrogen, and
 541 phosphorus cycles in three Rocky Mountain coniferous forests. *Canadian Journal of Forest*
 542 *Research-Revue Canadienne De Recherche Forestiere* 29, 1592-1603.

543 Levy-Varon, J.H., Schuster, W.S.F., Griffin, K.L., 2012. The autotrophic contribution to soil
 544 respiration in a northern temperate deciduous forest and its response to stand disturbance.
 545 *Oecologia* 169, 211-220.

546 Levy-Varon, J.H., Schuster, W.S.F., Griffin, K.L., 2014. Rapid rebound of soil respiration
 547 following partial stand disturbance by tree girdling in a temperate deciduous forest. *Oecologia*
 548 174, 1415-1424.

549 Lewandowski, T.E., Forrester, J.A., Mladenoff, D.J., Stoffel, J.L., Gower, S.T., D'Amato, A.W.,
 550 Balser, T.C., 2015. Soil microbial community response and recovery following group selection
 551 harvest: Temporal patterns from an experimental harvest in a US northern hardwood forest.
 552 Forest Ecology and Management 340, 82-94.

553 Liu, W.H., Bryant, D.M., Hutyra, L.R., Saleska, S.R., Hammond-Pyle, E., Curran, D., Wofsy,
 554 S.C., 2006. Woody debris contribution to the carbon budget of selectively logged and maturing
 555 mid-latitude forests. Oecologia 148, 108-117.

556 Luyssaert, S., Schulze, E.D., Borner, A., Knohl, A., Hessenmoller, D., Law, B.E., Ciais, P.,
 557 Grace, J., 2008. Old-growth forests as global carbon sinks. Nature 455, 213-215.

558 Marra, J.L., Edmonds, R.L., 1994. Coarse woody debris and forest floor respiration in an old-
 559 growth coniferous forest on the Olympic Peninsula, Washington, USA. Canadian Journal of
 560 Forest Research-Revue Canadienne De Recherche Forestiere 24, 1811-1817.

561 Moore, D.J.P., Trahan, N.A., Wilkes, P., Quaife, T., Stephens, B.B., Elder, K., Desai, A.R.,
 562 Negron, J., Monson, R.K., 2013. Persistent reduced ecosystem respiration after insect
 563 disturbance in high elevation forests. Ecology Letters 16, 731-737.

564 Nave, L.E., Gough, C.M., Maurer, K.D., Bohrer, G., Hardiman, B.S., Le Moine, J., Munoz, A.B.,
 565 Nadelhoffer, K.J., Sparks, J.P., Strahm, B.D., Vogel, C.S., Curtis, P.S., 2011. Disturbance and
 566 the resilience of coupled carbon and nitrogen cycling in a north temperate forest. Journal of
 567 Geophysical Research-Biogeosciences 116.

568 Pan, Y.D., Birdsey, R.A., Fang, J.Y., Houghton, R., Kauppi, P.E., Kurz, W.A., Phillips, O.L.,
 569 Shvidenko, A., Lewis, S.L., Canadell, J.G., Ciais, P., Jackson, R.B., Pacala, S.W., McGuire,
 570 A.D., Piao, S.L., Rautiainen, A., Sitch, S., Hayes, D., 2011. A Large and Persistent Carbon Sink
 571 in the World's Forests. *Science* 333, 988-993.

572 Pedlar, J.H., Pearce, J.L., Venier, L.A., McKenney, D.W., 2002. Coarse woody debris in relation
 573 to disturbance and forest type in boreal Canada. *Forest Ecology and Management* 158, 189-194.

574 Peterson, D.L., Wolken, J.M., Hollingsworth, T.N., Giardina, C.P., Littell, J.S., Joyce, L.A.,
 575 Swanston, C.W., Handler, S.D., Rustad, L.E., McNulty, S.G., 2014. Regional Highlights of
 576 Climate Change. In: Peterson, D.L., Vose, J.M., PatelWeynand, T. (Eds.), *Climate Change and*
 577 *United States Forests*, pp. 113-148.

578 Renninger, H.J., Carlo, N., Clark, K.L., Schaefer, K.V.R., 2014. Modeling respiration from snags
 579 and coarse woody debris before and after an invasive gypsy moth disturbance. *Journal of*
 580 *Geophysical Research-Biogeosciences* 119, 630-644.

581 Russell, M.B., Fraver, S., Aakala, T., Gove, J.H., Woodall, C.W., D'Amato, A.W., Ducey, M.J.,
 582 2015. Quantifying carbon stores and decomposition in dead wood: A review. *Forest Ecology and*
 583 *Management* 350, 107-128.

584 Russell, M.B., Woodall, C.W., Fraver, S., D'Amato, A.W., Domke, G.M., Skog, K.E., 2014.
 585 Residence Times and Decay Rates of Downed Woody Debris Biomass/Carbon in Eastern US
 586 Forests. *Ecosystems* 17, 765-777.

587 Stuart-Haentjens, E.J., Curtis, P.S., Fahey, R.T., Vogel, C.S., Gough, C.M., 2015. Net primary
 588 production of a temperate deciduous forest exhibits a threshold response to increasing
 589 disturbance severity. *Ecology* 96, 2478-2487.

590 Tang, J.W., Bolstad, P.V., Desai, A.R., Martin, J.G., Cook, B.D., Davis, K.J., Carey, E.V., 2008.
 591 Ecosystem respiration and its components in an old-growth forest in the Great Lakes region of
 592 the United States. *Agricultural and Forest Meteorology* 148, 171-185.

593 Teskey, R.O., Saveyn, A., Steppe, K., McGuire, M.A., 2008. Origin, fate and significance of
 594 CO₂ in tree stems. *New Phytologist* 177, 17-32.

595 Trahan, N.A., Dynes, E.L., Pugh, E., Moore, D.J.P., Monson, R.K., 2015. Changes in soil
 596 biogeochemistry following disturbance by girdling and mountain pine beetles in subalpine
 597 forests. *Oecologia* 177, 981-995.

598 Vanderhoof, M., Williams, C., Pasay, M., Ghimire, B., 2013. Controls on the rate of CO₂
 599 emission from woody debris in clearcut and coniferous forest environments. *Biogeochemistry*
 600 114, 299-311.

601 White, S.D., Hart, J.L., Schweitzer, C.J., Dey, D.C., 2015. Altered structural development and
 602 accelerated succession from intermediate-scale wind disturbance in *Quercus* stands on the
 603 Cumberland Plateau, USA. *Forest Ecology and Management* 336, 52-64.

604 Woodall, C.W., Russell, M.B., Walters, B.F., D'Amato, A.W., Fraver, S., Domke, G.M., 2015.
605 Net carbon flux of dead wood in forests of the Eastern US. *Oecologia* 177, 861-874.

606 Yatskov, M., Harmon, M.E., Krankina, O.N., 2003. A chronosequence of wood decomposition
607 in the boreal forests of Russia. *Canadian Journal of Forest Research-Revue Canadienne De*
608 *Recherche Forestiere* 33, 1211-1226.

609 Zhou, X., Wan, S.Q., Luo, Y.Q., 2007. Source components and interannual variability of soil
610 CO₂ efflux under experimental warming and clipping in a grassland ecosystem. *Global Change*
611 *Biology* 13, 761-775.

612

Figure Captions

Figure 1. Downed and standing woody debris carbon pools by decay class in a moderately disturbed upper Great Lakes forest in which all aspen and birch trees were stem girdled in 2008. Pool sizes reflect inputs from the experimental disturbance (35 Mg C ha^{-1}) plus those from naturally occurring tree senescence. Note the different y-axes and scales for downed (left) and standing (right) woody debris. Means $\pm \frac{1}{4}$ S.E. for visual clarity.

Figure 2. Means ± 1 S.E. of *in situ* coarse woody debris respiration (R_{CWD} ; A), wood gravimetric water content (ϕ_{CWD} ; B), and wood temperature (T_{CWD} ; C) of downed and standing woody debris by decay class during Summer 2014.

Figure 3. Standing dead wood sample height in relation to mean wood water content (ϕ_{CWD} ; A) and mean *in situ* coarse woody debris respiration (R_{CWD}) in relation to mean wood water content (ϕ_{CWD} ; B). An outlier (half-filled circle), greater than two standard deviations from the mean, was excluded from the regression analysis of mean R_{CWD} and mean ϕ_{CWD} . Means across sample dates ± 1 S.E.

Figure 4. Annual ecosystem CWD respiration, 2014, estimated using three different approaches in comparison to mean net ecosystem production (NEP) and ecosystem respiration (R_E) for a Great Lakes forest, 2009-2013 by CWD decay class. Note that NEP and R_E are expressed on a separate y-axis and scale. Error bars are ± 1 S.E. and represent cumulative uncertainty from model error and spatial variation in the case of ecosystem CWD respiration, or meteorological flux estimation for NEP and R_E (see Gough et al. 2013).

Figure 1

Figure 2

Figure 3

Figure 4