1 Regional validation and improved parameterization of the 3-PG model for *Pinus taeda* stands

2
3 Carlos A. Gonzalez-Benecke^{a,*}, Robert O. Teskey^b, Timothy A. Martin^c, Eric J. Jokela^c, Thomas R. Fox^d,
4 Michael B. Kane^b and Asko Noormets^e

5
6 ^a Department of Forest Engineering, Resources and Management, 269 Peavy Hall, Oregon State University,
7 Corvallis, OR 97331, USA

8 ^b Warnell School of Forestry and Natural Resources, 180 E. Green St., University of Georgia, Athens, GA 30601,
9 USA

10 ^c School of Forest Resources and Conservation, P.O. Box 110410, University of Florida, Gainesville, FL 32611,
11 USA

12 ^d Department of Forest Resources and Environmental Conservation, 319 Cheatham Hall, Virginia Polytechnic
13 Institute and State University, Blacksburg, VA 24061, USA

14 ^e Department of Forestry and Environmental Resources, P.O. Box 8008, North Carolina State University, Raleigh,
15 NC 27695, USA

16
17
18
19
20 * Corresponding author (carlos.gonzalez@oregonstate.edu), phone: 541-737-4952, fax: 541-737-4316

21
22 **Keywords:** Forest modeling, physiological process-based model, loblolly pine, ecophysiology, stand dynamics,
23 regional analysis.

Abstract

The forest simulation model, 3-PG, has the capability to estimate the effects of climate, site and management practices on many stand attributes using easily available data. The model, once calibrated, has been widely applied as a useful tool for estimating growth of forest species in many countries. Currently, there is an increasing interest in estimating biomass and assessing the potential impact of climate change on loblolly pine (*Pinus taeda* L.), the most important commercial tree species in the southeastern U.S. This paper reports a new set of 3-PG parameter estimates for loblolly pine, and describe new methodologies to determine important estimates. Using data from the literature and long-term productivity studies, we parameterized 3-PG for loblolly pine stands, and developed new functions for estimating NPP allocation dynamics, biomass pools at variable starting ages, canopy cover dynamics, effects of frost on production, density-independent and density-dependent tree mortality and the fertility rating. The model was tested against data from replicated experimental measurement plots covering a wide range of stand characteristics, distributed across the southeastern U.S. and also beyond the natural range of the species, using stands in Uruguay, South America. We used the largest validation dataset for 3-PG, and the most geographically extensive within and beyond a species' native range. Comparison of modeled to measured data showed robust agreement across the natural range in the U.S., as well as in South America, where the species is grown as an exotic. Across all tested sites, estimations of survival, basal area, height, quadratic mean diameter, bole volume and above-ground biomass agreed well with measured values, with R^2 values ranging between 0.71 for bole volume, and 0.95 for survival. The levels of bias were small and generally less than 13%. LAI estimations performed well, predicting monthly values within the range of observed LAI. The results provided strong evidence that 3-PG could be applied over a wide geographical range using one set of parameters for loblolly pine. The model can also be applied to estimate the impact of climate change on stands growing across a wide range of ages and stand characteristics.

1. Introduction

Loblolly pine (*Pinus taeda* L.) is one of the fastest growing pine species and has been planted on more than 10 million ha in the southeastern U.S. (Huggett et al., 2013). Its native range covers a wide area from the Atlantic coast to eastern Texas and from northern Florida to southern New Jersey (Figure 1). Loblolly pine has also been introduced into many countries, and large-scale plantations for timber production are found in Argentina, Brazil, China, New Zealand, South Africa and Uruguay (Borders and Bailey, 2001; Fassola et al., 2010).

Forest simulation models can be used to estimate forest productivity or biomass under diverse management and/or climate scenarios, and those estimates are of interest to landowners, managers and researchers. The semi-process-based simulation model, 3-PG (Physiological Processes Predicting Growth; Landsberg and Waring, 1997), has been extensively used to estimate stand attributes such as volume growth or biomass dynamics (Landsberg et al., 2001; Stape et al., 2004; Fontes et al., 2006; Sampson et al., 2006; Coops et al., 2010; Bryars et al., 2013; Gonzalez-Benecke et al., 2014a). The model uses species-specific empirical tree- and stand-level traits in combination with physiological attributes to quantify Net Primary Production (NPP, Mg ha^{-1}), allocation to the various biomass pools, population dynamics and soil water balance (Landsberg and Sands, 2011).

The 3-PG model can be used to estimate the effects on stand productivity of management, site characteristics and climate and has been parameterized for different tree species (Booth et al., 2000; Law et al., 2000; Waring, 2000; Dye, 2001; Rodriguez et al., 2002; Sands and Landsberg, 2002; Almeida et al., 2004; Flores and Allen, 2004; Coops et al., 2005; Waring et al., 2008; Rodriguez et al., 2009; Coops et al., 2010; Perez Cruzado et al., 2011; Gonzalez-Benecke et al., 2014a). Even though the model has been parameterized for loblolly pine (Landsberg et al., 2001 and 2003; Sampson et al., 2006; Bryars et al., 2013), questions have arisen about the generality of these parameterizations and the accuracy of model predictions across the southeastern U.S. and in other countries (Bryars et al., 2013). The modular implementation of the 3-PG model allows improvement of specific sub-routines. Several have been reported as requiring further refinement, including NPP partitioning, stand mortality, light interception, canopy closure and the fertility rating (FR) (Pinjuv et al., 2006; Almeida et al., 2010; Landsberg and Sands, 2011; Bryars et al., 2013). To improve model performance we used long-term datasets to obtain new species-specific parameters, and changed the structure of the model by introducing some new species-specific functions.

The objectives of this study were to parameterize the 3-PG model for loblolly pine, and validate it using data from measurement plots covering a wide range of age, productivity, management and geographical distribution

in the southeastern U.S. and Uruguay, South America. We used published data and long-term productivity studies from five university-forest industry research cooperatives in the southeastern U.S.: the Forest Biology Research Cooperative (FBRC) at the University of Florida, the Forest Productivity Cooperative (FPC) at North Carolina State University and Virginia Tech University, the Forest Modelling Research Cooperative (FMRC) at Virginia Tech University, the Western Gulf Forest Tree Improvement Program (WGFTIP) at Texas A&M University, and the Plantation Management Research Cooperative (PMRC) at the University of Georgia. This collaborative effort was part of the integrative research of the Pine Integrated Network: Education, Mitigation and Adaptation Project (PINEMAP, <http://www.pinemap.org/>). These shared datasets represented the outcome of frequently re-measured permanent plots established for a range of purposes, from monitoring plots of operational plantations to studies that included a variety of replicated treatments such as genetics, planting density, fertilization, weed control and thinning.

As we had access to the large datasets of PINEMAP, we were able to revise some important parameters of 3-PG using the alternative methods suggested by Gonzalez-Benecke et al. (2014a) to estimate species-specific parameters for FR, monthly needlefall rate, canopy cover, density-independent and density-dependent tree mortality, bole volume bark fraction, mean tree height, stemwood specific gravity, and initial biomass pools at any starting age. In addition, we incorporated new functions for estimating NPP partitioning, quadratic mean diameter and effect of frost temperature. We suggest that this new set of parameters could be applied across the native range of loblolly pine in the southeastern U.S.; however, certain parameters, when necessary, could be modified to improve predictions in other parts of the world where loblolly pine is grown.

2. Materials and methods

2.1. An overview of the 3-PG model

The 3-PG model (Landsberg and Waring, 1997) is a stand-level model that uses monthly weather data (e.g., global radiation, rainfall, number of rainy days, number of frost days and mean minimum and maximum temperatures) to predict growth of even-aged, mono-specific stands. The model also requires initial values of site characteristics such as soil texture class and upper and lower limits of available soil water (mm), as well as stand data about initial age, latitude of stand location, stocking (trees ha⁻¹) and biomass (Mg ha⁻¹) in roots (WR), foliage (WF) and stem (stemwood + bark + branches, WS). The 3-PG model has different sub-modules to estimate NPP,

biomass allocation, population dynamics and soil water balance at monthly intervals. A detailed description of the model can be found in Landsberg and Waring (1997) and Landsberg and Sands (2011). In this study we used the 3-PG version 3-PGpjs2.7 (Sands, 2010), implemented as a Microsoft Excel spreadsheet. The user-interface was modified allowing the user to change FR using SI, and initial biomass using tree density, age, mean dbh and mean tree height.

2.2. Parameter Estimation

Table 1 shows a summary of stand characteristics of the studies used for model fitting and parameter estimation. SI was estimated as the mean height of dominant and co-dominant trees at a base age of 25 years. For all sites in the U.S., SI was estimated with the height measurement at the age closest to 25 years using the equation reported by Diéguez-Aranda et al. (2006). For sites in Uruguay, SI was estimated with an operational function used by Cambium Forestal Uruguay S.A. and a reference age of 15 years (Daniel Ramirez, personal communication).

Table 1. Summary of data used for parameter estimation.

Project	Site	Institution	n	Lat	Long	AGE (yrs)	Dq (cm)	Nha (trees ha ⁻¹)	BA (m ² ha ⁻¹)	SI (m)	Parameters Estimated
Ameriflux	US-NC2	NCSU	4	35.80	-76.67	15 - 16	25.0 - 26.4	630 - 650	31.0 - 34.2	20.3 - 22.7	I
IMPAC	Gainesville	FBRC	12	29.76	-82.29	3 - 25	1.0 - 24.9	692 - 1538	0.1 - 50.1	13.9 - 27.2	II, IV, V, VI, VII, VIII
PPINES	Sanderson	FBRC	96	30.24	-82.33	2 - 14	2.0 - 20.7	988 - 2964	0.4 - 45.5	17.1 - 30.7	III, IV
PPINES	Waverly	FBRC	86	31.13	-81.75	2 - 14	2.2 - 21.5	988 - 2964	0.5 - 52.9	24.4 - 31.7	III, IV, VIII
SAGCD	Various (18)*	PMRC	250			2 - 12	1.1 - 32.1	205 - 1810	0.1 - 54.8	14.0 - 29.2	III, VII
CPCD	Various (14)*	PMRC	192			4 - 12	0.1 - 24.3	401 - 4485	0.1 - 46.3	16.8 - 33.2	II, III, VI
GSSS	Various (25)*	WGFTIP	437			5 - 20	0.1 - 27.0	232 - 4483	0.1 - 55.0	18.3 - 30.4	III, VII
RS2	Various (11)*	FPC	141			4 - 23	0.2 - 30.9	156 - 2050	0.1 - 48.2	15.5 - 26.8	III
RS5	Various (8)*	FPC	196			11 - 31	14.1 - 30.3	225 - 2032	9.0 - 47.6	15.6 - 25.6	III
RS8	Various (26)*	FPC	96			9 - 33	7.5 - 30.7	252 - 2590	4.7 - 52.1	15.1 - 24.3	III
N.A.	Tacuarembó	CAMBIUM	24			3 - 34	6.9 - 40.1	256 - 1381	5.1 - 55.3	20.9 - 29.8	V, VII
TOTAL			1506			2 - 34	0.1 - 40.1	156 - 4485	0.1 - 55.3	13.9 - 33.2	

n: number of plots; AGE: range of age (yrs); Dq: range of quadratic mean diameter (cm); BA: range of basal area (m² ha⁻¹), SI: range of site index (m).

I: canopy conductance; II: NPP partitioning; III: density-dependent tree mortality; IV: canopy cover; V: needlefall; VI: dbh-Ht relationship; VII: Dq-STEM relationship; VIII: FR-SI relationship.

*: multiple sites were used.

All model fitting and data analyses were performed using SAS 9.3 (SAS Inc., Cary, NC, USA). When boundary line fitting was performed, we used the quantile regression procedure with a quantile threshold of 0.99. When multiple variables were included in the fitted model, we used a logarithm transformation and a stepwise procedure with a threshold significance value of 0.15 as variable selection criteria. All variables included in the

model with a variance inflation factor (VIF) larger than 5 were discarded, as suggested by Neter et al. (1996). As non-linear model fitting was carried out, an empirical R^2 (Myers, 2000) was determined as:

$$R^2 = 1 - \frac{SSE/df_e}{SST/df_t} \quad [1]$$

where SSE and SST are the sum of squares of residuals and total, respectively, and df_e and df_t are the degrees of freedom of error and total, respectively.

2.2.1. Canopy conductance

Estimates of maximum (MaxCond, $m s^{-1}$) canopy conductance and stomatal response to VPD (CoeffCond, mb^{-1}) were obtained using two years of data from the US-NC2 long-term eddy covariance site, located in the lower Coastal Plain of North Carolina, USA (39°48' N, 76°40' W) (Noormets et al., 2010). The stand was 15-16 years old, with peak projected LAI of 4.0-4.3 $m^2 m^{-2}$ and mean tree height of 12-15 m. Further details of the study sites and measurement techniques can be found in Domec et al. (2009) and Noormets et al. (2010). Similar to Gonzalez-Benecke et al. (2014a), MaxCond and CoeffCond were estimated using meteorological measurements recorded with an automated weather station, latent heat fluxes from eddy-covariance measurements, and canopy conductance for water vapor was computed using an inverted form of the Penman-Monteith equation.

2.2.2. NPP partitioning

Previous versions of 3-PG allocated NPP to the three main tree components (foliage, stem and roots) using the ratio of foliage to stem mass (pFS) as a function of tree diameter. To better understand allocation pattern dynamics we used data from two long-term, replicated studies: CPCD (Coastal Plain Intensive Culture/Density Studies, from PMRC) and IMPAC (Intensive Management Practices Assessment Center, from FBRC). The CPCD dataset consisted of 192 plots that ranged in age from 4 to 12 years and the IMPAC dataset consisted of 12 plots that ranged in age from between 3 to 25 years (Table 1). For each study, we had access to the repeated measures raw inventory data. For all plots, above-ground biomass (foliage, branch, bark, stemwood) was calculated using the general functions reported in Gonzalez-Benecke et al. (2014b). Aboveground Net Primary Production (ANPP, $Mg ha^{-1} year^{-1}$) was calculated for each measurement interval as the net increment in woody biomass + foliage production during that period. Woody Biomass included increments in woody biomass (Iw , $Mg ha^{-1} year^{-1}$) and the

biomass of dead trees as suggested by Martin and Jokela (2004). For each plot and measurement, needlefall (NF, Mg ha⁻¹ year⁻¹), branchfall (BF, Mg ha⁻¹ year⁻¹) and litterfall (LF, Mg ha⁻¹ year⁻¹) were calculated using the functions reported by Gonzalez-Benecke et al. (2012). Foliage production (If, Mg ha⁻¹ year⁻¹) was assumed to be equal to the average of needlefall for the measurement interval. Woody biomass production (Iw) was assumed to be equal to net increment in woody biomass and included BF for the measurement interval.

We fitted a non-linear model to estimate the ratio between NPP allocation to foliage and NPP allocation to stem (pFS). The original equation used in 3-PG was an exponential decay to a non-zero asymptote function that correlated Age and pFS (Landsberg and Waring, 1997). That model was later modified by Gonzalez-Benecke et al. (2014a) using BA instead of Age. Upon further analysis we examined Age, BA and other stand attributes such as Nha, Dq and SDI. The model finally selected included stand Age and Dq to estimate pFS:

$$pFS = a1 \cdot Age^{a2} \cdot Dq^{a3} \quad [2]$$

where $a1$ to $a3$ are curve fit parameters (denoted in 3-PG as pFSC, pFSAge and pFSQMD, respectively).

2.2.3. Tree mortality and self-thinning

Parameter estimates for density-independent tree mortality (i.e. stochastic mortality that occurs prior to the onset of mortality due to intra-specific competition) were obtained after adapting the survival model reported by Harrison and Borders (1996). Similar to Gonzalez-Benecke et al. (2014a), we ran the model of Harrison and Borders (1996) under different conditions of planting density and SI, and then fitted the model of Sands (2004) to that dataset to maintain parsimony in the 3-PG model structure:

$$\gamma_{Nt} = \gamma_{N1} + (\gamma_{N0} - \gamma_{N1}) \cdot e^{\left(-\ln(2) \frac{Age}{Age_{\gamma}}\right)} \quad [3]$$

where e is the base of natural logarithm, γ_{N1} is the mortality rate of mature stands, γ_{N0} is the mortality rate at age=0 (seedling mortality rate), and Age_{γ} is the age at which $\gamma_{Nt} = \frac{1}{2} \cdot (\gamma_{N0} + \gamma_{N1})$.

Parameter estimates for density-dependent tree mortality (WSx1000, the single tree stem biomass at a stand density of 1000 trees ha⁻¹, and thinPower, the self-thinning rule parameter) were computed from permanent plot data (Table 1), after using a species-specific general biomass equation for WS reported by Gonzalez-Benecke et al. (2014b). The dataset used for model fitting consisted of 8842 observations (repeated plot x age measurements), including trees from 2 to 33 years old, with WS ranging between 0.1 to 367 kg tree⁻¹, growing in stands with Nha and SI ranging between 156 to 4485 trees ha⁻¹ and 14 to 33 m, respectively. Similar to Gonzalez-Benecke et al.

(2014a), value of thinPower was determined using linear fitting to the boundary line of the transformed data of mean plot WS and Nha for each year and each site; the value of WSx1000 was calculated after solving the fitted equation using Nha=1000.

2.2.4. Allometric relationships

Initial biomass pools (WF, WS and WR) are needed for model initialization. If the model user has no initial biomass estimations for the stand to be simulated, general biomass functions for foliage and stem that use dbh, Ht and age can be used as predictors. Using the dataset reported in Gonzalez-Benecke et al. (2014b), allometric relationships for WS, WF and branch and bark fraction (pBB) were obtained. The model for pBB was needed to estimate stemwood biomass by subtracting branch and stembark biomass from WS. The dataset consisted of a collection of several sources used previously for site-specific allometric functions (further details can be found in Gonzalez-Benecke et al., 2014b), including 744 trees measured at 25 sites, with age, dbh and height ranging between 2 to 30 years old, 1.3 to 32.6 cm and 1.5 to 22.9 m, respectively. For stands where dbh and Ht were known, WF and WS were estimated by the model:

$$W_{F,S} = w1 \cdot dbh^{w2} \cdot Ht^{w3} \cdot Age^{w4} \quad [4]$$

where $W_{F,S}$ is the dry mass of foliage (F) or stem (S), and $w1$ to $w4$ are curve fit parameters.

In order to estimate WR, a model was fitted to estimate the ratio between WR and WS (RFrac) using data from Kinerson et al. (1977), Gibson et al. (1985), Tuttle (1998), Adegbidi et al. (2002), and Roth et al. (2007). When RFrac was known, WR was determined as: $WR = RFrac \cdot WS$. The dataset consisted of 168 trees from 2 to 27 years old, with dbh and height ranging between 0.3 to 26.7 cm and 2.0 to 18.6 m, respectively. The data were collected across the natural range of the species distribution, under different management and stand development conditions. The root systems were excavated to a depth of 40 cm in a 1 m² pit around the stump of each selected tree, and all live pine roots larger than a 2 mm diameter were weighed. We determined RFrac as a function of age as follows:

$$RFrac = r0 + r1 \cdot e^{(r2 \cdot Age)} \quad [5]$$

where e is the base of natural logarithm, and $r1$ to $r3$ are curve fit parameters. Other predictors were tested, such as dbh, height and Nha, but this model showed the better goodness of fit.

Similar to Gonzalez-Benecke et al. (2014a), alternative allometric models were developed to estimate initial WS, WF and WR for young stands when dbh was not available, using total tree height (Ht, m) and age as the main

predictors. For WS and WF, the data consisted of 338 trees measured at 10 sites, including trees from 1 to 4 years old, with Ht ranging between 0.9 to 6.0 m (Colbert, 1990; Roth et al., 2007; Samuelson et al. 2004; Maier et al. 2012). For WR we used the same relationship described previously. For young stands, when dbh data were not available, the model selected was:

$$W_{F,S} = w1 \cdot Ht^{w2} \cdot Age^{w3} \quad [6]$$

where $W_{F,S}$ is the dry mass of foliage (F) or stem (S), and $w1$ to $w3$ are curve fit parameters.

The relationship between age and pBB was fitted using an exponential decay to a non-zero asymptote function (Sands and Landsberg, 2002):

$$pBB = pBB1 + (pBB0 - pBB1) \cdot e^{\left(-\ln(2) \cdot \frac{Age}{Age_{BB}}\right)} \quad [7]$$

where e is the base of natural logarithm, pBB1 is the branch and bark fraction of mature stands, pBB0 is the branch and bark fraction at age=0 (planting), and Age_{BB} is the age at which $pBB = \frac{1}{2} \cdot (pBB0 + pBB1)$. Similar to allometric relationships described previously, we used the dataset reported by Gonzalez-Benecke et al. (2014b). This dataset included 427 trees measured at 11 sites, with age, dbh and height ranging between 2 to 25 years old, 1.3 to 30.1 cm and 1.5 to 21.3 m, respectively.

The relationship between Dq and mean tree height (H, m) was obtained from permanent plot data. We fitted separate models for stands growing in the southeastern U.S. (7334 paired Dq-H data were used, including trees from 2 to 25 years old, with Dq and H ranging between 0.3 to 37.8 cm and 1.4 to 26.3 m, respectively) and stands growing in Uruguay (175 paired Dq-H data were used, including trees from 3 to 34 years old, with Dq and H ranging between 6.9 to 40.1 cm and 4.5 to 26.6 m, respectively) (Table 1). The relationship between Dq and H was fitted using several stand-level variables as covariates. The variables considered were Age, Nha and BA, which represented different characteristics of the stands, such as stocking, productivity and competition, which could affect the height-diameter relationships. The model finally selected to estimate mean height was:

$$H = h1 \cdot Dq^{h2} \cdot Age^{h3} \cdot Nha^{h4} \quad [8]$$

where $h1$ to $h4$ are curve fit parameters (denoted in 3-PG as aH, nHD, nHAge and nHN, respectively).

The original equation in 3-PG used the relationships between WS and dbh to estimate dbh from a known WS (Landsberg and Waring, 1997). As the model used stem diameter to estimate BA, and considering that the model estimated WS (in Mg ha⁻¹) directly from NPP, we decided to estimate Dq (the dbh of the tree of mean BA) from stand-level WS, including age and stand density (Nha) as covariates. The model finally selected was:

$$Dq = b1 + b2 \cdot WS^{b3} \cdot Age^{b4} \cdot Nha^{b5} \quad [9]$$

where $b1$ to $b5$ are curve fit parameters (denoted in 3-PG as $a11Ws$, $a1Ws$, $n1Ws$, $n2Ws$ and $n3Ws$, respectively).

For stands growing in the southeastern U.S. we computed WS using the general biomass equation reported by Gonzalez-Benecke et al. (2014b); for stands growing in Uruguay, South America, we computed WS using the biomass equations reported by Fassola et al. (2010). Table 1 describes the data used for the Dq-STEM analysis.

After bole volume inside bark (VIB , $m^3 ha^{-1}$) was computed, we estimated bole volume outside bark (VOB , $m^3 ha^{-1}$) from VIB using the term $Vratio$, which is the ratio between VOB and VIB using the same approach described by Gonzalez-Benecke et al. (2014a). We created the dataset needed for model fitting by running the growth and yield model reported by Harrison and Borders (1996) for a rotation length of 30 years under different conditions of planting density (500, 1500 and 2500 trees ha^{-1}) and SI (15, 23 and 30 m). The relationship between $Vratio$ and VIB was also fitted using several stand-level variables as covariates. The model finally selected to estimate $Vratio$ was:

$$Vratio = r1 \cdot VIB^{r2} \cdot Nha^{r3} \cdot Age^{r4} \quad [10]$$

where $r1$ to $r4$ are curve fit parameters (denoted in 3-PG as aVR , $nVRVi$, $nVRN$ and $nVRAge$, respectively).

2.2.5. Wood specific gravity

Wood specific gravity (SG) was needed to convert stemwood mass ($Mg ha^{-1}$) to VIB . As it has been documented that SG differs between trees growing in the United States and South America (Higa et al., 1973; Barrichelo et al., 1977), we developed separate models for each geographic location. For trees growing in the southeastern U.S. we used the data reported by Gonzalez-Benecke et al. (2011) and fitted a new model that maintained parsimony with the 3-PG model structure. For trees growing in South America (Argentina, Brazil and Uruguay), we used data reported by Higa et al. (1973), Barrichelo et al. (1977), Pereyra and Gelid (2002), Weber (2005), Von Wallis et al. (2007), Pezzutti (2011) and Barth et al. (2013). The relationships between age and wood specific gravity (SG) were determined by fitting the model proposed by Sands (2010):

$$SG = \rho_1 + (\rho_0 - \rho_1) \cdot e^{\left(-\ln(2) \cdot \frac{Age}{Age_p}\right)} \quad [11]$$

where e is the base of natural logarithm, ρ_1 is the SG of mature stands, ρ_0 is the SG at age=0, and Age_p is the age at

which $SG = \frac{1}{2} \cdot (\rho_0 + \rho_1)$.

2.2.6. Specific needle area

Data used to determine the relationships between age and specific needle area (SNA, m² kg⁻¹) were obtained from the literature review (see Figure 8 for list of references used). We fitted the model proposed by Sands (2010):

$$\text{SNA} = \sigma_1 + (\sigma_0 - \sigma_1) \cdot e^{\left(-\ln(2) \cdot \left(\frac{\text{Age}}{\text{Age}_\sigma}\right)^2\right)} \quad [12]$$

where e is the base of natural logarithm, σ_1 is the SNA of mature stands σ_0 is the SNA at age=0; and Age_σ is the age at which $\text{SNA} = \frac{1}{2} \cdot (\sigma_0 + \sigma_1)$.

2.2.7. Canopy Cover

We used data from 182 plots installed in two PPINES (Pine Productivity Interactions on Experimental Sites, from FBRC) studies in FL and GA (Table 1) to analyze canopy cover dynamics of young loblolly pine stands. These studies were selected because they provided long-term repeated measurements of canopy cover development under contrasting conditions. The studies included the combinations of two contrasting silvicultural treatments (operational and high intensity), two contrasting planting densities (1,334 and 2,990 trees ha⁻¹), and seven different loblolly pine full-sib genetic families. Further details can be found in Roth et al. (2007). The dataset included yearly measurements of dbh and Ht, from age 2 to 14 years, and live crown widths at ages 3, 4 and 5 years. For each measured tree, live crown area (CA, m²) was determined assuming an elliptical crown shape. Following the approach of Gonzalez-Benecke et al (2014a), for each site and plot (that included the combination of planting density, culture and genetic family), a model was fitted to estimate CA as a function of dbh:

$$\text{CA} = a \cdot \text{dbh}^b \quad [13]$$

Only the effect of genetic family was significant in the allometry of CA and no effect of site, planting density and culture was detected ($P > 0.18$, data not shown). Using family-specific models, CA was calculated for all measured trees. Following Gonzalez-Benecke et al. (2014a), the sum of CA for each plot was expressed as a proportion of the area of the plot and the variable CanCover was determined for each age. After canopy closure the relationship used in this study would not be adequate as the allometry of crown width changes (Pretzsch et al., 2012). As 3-PG uses a maximum value of CanCover of 1 (not accounting for overlapping branches), values of CanCover greater than 1 were assumed to be 1. A function to describe the dynamics of CanCover prior to reaching

full canopy closure was fitted using age and other stand attributes such as Dq, SDI, BA and Nha. The model finally selected to estimate mean CanCover was:

$$\text{CanCover} = c1 \cdot \text{BA}^{c2} \cdot \text{Age}^{c3} \quad [14]$$

where $c1$ to $c3$ are curve fit parameters (denoted in 3-PG as aCan, nCanBA, nCanAge, respectively).

2.2.8. Needlefall, litterfall and forest floor accumulation

We followed the approach of Gonzalez-Benecke et al. (2014a) to analyze the dynamics of monthly fractional rate of needlefall (γ_N , month⁻¹), using needlefall (NF, Mg ha⁻¹ month⁻¹) data from the IMPAC study (Dalla-Tea and Jokela, 1991; Jokela and Martin, 2000). Phenological month for needlefall (NMonth) was defined as starting in May (NMonth=1) and ending in April (Nmonth=12). After expressing γ_N as a proportion of annual maximum γ_N (γ_{Nx} , month⁻¹), a non-linear model was fitted to the relationship between NMonth and the monthly average γ_N . The final model was:

$$\frac{\gamma_N}{\gamma_{Nx}} = \frac{\gamma_{N1} + \gamma_{N2} \cdot \text{NMonth}}{1 + \gamma_{N3} \cdot \text{NMonth} + \gamma_{N4} \cdot \text{NMonth}^2 + \gamma_{N5} \cdot \text{NMonth}^3} \quad [15]$$

where γ_{N1} to γ_{N5} are curve fit parameters.

To estimate litterfall (LF, Mg ha⁻¹ month⁻¹), we used the model ratio between NF and LF reported by Gonzalez-Benecke et al. (2014a). Using LF and a litter decay rate =0.15 (Binkley, 2002), we incorporated into 3-PG the calculation of forest floor accumulation (Mg ha⁻¹).

2.2.9. Effect of frost on production

Previous versions of 3-PG have incorporated the effects of frost on canopy conductance by using a factor called kF, the number of days of production lost per frost day. The frost-dependent growth modifier fFrost depended on kF and the number of frost days (FrostDay) and was calculated as fFrost=1-kF·FrostDay/30. The effect of frost on stand production was assumed to be independent of frost intensity. For example, Sands and Landsberg (2002) used kF=0 for *E. globulus*, assuming that there is no effect of frost on production, or Bryars et al. (2013) used kF=1 for loblolly pine, assuming that there is one day of production lost per each frost day. Based on observations of Teskey et al. (1987) and Polster and Fuchs (1963), as redrawn in Larcher (1995), the reduction in photosynthesis or leaf conductance depended on the intensity of frost. Thus, we modified the fFrost function to account for the effect of frost intensity. The impact of frost on growth reduction was conducted using data reported in Teskey et al. (1987),

where maximum leaf conductance during the day following a frost night was correlated with minimum night temperature on 8 year-old trees. After expressing conductance relative to conductance at 0° Celsius (fractional conductance, kF), the model used was:

$$kF = e^{(tF \cdot T_{min})} \quad [16]$$

where e is the base of natural logarithm, tF is the rate of production loss per degree Celsius below zero and T_{min} is the minimum temperature of each frost day. For days when $T_{min} > 0$, $fFrost$ was set equal to 1. Using this parameter, $fFrost$ was calculated as:

$$fFrost = 1 - (1 - e^{(tF \cdot T_{min})}) \cdot FrostDay/30 \quad [17]$$

where e is the base of natural logarithm and $FrostDay$ is the number of frost days of each month.

2.2.10. Fertility rating

Following Gonzalez-Benecke et al. (2014a), we used the approach of correlating FR with changes in site index (SI, m). Using this approach, FR was the only parameter obtained from calibration and not from observed/reported data. The relationship between FR and SI was analyzed using data from 47 permanent plots from the PINEMAP dataset, installed at 12 sites, one site per state: two studies from FBRC, one from FMRC, six from FPC, one from PMRC and two from WGFTIP. The first selection criteria was that all sites would have at least 10 years of measurement interval. Then, all sites were randomly selected to account for variability in geographic location within each state (avoid two sites in same county). On each site, 3-4 plots were randomly selected. We also included data from 16 permanent plots from the CAPPS study (Consortium for Accelerated Pine Production Studies, from PMRC) installed in GA. The treatments applied in each of the 63 plots created a wide range in productivity, similar to the range in productivity found in operational and experimental plots in the southeastern United States (Fox et al., 2007). The dataset consisted of 510 plot-level data points, including stands from 2 to 27 years old, with N_{ha} and SI ranging between 302 to 4,434 trees ha^{-1} and 13.9 to 33.1 m, respectively (Table 2). On each plot, total above-ground biomass (AGB, Mg ha^{-1}) was determined using the general biomass function reported by Gonzalez-Benecke et al. (2014b).

Similar to Gonzalez-Benecke et al. (2014a), after obtaining all parameter estimates required by 3-PG, we determined the value of FR that minimized the error of AGB, by recording for each plot the value that had the minimum mean square error of the fitting between the observed and predicted AGB (including all measurements).

Finally, after pooling all paired data from all 63 plots, SI was correlated with the optimum FR. The following sigmoidal curve was finally selected to estimate FR:

$$FR = \frac{f1}{1+f2 \cdot e^{(-f3 \cdot SI)}} \quad [18]$$

Where e is the base of natural logarithm $f1$ to $f3$ are curve fit parameters. Once the FR function had been developed by calibration, it was applied unchanged to the validation data set. This is in contrast to most applications of the model, in which the FR parameter is used as a “tuning” parameter.

2.2.10. Parameters obtained from literature review

All other parameter estimates shown in Table 4 were obtained from previous reports of 3-PG parameterizations for loblolly pine (Table 4). From Sampson et al. (2006), canopy quantum yield $\alpha_c = 0.053 \text{ mol C mol}^{-1} \text{ photon}$, and the age modifiers MaxAge, nAge and rAge = 200, 1.5 and 0.75. From Bryars et al. (2013), maximum (pRx) and minimum (pRn) fraction of NPP to roots = 0.4 and 0.2; temperature modifiers Tmin, Topt and T max = 4, 25 and 38 °C; fertility effects factors m0, fNo and fNn = 0, 0.3 and 1; maximum proportion rainfall canopy interception Maxintcptn = 0.2; LAI for maximum rainfall interception LAImaxIntcp = 5; light extinction coefficient $k = 0.57$; monthly root turnover = 0.0168.

2.3. Model evaluation

The independent validation dataset included data from 91 permanent plots distributed in 24 sites in 12 states in the southeastern U.S. (two sites per state). The model was also validated against data from 10 permanent plots growing in operational stands in Tacuarembó, Uruguay (properties of Cambium Forestal Uruguay S.A.). Figure 1 shows the location of all validation sites. An additional validation was conducted on projected LAI ($\text{LAI, m}^2 \text{ m}^{-2}$) estimates using data from IMPAC study (Jokela and Martin, 2000), where monthly LAI was estimated from needlefall collected monthly from age 6 to 19 years, using six circular litter traps (1 m^2) installed in each of the 12 study plots (see section 2.2.8). Further details of LAI calculations can be found in Jokela and Martin (2000). On each plot, we modified FR using the observed SI and the FR-SI relationship reported in this study.

FIGURE 1

The performance of 3-PG for loblolly pine was compared against independent data not used in model development. The goodness-of-fit between the observed and predicted values was evaluated using three measures of accuracy: (i) root mean square error (RMSE); (ii) mean bias error (Bias, the difference between observed and predicted values); and (iii) coefficient of determination (R^2). Variables evaluated included BA, Nha, H, AGB and bole volume outside bark (VOB, $m^3 ha^{-1}$). For each plot, observed VOB was computed with the function reported by Van Deusen et al. (1981); observed AGB was computed using the general biomass function reported by Gonzalez-Benecke et al. (2014a). For sites in Uruguay, functions to estimate VOB were not available and VIB was used instead, and was computed with the function reported by Rachid et al. (2014). Observed AGB for the Uruguay sites was computed using the function reported by Fassola et al. (2010). For each variable, we used F-tests to determine if the relationship between predicted and observed values had a slope and intercept different than one and zero, respectively. All statistical analyses were performed using SAS 9.3 (SAS Inc., Cary, NC, USA).

Model validation was conducted by running the model from age of first measurement to the age of last measurement. Initial biomass pools to initialize the model were determined for each plot using the equations for WF, WS and WR reported in this study. Table 2 shows a summary of stand characteristics for each site used for model validation and FR/SI calibration.

Table 2. Summary of data used for model validation and FR/SI calibration.

State	County	Institution	n	Lat	Long	AGE (yrs)	Dq (cm)	Nha (trees ha^{-1})	BA ($m^2 ha^{-1}$)	SI (m)	Reference
AL	Bibb	FPC	4*	33.03	-87.21	18 - 26	18.8 - 28.8	302 - 336	8.4 - 21.9	17.8 - 19.8	1
	Butler	FPC	4	31.57	-86.67	4 - 18	2.7 - 20.8	923 - 1369	0.5 - 36.2	20.8 - 23.7	2
	St. Clair	PMRC	4	33.84	-86.30	2 - 12	0.1 - 24.1	355 - 1494	0.1 - 38.3	21.6 - 25.4	3
AR	Bradley	FPC	4*	33.59	-92.17	19 - 27	21.9 - 29.4	358 - 494	15.6 - 30	18.3 - 20.1	1
	Hempstead	WGFTIP	4	33.96	-93.75	5 - 20	6.6 - 27.8	333 - 969	2.6 - 34.4	20.5 - 21.6	4
	Perry	FMRC	3	34.88	-93.07	15 - 30	15.8 - 29.2	652 - 1265	17.8 - 47	17.2 - 18.5	5
FL	Alachua	FBRC	4*	29.76	-82.29	4 - 25	2.4 - 21.8	1010 - 1497	0.6 - 44.3	15.0 - 23.0	6
	Santa Rosa	FPC	4	30.77	-86.97	5 - 13	4 - 13.7	1262 - 1441	1.6 - 20.9	15.5 - 19.9	7
	Taylor	FBRC	4	30.16	-83.75	2 - 14	2.7 - 25.4	1070 - 2902	0.7 - 54.8	27.2 - 32.9	8
GA	Camden	FBRC	4*	31.13	-81.75	2 - 14	3.3 - 22.3	1098 - 2779	1.1 - 48.6	27.1 - 28.8	8
	Jones	FPC	4	33.10	-83.60	25 - 33	19.8 - 30.3	333 - 428	12.9 - 24.7	15.5 - 18.5	1
	Stewart	FMRC	3	32.12	-84.66	6 - 17	7.1 - 17.1	1532 - 1775	7.0 - 37.7	17.7 - 18.8	5
	Clarke	PMRC	4*	33.60	-83.38	2 - 15	0.5 - 21.0	1230 - 1580	0.1 - 45.2	24.5 - 29.5	9
	Jasper	PMRC	4*	33.32	-83.39	2 - 15	0.7 - 20.5	1200 - 1588	0.2 - 42.2	23.7 - 29.3	9
	Tift	PMRC	4*	31.30	-83.51	3 - 15	1.6 - 22.1	1070 - 1440	0.5 - 40.9	27.6 - 29.0	9
	Pierce	PMRC	4*	31.10	-82.42	2 - 15	0.8 - 20.9	1356 - 1660	0.2 - 46.6	26.1 - 32.8	9
	Allen	WGFTIP	4*	30.65	-92.82	5 - 20	2.1 - 22.6	890 - 1268	0.4 - 40.5	18.2 - 20.1	4
LA	Bienville	FPC	4	32.32	-92.83	16 - 24	14.6 - 20.2	1161 - 1482	23.9 - 40.6	16.8 - 17.7	10
	Washington	FPC	4	30.85	-90.04	14 - 24	12.5 - 17.2	1418 - 1912	21.6 - 36.5	14.1 - 17.7	10
	Kemper	FPC	4	32.75	-88.45	11 - 19	15.1 - 22.5	787 - 896	14.2 - 35.8	20.3 - 22.4	10
MS	Marion	FPC	4*	31.38	-90.27	4 - 14	4 - 18.8	914 - 1778	1.8 - 30.7	22.0 - 24.6	2
	Pontotoc	WGFTIP	4	34.28	-88.92	5 - 20	7.3 - 24.1	458 - 1339	4.2 - 37.4	21.3 - 22.6	4

NC	Bertie	FPC	4*	36.21	-76.95	10 - 20	12.3 - 20.5	1428 - 1517	17.2 - 50.2	19.7 - 20.5	10
	Bladen	FPC	4	34.60	-78.60	4 - 23	3 - 26.9	726 - 1508	1.0 - 42.8	23.1 - 26.8	2
	Chatham	FMRC	3	35.63	-79.08	24 - 39	19.1 - 25.6	396 - 1260	17.4 - 39.9	15.4 - 17.1	5
OK	McCurtain	FMRC	4	34.20	-95.05	17 - 26	18.4 - 36.6	226 - 1252	13.3 - 45.5	16.2 - 18.1	5
	Pushmataha	WGFTIP	4*	34.43	-95.22	5 - 15	3.7 - 20.5	1041 - 1627	1.4 - 47.6	18.7 - 20.2	4
	Pushmataha	WGFTIP	4	34.47	-95.20	5 - 20	4.5 - 21.6	1145 - 1517	2.2 - 52	19.8 - 21.4	4
SC	Hampton	PMRC	4	32.79	-80.95	2 - 12	2.8 - 24.2	672 - 1476	0.4 - 39.2	28.5 - 30.3	11
	Laurens	PMRC	4*	34.40	-81.90	4 - 12	2.7 - 23.2	486 - 4434	0.3 - 46.9	21.3 - 23.4	11
	Williams	FPC	4	33.59	-79.48	4 - 22	0.4 - 23.2	736 - 1340	0.1 - 39.0	18.6 - 20.9	2
TN	Bradley	FPC	4	35.02	-84.84	12 - 22	12 - 18.5	1468 - 2040	19.1 - 48.3	17.0 - 19.3	10
	Hardin	FPC	4	35.15	-88.01	5 - 17	4.8 - 18.2	1366 - 1655	2.6 - 36.2	19.3 - 21.7	2
	Rhea	FPC	4*	35.72	-84.77	12 - 22	14 - 20.9	1330 - 1411	21 - 45.7	16.9 - 18.4	10
TX	Polk	FMRC	4*	30.77	-94.82	4 - 14	7.6 - 20.6	907 - 1072	4.4 - 30.4	22.3 - 23.5	5
	San Augustine	PMRC	4	31.40	-94.05	2 - 8	0.9 - 20.7	469 - 2938	0.1 - 35.0	23.8 - 24.6	1
	Walker	FMRC	4	31.00	-95.48	19 - 31	16.8 - 25.3	693 - 1502	24 - 37.3	15.2 - 17.0	5
VA	King & Queen	FPC	4*	37.62	-76.78	4 - 18	2.2 - 22.3	999 - 1655	0.6 - 41	18.2 - 23.5	2
	Prince Edward	FMRC	3	37.13	-78.40	19 - 34	15.2 - 29.8	516 - 1729	19.7 - 46.1	18.8 - 19.3	5
	Sussex	FPC	4	36.88	-77.07	4 - 18	1.5 - 18.6	1314 - 1550	0.2 - 38.4	19.4 - 21.2	2
URUGUAY	Tacuarembó	CAMBIUM	10	-31.48	-55.99	7 - 17	15.8 - 40.1	255 - 1000	8.8 - 55.3	20.9 - 29.8	1
TOTAL				148		2 - 39	0.1 - 30.3	226 - 4434	0.1 - 55.3	14.0 - 32.9	

n: number of plots; AGE: range of age (yrs); Dq: range of quadratic mean diameter (cm) across AGE and plots; BA: range of basal area ($\text{m}^2 \text{ha}^{-1}$) across AGE and plots, SI: range of site index at base age=25 years (m) across plots.

FBRC: Forest Biology Research Cooperative; FMRC: Forest Modeling Research Cooperative; FPC: Forest Productivity Cooperative; PMRC: Plantation Management Research Cooperative; WGFTIP: Western Gulf Forest Tree Improvement Program; CAMBIUM: Cambium S.A.

*: Site used for FR/SI analysis.

References: 1: Carlson et al., 2014; 2: Nilsson and Allen, 2003; 3: Zhao et al., 2012; 4: Koralewski et al., 2015; 5: Russell et al., 2010; 6: Jokela and Martin, 2000; 7: Leggett and Kelting, 2006; 8: Roth et al., 2007; 9: Borders et al., 2004; 10: Zhao et al., 2011; 11: Hynynen et al., 1998.

AL: Alabama; AR: Arkansas; FL: Florida; GA: Georgia; LA: Louisiana; MS: Mississippi; NA: North Carolina; OK: Oklahoma; SC: South Carolina; TN: Tennessee; TX: Texas; VA: Virginia.

2.4. Climate and soil data

The weather data consisted of monthly average daily maximum (T_{max} , $^{\circ}\text{C}$) and minimum (T_{min} , $^{\circ}\text{C}$) temperature, monthly total rainfall (Rain, mm month^{-1}), monthly average daily total solar radiation ($\text{MJ m}^{-2} \text{day}^{-1}$), number of rainy days (the number days with rainfall $> 1 \text{ mm, month}^{-1}$) and number of frost days (the number of days with $T_{\text{min}} < 0 \text{ }^{\circ}\text{C, month}^{-1}$). For the AMERIFLUX study, weather data were collected from an automatic weather station installed at the site (Noormets et al., 2010). For all other sites in the U.S., daily weather data were obtained online from the University of Idaho Gridded Surface Meteorological Dataset (<http://climate.nkn.uidaho.edu/METDATA/>), and selecting the weather station nearest to each study site. For sites from Uruguay, daily weather data were obtained online from the Instituto Nacional de Investigación Agropecuaria (<http://www.inia.org.uy/online/site/gras.php>), and selecting the weather station at INIA-Tacuarembó. The soils data collected were texture class (s: sandy; sl: sandy-loam; cl: clay), and maximum and minimum available soil water (mm). For all sites in the U.S., soils data were obtained online from the USDA's National Resources Conservation

Service (<http://websoilsurvey.sc.egov.usda.gov/App/WebSoilSurvey.aspx>), using site coordinates. For sites in Uruguay, soils data were available from soil classification maps for each site using the soil classification of CONEAT (Comisión Nacional de Estudio Agro económico de la Tierra; www.prenader.gub.uy/coneat/). A summary of soil and weather data of all sites used for model validation and FR/SI analysis is presented in Table 3.

Table 3. Summary of soil and weather data of sites used for model validation and FR/SI calibration (§).

State	County	Institution	Soil Class	ASW	Tmin-w	Tmax-s	Rad-w	Rad-s	Rain	Nrain	Nfrost
AL	Bibb	FPC §	SL	187	3.0	33.0	11.0	19.4	1,460	150	31
	Butler	FPC	SL	227	1.0	32.6	10.4	20.0	1,421	148	47
	St. Clair	PMRC	SL	155	0.4	32.3	10.1	20.2	1,390	148	50
AR	Bradley	FPC §	SL	165	0.9	32.9	10.1	20.6	1,380	129	44
	Hempstead	WGFTIP	SL	164	0.0	32.8	10.1	20.7	1,330	131	52
	Perry	FMRC	SL	140	-1.7	32.5	10.0	20.8	1,332	129	68
FL	Alachua	FBRC §	S	116	6.8	32.8	11.8	18.0	1,310	138	11
	Santa Rosa	FPC	SL	150	4.8	32.9	11.0	18.7	1,662	156	20
	Taylor	FBRC	S	125	5.7	33.2	11.5	18.3	1,442	152	17
GA	Camden	FBRC §	SL	213	5.3	32.7	11.2	18.5	1,284	144	14
	Jones	FPC	CL	218	0.5	32.6	10.5	19.8	1,207	150	42
	Stewart	FMRC	SL	190	2.0	32.8	10.9	19.6	1,271	145	33
	Clarke	PMRC §	CL	205	4.1	36.7	12.2	22.2	1,236	131	27
	Jasper	PMRC §	C	205	4.1	35.9	11.6	22.5	1,198	130	25
	Tift	PMRC §	CL	198	1.1	36.4	10.9	22.8	1,175	119	44
LA	Pierce	PMRC §	SL	178	1.3	36.7	10.7	22.8	1,181	123	42
	Allen	WGFTIP §	SL	214	4.2	33.1	11.0	19.2	1,617	153	23
	Bienville	FPC	CL	211	5.3	33.3	10.8	19.4	1,600	140	15
MS	Washington	FPC	CL	241	2.4	33.6	10.5	20.3	1,379	131	33
	Kemper	FPC	CL	173	3.1	32.8	11.0	19.7	1,543	149	30
	Marion	FPC §	CL	256	1.9	33.0	10.4	19.8	1,362	147	40
NC	Pontotoc	WGFTIP	SL	175	0.6	32.5	9.8	20.4	1,470	144	48
	Bertie	FPC §	S	169	1.2	31.6	9.9	19.4	1,259	133	44
	Bladen	FPC	SL	203	-0.3	31.2	9.2	19.8	1,217	148	56
OK	Chatham	FMRC	SL	228	-1.5	31.6	9.7	20.2	1,129	140	72
	McCurtain	PMRC	S	128	-0.1	33.1	10.3	21.1	1,290	131	53
	Pushmataha	WGFTIP §	SL	146	-0.8	32.9	10.3	21.2	1,271	127	61
SC	Pushmataha	WGFTIP	SL	184	-0.8	33.1	10.2	21.1	1,278	127	62
	Hampton	PMRC	SL	148	2.4	32.2	10.2	18.9	1,263	151	31
	Laurens	PMRC §	S	166	2.9	32.9	10.6	19.1	1,258	149	31
TN	Williams	FPC	SL	176	-0.6	32.3	10.1	20.2	1,158	140	60
	Bradley	FPC	CL	247	-1.7	31.6	9.5	20.7	1,430	160	72
	Hardin	FPC	SL	184	-0.8	31.2	9.6	20.3	1,415	155	62
TX	Rhea	FPC §	S	123	-1.8	30.7	9.2	20.3	1,391	163	71
	Polk	FMRC §	SL	254	2.2	33.8	10.9	20.7	1,329	127	36
	San Augustine	PMRC	SL	166	3.8	34.5	11.1	20.9	1,118	120	22
VA	Walker	FMRC	SL	161	3.5	33.8	11.1	20.7	1,276	128	31
	King & Queen	FPC §	CL	241	-1.2	31.1	9.0	19.9	1,204	144	67
	Prince Edward	FMRC	SL	221	-1.4	31.0	8.7	20.0	1,149	134	69
	Sussex	FPC	SL	202	-2.6	30.6	9.2	20.4	1,117	141	81
URUGUAY	Tacuarembó	CAMBIUM	SL	195	4.2	33.1	11.0	19.2	1,617	153	23

Soil Class: Soil texture class (s: sandy; sl: sandy-loam; cl: clay); ASW: Available soil water, the difference between maximum and minimum ASW (mm); Tmin-w: average daily minimum temperature of winter months (°C); Tmax-s: average daily maximum temperature of summer months (°C); Rad-w: average daily total solar radiation of winter months ($\text{MJ m}^{-2} \text{ day}^{-1}$); Rad-s: average daily total solar radiation of summer months ($\text{MJ m}^{-2} \text{ day}^{-1}$); Rain: average yearly total rainfall (mm year^{-1}), Nrain: average yearly total number of rainy days, Nfrost: average yearly total number of frost days.

Winter months: December 1 to February 28 in U.S., June 1 to August 31 in Uruguay; summer months: June 1 to August 31 in U.S., December 1 to February 28 in Uruguay.

AL: Alabama; AR: Arkansas; FL: Florida; GA: Georgia; LA: Louisiana; MS: Mississippi; NA: North Carolina; OK: Oklahoma; SC: South Carolina; TN: Tennessee; TX: Texas; VA: Virginia.

§: Site used for FR/SI calibration.

Weather data from years used for validation or FR/SI analysis.

3. Results

3.1. Model Fitting

The parameter estimates used by 3-PG for loblolly pine are reported in Table 4. All parameter estimates from model fitting were significant at $P < 0.05$.

Table 4. Description of 3-PG parameters and values for loblolly pine.

Meaning / Comment	3-PG symbol	Unit	Value	Sources
Biomass partitioning and turnover				
Allometric relationships & partitioning				
Constant in the foliage : stem partitioning ratio relationship	pFSC	-	0.406	This study
Power of Age in the foliage: stem partitioning ratio relationship	pFSAge	-	0.311	This study
Power of Dq in the foliage : stem partitioning ratio relationship	pFSD	-	-0.288	This study
Intercept in the Dq v. stem mass relationship	a11Ws	-	-3.707*	This study
Constant in the Dq v. stem mass relationship	a1Ws	-	54.449*	This study
Power in the Dq v. stem mass relationship	n1Ws	-	0.253*	This study
Power of Age in the Dq v. stem mass relationship	n2Ws	-	0.037*	This study
Power of Nha in the Dq v. stem mass relationship	n3Ws	-	-0.306*	This study
Maximum fraction of NPP to roots	pRx	-	0.40	1
Minimum fraction of NPP to roots	pRn	-	0.20	1
Needlefall, litterfall, litter decay & root turnover				
Maximum needlefall rate	γFx	month ⁻¹	0.157	This study
Month at which needlefall rate has maximum value	$t\gamma Fx$		11	This study
Average yearly decay rate of litter		year ⁻¹	0.15	This study
Needlefall to litterfall ratio at age 0	NF ₀	-	0.733	This study
Needlefall to litterfall ratio for mature stands	NF ₁	-	1.0	This study
Age at which Needlefall to litterfall ratio = ($\sigma_0 + \sigma_1$)/2	Age _{NLR}	year	21.5	This study
Average monthly root turnover rate	γR	month ⁻¹	0.0168	1
NPP & conductance modifiers				
Temperature modifier (fT)				
Minimum temperature for growth	Tmin	°C	4	1
Optimum temperature for growth	Topt	°C	25	1
Maximum temperature for growth	Tmax	°C	38	1
Frost modifier (fFrost)				
Reduction rate of production per degree Celsius below zero	tF	Day °C ⁻¹	0.178	This study
Soil water modifier (fSW)				
Moisture ratio deficit for $f_q = 0.5$	SWconst	-	0.7	1
Power of moisture ratio deficit	SWpower	-	9	1
Fertility effects				
Value of 'm' when FR = 0	m0	-	0	1
Value of 'fNutr' when FR = 0	fN0	-	0.3	1
Power of (1-FR) in 'fNutr'	fNn	-	1	1
Age modifier (fAge)				
Maximum stand age used in age modifier	MaxAge	year	200	2
Power of relative age in function for fAge	nAge	-	1.5	2
Relative age to give fAge = 0.5	rAge	-	0.5	2
Stem mortality & self-thinning				
Mortality rate for large t	γNx	% year ⁻¹	0.392	This study
Seedling mortality rate (t = 0)	$\gamma N0$	% year ⁻¹	2.320	This study

Age at which mortality rate has median value	$t\gamma N$	year	10.853	This study
Shape of mortality response	$n\gamma N$	-	1	This study
Max. stem mass per tree @ 1000 trees/hectare	$wSx1000$	kg tree ⁻¹	230	This study
Power in self-thinning rule	$thinPower$	-	1.174	This study
Fraction mean single-tree foliage biomass lost per dead tree	mF	-	0	1
Fraction mean single-tree root biomass lost per dead tree	mR	-	0.2	1
Fraction mean single-tree stem biomass lost per dead tree	mS	-	0.4	1

Canopy structure and processes

Specific needle area (σ)

Specific needle area at age 0	$\sigma 0$	m ² kg ⁻¹	5.529	This study
Specific leaf area for mature leaves	$\sigma 1$	m ² kg ⁻¹	3.875	This study
Age at which specific needle area = ($\sigma 0 + \sigma 1$)/2	$t\sigma$	year	5.971	This study

Light interception

Extinction coefficient for absorption of PAR by canopy	k	-	0.57	1
Constant in the CanCover relationship	$aCan$	-	0.258	This study
Power of BA in the CanCover relationship	$nCanBA$	-	0.688	This study
Power of Age in the CanCover relationship	$nCanAge$	-	-0.198	This study
Maximum proportion of rainfall evaporated from canopy	$MaxIntcptn$	-	0.2	1
LAI for maximum rainfall interception	$LAI_{maxIntcptn}$	-	5	1

Production and respiration

Canopy quantum efficiency	α_c	mol C mol PAR ⁻¹	0.053	2
Ratio NPP/GPP	Y	-	0.47	1

Canopy Conductance (g_c)

Minimum canopy conductance	$MinCond$	m s ⁻¹	0	This study
Maximum canopy conductance	$MaxCond$	m s ⁻¹	0.0118	This study
LAI for maximum canopy conductance	LAI_{g_c}	-	3	1
Defines stomatal response to VPD	$CoeffCond$	mb ⁻¹	0.0408	This study
Canopy boundary layer conductance	$BLcond$	m s ⁻¹	0.1	1

Wood and stand properties

Branch and bark fraction (pBB)

Branch and bark fraction at age 0	$pBB0$	-	1.198	This study
Branch and bark fraction for mature stands	$pBB1$	-	0.235	This study
Age at which pBB = ($pBB0 + pBB1$)/2	tBB	year	1.737	This study

Wood basic specific gravity

Minimum basic density - for young trees	$\rho 0$	-	0.358 **	This study
Maximum basic density - for older trees	$\rho 1$	-	0.482 **	This study
Age at which $\rho = (\rho_{min} + \rho_{max})/2$	$t\rho$	year	7.054 **	This study

Stem height

Constant in the D_q v. height relationship	aH	-	0.230 ***	This study
Power of D_q in the D_q v. height relationship	nHD	-	0.91 ***	This study
Power of Age in the D_q v. height relationship	$nHAge$	-	0.261 ***	This study
Power of Nha in the D_q v. height relationship	nHN	-	0.110 ***	This study

Volume Ratio

Constant in the bole volume ratio relationship	aVR	-	1.232	This study
Power of VIB in the bole volume ratio relationship	$nVRVi$	-	-0.017	This study
Power of Nha in the bole volume ratio relationship	$nVRN$	-	0.025	This study
Power of Age in the bole volume ratio relationship	$nVRAge$	-	-0.030	This study

*: For Uruguay $a1Ws = -0.143$; $a1Ws = 43.721$; $n1Ws = 0.353$; $n2Ws = -0.0099$; $n3Ws = -0.342$

**: For Uruguay $\rho 0 = 0.328$; $\rho 1 = 0.478$; $t\rho = 6.913$

***: For Uruguay $aH = 0.080$; $nHD = 0.923$; $nHAge = 0.308$; $nHN = 0.244$

References: 1: Bryars et al., 2013; 2: Sampson et al., 2001.

There was a negative relationship between canopy conductance (G_c , m s⁻¹) and mean daily VPD ($n=132$; $P<0.001$; $R^2=0.96$). The model fitted to estimate G_c parameters is shown in Figure 2. Maximum ($MaxCond$) and

minimum (MinCond) canopy conductance, and the response of canopy conductance to VPD (CoeffCond) were 0.0188 m s⁻¹, 0 m s⁻¹ and 0.0408 mb⁻¹, respectively (Table 4).

FIGURE 2

NPP partitioning was set as a function of Dq and age (n=924; P<0.001; R²=0.95). NPP allocation to stem (pS) increased rapidly until reaching values ranging between 0.6 - 0.8 at Dq larger than about 2 cm. Conversely, NPP allocation to foliage (pF) decreased sharply until reaching values ranging between 0.2 - 0.4 at Dq larger than about 3 cm (Figure 3a). The ratio between pF and pS (pFS) ranged between 0.25 and 0.6 for Dq larger than 5 cm and there was a good agreement between observed and predicted values (Figure 3b). The parameter estimates of the new pFS function were 0.406, 0.311 and -0.288, for pFSC, pFSAge and pFSD, respectively (Table 4).

FIGURE 3

Allometric relationships for Dq as a function of stem biomass (WS), mean height (H) as a function of Dq, pBB as a function of age, and volume ratio (Vratio) as a function of bole volume inside bark (VIB) are shown in Figure 4. The model to estimate Dq (Figure 4a) was dependent on WS (Mg ha⁻¹), age and Nha. We fitted separate models for stands growing in the southeastern U.S. (Dq_{US}) and Uruguay (Dq_{UR}):

$$Dq_{US} = -3.707 + 54.449 \cdot WS^{0.253} \cdot Age^{0.0374} \cdot Nha^{-0.3065} \quad (n=4380; P<0.001; R^2=0.97)$$

$$Dq_{UR} = -0.142 + 43.721 \cdot WS^{0.353} \cdot Age^{0.0099} \cdot Nha^{-0.3424} \quad (n=181; P<0.001; R^2=0.98)$$

In this model, the parameter estimate associated with Nha was negative, indicating that for the same WS and age, stands with higher density would have smaller diameters. Differences in model estimations reflect differences in allocation and/or wood specific gravity between geographic areas. For example, a 15-year-old stand with 1000 trees per ha and a WS of 200 Mg ha⁻¹, would have, on average, Dq= 24.0 or 29.9 cm, if growing in the southeastern U.S. or Uruguay, respectively.

The model to estimate mean height (Figure 4b) was also dependent on Dq, age and Nha. We fitted separate models for stands growing in the southeastern U.S. (H_{US}) and Uruguay (H_{UR}):

$$H_{US} = 0.2304 \cdot Dq^{0.9171} \cdot Age^{0.2616} \cdot Nha^{0.1098} \quad (n=7334; P<0.001; R^2=0.99)$$

$$H_{UR} = 0.0799 \cdot Dq^{0.9226} \cdot Age^{0.3080} \cdot Nha^{0.2439} \text{ (n=175; } P<0.001; R^2=0.98).$$

The model to estimate fraction of branch and bark to stem biomass (Figure 4c) was dependent on age. In this model the average pBB for seedlings was about 1, decreasing to about 0.235 as the trees aged:

$$pBB = 0.235 + (1.198 - 0.235) \cdot e^{(-\ln(2) \cdot \frac{Age}{1.737})} \text{ (n=114; } P<0.001; R^2=0.51)$$

Bole volume ratio (Figure 4d) was dependent on age and Nha. In this model, too, the parameter estimate associated with age was negative, indicating that older trees were likely to have a larger bark fraction:

$$Vratio = 1.232 \cdot VIB^{-0.0166} \cdot Nha^{0.0248} \cdot Age^{-0.0299} \text{ (n=324; } P<0.001; R^2=0.99).$$

FIGURE 4

Parameter estimates of models needed to estimate initial biomass pools for stands growing in the southeastern U.S. are shown in Table 5. For WF when dbh data were available ($Ht > 3$ m), the parameter estimate associated with Ht was negative, indicating that for the same dbh and Age, taller trees had less living needle biomass than shorter trees. In all cases, the parameter estimate associated with Age was negative, indicating that for the same size (dbh and/or H), older trees had less WS or WF. For young trees ($Ht < 3$ m), WF was dependent on Ht and age, and WS was dependent only on Ht. For RFrac, the parameter estimate associated with Age was negative, indicating that as trees aged, the ratio between WR and WS decreased, reaching a value of about 0.233 for trees older than about 15 years.

Table 5. Parameter estimates and fitted statistics of equations for predicting initial WF, WS and RFrac for stands growing in the southeastern U.S.

	Model	Equation	Parameter	Parameter estimate	SE	R ²	RMSE
Ht > 3 m	WF = $w_1 \cdot dbht^{w2} \cdot Ht^{w3} \cdot Age^{w4}$	5	w1	0.0997	0.0125	0.876	1.66
			w2	2.2416	0.0886		
			w3	-0.591	0.0866		
			w4	-0.3184	0.0422		
	WS = $w_1 \cdot dbht^{w2} \cdot Ht^{w3} \cdot Age^{w4}$	5	w1	0.015	0.000965	0.986	8.74
			w2	2.0449	0.0303		
			w3	1.2165	0.039		
			w4	-0.2078	0.0159		
Ht < 3 m	WF = $w_1 \cdot Ht^{w2} \cdot Age^{w3}$	6	w1	0.3253	0.034	0.926	1.80
			w2	2.5944	0.1336		
			w3	-0.6389	0.1699		
	WS = $w_1 \cdot Ht^{w2}$	6	w1	0.0904	0.00797	0.954	0.57
			w2	2.4223	0.0559		
			w3	-0.6389	0.1699		
All	RFrac = $r0 + r1 \cdot e^{(r2 \cdot Age)}$	7	r0	0.2333	0.018	0.913	0.12
			r1	10.6424	1.1156		
			r2	-1.851	0.0994		

WF: foliage dry mass (kg tree⁻¹); WS: stem dry mass (kg tree⁻¹); dbh: diameter outside-bark at 1.37 m height (cm); Ht: total tree height (m); RFrac: ratio between WR and WS; SE: standard error; R²: coefficient of determination; RMSE: root mean square error. For all parameter estimates: P-value < 0.001.

Figure 5 shows the tree mortality relationships. The model of Harrison and Borders (1996) was used to calculate mortality rate (Figure 5a). The parameter estimates for the model fitted for density-independent mortality are shown in Table 4. When using boundary line analysis for density-dependent mortality (Figure 5b), the slope of the self-thinning line (thinPower) was -1.174 and the maximum stem mass per tree at 1000 trees ha⁻¹ (WSx1000) was 230 kg (Table 4).

FIGURE 5

Family-specific allometric relationships were used to estimate crown area for each plot (models not shown). Using these relationships, the fractional canopy cover for each plot was calculated for both PPINES studies. Figure 6a shows the relationship between age and canopy cover development, which affected the timing to reach full canopy closure (fractional canopy cover =1). In the PPINES studies, plots with a narrow planting density and high culture (N-H) reached full canopy cover at about 3 years, while plots with a wide planting density and low culture (WL) reached full canopy cover at ages older than 10 years. After canopy closure the allometry of branches changes (Valentine et al., 2012) and the relationship used in this study may not be adequate, but for 3-PG any value of CanCover > 1 is assumed to be 1. When the fractional canopy cover was plotted against BA, a suite of curves was observed, and those curves were expressed as:

$$\text{CanCover} = 0.258 \cdot \text{BA}^{0.6883} \cdot \text{Age}^{-0.1986} \text{ (n=559; P<0.001; R}^2\text{=0.98).}$$

FIGURE 6

Figure 7 shows average monthly WF (Figure 7a), NF (Figure 7b) and γN (Figure 7c) for the IMPAC study, where fertilization (F) and weed control (W) treatments created a wide range in foliage biomass and needlefall (Jokela and Martin, 2000). In general, across all 14 years of measurements, control plots (C) that did not receive fertilization and weed control, showed lower monthly WF and NF. Maximum and minimum WF was reached in July

(NMonth=3) and February (NMonth=10), respectively, and maximum and minimum NF was attained in November (NMonth=7) and March (NMonth=11). All treatments showed similar γN within each month, reaching maximums and minimums in November (NMonth=7) and April (NMonth=12), respectively. The final model fitted was:

$$\frac{\gamma N}{\gamma N_x} = \frac{0.1183 + 11.9827 \cdot NMonth}{1 + 1.1522 \cdot NMonth - 0.8833 \cdot NMonth^2 + 0.0713 \cdot NMonth^3} \quad (n=144; P<0.001; R^2=0.96)$$

We left the model flexible to account for site-specific needlefall dynamics, allowing the user to change γN_x and the month when γN reached the maximum ($t\gamma N_x$). As default values, we determined a mean γN_x of 0.157 month⁻¹ (n=12; SE=0.031) and November ($t\gamma N_x=11$) as the month when γN_x peaked (Table 3). Mean monthly γN was 0.062 month⁻¹.

FIGURE 7

Figure 8 shows the age-dependent relationships for SNA (Figure 8a) and whole-tree SG for trees growing in the southeastern U.S. (SG_{US}) and Uruguay (SG_{UR}) (Figure 8b). For both variables, an exponential decay to a non-zero asymptote was fitted from the data:

$$SNA = 3.875 + (5.529 - 3.875) \cdot e^{(-\ln(2) \frac{Age}{5.971})} \quad (n=94; P<0.001; R^2=0.37)$$

$$SG_{US} = 0.482 + (0.358 - 0.482) \cdot e^{(-\ln(2) \frac{Age}{7.054})} \quad (n=30; P<0.001; R^2=0.98)$$

$$SG_{UR} = 0.478 + (0.328 - 0.478) \cdot e^{(-\ln(2) \frac{Age}{6.913})} \quad (n=20; P<0.001; R^2=0.86)$$

Average SNA for 1-year-old trees was about 5.5 m² kg⁻¹, decreasing as trees aged to values of about 3.9 m² kg⁻¹ (Figure 8a). The models for whole-tree SG indicated that for trees growing in the southeastern U.S. (SG_{US}) and South America (SG_{SA}), SG of seedlings at planting was about 0.36 and 0.32, respectively, increasing as the trees aged to values of about 0.48 and 0.47, respectively (Table 3). The models predict that at an age of 20 years, $SG_{US} = 0.48$ and $SG_{SA} = 0.45$ (Figure 8b).

FIGURE 8

The curve that showed the best fit between minimum daily temperature (Tmin) and fractional leaf conductance (kF) is was:

$$kF = e^{(0.178 \cdot T_{min})} \quad (n=8, P<0.001; R^2=0.96).$$

On frost days, for each degree Celsius below zero during the night before, leaf conductance during the daytime was reduced at an exponential rate of 17.8%. For example, if a month had Frostday=5 and $T_{min}=-3^{\circ}\text{C}$, productivity would be reduced by $F_{frost}=0.902$ ($F_{frost}=1-(1-e^{(t_F \cdot T_{min})}) \cdot \text{Frostday} / 30$).

3.2. Iterative calibration of FR

We analyzed the relationship between FR and SI using iterative calibration on 63 plots randomly selected across the distribution range of loblolly pine. The silvicultural treatments applied in each of the selected plots created a wide span in productivity, resulting in SI ranging between 13.9 to 33.1 m (Table 2). Figure 9 shows the relationship between SI and FR. The model predicts a $FR=0.35$ for stands with $SI=15$ m and a $FR=0.96$ for stands with $SI=35$ m. The curve that showed the best fit and biological meaning was:

$$FR = \frac{1.1272}{1+14.9144 \cdot e^{(-0.1277 \cdot SI)}} \quad (n=63; P<0.001; R^2=0.68).$$

FIGURE 9

3.3. Model evaluation

There was agreement between observed and predicted values, with no clear tendencies to over or underestimate for any of the variables tested. Across all sites, including stands growing in Uruguay, the slope and the intercept of the relationship between predicted and observed values were not statistically different from one ($P>0.13$) and zero ($P>0.09$), respectively (Figure 10).

FIGURE 10

The LAI estimations performed well, predicting monthly values within the range of observed LAI reported for the IMPAC study. For the control plots that did not receive fertilization or herbicide treatments (Figure 11a), predicted and observed LAI followed similar trends, continuing to increase after age 14 years. The model also accurately estimated the amplitude and timing of seasonal LAI variation. For plots that received sustained fertilization (Figure 11b), herbicide (Figure 11c) and the combination (Figure 11d) of both treatments, the model

showed adequate predictions also following similar seasonal trends, showing a mean bias of about 2.3%. On the other hand, maximum monthly LAI (LAI-peak) was not well predicted by the model, having a mean underestimation of about 11% (Table 6).

FIGURE 11

Using monthly values of LAI shown previously, mean annual LAI (LAI-mean) was calculated for each plot and year of observation (Figure 12). There was a strong correlation between observed and predicted values ($P < 0.001$; $R^2 = 0.85$), showing a mean bias of 5.6% (data not shown).

FIGURE 12

All model performance tests showed that AGB, VOB, BA, Nha, Dq, H, LAI-mean, LAI-min and LAI-peak estimations agreed with measured values (Table 6). Across all sites and for all estimations, the RMSE ranged between 3% for Dq to 31% for VOB, both for stands growing in Uruguay. The Bias ranged between 14% under-estimations for LAI-peak at the IMPAC study and 14% over-estimations for VOB in Uruguay, both on stands with $SI > 25$ m. Estimated and observed values were highly correlated, with R^2 values ranging between 0.47 and 0.96. There was no clear trend indicating that there was different model behavior for stands with different SI. Even though the Bias percentage of AGB estimations increased from -2.6% for stands with $SI < 20$ m, to -16.3% for stands with $SI > 25$ m, the Bias percentage of VOB did not change. Furthermore, for LAI-min and LAI-mean, the Bias percentage was reduced as SI increased.

Table 6. Summary of model evaluation statistics for AGB, VOB, BA, Nha, Dq, H and LAI estimations for loblolly pine stands growing in sites with different SI in U.S. and Uruguay.

Variable	Location	SI class	$\bar{O}$	$\bar{P}$	n	RMSE	Bias	R^2
AGB	U.S.	15 - 20	89.7	92.0	184	16.9 (18.9)	-2.3 (-2.6)	0.854
		20 - 25	68.3	66.9	222	17.1 (25.1)	1.4 (2.1)	0.874
		25 - 30	71.2	82.9	104	16.9 (23.8)	-11.6 (-16.3)	0.939
	URUGUAY	20 - 25	130.3	132.4	30	15.5 (11.9)	-2.1 (-1.6)	0.866
		25 - 30	164.3	166.0	32	14.4 (8.8)	-1.7 (-1.0)	0.891
VOB*	U.S.	15 - 20	184.6	176.9	184	44.2 (23.9)	15.9 (8.6)	0.729
		20 - 25	133.4	131.3	222	41.1 (30.8)	13.6 (10.2)	0.814
		25 - 30	132.7	162.8	104	25.3 (19.1)	-11.4 (-8.6)	0.967

BA	URUGUAY	20 - 25	191.6	214.8	30	37.5 (19.6)	-23.2 (-12.1)	0.722
		25 - 30	239.4	272.1	32	46.1 (19.2)	-32.7 (-13.7)	0.602
	U.S.	15 - 20	25.6	27.2	184	5.4 (21.1)	-1.5 (-6.0)	0.767
		20 - 25	20.8	20.7	222	4.1 (19.5)	0.1 (0.4)	0.889
		25 - 30	20.9	23.2	104	4.5 (21.7)	-2.2 (-10.7)	0.917
Nha	URUGUAY	20 - 25	30.1	31.5	30	2.8 (9.4)	-1.5 (-4.8)	0.911
		25 - 30	33.8	35.1	32	3.0 (9.0)	-1.3 (-3.8)	0.915
	U.S.	15 - 20	1,045	1,042	184	117 (11.2)	3 (0.3)	0.923
		20 - 25	1,218	1,156	222	95 (7.8)	62 (5.1)	0.945
		25 - 30	1,471	1,437	104	147 (10.0)	34 (2.3)	0.956
Dq	URUGUAY	20 - 25	646	639	30	67 (10.3)	8 (1.2)	0.931
		25 - 30	508	518	32	48 (9.5)	-10 (-1.9)	0.961
	U.S.	15 - 20	18.5	19.1	184	1.8 (9.7)	-0.6 (-3.2)	0.912
		20 - 25	14.4	14.8	222	1.4 (9.7)	-0.5 (-3.3)	0.948
		25 - 30	13.3	14.2	104	1.8 (13.8)	-1.0 (-7.2)	0.925
H	URUGUAY	20 - 25	25.3	26.4	30	2.0 (7.7)	-1.1 (-4.3)	0.889
		25 - 30	30.6	31.2	32	1.0 (3.3)	-0.5 (-1.7)	0.966
	U.S.	15 - 20	13.9	16.6	184	2.3 (16.5)	-2.0 (-14.3)	0.763
		20 - 25	10.6	12.7	222	1.2 (11.2)	-0.7 (-6.4)	0.921
		25 - 30	10.0	11.2	104	1.2 (12.5)	0.1 (0.7)	0.920
	URUGUAY	20 - 25	15.7	17.2	30	1.5 (9.6)	-1.1 (-6.9)	0.682
		25 - 30	18.2	18.9	32	0.8 (4.3)	-0.1 (-0.8)	0.889
LAI-mean	IMPAC	13 - 20	1.59	1.59	33	0.35 (22.1)	-0.001 (-0.13)	0.862
		20 - 25	2.90	2.90	55	0.48 (16.7)	-0.001 (-0.04)	0.511
		25 - 30	3.13	2.94	44	0.43 (13.9)	0.184 (5.9)	0.587
LAI-min	IMPAC	13 - 20	1.16	1.30	33	0.30 (26.2)	-0.143 (-12.3)	0.814
		20 - 25	2.11	2.39	55	0.47 (22.0)	-0.279 (-13.2)	0.528
		25 - 30	2.30	2.44	44	0.35 (15.3)	-0.135 (-5.9)	0.540
LAI-peak	IMPAC	13 - 20	2.01	1.83	33	0.51 (25.0)	0.18 (8.8)	0.809
		20 - 25	3.65	3.32	55	0.72 (19.6)	0.34 (9.2)	0.473
		25 - 30	3.92	3.36	44	0.75 (19.3)	0.56 (14.2)	0.563

AGB: above-ground biomass (Mg ha^{-1}); VOB: stand bole volume outside bark ($\text{m}^3 \text{ha}^{-1}$); BA: stand basal area ($\text{m}^2 \text{ha}^{-1}$); Nha: trees per hectare (ha^{-1}); Dq: quadratic mean diameter (cm); H: mean tree height (m); LAI-mean: mean annual projected leaf area index ($\text{m}^2 \text{m}^{-2}$); LAI-min: minimum annual projected leaf area index ($\text{m}^2 \text{m}^{-2}$); LAI-peak: maximum annual projected leaf area index ($\text{m}^2 \text{m}^{-2}$); SI: site index class range (m); $\bar{O}$: mean observed value; $\bar{P}$: mean predicted value; n : number of observations; RMSE: root of mean square error (same unit as observed value); Bias: absolute bias (observed-predicted; same unit as observed value); R^2 : coefficient of determination. Values in parenthesis are percentage relative to observed mean.

*: For sites in Uruguay: VIB.

4. Discussion

The 3-PG model has broad potential for application e.g., for regional analysis of loblolly pine stand dynamics or assessing the impact of future climate scenarios on stand productivity across a wide range of ages and stand characteristics. Because we had access to a large, previously-unavailable set of data, we were able to revise some important parameter estimates using the alternative methods suggested by Gonzalez-Benecke et al. (2014) across a far broader range of climate, soils, and silvicultural intensity than previous researchers. The new approaches presented in this study provided new algorithms for canopy cover dynamics, fertility rating, needlefall dynamics, NPP allocation dynamics, mortality, frost intensity effects on production, quadratic mean diameter (and therefore, basal area), mean height and initial biomass estimations.

A critical variable used in 3-PG is FR, the empirical index that ranks soil fertility on a scale from 0 (extremely infertile) to 1 (optimum), and modifies canopy quantum efficiency and root allocation (Dye et al., 2004; Swenson et al., 2005; Fontes et al., 2006; Almeida et al., 2010; Gonzalez-Benecke et al., 2014a). Most previous applications of the 3-PG model relied on using arbitrary FR values determined by calibration which gave adequate results, but lacked a mechanistic basis and independence. Working with Douglas-fir (*Pseudotsuga menziesii*) stands, Swenson et al. (2005) indirectly correlated FR with SI by using 3-PG model to estimate SI, after computing FR from soil nitrogen content. The authors provided the logarithmic function but did not show any index of correlation or goodness of fit of the model fitted. Coops et al. (2011), working with five conifer species in Pacific Northwest of the U.S., also used 3-PG model to estimate SI, but used a constant value of FR=0.7 in all their calculations.

Dye et al. (2004), also correlated FR with SI, but estimated FR from known SI. The authors showed an alternative approach to estimate FR. This method was successfully tested by Gonzalez-Benecke et al. (2014a) for slash pine (*Pinus elliotii* var. *elliotii* Engelm.), supporting that FR was positively correlated with changes in SI; SI integrated a variety of factors including nutrient dynamics and site water balance. The model reported in this study predicted a maximum FR of 1 for stands with a SI of 37 m, a value that was above the range of observed plots (in our dataset consisting of 148 plots, only 8 plots had SI larger than 32 m, and those plots corresponded to studies with sustained weed control and fertilization). It is expected that improved genetics and silviculture can lead to higher values than those observed in our current database (Fox et al., 2007). The use of SI as a determinant of FR allows the model to be applied across a range of sites without resorting to model “tuning” to estimate FR. Nevertheless, further research is needed to improve the goodness of fit of that relationship. In a recent publication, Subedi et al. (2015) used SI to estimate FR for loblolly pine stands. The authors determined the relationship between stem volume at age 11 years and SI for each plot and then assigned a value of FR = 0 for SI=10.7 m and FR = 1 for SI of 30.5 m (plots with minimum and maximum volume at age 11 years observed in their dataset). Finally the authors arbitrarily assigned evenly the steps of FR associated to stem volume. We consider that our method is stronger as does not assume a linear relationship between FR and stem volume at age 11 years. Furthermore, instead of using one target age (11 years), we used all years of observations on each plot finding the value of FR that minimized the overall prediction error.

Assessments of mortality occurring prior to the onset of intraspecific competition (density-independent tree mortality) were not included in the original version of the model (Landsberg and Waring, 1997), that assumed no

mortality until self-thinning was triggered (density-dependent tree mortality). Several authors reported inadequate results in model performance, especially in stand density estimations (Sands and Landsberg, 2002; Pinjuv et al., 2006; Bryars et al., 2013). Following Sands (2004), who first proposed the inclusion of the density-independent model shown in eq. 4, Pérez-Cruzado et al. (2011) and Gonzalez-Benecke et al. (2014a) included, successfully, new species-specific parameter estimates for *Eucalyptus nitens* H. Deane and Maiden and slash pine, respectively, allowing for the estimation of random or stress-induced mortality observed under field conditions. Gonzalez-Benecke et al. (2014a) reported an overall bias of about 1% on stand density estimations. The inclusion of the mortality function reported by Harrison and Borders (1996) in the form of the model proposed by Sands (2004), greatly improved the estimations of stand density. In our study the overall bias was less than 3%.

In general, the model assumes that the fractional ground covered by the canopy (CanCover) is proportional to stand age until the age of full canopy cover (fullCanAge, years). This parameter is uncertain, as the age for full canopy cover depends on genetics, stand density and levels of productivity (Radke and Burkhardt, 1999). Bryars et al. (2013) reported a fullCanAge of 2 years, based on data reported by Burkes et al. (2003) for stands with high silviculture intensity and high planting density (between 3700 and 4400 trees ha⁻¹). Similar to our earlier findings (Gonzalez-Benecke et al., 2014a), we concluded that the use of a single age to determine the year to reach full canopy cover was not satisfactory. A relationship between fractional canopy cover, age and BA improved canopy cover estimations under different conditions of stand age, genetics, density and productivity in loblolly pine stands. This relationship was in agreement with our rationale that the moment when the stand reaches full canopy closure was correlated with stand density and productivity. With our new approach, the estimations of absorbed PAR are less dependent on arbitrary tuning of the CanCover parameter.

The model needs as inputs initial biomass allocations to roots (WR), foliage (WF) and stem (stemwood + bark + branches, WS). Sands and Landsberg (2002) emphasized the importance of the initial biomass conditions, especially to initial canopy development. If the model user has no initial biomass estimations for the stand to be simulated, we provide general individual-tree biomass functions for loblolly pine that use dbh, Ht and age as predictors. We also provide new biomass functions that can be used for young stands, where individual-tree biomass can be estimated using Ht and age. Using the raw data from 764 measured trees, we adapted the models reported by Gonzalez-Benecke et al. (2014b) and produced alternative functions to estimate initial biomass pools. The parameter estimates and relationships presented in this study have a wide range of applicability, and not only for 3-PG

initialization, but also life-cycle analysis (Puettmann et al., 2010), and estimation of stand biomass and carbon sequestration dynamics (Gonzalez-Benecke et al., 2011).

Landsberg and Sands (2011) remarked that relationships used for biomass allocation deserved further research. Previous versions of the model estimated NPP allocation to stem and foliage (pFS) based on individual-tree stem mass (kg tree^{-1}) and dbh. Gonzalez-Benecke et al. (2014a) introduced a new function to estimate pFS using BA rather than using dbh. For stands with a Dq of 20 cm, the modified model predicted pFS values between 0.35 and 0.45 (covering most of the distribution of observed values), and were larger than the comparable value of 0.25 reported by Bryars et al. (2013). Our new model used to estimate pFS included Dq and age, supporting our rationale that stand productivity and competition were correlated with changes in NPP allocation. These modifications contributed to our accurate predictions of stand biomass.

The model performed well when describing LAI dynamics, and mean annual LAI (LAI-mean) was well estimated (mean bias of 2.3% across all SI classes) in most cases. However, the model could not accurately reproduce the peak (mean bias of +11% across all SI classes) and minimum (mean bias of -10%, across all SI classes) LAI. This could be an effect of inadequacy of the model to capture the impacts of nutrition on foliage allocation and retention (Albaugh et al., 1998). Nevertheless, it is highly unlikely that stands receiving extremely high management inputs, such as those in some experimental plots, would be found under regular field conditions. An important area for future research is to better understand how intensive management and genetic selection affects LAI phenology.

Needlefall rate was a key factor that affected LAI estimations. Previous versions of the model used a function that described the dynamics of needlefall as trees developed, assuming a constant needlefall rate after canopy closure (Sands and Landsberg, 2002). Gonzalez-Benecke et al. (2014a) introduced equations to better describe the seasonal dynamics of needlefall for slash pine. We applied these equations to loblolly pine and found that the model was appropriate for the species, with needlefall also peaking in November; however, the maximum needlefall rate was 36% larger than for slash pine. Similar differences between species in total yearly needlefall were reported by Dalla-Tea et al. (1991). Our approach, that estimated a variable monthly needlefall rate, produced more realistic LAI dynamics than previous versions of the model, both in terms of average annual LAI, as well as within-year amplitude.

The dataset used for model validation includes plots with different treatments of fertilization and vegetation management. We used a simple approach to address the effects of those treatments by changing the FR reflected on observed SI on each plot. Wei et al. (2014) included the effects of competing vegetation on water and light availability in a recent publication on *P. ponderosa* trees. We recognize the great contribution of their study in including explicitly understory biomass into light interception and water balance modules of the model, but we also have to mention that the authors used a value of FR calibrated for each plot assuming no effect of FR on canopy quantum yield and only effects on biomass allocation. We are aware that competing vegetation has not only effects on nutrient availability, but our approach showed adequate results for loblolly pine stands on a variety of sites.

No other study using 3-PG with any species has shown the range of validation sites as reported here. Previous versions used with loblolly pine were validated using datasets restricted to single U.S. state, i.e., Georgia for Bryars et al. (2013) and North Carolina for Landsberg et al. (2001) and Sampson et al. (2006). In this study, the model was regionally validated under different stand and site conditions, covering zones beyond the native range for loblolly pine, including stands growing in Uruguay, South America. Even though other factors such as storms, disease and pests could be significant (VanderShaaf and Priesley, 2006; Aubrey et al., 2007; Coyle et al., 2015), the model showed adequate estimates over a range of climate and soil conditions. Interestingly, the validation results for stands in Uruguay were satisfactory, showing good agreement for all variables tested, and supported the robustness of the model. Even though the use of specific parameters for the Dq-WS, H-Dq and SG-Age functions greatly improved model performance, we consider that estimates of stem volume and LAI-peak could be improved as new data becomes available e.g., whole-tree SG for older trees, seasonal phenology of SNA and NPP allocation or soil/site variables that can improve the FR-SI relationship. These results have important implications, as the model can be used to capture the variation in loblolly pine productivity across the southeastern U.S., including sites in South America.

5. Conclusions

This study reports a regional validation of and alternative parameterization for the 3-PG model for loblolly pine. The new model was successfully tested against data from stands of varying characteristics that were distributed across the species' native range in the southeastern U.S., as well as in Uruguay, South America. We used the largest and most geographically extensive validation dataset across a species' range with 3-PG. In addition, data from the

literature and long-term studies were used to develop new functions for estimating important parameters for the model. Similar to previous work on slash pine, we estimated fertility rating based on a positive correlation with site index. The new parameterization and validation represents an improvement to previous published versions, allowing for improved assessments of stand productivity across a wide range of ages and stand characteristics for this species. The results in this study provide strong evidence that 3-PG can be applied over a wide geographical range using the same set of physiological parameters.

Acknowledgements

This research was supported by the U.S. Department of Defense, through the Strategic Environmental Research and Development Program (SERDP), the Pine Integrated Network: Education, Mitigation and Adaptation project (PINEMAP), a Coordinated Agricultural Project funded by the USDA National Institute of Food and Agriculture, Award #2011-68002-30185, the University of Florida's Carbon Resources Science Center and the Forest Biology Research Cooperative. Special thanks goes to Dr. Harold Burkhart at Virginia Tech University, Dr. Thomas Byram at Texas A&M University and to Mr. Daniel Ramirez and Mr. Juan P. Burgos at Cambium Forestal Uruguay S.A., for providing valuable data for parameter estimation and model validation.

References

- Adegbidi, H.G., Comerford, N.B., Jokela, E.J., Barros, N.F., 2002. Root development of young loblolly pine in Spodosols in southeast Georgia. *Soil Sci. Soc. Am. J.* 68, 596-604.
- Albaugh, T.J., Allen, H.L., Dougherty, P.M., Kress, L.W., King, J.S., 1998. Leaf area and above- and belowground growth responses of loblolly pine to nutrient and water additions. *Forest Sci.* 44, 317-328.
- Almeida, A.C., Landsberg, J.J., Sands, P.J., 2004. Parameterization of the 3-PG model for fast-growing *Eucalyptus grandis* plantations. *Forest Ecol. Manage.* 193, 179-195.
- Almeida, A.C., Siggins, A., Batista, T.R., Beadle, C., Fonseca, S., Loos, R., 2010. Mapping the effect of spatial and temporal variation in climate and soils on Eucalyptus plantation production with 3-PG, a process-based growth model. *Forest Ecol. Manage.* 259, 1730-1740.
- Aubrey, D.P., Coleman, M.D., Coyle, D.R. 2007. Ice damage in loblolly pine: understanding the factors that influence susceptibility. *Forest Sci.* 53, 5880-589.
- Aylott, M.J., Casella, E., Tubby, I., Street, N.R., Smith, P., Taylor, G., 2008. Yield and spatial supply of bioenergy poplar and willow short-rotation coppice in the UK. *New Phytol.* 178, 358-370.
- Barrichelo, L.E.G., Kageyama, P.Y., Speltz, R.M., Bonish, H.J., Brito, J.O., Ferreira, M., 1977. Estudos de procedências de *Pinus taeda* visando seu aproveitamento industrial. IPEF, Piracicaba, 15, 1-14.
- Barth, S.R., Fassola H.E., Crechi, E., Keller, A., Winck, R.A., Figueredo, E.I., Knebel O.E., 2013. Wood density of *Pinus taeda* L. implanted in Misiones-NE Corrientes, Argentina, on longitudinal and transverse directions. 15as Jornadas Técnicas Forestales y Ambientales. Facultad de Ciencias Forestales, UNaM - EEA Montecarlo, INTA. 7, 8 y 9 de Junio de 2012 -Eldorado, Misiones, Argentina.
- Binkley, D., 2002. Ten-year decomposition in a loblolly pine forest. *Can J. Forest Res.* 32, 2231-2235.
- Bryars, C., Maier, C., Zhao, D., Kane, M., Borders, B., Will, R., Teskey, R., 2013. Fixed physiological parameters in the 3-PG model produced accurate estimates of loblolly pine growth on sites in different geographic regions. *Forest Ecol. Manage.* 289, 501-514.
- Booth, T.H., Hong Hai, P., Khac Hieu, N., Jovanovic, T., Landsberg, J., Parsons, M., 2000. Increasing Carbon Sequestration in Planted Forests in Vietnam Through the Use of Genetically Improved Planting Stock, and Modeling to Quantify the Benefits Achieved. Component 2: Carbon Inventory and Growth Prediction. CSIRO Forestry and Forest Products Client Report for the International Greenhouse Partnerships Office, Canberra.
- Borders, B.E., Bailey, R.L., 2001. Loblolly pine - pushing the limits of growth. *South. J. Appl. For.* 25:69-74.
- Carlson, C.A., Fox, T.R., Allen, H.L., Albaugh, T.J., Rubilar, R.A., Stape, J.L., 2014. Growth responses of loblolly pine in the southeast United States to midrotation applications of nitrogen, phosphorus, potassium, and micronutrients. *Forest Sci.* 60, 157-169.
- Colbert, S.R., Jokela, E.J., Neary, D.G., 1990. Effects of annual fertilization and sustained weed control on dry matter partitioning, leaf area, and growth efficiency of juvenile loblolly and slash pine. *Forest Sci.* 36, 995-1014.
- Coops, N.C., Waring, R.H., 2001. Estimating forest productivity in the eastern Siskiyou Mountains of southwestern Oregon using satellite driven process model, 3-PGS. *Can. J. Forest Res.* 13, 143-154.

786 Coops, N.C., Waring, R.H., Law, B.E., 2005. Assessing the past and future distribution and productivity of
787 ponderosa pine in the Pacific Northwest using a process model, 3-PG. *Ecol. Model.* 183, 107-124.

788 Coops, N.C., Hember, R.A., Waring, R.H., 2010. Assessing the impact of current and projected climates on
789 Douglas-Fir productivity in British Columbia, Canada using a process-based model (3-PG). *Can. J. Forest Res.*
790 40, 511-524.

791 Coops, N.C., Gaulton, R., Waring, R.H., 2011. Mapping site indices for five Pacific Northwest conifers using a
792 physiologically based model. *Applied Veg. Sci.* 14, 268-276.

793 Coyle, D.R., Klepzig, K.D., Koch, F.H., Morris, L.A., Nowak, J.T., Oak, S.W., Orosina, W.J., Smith, W.D.,
794 Gandhi, K.J.K., 2015. A review of southern pine decline in North America. *Forest Ecol. Manage.* 349, 134-148.

795 Dalla-Tea, F., Jokela, E.J., 1991. Needlefall, canopy light interception, and productivity of young intensively
796 managed slash and loblolly pine stands. *Forest Sci.* 37, 1298-1313.

797 Diéguez-Aranda, U., Burkhart, H.E., Amateis, R.L., 2006. Dynamic site model for loblolly pine (*Pinus taeda* L.)
798 plantations in the United States. *Forest Sci.* 52:262-272.

799 Domec, J.-C., Noormets, A., King, J.S., Sun, G., McNulty, S.G., Gavazzi, M.J., Boggs J.L., Treasure E.A., 2009.
800 Decoupling the influence of leaf and root hydraulic conductances on stomatal conductance and its sensitivity to
801 vapour pressure deficit as soil dries in a drained loblolly pine plantation. *Plant Cell Environ.* 32, 980-991.

802 Dougherty, P.M., Hennessey, T.C., Zarnoch, S.J., Stenberg, P.T., Holeman, R.T., Wittwer, R.F. 1995. Effects of
803 stand development and weather on monthly leaf biomass dynamics of a loblolly pine (*Pinus taeda*) stand. *Forest*
804 *Ecol. Manage.* 72: 213-227.

805 Dye, P.J. 2001. Modelling growth and water use in four *Pinus patula* stands with the 3-PG model. *South Afr.*
806 *Forestry J.* 191, 53-64.

807 Dye, P.J., Jacobs, S., Drew, D., 2004. Verification of 3-PG growth and water-use prediction in twelve Eucalyptus
808 plantation stands in Zululand, South Africa. *Forest Ecol. Manage.* 193, 197-218.

809 Fassola, H. E., Crechi, E. H., Barth, S. R., Keller, A.E., Winck, R.A., Martiarena, R., Von Wallis, A., Pinazo, M. A.,
810 Knebel, O., 2012. Regional prediction models for aboveground biomass estimation of *Pinus taeda* in misiones
811 and NE corrientes region, Argentina. In 14as Jornadas Técnicas Forestales y Ambientales. Facultad de Ciencias
812 Forestales, UNaM - EEA Montecarlo, INTA. 10-12 de Junio de 2010 - Eldorado, Misiones, Argentina.

813 Flores, F.J., Allen, H.L., 2004. Efectos del clima y capacidad de almacenamiento de agua del suelo en la
814 productividad de rodales de pino radiata en Chile: un análisis utilizando el modelo 3-PG. *Bosque* 25, 11-24.

815 Fontes, L., Landsberg, J., Tomé, J., Tomé, M., Pacheco, C.A., Soares, P., Araujo, C., 2006. Calibration and testing
816 of a generalized process-based model for use in Portuguese Eucalyptus plantations. *Can. J. Forest Res.* 36,
817 3209-3221.

818 Fox, T.R., Jokela, E.J., Allen, H.L., 2007. The development of pine plantation silviculture in the southern United
819 States. *J. Forestry* 105, 337-347

820 Gibson, M.O., McMillin, C.W., Shoulders, E., 1985. Preliminary results for weight and volume of even-aged,
821 unthinned, planted southern pines on three sites in Louisiana. In: Saucier, J.R. (Ed.), *Proceeding of Southern*
822 *Forest Biomass Workshop* Athens, GA., June 5-7, 1984, pp. 69-74.

823 Gonzalez-Benecke, C.A., Martin, T.A., Jokela, E.J., 2011. A flexible hybrid model of life cycle carbon balance of
824 loblolly pine (*Pinus taeda* L.) management systems. *Forests* 2, 749-776.

825 Gonzalez-Benecke, C.A., Jokela, E.J., Martin, T.A., 2012 Modelling the effects of stand development, site quality
826 and silviculture on leaf area index, litterfall, and forest floor accumulation in loblolly and slash pine plantations.
827 Forest Sci. 58, 457-471.

828 Gonzalez-Benecke, C.A., Jokela, E.J., Cropper Jr., W.P., Bracho, R., Leduc, D.J., 2014a. Parameterization of model
829 3-PG for *Pinus elliottii* stands using alternative methods to estimate fertility rating, biomass partitioning and
830 canopy closure. Forest Ecol. Manage. 327:55-75.

831 Gonzalez-Benecke, C.A., Gezan, S.A., Albaugh, T.J., Burkhardt, H.E., Jokela, E.J., Maier, C.A., Martin, T.A.,
832 Rubilar, R.A., Samuelson, L.J., 2014b. Local and general above-stump biomass functions for loblolly and slash
833 pine trees. Forest Ecol. Manage. 334, 254-276.

834 Valentine, H.T., Mäkelä, A., Green, E.J. Amateis, R.L., Mäkinen, H., Ducey, M.J. 2012. Models relating stem
835 growth to crown length dynamics: application to loblolly pine and Norway spruce. Trees 26, 469-478.

836 Harrison, W.M.; Borders, B.E. 1996. Yield Prediction and Growth Projection for Site-Prepared Loblolly Pine
837 Plantations in the Carolinas, Georgia, Alabama and Florida; PMRC Technical Report 1996-1; University of
838 Georgia: Athens, GA, USA, p. 66.

839 Higa, A., Kageyama, P., Ferreira, M., 1973. Variação da densidade básica da madeira de *P. elliottii* var. *elliottii* e *P.*
840 *taeda*. IPEF, Piracicaba, 7, 79-91.

841 Huggett, R., Wear, D.N., Li, R., Coulston, J., Liu, S., 2013. Forest forecasts. In: Wear, D.N., Greis, J.G., (Eds.), The
842 Southern Forest Futures Project: technical report. Gen. Tech. Rep. SRS-GTR-178. Asheville, NC: USDA-Forest
843 Service, Southern Research Station, 542p.

844 Hynynen, J., Burkhardt, H.E., Allen, H.L., 1998. Modeling tree growth in fertilized midrotation loblolly pine
845 plantations. Forest Ecology Manage. 107, 213-229.

846 Jokela, E.J., Martin, T.A., 2000. Effects of ontogeny and soil nutrient supply on production, allocation and leaf area
847 efficiency in loblolly and slash pine stands. Can. J. Forest Res. 30, 1511-1524.

848 Kinerson, R., Higginbotham, K.O., Chapman, R.C. 1974. The dynamics of foliage distribution within a forest
849 canopy. J. Appl. Ecol. 11, 347-353.

850 Kinerson, R.S., Ralston, C.W., Wells, C.G., 1977. Carbon cycling in a loblolly pine plantation. Oecologia 29, 1-10.

851 Koralewski T.E., Wang, H-H., Grant, W.E., Byram, T.D., 2015. Plants on the move: Assisted migration of forest
852 trees in the face of climate change. Forest Ecol. Manage. 344, 30-37.

853 Landsberg, J.J., Waring, R.H., 1997. A generalized model of forest productivity using simplified concepts of
854 radiation-use efficiency, carbon balance and partitioning. Forest Ecol. Manage. 95, 209-228.

855 Landsberg, J.J., Johnsen, K.H., Albaugh, T.J., Allen, H.L., McKeand, S.E., 2001. Applying 3-PG, a simple process-
856 based model designed to produce practical results, to data from loblolly pine experiments. Forest Sci. 47, 43-51.

857 Landsberg, J.J., Waring, R.H., Coops, N.C., 2003. Performance of the productivity model 3-PG applied to a wide
858 range of forest types. Forest Ecol. Manage. 172, 199-214.

859 Landsberg J.J., Sands, P.J., 2011. Physiological Ecology of Forest Production: Principles, Processes and Modelling.
860 Oxford, UK: Elsevier Inc., pp 241-282.

861 Larcher, W. 1995. Physiological Plant Ecology, Third Edition. Springer, Berlin. 506 p.

862 Law, B.E., Waring, R.H., Anthoni, P.M., Aber, J.D., 2000. Measurements of gross and net productivity and water
863 vapor exchange of a *Pinus ponderosa* ecosystem, and an evaluation of two generalized models. *Global Change*
864 *Biol.* 6, 155-168.

865 Leggett, Z.H., Kelting, D.L., 2006. Fertilization effects on carbon pools in loblolly pine plantations on two upland
866 sites. *Soil Sci. Soc. Am. J.* 70, 279-286

867 Maier, C.A., Johnsen, K.H., Dougherty, P., McInnis, D., Anderson, P., Patterson, S., 2012. Effect of harvest residue
868 management on tree productivity and carbon pools during early stand development in a loblolly pine plantation.
869 *Forest Sci.* 58, 430-445.

870 Martin, T.A., Jokela, E.J., 2004. Developmental patterns and nutrition impact radiation use efficiency components in
871 southern pine stands. *Ecol. Appl.* 14, 1839-1854.

872 Myers R.H., 2000. Classical and modern regression with applications. PWS and Kent Published, 2nd edition.
873 Boston, 488 p.

874 Neter, J., Kutner, M.H., Nachtsheim, C.J., Wasserman, W., 1996 Applied linear statistical models. Irwin, 4th
875 edition, 770 p.

876 Nilsson, U., Allen, H.L., 2003. Short- and long-term effects of site preparation, fertilization and vegetation control
877 on growth and stand development of planted loblolly pine. *Forest Ecol. Manage.* 175, 367-377.

878 Noormets, A., Gavazzi, M.J., McNulty, S.G., Domec, J.-C., Sun, G., King, J.S., Chen, J., 2010. Response of carbon
879 fluxes to drought in a coastal plain loblolly pine forest. *Global Change Biol.* 16, 272-287.

880 Pereyra, O., Gelid, M., 2002. Estudio de la variabilidad de la densidad básica de la madera de *Pinus taeda* para
881 plantaciones de misiones y norte de corrientes. *Floresta* 33, 3-19.

882 Pérez-Cruzado, C., Muñoz-Sáez, F., Basurco, F., Riesco, G., Rodríguez-Soalleiro, R., 2011. Combining empirical
883 models and the process-based model 3-PG to predict *Eucalyptus nitens* plantations growth in Spain. *Forest*
884 *Ecol. Manage.* 262, 1067-1077.

885 Pezzutti R.V., 2011. Efeitos biológicos e econômicos de tratamentos silviculturais em plantios de *Pinus taeda* L. no
886 nordeste argentino. Tesis de Doctorado. Universidade Federal de Santa Maria, Santa Maria, RS, Brasil, pp. 179.

887 Pinjuv, G., Mason, E.G., Watt, M., 2006. Quantitative validation and comparison of a range of forest growth model
888 types. *Forest Ecol. Manage.* 236, 37-46.

889 Polster, H., Fuchs, S., 1963. Winterassimilation und -Atmung der Kiefer (*Pinus silvestris* L.) im mitteldeutschen
890 Binnenlandklima. *Archiv für Forstwesen* 12, 1011-1024.

891 Powers, R.F., Scott, D.A., Sanchez, F.G., Voldseth, R.A., Page-Dumroese, D., Elliof, J.D., Stone, D.M., 2005. The
892 North American long-term soil productivity experiment: findings from the first decade of research. *Forest Ecol.*
893 *Manage.* 220, 31-50.

894 Pretzsch, H., Matthew, C., Dieler, J., 2012. Allometry of tree crown structure. Relevance for space occupation at the
895 individual plant level and for self-thinning at the stand level. In *Growth and Defence in Plants*, Matyssek et al.
896 (eds.), Springer, pp 287-310.

897 Puettmann, M.E., Bergman, R., Hubbard, S., Johnson, L., Lippke, B., Oneil, E., 2010. Cradle-to-Gate Life-Cycle
898 Inventory of the US Wood Products Production: CORRIM Phase I and Phase II products. *Wood Fiber Sci.* 42,
899 15-28.

900 Rachid C.C., Mason E.G., Woollons, R., Resquin, F., 2014. Volume and taper equations for *P. taeda* (L.) and *E.*
901 *grandis* (Hill ex. Maiden). Agrociencia Uruguay 18, 47-60.

902 Radtke, P.J., Burkhart, H.E., 1999. Basal area growth and crown closure in a loblolly pine spacing trial. For. Sci. 45,
903 35-44

904 Rodríguez, R., Espinosa, M., Real P., Inzunza, J., 2002. Productivity analysis of radiata pine plantations under
905 different silvicultural regimes using the 3-PG process-based model. Austral. For. J. 65, 165-172.

906 Rodríguez, R., Real P., Espinosa, M., Perry, D.A., 2009. A process-based model to evaluate site quality for
907 *Eucalyptus nitens* in the Bio-Bio Region of Chile. Forestry 82, 149-162.

908 Roth, B.E., Jokela, E.J., Martin, T.A., Huber, D.A., White, T.L., 2007. Genotype x environment interactions in
909 selected loblolly and slash pine plantations in the Southeastern United States. Forest Ecol. Manage. 238, 175-
910 188.

911 Russell, M.B., Amateis, R.L., Burkhart, H.E., 2010. Implementing regional locale and thinning response in the
912 loblolly pine height-diameter relationship. South. J. Appl. For. 34: 21-27.

913 Sampson, D.A., Waring, R.H., Maier, C.A., Gough, C.M., Ducey, M.J., Johnsen, K.H., 2006. Fertilization effects on
914 forest carbon storage and exchange, and net primary production: A new hybrid process model for stand
915 management. Forest Ecol. Manage. 221, 91-109.

916 Samuelson, L.J., Johnsen, K., Stokes, T., 2004. Production, allocation, and stemwood growth efficiency of *Pinus*
917 *taeda* L. stands in response to 6 years of intensive management. Forest Ecol. Manage. 192, 59-70.

918 Sanchez, F.G., Scott, D.A., Ludovici, K.H., 2006. Negligible effects of severe organic matter removal and soil
919 compaction on loblolly pine growth over 10 years. Forest Ecol. Manage. 227, 145-154.

920 Sands, P.J. and Landsberg, J.J., 2002. Parameterization of 3-PG for plantation-grown *Eucalyptus globulus*. Forest
921 Ecol. Manage. 163, 273-292.

922 Sands, P.J., 2004. Adaptation of 3-PG to novel species: guidelines for data collection and parameter assignment.
923 Technical Report 141, CRC for Sustainable Production Forestry, Hobart, Australia, 34 p.

924 Sands, P.J., 2010. 3PG_{PJS} user manual, 27 p.

925 Shan, J., Morris, L.A., Hendrick, R.L., 2001. The effects of management on soil and plant carbon sequestration in
926 slash pine plantations. J. Appl. Ecol. 38, 932-941.

927 Stape, J.L., Binkley, D., Ryan, M.G., 2004. Testing the utility of the 3-PG model for growth of *Eucalyptus grandis* x
928 *urophylla* with natural and manipulated supplies of water and nutrients. Forest Ecol. Manage. 193, 219-234.

929 Subedi, S., Fox, T.R., Wynne, R.H., 2015. Determination of fertility rating (FR) in the 3-PG model for loblolly pine
930 plantations in the Southeastern United States based on site index. Forests 6, 3002-3027.

931 Swenson, J. J., Waring, R. H., Fan, W., Coops, N., 2005. Predicting site index with a physiologically based growth
932 model across Oregon, USA. Can. J. Forest Res. 35, 1697-1707.

933 Teskey, R.O., Bongarten, B.C., Cregg, B.M., Dougherty, P.M., Hennessey, T.C., 1987. Physiology and genetics of
934 tree growth response to moisture and temperature stress: an examination of the characteristics of loblolly pine
935 (*Pinus taeda* L.). Tree Physiol. 3, 41-61.

- 936 Tuttle, C.L. 1978. Root biomass and nutrient content of a 25-yearold loblolly pine (*Pinus taeda* L.) plantation in east
937 Texas. M.Sc. thesis, Texas A&M Univ., College Station, TX, 69 p,
- 938 VanderSchaaf, C.L. and Prisley, S.P., 2006. Factors affecting site productivity of loblolly pine plantations across the
939 Southeastern United States. In Prisley, S., Bettinger, P., Hung, I-K., Kushla, J. eds. Proceedings of the 5th
940 Southern Forestry and Natural Resources GIS Conference, June 12-14, 2006, Asheville, NC. Warnell School of
941 Forestry and Natural Resources, University of Georgia, Athens, GA.
- 942 Van Deusen, P.C., Sullivan, A.D., Matney, T.G., 1981. A prediction system for cubic foot volume of loblolly pine
943 applicable through much of its range. South. J. Appl. For. 5, 186-189.
- 944 Von Wallis, A., Martiarena, R.A., Pinazo, M.A., 2007. Densidad básica de la madera de *Pinus taeda* L. Marion en
945 función de la densidad arbórea del rodal, altura y posición transversal en el fuste. III Congreso Iberoamericano
946 de Productos Forestales. IBEROMADERA 2007, 3 al 5 de Julio 2007. Buenos Aires. Argentina.
- 947 Waring, R.H., 2000. A process model analysis of environmental limitations on growth of Sitka spruce plantations in
948 Great Britain. Forestry 73, 65-79.
- 949 Waring, R., Nordmeyer, A., Whitehead, D., Hunt, J., Newton, M., Thomas, C., Irvine, J. 2008. Why is the
950 productivity of Douglas-fir higher in New Zealand than in its native range in the Pacific Northwest, USA? For.
951 Ecol. Manage. 255, 4040-4046.
- 952 Weber, E.M., 2005. Densidad básica de madera de *Pinus taeda* L. Marion de diferentes edades, Misiones,
953 Argentina. Floresta 35, 487-494.
- 954 Wei, L., Marshall, J.D., Zhang, J., Zhou, H., Powers, R.F. 2014. 3-PG simulations of young ponderosa pine
955 plantations under varied management intensity: Why do they grow so differently? For. Ecol. Manage. 255,
956 4040-4046.
- 957 Zhao, M.F., Xiang, W.H., Peng, C.H., Tian, D.L., 2009. Simulating age-related changes in carbon storage and
958 allocation in a Chinese fir plantation growing in southern China using the 3-PG model. Forest Ecol. Manage.
959 257, 1520-1531.
- 960 Zhao, D., Kane, M. and Borders, B., 2011. Growth response to planting density and management intensity and
961 planting density in loblolly pine plantations in the southeastern United States. Annals Forest Sci. 68, 625-635.
- 962 Zhao, D., Kane, M., Borders, B., Subedi, S., Akers, M., 2012. Effects of cultural intensity and planting density on
963 stand-level aboveground biomass production and allocation of 12-year old loblolly pine plantations in the Upper
964 Coastal Plain and Piedmont of the southeastern United States. Can. J. Forest Res. 42, 111-122.

Figure legends.

Fig. 1. a) Location of sites used for validation (black circle, n=91 plots in 24 sites) or FR/SI analysis (grey triangle, n=63 plots in 16 sites) in the U.S.: the species natural distribution range is the shaded area; b) Location of sites used for validation in Uruguay (n=10 plots). AL: Alabama; AR: Arkansas; FL: Florida; GA: Georgia; LA: Louisiana; MS: Mississippi; NA: North Carolina; OK: Oklahoma; SC: South Carolina; TN: Tennessee; TX: Texas; VA: Virginia.

Fig 2. Model fitting for canopy conductance sensitivity to VPD. Data from a 15-16 year-old eddy-covariance site located in the Lower Coastal Plain of North Carolina, U.S.

Fig. 3. Figure 3. a) Relationship between observed values of Dq and NPP allocation to foliage (pF), stem (pS) and stem to foliage ratio (pFS) for loblolly pine stands ranging in age from 2 to 25 years old (all data points are observed values); b) Observed and predicted values of pFS. Panels c) and d) shows inserts for Dq smaller than 5 cm.

Fig 4. Allometric relationships for a) stem biomass (WS) and Dq, b) Dq and mean height (H), c) age and branch and bark fraction (pBB), and d) bole volume inside bark (VIB) and bole volume ratio (Vratio) for different planting density (PD) and site index (not labelled). Panels a) and b) show observed (filled symbol) and predicted (open symbol)) values for stands growing in U.S. (circles) and Uruguay (triangles).

Fig. 5. Tree mortality relationships. a) Relationship between age density-independent tree mortality (γNt) using the model of Harrison and Borders (1996); b) Density-dependent tree mortality, based on the relationship between stem biomass (WS, kg tree⁻¹) and stand density (both in natural logarithm scale). The self-thinning line is the theoretical upper limit.

Fig 6. Figure 6. Relationship between canopy development (expressed as fractional canopy cover, CanCover) and a) age and b) basal area for loblolly pine stands of PPINES study. Plots that received combinations of two contrasting silvicultural treatments (operational, O, and high intensity, H) and two contrasting planting densities (1,334 trees ha⁻¹, W, and 2,990 trees ha⁻¹, N). Panel a) show mean values for stands growing in GA. Panel b) show observed and

predicted values for all plots growing on sites in FL and GA. Dashed line represents the point when the stand reach canopy closure (CanCover=1).

Fig. 7. Seasonal dynamics of foliage biomass and needlefall for the IMPAC study, showing average monthly a) foliage dry mass (WF, Mg ha^{-1}), b) needlefall (NF, $\text{Mg ha}^{-1} \text{ month}^{-1}$) and c) fractional needlefall (γ_N as a proportion of annual maximum).

Fig 8. Model fitted for age-dependent relationship between for a) specific needle area (SNA) and b) whole-tree wood specific gravity (SG) for trees growing in southeast U.S. (SG_{US} , solid line; adapted from Harrison and Borders, 1996) and Uruguay (SG_{UR} , dashed line).

Fig. 9. Relationship between site index (SI, m) and FR after iterative calibration for 63 plots in 12 states in southeastern U.S. AL: Alabama; AR: Arkansas; FL: Florida; GA: Georgia; LA: Louisiana; MS: Mississippi; NA: North Carolina; OK: Oklahoma; SC: South Carolina; TN: Tennessee; TX: Texas; VA: Virginia.

Fig. 10. Model validation for 34 tested sites (101 plots total; 91 in the southeastern U.S.; 10 in Uruguay, South America) for different site index. Observed versus predicted (simulated with 3-PG) values of a) total above ground biomass (AGB, Mg ha^{-1}), b) stand density (Nha, trees ha^{-1}); c) bole volume over-bark (VOL, $\text{m}^3 \text{ ha}^{-1}$); d) stand basal area (BA, $\text{m}^2 \text{ ha}^{-1}$); e) quadratic mean diameter (Dq, cm), and f) mean tree height (H, m). The dotted line corresponds to the 1-to-1 relationship.

*: For sites in Uruguay: VIB.

Fig. 11. Validation of monthly projected leaf area index (LAI, $\text{m}^2 \text{ m}^{-2}$). Observed (filled circle) and predicted (open circle) values for loblolly pine stands grown at the IMPAC study under the following silvicultural treatments a) Control, b) Fertilization, c) Herbicide, and d) Fertilization + Herbicide (Jokela and Martin, 2000).

1019 **Fig. 12.** Relationship between observed and predicted mean annual projected leaf area index (LAI, $\text{m}^2 \text{m}^{-2}$) for
1020 loblolly pine stands grown at the IMPAC study under the following silvicultural treatments: control (C), herbicide
1021 (H), fertilization (F), and fertilization + herbicide (FH) (Jokela and Martin, 2000).
1022

Figure 1. a) Location of sites used for validation (black circle, n=91 plots in 24 sites) or FR/SI analysis (grey triangle, n=63 plots in 16 sites) in the U. S.: the species natural distribution range is the shaded area; b) Location of sites used for validation in Uruguay (n=10 plots). AL: Alabama; AR: Arkansas; FL: Florida; GA: Georgia; LA: Louisiana; MS: Mississippi; NA: North Carolina; OK: Oklahoma; SC: South Carolina; TN: Tennessee; TX: Texas; VA: Virginia.

Figure 2. Model fitting for canopy conductance sensitivity to VPD. Symbols are mean daily data from eddy covariance measurements in a 15-16 year-old stand located in the Lower Coastal Plain of North Carolina, U.S.

Figure 3. a) Relationship between observed values of Dq and NPP allocation to foliage (pF), stem (pS) and stem to foliage ratio (pFS) for loblolly pine stands ranging in age from 2 to 25 years old (all data points are observed values); b) Observed and predicted values of pFS. Panels c) and d) shows inserts for Dq smaller than 5 cm.

Figure 4. Allometric relationships for a) stem biomass (WS) and Dq, b) Dq and mean height (H), c) age and branch and bark fraction (pBB), and d) bole volume inside bark (VIB) and bole volume ratio (Vratio) for different planting density (PD) and site index (not labelled). Panels a) and b) show observed (filled symbol) and predicted (open symbol) values for stands growing in U.S. (circles) and Uruguay (triangles).

1071

1072

1073 Figure 5. Tree mortality relationships. a) Relationship between age density-independent tree mortality (γNt)
 1074 using the model of Harrison and Borders (1996); b) Density-dependent tree mortality, based on the relationship
 1075 between stem biomass (WS, kg tree⁻¹) and stand density (both in natural logarithm scale). The self-thinning line is
 1076 the theoretical upper limit.

1077

1078

1079

1080

1081

Figure 6. Relationship between canopy development (expressed as fractional canopy cover, CanCover) and a) age and b) basal area for loblolly pine stands of PPINES study. Plots that received combinations of two contrasting silvicultural treatments (operational, O, and high intensity, H) and two contrasting planting densities (1,334 trees ha^{-1} , W, and 2,990 trees ha^{-1} , N). Panel a) show mean values for stands growing in GA. Panel b) show observed and predicted values for all plots growing on sites in FL and GA. Dashed line represents the point when the stand reach canopy closure (CanCover=1).

Figure 7. Seasonal dynamics of foliage biomass and needlefall for the IMPAC study, showing average monthly a) foliage dry mass (WF, Mg ha^{-1}), b) needlefall (NF, $\text{Mg ha}^{-1} \text{ month}^{-1}$) and c) fractional needlefall (γN as a proportion of annual maximum).

Figure 8. Model fitted for age-dependent relationship between for a) specific needle area (SNA) and b) whole-tree wood basic specific gravity (SG) for trees growing in southeast U.S. (SG_{US} , solid line; adapted from Harrison and Borders, 1996) and Uruguay (SG_{UR} , dashed line).

Figure 9. Relationship between site index (SI, m) and FR after iterative calibration for 63 plots in 12 states in southeastern U.S. AL: Alabama; AR: Arkansas; FL: Florida; GA: Georgia; LA: Louisiana; MS: Mississippi; NA: North Carolina; OK: Oklahoma; SC: South Carolina; TN: Tennessee; TX: Texas; VA: Virginia.

Figure 10. Model validation for 34 tested sites (101 plots total; 91 in the southeastern U.S.; 10 in Uruguay, South America). Observed *versus* predicted (simulated with 3-PG) values of a) total above ground biomass (AGB, Mg ha⁻¹), b) stand density (Nha, trees ha⁻¹); c) bole volume over-bark (VOL, m³ ha⁻¹); d) stand basal area (BA, m² ha⁻¹); e) quadratic mean diameter (Dq, cm), and f) mean tree height (H, m). The dotted line corresponds to the 1-to-1 relationship.

*: For sites in Uruguay: VIB.

Figure 11. Validation of monthly projected leaf area index (LAI, $\text{m}^2 \text{m}^{-2}$). Observed (filled circle) and predicted (open circle) values for loblolly pine stands grown at the IMPAC study under the following silvicultural treatments a) Control, b) Fertilization, c) Herbicide, and d) Fertilization + Herbicide (Jokela and Martin, 2000).

Figure 12. Relationship between observed and predicted mean annual projected leaf area index (LAI-mean, $\text{m}^2 \text{m}^{-2}$) for loblolly pine stands grown at the IMPAC study under the following silvicultural treatments: control (C), herbicide (H), fertilization (F), and fertilization + herbicide (FH) (Jokela and Martin, 2000).