Biotic and abiotic impacts of Double-crested Cormorant breeding colonies on forested islands in the southeastern United States

Diana J. R. Lafferty^{ae1}, Katie C. Hanson-Dorr^{b2}, Amanda M. Prisock^{c3}, Brian S. Dorr^{b*}

^a*Carnivore Ecology Laboratory, Department of Wildlife, Fisheries and Aquaculture, Mississippi State University, Mississippi State, MS, USA*

^b*US Department of Agriculture, Wildlife Services National Wildlife Research Center, Mississippi Field Station, P.O. Box 6099, Mississippi State, Mississippi, 39762, USA*

^c*Weyerhaeuser Company, 29 Tom Rose Road, Columbus, MS 39704, USA*

^e*Department of Forestry and Environmental Resources, Program in Fisheries, Wildlife and Conservation Biology, David Clark Labs, North Carolina State University, Raleigh, NC 27695, USA*

* Corresponding author. Tel.: +1 662 325 8216

E-mail addresses: djr.lafferty@gmail.com (DJR Lafferty), katie.c.hanson-dorr@aphis.usda.gov (KC Hanson-Dorr), amanda.prisock@weyerhaeuser.com (AM Prisock), brian.s.dorr@aphis.usda.gov (BS Dorr).

¹ Tel.: +1 919 513 3312

² Tel.: +1 662 325 5489

³ Tel.: +1 662 245 5250

Author Contributions: BD and KCHD conceived of the idea; BD, KCHD and AMP conducted fieldwork; DJRL conducted statistical analyses; DJRL, wrote the manuscript; BD, KCHD, AMP reviewed the manuscript and provided editorial advice.

ABSTRACT: Double-crested cormorant (*Phalacrocorax auritus*) numbers have increased in North America, and concomitantly cormorants appear to be expanding their nesting range in the southeastern United States. Because colonial nesting waterbirds can impact water quality, soil chemistry and subsequent vegetation succession patterns, our goal was to assess the extent to which cormorant breeding colonies are influencing the biotic and abiotic attributes of forested islands in the southeastern United States. Our objectives were to 1) compare water quality characteristics in near-shore surface waters around forested islands with and without nesting cormorants during the peak-nesting/fledgling period and post-fledgling period, 2) measure soil chemistry parameters for forested islands with and without nesting cormorants, and 3) compare tree health metrics on forested islands with and without nesting cormorants. Our results indicate that cormorant colonies are not significant contributors to general coliforms or *E. coli* levels in waters surrounding southern breeding colonies. Cormorants also do not appear to have significant direct effects on water chemistry. We did find that cormorant colonies are affecting soil chemistry. Soil from within the nesting colony was more acidic and had greater concentrations of phosphorous than soils on reference islands. In addition, we found evidence that cormorants are negatively affecting tree health within nesting colonies as evidenced by a greater number of trees of lower vigor class within the nesting colonies compared to reference sites. While cormorants do cause abiotic and biotic changes, these are part of the natural ecological processes that occur following waterbird colonization. Management to reduce unwanted impacts that nesting cormorants are having on forested island habitats should be considered within a framework that allows for natural ecological processes, including changes in soil chemistry and subsequent vegetation succession.

Key words: double-crested cormorant, forest health, human-wildlife conflict, soil chemistry, waterbirds, water quality

1. Introduction

In North America, the Double-crested cormorant (*Phalacrocorax auritus*; hereafter cormorant) has substantially increased in abundance from relatively low numbers in the early 1970's (Dorr et al. 2012). Cormorants have historically wintered in the southeastern United States, but nesting cormorants in the region were relatively rare compared to their northern breeding regions and occurred primarily along the gulf coast and major rivers (Wires and Cuthbert 2006, Hanson et al. 2010). More recently cormorants appear to have expanded their southeastern nesting range to include man-made sloughs, lakes, and reservoirs not historically available or reported (Reinhold et al. 1998, Hatch and Weseloh 1999, Wires and Cuthbert 2006, Hanson et al. 2010). The increased numbers of cormorants throughout their range, while a conservation success story, has also come with increased human-cormorant conflicts (Taylor and Dorr 2003, Dorr et al. 2012). Conflicts associated with nesting cormorants differ from those associated with wintering cormorants. Issues with wintering cormorants typically are associated with direct predation on aquaculture or recreational fisheries (Taylor and Dorr 2003), whereas cormorant breeding colonies may have broader impacts due to increased competition with co-nesting species, vegetation damage, and changes in soil and water quality associated with the release of nutrient-rich waste due to guano deposition, regurgitated food, carcasses and feathers (Jarvie et al. 1999, Shieldcastle and Martin 1999, Taylor and Dorr 2003, Ayers et al. 2015, Stewart et al. 2015). Jarvie et al. (1999) found that Black-crowned Night Herons (*Nycticorax nycticorax*) nesting pairs declined with increasing cormorant nesting. Herbert et al. (2005)

documented significant damage to trees on forested islands associated with cormorant nesting and that vegetation damage can negatively impact co-nesting avian species that are obligate tree nesters. Fish-eating avian species, such as cormorants, also can affect soil quality (Ishida 1996, Ligeza and Smal 2003, Breuning-Madsen et al. 2010, Ayers et al. 2015). Ishida (1996) found that changes in soil nutrients caused by nesting great cormorants (*Phalacrocorax carbo*) simplified forest structure and reduced tree species diversity. Breuning-Madsen et al. (2010) found that great cormorant colonies significantly impacted soil nutrient profiles that affected plant community composition by limiting plant diversity to salt-tolerant plants that can grow in extremely nutrient-rich soils.

In addition to the potential terrestrial ecological impacts of nesting cormorant colonies, waterfowl feces may be an important source of fecal contamination in water bodies (Standridge et al. 1979, Makino et al. 2000, Lu et al. 2008). High levels of coliform and *E. coli* bacteria are a concern for many recreational water environments because of the known association between fecal matter and human health risks, and are commonly used as indicator species to measure the quality of recreational and drinking waters (USEPA 1986, 1999, WHO 2006). For example, McLellan (2004) found that waterbird feces was an important source of bacterial pollution in some beach areas and contributed to beach closures in efforts to limit potential human health risks. Although research has been conducted on impacts of cormorants on co-nesting species, vegetation, and soil and water quality on their northern breeding grounds (Bédard et al. 1995, Jarvie et al. 1999, Shieldcastle and Martin 1999, Hebert et al. 2005, Ayers et al. 2015), no research has been conducted on the impact of cormorant breeding colonies in southeastern United States in this regard.

Within the last 10 – 15 years cormorants began nesting on islands in Guntersville, Lake, AL, resulting in concerns regarding impacts to fisheries, vegetation, and soil and water quality (Barras 2004). Impacts to several islands in Guntersville Lake have been observed including loss of vegetation and possibly increased shoreline erosion (Barras 2004), presumably due to nesting cormorants. Given the diversity of soil and vegetation types, longer growing season, and generally more eutrophic water-bodies, the ecological impacts of cormorants at southern breeding colony sites may differ substantially from impacts documented at northern colonies. Thus, our goal for this study is to understand if and how cormorant breeding colonies influence biotic and abiotic attributes of forested islands in the southeastern United States. Our objectives were to 1) compare water quality characteristics in near-shore surface waters around forested islands with and without nesting cormorants during the peak-nesting/fledgling period and post-fledgling period, 2) measure soil chemistry parameters for forested islands with and without nesting cormorants, and 3) compare tree health metrics on forested islands with and without nesting cormorants.

2. Methods

2.1. Study site – Lake Guntersville Alabama, USA.

Lake Guntersville (34° 32' 19.82 N, 86° 07' 05.14 W) is the largest lake (27,964 ha) in Alabama, U.S.A. (Fig. 1) and spans 121 km from Nickajack Dam, Tennessee, to Guntersville Dam, Alabama. The reservoir is managed by the Tennessee Valley Authority (TVA) on the Tennessee River in northeastern Alabama and southeastern Tennessee. Cormorants currently nest at three primary locations in Lake Guntersville: Conner's Island (CON), South Sauty Island (SSS), and North Sauty Island (NS); hereafter, these islands will be referred to as colony islands. Reference islands that lacked nesting cormorant colonies were selected based on similar size and

proximity (adjacency) to colony islands: southeast of Connor's Island (SEC), north of South Sauty (NSS), west of South Sauty (WSS). When more than one reference island was available, the reference island was selected at random.

2.2. Study Design

2.2.1 Water sampling

Water samples for microbial and chemical analyses were collected at each of the colony islands and the reference islands during the period of peak nesting/fledging of chicks (mid-May) and during the post fledging period when cormorant numbers are at a minimum (mid-August), 2010. Island shorelines were mapped using digital orthoquad imagery ArcGIS ArcMap (ESRI® 9.1, Esri Inc., Redlands, CA, U.S.A).

For microbial analysis, the largest islands shoreline was divided into 10 equal length sample sections. The beginning of each section represents a shoreline starting point for obtaining water samples. Shoreline sections of smaller islands were sampled in proportion to their length relative to the shoreline of the largest island. If section lengths were less than 100 m, a minimum of four sample points were selected, one in each cardinal direction. Water samples were collected at each shoreline section starting point at intervals of 5 m, 15 m and 25 m from the shoreline. Water samples were placed in sterile Whirl-Pak® bags and placed on ice until all water samples were collected. Samples were plated within 4 h of collection (Vail et al. 2003) on 3M Petrifilm™ *E. coli*/Coliform plates (Petrifilm™) for enumerating general coliform and *E. coli* bacteria (colony forming units [CFU]). Petrifilm™ count plates are a simple, safe, reliable and low cost method for monitoring environmental water samples in the field (Vail et al. 2003). Prior to plating, water samples were allowed to increase to ambient temperature. Plates were inoculated with 1 ml of water using sterile, disposable pipets and allowed to incubate at $35 \pm 1^\circ$

C for 24 ± 2 h prior to enumeration. Used plates were disinfected before disposal by soaking in 1% hypochlorite solution for 1-hour.

Due to logistic constraints associated with the amount of time to run water chemistry tests on-site, samples were taken only at 5 m from shore and at 4 sample locations, one in each cardinal direction from each study island during both the peak nesting/fledgling period (mid-May) and post fledgling period (mid-August). Four water quality parameters were tested: pH, nitrate (NO_3 mg/L), ammonia (NH_3 mg/L), and phosphate (PO_4 mg/L). A digital meter was used to measure pH (± 0.1 pH at 20°C). A Hach multi-parameter surface water test kit was used to measure water chemistry parameters.

2.2.2 Soil sampling

Soil samples were collected from each of the colony islands and reference islands. To establish sampling plots, each of the six islands was overlain with a $10\text{ m} \times 10\text{ m}$ (0.10 ha) grid using the most recent digital orthoquad imagery available for the area via ArcGIS ArcMap (ESRI[®] 9.1, Esri Inc., Redlands, CA, U.S.A). Sampling of 0.10 ha plots was conducted in proportion to island area by randomly sampling 20% of the total available plots on each island. Soil core samples were collected with a soil auger (9 cm diameter $\times$ 20 cm deep) at the center of each plot, excluding surface debris. All soil samples were assigned a treatment level based on location relative to nesting colonies that ranged from 1 to 3 (1 = colony island within a nesting colony; 2 = colony island outside a nesting colony; 3 = reference island). Soil samples were labeled and placed in ice chests in doubled paper bags for transport to Mississippi State University Extension Service, Soils Testing Laboratory (Mississippi State, MS, U.S.A.) for analyses. Extractable nutrient concentrations (kg/ha), milli-equivalents/100g, and base saturation were determined for percent organic material (%OM), nitrate ($\text{NO}_3\text{-N}$), phosphorous (P),

potassium (K), calcium (Ca), magnesium (Mg), zinc (Zn), sulfur (S), sodium (Na), and soil pH for each sample. Nutrients were selected based on importance in biogeochemical circulation and plant physiology (Ligeza and Smal 2003, Ishida 1996, Breuning-Madsen et al. 2010). Soil samples at all locations were collected during the peak cormorant nesting/fledging period (mid-May).

2.2.3 Tree surveys

Randomly selected 10 m diameter circular plots (e.g. 0.02 ha) were used to survey trees on colony and reference islands. During surveys, species composition, tree diameter at breast height (DBH) in cm, tree health status (e.g., vigor class described below), the number of cormorant nests and the number of nests of other co-nesting waterbird species were recorded. All surveyed trees were assigned a treatment level based on location that ranged from 1 to 3 (1 = colony island within a nesting colony; 2 = colony island outside a nesting colony; 3 = reference island).

Tree health status criteria were adapted from the forest inventory and analysis national core field guide (USFS 2007). We added 2 categories for standing dead trees to provide a Likert Scale measure (5 categorical values) ranging from healthy to standing dead. Vigor classes are defined as follows: Vigor Class 1 □ uncompact live crown ratio of 35% or higher, < 5% dieback, and > 80% of the foliage is normal or > 50 percent of each leaf is not damaged or missing. Twigs and branches that are dead because of normal shading are not included. Vigor Class 2 □ saplings do not meet Class 1 or 3 criteria. May have uncompact live crown ratio, may or may not have dieback and may have between 21 and 100% of the foliage classified as normal. Vigor Class 3 □ saplings may have uncompact live crown ratio and have 1 □ 20% normal foliage or the percent of foliage missing combined with the percent of leaves that are over 50%

183 damaged or missing equal 80% or more of the live crown. Twigs and branches that are dead
184 because of normal shading are not included. Vigor Class 4 □ fresh standing dead tree, limbs
185 almost all present and top pointed or not broken and 90 □ 100% of bark remaining. Vigor Class
186 5 □ old standing dead tree, some or almost all limbs broken and/or top broken, and 10 □ 100%
187 bark slippage (loss).

188 *2.2.4 Cormorant nesting and co-nesting species surveys*

189 Cormorant and co-nesting waterbird species surveys were conducted by an observer
190 using binoculars from a boat that circled the island at < 100 m. During this survey, the number of
191 cormorants and co-nesting species nests observed in each tree were documented and the presence
192 or absence of chicks in each nest were recorded. When nests were observed, the tree species was
193 identified and classified as alive or dead.

194 *2.3. Statistical analyses*

195 *2.3.1 General approach*

196 We conducted all statistical analyses in R 3.0.2 (R Development Core Team 2014).
197 Following Zuur et al. (2009), we began model building by discerning appropriate model
198 distributions (e.g., Poisson). For example, if the data were overdispersed, we examined the
199 application of negative binomial or zero-inflated Poisson models. We also assess the support for
200 random effects and for the inclusion of spatial correlation structures and alternative variance
201 structures when appropriate. We used likelihood ratio tests to test support for the most complex
202 models and for sequentially more parsimonious models until dropping additional covariates did
203 not improve model fits (Zuur et al. 2009). Model validation was conducted following Zuur et al.
204 (2009, 2012). We examined model coefficients with 95% confidence intervals for interpretation
205 of covariate effects if intervals excluded zero.

2.3.2 Water quality

To test for potential effects of nesting cormorants on general coliform and *E. coli* levels in near shore island waters, we used general linear mixed models from the package *nlme* (Pinheiro et al. 2014). We specified treatment (colony or reference island) as a categorical covariate and included distance from shoreline (5m, 15m, 25m) and time (peak nesting/fledgling period [mid-May], post nesting/fledgling period [mid-August]) as fixed effects and we included a treatment $\times$ time interaction term when appropriate. In addition, we tested whether the random effect of colony-reference island pairs or the random effect of island alone was warranted. To test for the influence of nesting cormorants on surface water chemistry in near shore water, we used linear models and generalized linear mixed models in the package *lme4* (Bates et al. 2014). Again we specified treatment and time as fixed effects and tested whether inclusion of random effects for colony-reference island pairs or the random effect of island alone improved model fit.

2.2.3 Soil chemistry

We used general linear mixed models and linear models to test for potential effects of nesting cormorants on soil chemistry parameters (%OM, NO₃-N, P, K, Ca, Mg, Zn, S, Na, pH) on the log-scale. We specified treatment as a categorical fixed effects ranging from 1 to 3 (1 = colony island within a nesting colony; 2 = colony island outside a nesting colony, 3 = reference island) and tested whether the random effect of colony-reference island pairs or island alone was warranted, while accounting for spatial correlation and alternative variance structures as needed to improve model fit.

2.2.4 Tree health

To examine the potential effects of nesting cormorants on tree health, we used ordered logistic regression with tree vigor class as the response variable. We included treatment as a

categorical fixed effect ranging from 1 to 3 (1 = colony island within a nesting colony; 2 = colony island outside a nesting colony, 3 = reference island). Tree species also was included as a fixed effect, whereas DBH was included as a continuous fixed effects. We also tested whether the random effects of colony-reference island pairs or island alone was warranted.

3. Results

3.1 Water quality

3.1.1 General coliform and *E. coli* levels

The likelihood ratio test from Poisson distributed models used to explain general coliform levels indicated support for inclusion of a random effect for treatment-reference island pairs ($L = 39.76$, $df = 6$, $P < 0.001$). Inclusion of a treatment $\times$ time interaction was not supported ($L = 1.27$, $df = 1$, $P = 0.26$), nor was the inclusion of treatment alone ($L = 3.55$, $df = 1$, $P = 0.06$). Dropping distance ($L = 16.24$, $df = 1$, $P < 0.001$) or time ($L = 791.95$, $df = 1$, $P < 0.001$) effects did not improve model fit. Overall, general coliform levels decreased only slightly with increasing distance from island shorelines ($\beta = -0.009$, $SE = 0.002$, $P < 0.001$, $CI = -0.013$ to -0.005) but increased substantially from mid-May to mid-August ($\beta = 1.23$, $SE = 0.05$, $P < 0.001$, $CI = 1.14$ to 1.33) (Fig. 2).

E. coli data were overdispersed, likely due to the number of zeros in the data (84% of the data were zeros), thus we used a zero-inflated Poisson model. The likelihood ratio test indicated lack of support for inclusion of the random effect for treatment-reference island pairs ($L = 3.16$, $df = 6$, $P = 0.08$) and for the random effect of island alone ($L = 2.75$, $df = 7$, $P = 0.10$). Inclusion of a treatment $\times$ time interaction was supported ($L = 6.02$, $df = 2$, $P < 0.001$), although treatment alone had no effect ($\beta = -1.17$, $SE = 0.08$, $P = 0.14$, $CI = -2.75$ to 0.40). Distance to shore also had no measurable effect on *E. coli* levels ($\beta = -0.04$, $SE = 0.03$, $P = 0.11$, $CI = -0.10$ to 0.01) but

dropping the effect of distance ($L = 6.95$, $df = 2$, $P = 0.03$) did not improve model fit. Overall, *E. coli* levels decreased between mid-May and mid-August ($\beta = -3.76$, $SE = 0.75$, $P < 0.001$, $CI = -5.23$ to -2.29), although the treatment $\times$ time interaction indicated that *E. coli* levels decreased substantially between mid-May and mid-August at islands with nesting cormorants (treatment islands), whereas *E. coli* levels increased slightly between mid-May and mid-August at reference islands without nesting cormorants ($\beta = 4.22$, $SE = 1.18$, $P < 0.001$, $CI = 1.83$ to 6.61) (Fig. 2).

3.1.2 Water chemistry

The likelihood ratio tests from the linear models used to explain variation in surface water pH level 5m from island shorelines indicated the random effects of treatment-reference island pairs ($L = 0.23$, $df = 6$, $P = 0.63$) and islands alone ($L = 0.30$ $df = 6$, $P = 0.59$) were unwarranted. Inclusion of a treatment $\times$ time interaction was supported ($L = 12.38$, $df = 1$, $P < 0.001$) but time alone had no measurable effect on pH levels ($\beta = -0.004$, $SE = 0.24$, $P = 0.98$, $CI = -0.04$ to 0.99). Treatment, however, did have an influence on pH levels with higher pH measured in nearshore waters around reference islands ($\beta = 0.78$, $SE = 0.22$, $P < 0.001$, $CI = 0.36$ to 1.21) compared to treatment islands, with treatment $\times$ time interaction resulting in lower pH levels in mid-August in nearshore waters around reference islands compared to treatment islands ($\beta = -1.15$, $SE = 0.32$, $P < 0.001$, $CI = -1.77$ to -0.52) (Fig. 2).

The likelihood ratio tests from the linear mixed models used to explain variation in ammonia (NH_3) levels in surface water 5m from island shorelines indicated that the random effect of treatment-reference island pairs ($L = 0.08$, $df = 5$, $P = 0.78$) was unwarranted, however, the random effect of islands alone ($L = 5.54$ $df = 6$, $P = 0.02$) improved model fit. We found no evidence of a treatment $\times$ time interaction ($L = 2.12$ $df = 1$, $P = 0.16$) and dropping treatment improved model fit ($L = 27.34$ $df = 1$, $P = 0.98$). Inclusion of time was merited ($L = 29.11$ $df = 1$,

$P < 0.001$) and overall, ammonia levels decreased between mid-May and mid-August ($\beta = -0.08$ [mg/L], $SE = 0.01$, $P < 0.001$, $CI = -0.10$ to -0.05 [mg/L]) (Fig. 2).

We fit linear mixed models on the log-scale to assess variation in phosphate (PO_4) levels in surface water 5m from island shorelines. The likelihood ratio tests indicated there was no support for inclusion of the random effect of treatment-reference island pairs ($L = 1.05$, $df = 6$, $P = 0.99$) or little support for islands alone ($L = 3.16$, $df = 6$, $P = 0.06$). Inclusion of a treatment $\times$ time interaction also was not supported ($L = 0.66$, $df = 1$, $P = 0.42$) and dropping treatment ($L = 0.64$, $df = 1$, $P = 0.42$) and time ($L = 1.75$, $df = 1$, $P = 0.18$) improved model fit. As such, the null model was the best supported model.

Because only two of the 48 water samples yielded non-zero nitrate (NO_3-N) values, we were unable to assess the effects of treatment and time on nitrate levels (mg/L) in nearshore waters around islands with and without nesting cormorants. Thus, nesting cormorants have no measurable effect on nitrate levels in surface waters near islands that host cormorant nesting colonies.

3.2 Soil chemistry

All data were log-transformed to improve normality of the residuals except pH. With the exception of Na ($L = 3.31$ $df = 5$, $P = 0.06$) and NO_3-N ($L = 2.00$, $df = 5$, $P = 0.16$), likelihood ratio tests indicated that the inclusion of the random effect of island was supported for all other analyses to evaluate the influence of cormorant nesting colonies on soil chemistry parameters. Likelihood ratio tests also indicated the effect of treatment (1 = colony island within a nesting colony; 2 = colony island outside a nesting colony, 3 = reference island) was significant for P ($L = 5.39$, $df = 2$, $P = 0.06$), K ($L = 31.26$, $df = 2$, $P < 0.001$), Mg ($L = 7.52$, $df = 2$, $P = 0.02$) and pH ($L = 5.89$, $df = 2$, $P = 0.05$). Post hoc tests revealed significant differences in P ($P = 0.05$), K

($P = 0.03$) and pH ($P = 0.05$) between soil samples taken from colony islands within a nesting colony (treatment 1) and on reference islands (treatment 3) (Fig. 3). Mg levels were not different within nesting colonies (treatment 1) and on reference islands (treatment 3) although Mg levels were higher on colony islands outside the nesting colony (treatment 2) compared to the other treatments ($P = 0.02$) (Fig. 3).

3.3 Tree health

Although we documented differences in the number of tree species identified during tree surveys across the six study islands (Table 1), tree species was not an important factor in regards to tree vigor class. However, variation in tree vigor class was influenced by a combination of DBH and treatment (1 = colony island within a nesting colony; 2 = colony island outside a nesting colony, 3 = reference island) (Fig. 4a). For a unit increase in DBH the odds of a tree being categorized as vigor class 1 versus vigor class 2, 3, 4 or 5 was multiplied by 0.98 (CI = 0.97 to 1.00) (Fig. 4). In addition, the odds of a tree being classified as vigor class 1 versus vigor class 2, 3, 4 or 5 was 0.51 times greater (CI = 0.28 to 0.93) in treatment 2 compared to treatment 1, whereas the odds of a tree being classified in vigor class 1 versus vigor class 2, 3, 4 or 5 was 0.30 times greater (CI = 0.09 to 1.00) in treatment 3 versus treatment 1. We also found a more even distribution of tree vigor classes inside cormorant nesting colonies than outside nesting colonies or on reference islands (Fig. 4b). Tukey post hoc tests confirmed significant differences in tree vigor class between all pairwise treatment combinations (2 $\square$ 1: $P < 0.001$; 3 $\square$ 1: $P < 0.001$; 3 $\square$ 2: $P = 0.021$) (Fig. 4b), with the greatest difference observed between treatments 1 and 2.

3.4 Cormorant and co-nesting species

During nesting surveys, we documented 308 cormorant nests and 92 great blue heron (*Ardea herodias*) nests on colony islands as well as 16 great blue heron nests on one reference island (Table 2). Cormorant chicks were present in 100% of cormorants nests documented during nesting surveys and > 90% of great blue heron nests contained chicks (Table 2). Of the cormorant nests documented during nesting surveys, 90% occurred in Loblolly Pine (*Pinus taeda*), while only 6% occurred Yellow Poplar (*Liriodendron tulipifera*), 3% in Bald Cypress (*Taxodium distichum*) and 1% in Sycamore (*Plantus occidentalis*). The status (i.e., live/dead) of cormorant nesting trees varied across islands with approximately 50% of cormorant nesting trees were identified as dead (Table 3). Although not included in nest surveys, during tree surveys, additional cormorant nests were observed in Sweetgum (*Liquidambar styraciflua*; n = 6) and Silver Maple (*Acer saccharinum*; n = 1) on each island (section 2.2.3 above). Of the great blue heron nests documented during nesting surveys, 67% occurred in Loblolly Pine, 32% in Yellow Poplar, and < 1% in Sycamore, with approximately 50% of great blue heron nesting trees identified as dead (Table 3). Again, although not included in nest surveys, additional great blue heron nests were observed in Sweetgum (n = 1), Silver Maple (n = 1), Persimmon (*Diospyros virginiana*; n = 1) and Willow Oak (*Quercus phellos*; n = 1) during tree surveys (section 2.2.3 above).

4. Discussion

We found significant differences in general coliform and *E. coli* levels, with general coliform levels increasing substantially from mid-May to mid-August and to a lesser degree with proximity to island shorelines regardless of presence of cormorant colonies. *E. coli* levels had the opposite trend compared to general coliforms, decreasing between mid-May and mid-August with a greater declines occurring on islands with nesting cormorants compared to reference

islands. Our results also indicated few differences in water chemistry between islands with and without cormorant colonies. Treatment, however, did have an influence on pH levels with higher pH measured in near shore waters around reference islands compared to treatment islands, and lower pH levels in mid-August in near shore waters around reference islands compared to treatment islands. We also found significant differences in pH, P and K between soil samples taken from colony islands within a nesting colony and on reference islands. Differences in Mg were only significant between samples taken from colony islands within a nesting colony and colony islands outside a nesting colony (Fig. 3).

Loblolly pine and maple spp. were the most common tree types on most islands (Table 1) although the majority of cormorants (90%) nested in loblolly pine, suggesting a preference for this species. Similar to soil chemistry data we found tree vigor class was influenced by the presence of cormorant colonies and interestingly tree DBH. For each unit increase in DBH, the odds of a tree being categorized as vigor class 1 versus a higher vigor class increased by a factor of 0.98, which suggests that cormorants are selecting larger trees. In addition, the odds of a tree being classified as vigor class 1 versus higher vigor classes was 0.51 times greater outside of a colony versus within a colony and 0.30 times greater to have a vigor class 1 on reference islands versus colony islands, which suggests that cormorants are affecting tree health as indexed by vigor class regardless of differences among tree species.

Despite some spatial and temporal difference in water parameters, the patterns we observed suggested that water quality and chemistry of near shore waters surrounding cormorant colonies may have been influenced to a greater degree by factors other than the presence of cormorant colonies. For example, if cormorants were a significant contributor to *E. coli* levels in near shore waters, we expected to see an increase in *E. coli* levels from mid-May to mid-August,

yet we observed a slight decrease in *E. coli* over this timeframe. However, observed decrease in *E. coli* levels from mid-May to mid-August may simply reflect a reduction in *E. coli* levels following a rapid decrease in cormorant numbers during the post fledging period. As expected, general coliform levels increased from mid-May to mid-August in near shore waters, although this may have been a consequence of increasing water temperature from mid-May to mid-August. We also found no measurable differences in nitrate levels in surface waters around colony and reference islands. Further, despite slightly lower pH levels in surface waters surrounding colony islands relative to references islands, pH levels were still alkaline, thus showing no evidence of water acidification. Although cormorant colonies may be a source of bacterial and chemical pollution on islands and in near shore waters (Klimaszyk 2012, Stewart et al. 2015), cumulatively our data suggest nesting cormorants had little effect on water quality and chemistry in Guntersville, Lake.

Our results provide an interesting contrast to other studies that have found that waterbirds such as gulls (*Larus* sp.), Canada geese (*Branta canadensis*) and great cormorants occurring at high densities can be drivers of changes in water quality. For instance, several studies have demonstrated that waterbirds can be a significant source of bacterial contamination in inland water bodies (Hussong et al. 1979, Benton et al. 1983, Lévesque et al. 2000, Kirschner et al. 2004, Meerburg et al. 2011, Klimaszyk and Rzymiski 2013). Additionally, guano inputs to surface waters can accelerate the process of eutrophication (Niewolak 1999, Nakamura et al. 2010, Klimaszyk and Rzymiski 2013) and lead to hyper-eutrophic and polluted systems (Stewart et al. 2015). However, factors including the density and diversity of waterbirds, their diet and feeding habits as well as the size of the waterbody all influence the level of bacterial contamination in the water (Hussong et al. 1979). For instance, the highest density of cormorant

389 nests recorded on any of our study islands (South Sauty Island [SSS]) was 97.1 cormorants/ha²,
390 whereas cormorant nest densities during the same sampling year (i.e., 2010) in the Great Lakes
391 region on some islands was substantially higher (e.g., > 500 cormorants/ha²), although islands in
392 the Great Lakes also host large populations of other waterbird species (Ayers et al. 2015, Stewart
393 et al. 2015). In addition, the movement of water, variable nutrient inputs and the suspension of
394 inorganic and organic matter in the water column also may influence the density of general
395 coliforms and *E. coli* in near shore waters thereby affecting water quality (Gwiazda et al. 2014).
396 Thus, although we did not find evidence that cormorants were having a direct effect on water
397 quality or chemistry, cormorants may have indirect effects due to interactions among nutrient
398 flow, aquatic plant growth and slight differences in pH associated with cormorant colonies.

399 Further, we found that soils within cormorant nesting colonies had lower pH and higher
400 accumulations of P and K compared to soils on reference islands. In addition, soils on islands
401 with cormorant colonies but outside the spatial extent of the colonies exhibited intermediate pH
402 levels and intermediate concentrations of P and K, thus showing a stepwise pattern of pH levels
403 and P and K concentrations across the three natural treatments (Fig 3). Several studies have
404 reported significant changes in soil chemistry associated with colonial nesting birds (e.g., Mun
405 1997, Ligeza and Smal 2003, Hobara et al. 2005, Ayers et al. 2015, Borkowska et al. 2015).
406 Specifically, soils tend to be more acidic within nesting colonies compared to reference sites,
407 suggesting that deposition of avian guano results in soil acidification (Ishida 1996, Cuthbert et al.
408 2002, Breuning-Madsen et al. 2010, Ayers et al. 2015). Soil acidification can result in some
409 nutrients (e.g., Al) becoming noxious and potentially toxic to plants depending on the nutrients'
410 reactivity at low pH, whereas other nutrients (e.g., Ca, K, Mg) may be depleted or become
411 unavailable for uptake by plants in acidic soils (Ashman and Puri 2013). Further, high

concentrations of P can be toxic and limit plant uptake of some macronutrients (e.g., iron, zinc), thereby effecting plant growth (Pitt and Provin 2008), whereas high concentrations of K are neither noxious nor toxic but may influence osmotic pressure caused by the number of K ions in the soil thereby affecting synergistic action with other soil nutrients (Ligeza and Smal 2003). In addition, avian guano, regurgitation, carcasses and feather deposits and subsequent changes in soil chemistry also can lead to loss of native vegetation, diminished plant diversity and facilitation of nutrient-tolerant exotic species (Ishida 1997, Mun 1997, Rippey 2002, Weseloh et al. 2002, Ayers et al. 2015). However, the negative effects of cormorants and other avian species on biotic and abiotic environments are conditioned by the number of waterbirds contributing to the system, which, compared to northern ecosystem, cormorant inputs in Lake Guntersville is relatively limited.

Although the general pattern of differences in soil chemistry parameters across the three treatments in our study area were similar to what has been documented in northern ecosystems, our results were less dramatic likely because our study system has substantially fewer nesting cormorants. In northern ecosystems the impacts of cormorants and other colonial nesting birds on soil chemistry includes significant differences in additional nutrients (e.g., $\text{NO}_3\text{-N}$, Zn) that extends well-beyond the spatial extent of the nesting colony, thereby impacting adjacent soils and vegetation (e.g., Ayers et al. 2015). Greater spatial extent of the impact of colonial nesting birds on soils in northern ecosystems may be a due to multiple factors. For example, in northern ecosystems the additive effects of high numbers of co-nesting species (e.g., gulls [*Larus* spp.], American White Pelicans [*Pelecanus erythrorhynchos*]) that only nest on the ground may result in more direct guano inputs to the soil (Ayers et al. 2015), whereas in our southeastern study area, we observed high numbers of cormorants but relatively few co-nesting species and no

ground nesting species. Moreover, the density of nesting cormorants in northern systems is substantially greater than what we observed in our study system (see Ayers et al. 2015, Stewart et al. 2015).

Northern and southeastern ecosystems also have different precipitation regimes, microsite characteristics, soil properties (e.g., texture, substrate [sand, clay loam]), different plant assemblages and subsequent differences in nutrient uptake by different plant species, all of which could affect soil chemistry. Further, differences in spatio-temporal patterns of cormorant colonization, subsequent population increases as well as differences in nest densities may account for some of the observed differences in the magnitude of soil chemistry parameters among samples within colonies, adjacent to colonies and on reference islands in northern and southeastern ecosystems.

Our result also indicated that tree vigor class was negatively affected by nesting cormorant colonies, although impacts were mainly to loblolly pine due to an apparent preference of this tree species by cormorants. The greatest impacts were found within the spatial extent of the nesting colonies relative to trees on reference islands, which is a similar stepwise pattern to what we observe in the soil data across the three treatments. Previous studies have reported substantial tree damage and tree cover loss on islands with nesting cormorant colonies (Bédard 1988, Chapdelaine and Bédard 1995, Rippey 2002, Boutin et al. 2011). Cormorants can cause physical damage to trees by removing foliage for nesting material and by breaking branches (Lemmon et al. 1994, Aderman and Hill 1995, Hobara et al. 2001, Hebert et al. 2005). In addition, as trees die, branches become less stable for nesting and cormorants often abandon unstable trees in favor of nearby sturdier trees with remaining foliage (Rippey 2002, Hebert et al. 2005). As such, it is not surprising that tree damage caused by cormorants often expands across

islands habitats as dead and dying trees are abandoned and new trees are selected as nesting sites (Hebert et al. 2005). Interestingly, we also found that trees with smaller DBH appeared to be more affected by nesting cormorants. This result may be a consequence of multiple interacting factors. For instance, large increases in tree debris (i.e., litterfall) resultant from cormorant damage has been reported (Hobara et al. 2001) and this debris may negatively affect smaller trees via physical damage as well as changing microsite conditions. Also, elevated ammonium levels in the soil under cormorant colonies due to guano can result in toxicity to many plants, particularly in areas where soil pH is low (Britto and Kronzucker 2002). Thus, cormorant guano may have a more immediate impact on understory vegetation that includes smaller trees within the colony compared to larger, well-established trees. However, the impacts of nesting cormorants on large healthy trees in our study area was apparent, although the effects may be more evident in subsequent years as trees die and are subsequently abandoned by nesting cormorants in favor of healthier trees with remaining foliage (Rippey 2002, Hebert et al. 2005).

5. Management and Research Implications

Our study indicates that cormorants are having an effect on forested island habitats in Lake Guntersville, AL, USA, particularly changes in soil characteristics and damage to tree health. Due to a possible preference for loblolly pine for nesting, if restoration of island trees is a management goal, use of non-pine species may reduce attractiveness of these islands to cormorants. Overall, the biotic and abiotic impacts of cormorants nesting in this watershed appears to be less dramatic than findings in northern ecosystems. For example, we found little evidence that cormorants are impacting water quality and chemistry at current cormorant nesting densities. However, large numbers of waterfowl do not always affect water quality (Brierley et al. 1975). As such, it is likely that the impacts of nesting cormorants on water quality/chemistry

and forest structure and composition may be uncoupled. Despite quantifiable impacts, colonization and subsequent succession of forested islands following waterbird colonization is a natural process (Wires and Cuthbert 2006). Stewart et al. (2015) used paleoecological techniques to evaluate spatial and temporal patterns of colonial nesting waterbirds use of islands over relatively long time scales (100-150 years) in the Great Lakes region, providing greater historical ecological context for current patterns of cormorant and other waterbird use of island habitats. Thus, any consideration of management action to reduce unwanted abiotic and biotic impacts that nesting cormorants are having on forested island habitats for socio-economic reasons should be considered within a framework that allows for natural ecological processes, including vegetation succession following cormorant colonization.

6. Acknowledgements

This research was funded by the U.S. Department of Agriculture, Wildlife Services, National Wildlife Research Center. We thank Robert Ritchey and Jerry Feist of U.S. Department of Agriculture, Wildlife Services and Paul Fioranelli with the National Wildlife Research Center for logistical and field support. We thank the Keith Crouse of Mississippi State University Extension Service Soil Testing Laboratory for conducting soil nutrient analyses and Tara Conkling for advice regarding statistical analysis.

7. References

- Aderman, A. R., and E. P. Hill. 1995. Locations and numbers of Double-crested Cormorants using winter roosts in the Delta Region of Mississippi. *Colonial Waterbirds*:143-151.
- Ashman, M., and G. Puri. 2013. *Essential soil science: a clear and concise introduction to soil science*. John Wiley & Sons.
- Ayers, C. R., K. C. Hanson-Dorr, S. O'Dell, C. D. Lovell, M. L. Jones, J. R. Suckow, and B. S.

504 Dorr. 2015. Impacts of colonial waterbirds on vegetation and potential restoration of
505 island habitats. *Restoration Ecology* 23:252-260.

506 Barras, S. C. 2004. Double-Crested Comorants in Alabama. USDA National Wildlife Research
507 Center-Staff Publications:77.

508 Bates, D., M. Maechler, B. Bolker, and S. Walker. 2014. lme4: Linear mixed-effects models
509 using Eigen and S4. R package version 1.1-7. This is computer program (R package). The
510 URL of the package is: <http://CRAN.R-project.org/package=lme4>.

511 Bédard, H. 1988. Les Montagnais et la réserve de Betsiamites, 1850-1900. Institut québécois de
512 recherche sur la culture; Ville Saint-Laurent, Québec: Distribution, Diffusion Prologue.

513 Bédard, J., A. Nadeau, and M. Lepage. 1995. Double-crested cormorant culling in the St.
514 Lawrence River estuary. *Colonial Waterbirds*:78-85.

515 Benton, C., F. Khan, P. Monaghan, W. Richards, and C. Shedden. 1983. The contamination of a
516 major water supply by gulls (*Larus* sp.): a study of the problem and remedial action
517 taken. *Water Research* 17:789-798.

518 Borkowska, L., E. Królak, Z. Kasprzykowski, and P. Kaczorowski. 2015. The influence of
519 *Corvus frugilegus* nesting on soil parameters and plant composition in poor and fertile
520 habitats. *Landscape and Ecological Engineering* 11:161-167.

521 Boutin, C., T. Dobbie, D. Carpenter, and C. E. Hebert. 2011. Effects of Double-Crested
522 Cormorants (*Phalacrocorax auritus* Less.) on Island Vegetation, Seedbank, and Soil
523 Chemistry: Evaluating Island Restoration Potential. *Restoration Ecology* 19:720-727.

524 Breuning-Madsen, H., C. Ehlers-Koch, J. Gregersen, and C. L. Løjtnant. 2010. Influence of
525 perennial colonies of piscivorous birds on soil nutrient contents in a temperate humid
526 climate. *Geografisk Tidsskrift-Danish Journal of Geography* 110:25-35.

527 Brierley, J. A., D. K. Brandvold, and C. J. Popp. 1975. Waterfowl refuge effect on water quality:
 528 I. Bacterial populations. *Journal (Water Pollution Control Federation)*:1892-1900.
 529 Britto, D. T., and H. J. Kronzucker. 2002. NH₄⁺ toxicity in higher plants: a critical review.
 530 *Journal of Plant Physiology* 159:567-584.
 531 Chapdelaine, G., and J. Bédard. 1995. Recent changes in the abundance and distribution of the
 532 Double-crested Cormorant in the St. Lawrence River, Estuary and Gulf, Quebec, 1978-
 533 1990. *Colonial Waterbirds*:70-77.
 534 Cuthbert, F. J., L. R. Wires, and J. E. McKearnan. 2002. Potential impacts of nesting double-
 535 crested cormorants on great blue herons and black-crowned night-herons in the US Great
 536 Lakes region. *Journal of Great Lakes Research* 28:145-154.
 537 Dorr, B. S., L. W. Burger, S. C. Barras, and K. C. Godwin. 2012. Double-crested cormorant
 538 distribution on catfish aquaculture in the Yazoo River Basin of Mississippi. *Wildlife*
 539 *Society Bulletin* 36:70-77.
 540 Gwiazda, R., A. Woźnica, B. Łozowski, M. Kostecki, and A. Flis. 2014. Impact of waterbirds on
 541 chemical and biological features of water and sediments of a large, shallow dam
 542 reservoir. *Oceanological and hydrobiological studies* 43:418-426.
 543 Hanson, K. C., T. L. D. Vault, and S. J. Dinsmore. 2010. Increased abundance and first breeding
 544 record of the Neotropic Cormorant (*Phalacrocorax brasilianus*) on the Alluvial Plain of
 545 Mississippi. *Southeastern naturalist* 9:385-394.
 546 Hatch, J. J., and D. V. Weseloh. 1999. Double-crested Cormorant(*Phalacrocorax auritus*). *The*
 547 *Birds of North America*:36.
 548 Hebert, C. E., J. Duffe, D. C. Weseloh, E. T. Senese, and G. D. Haffner. 2005. Unique island

549 habitats may be threatened by double-crested cormorants. *Journal of Wildlife*
550 *Management* 69:68-76.

551 Hobara, S., K. Koba, T. Osono, N. Tokuchi, A. Ishida, and K. Kameda. 2005. Nitrogen and
552 phosphorus enrichment and balance in forests colonized by cormorants: implications of
553 the influence of soil adsorption. *Plant and Soil* 268:89-101.

554 Hobara, S., T. Osono, K. Koba, N. Tokuchi, S. Fujiwara, and K. Kameda. 2001. Forest floor
555 quality and N transformations in a temperate forest affected by avian-derived N
556 deposition. *Water, Air, and Soil Pollution* 130:679-684.

557 Hussong, D., J. Damare, R. J. Limpert, W. Sladen, R. Weiner, and R. Colwell. 1979. Microbial
558 impact of Canada geese (*Branta canadensis*) and whistling swans (*Cygnus columbianus*
559 *columbianus*) on aquatic ecosystems. *Applied and environmental microbiology* 37:14-20.

560 Ishida, A. 1996. Effects of the common cormorant, *Phalacrocorax carbo*, on evergreen forests in
561 two nest sites at Lake Biwa, Japan. *Ecological Research* 11:193-200.

562 _____. 1997. Seed germination and seedling survival in a colony of the common cormorant,
563 *Phalacrocorax carbo*. *Ecological Research* 12:249-256.

564 Jarvie, S., H. Blokpoel, and T. Chipperfield. A geographic information system to monitor nest
565 distributions of Double-crested Cormorants and Blackcrowned Night-herons at shared
566 colony sites near Toronto, Canada. 1999.

567 Kirschner, A. K., T. C. Zechmeister, G. G. Kavka, C. Beiwl, A. Herzig, R. L. Mach, and A. H.
568 Farnleitner. 2004. Integral strategy for evaluation of fecal indicator performance in bird-
569 influenced saline inland waters. *Applied and environmental microbiology* 70:7396-7403.

570 Klimaszuk, P. 2012. May a cormorant colony be a source of coliform and chemical pollution in a
571 lake? *Oceanological and hydrobiological studies* 41:67-73.

572 Klimaszyk, P., and P. Rzymiski. 2013. Impact of cormorant (*Phalacrocorax carbo sinensis* L.)
573 colonies on microbial pollution in lakes. *Limnological Review* 13:139-145.

574 Lemmon, C., G. Bugbee, and G. Stephens. 1994. Tree damage by nesting double-crested
575 cormorants in Connecticut. *Connecticut Warbler* 14:27-30.

576 Lévesque, B. t., P. Brousseau, F. Bernier, É. Dewailly, and J. Joly. 2000. Study of the bacterial
577 content of ring-billed gull droppings in relation to recreational water quality. *Water*
578 *Research* 34:1089-1096.

579 Ligeza, S., and H. Smal. 2003. Accumulation of nutrients in soils affected by perennial colonies
580 of piscivorous birds with reference to biogeochemical cycles of elements. *Chemosphere*
581 52:595-602.

582 Lu, J., J. W. Santo Domingo, R. Lamendella, T. Edge, and S. Hill. 2008. Phylogenetic diversity
583 and molecular detection of bacteria in gull feces. *Applied and environmental*
584 *microbiology* 74:3969-3976.

585 Makino, S.-i., H. Kobori, H. Asakura, M. Watarai, T. Shirahata, T. Ikeda, K. Takeshi, and T.
586 Tsukamoto. 2000. Detection and characterization of Shiga toxin-producing *Escherichia*
587 *coli* from seagulls. *Epidemiology and infection* 125:55-61.

588 McLellan, S. L. 2004. Genetic diversity of *Escherichia coli* isolated from urban rivers and beach
589 water. *Applied and environmental microbiology* 70:4658-4665.

590 Meerburg, B. G., M. G. Koene, and D. Kleijn. 2011. *Escherichia coli* concentrations in feces of
591 geese, coots, and gulls residing on recreational water in The Netherlands. *Vector-Borne*
592 *and Zoonotic Diseases* 11:601-603.

593 Mun, H. 1997. Effects of colony nesting of *Adrea cinerea* and *Egretta alba modesta* on soil
594 properties and herb layer composition in a *Pinus densiflora* forest. *Plant and Soil* 197:55-

595 59.

596 Nakamura, M., T. Yabe, Y. Ishii, K. Kamiya, and M. Aizaki. 2010. Seasonal changes of shallow
597 aquatic ecosystems in a bird sanctuary pond. *Journal of Water and Environment*
598 *Technology* 8:393-401.

599 Niewolak, S. 1999. Evaluation of Pollution and the Sanitary-Bacteriological State of Lake
600 Wigry, Poland. *Polish Journal of Environmental Studies* 8:89-100.

601 Pinheiro, J., D. Bates, S. DebRoy, and D. Sarkar. 2014. R Core Team (2014). nlme: linear and
602 nonlinear mixed effects models. R package version 3.1–117. URL: [http://cran.r-project.](http://cran.r-project.org/web/packages/nlme/index.html)
603 [org/web/packages/nlme/index.html](http://cran.r-project.org/web/packages/nlme/index.html).

604 Pitt, J. L., and T. Provin. 2008. Phosphorus--Too Much and Plants May Suffer. Texas FARMER
605 Collection.

606 R Development Core Team. 2014. R: a language and environment for statistical computing.
607 Vienna, Austria: R Foundation for Statistical Computing. Retrieved from: [http://www.R-](http://www.R-project.org)
608 [project.org](http://www.R-project.org)

609 Reinhold, D. S., A. J. Mueller, and G. Ellis. 1998. Observations of nesting Double-crested
610 Cormorants in the delta region of Mississippi. *Colonial Waterbirds* 21:450-451.

611 Rippey, E. 2002. 10. Increasing numbers of pied cormorants breeding on the islands off Perth,
612 Western Australia and consequences for the vegetation. *Sort* 50:500.

613 Shieldcastle, M. C., and L. Martin. 1999. Colonial Waterbird Nesting on West Sister Island
614 National Wildlife Refuge and the Arrival of. *Technical Bulletin*:115.

615 Standridge, J. H., J. J. Delfino, L. B. Kleppe, and R. Butler. 1979. Effect of waterfowl (*Anas*
616 *platyrhynchos*) on indicator bacteria populations in a recreational lake Madison,
617 Wisconsin. *Applied and environmental microbiology* 38:547-550.

618 Stewart, E. M., Michelutti, N., Shenstone-Harris, S., Grooms, C., Weseloh, C., Kimpe, L. E.,
 619 Blais, J. M., and P. J. Smol. 2015. Tracking the History and Ecological Changes of
 620 Rising Double-Crested Cormorant Populations Using Pond Sediments from Islands in
 621 Eastern Lake Ontario. PloS One 10:e0134167.

622 Taylor, J. D., and B. S. Dorr. 2003. Double-crested cormorant impacts to commercial and natural
 623 resources. USDA National Wildlife Research Center-Staff Publications:278.

624 United States Department of Agriculture Forest Service (USFS). 2007. Forest inventory and
 625 analysis national core field guide volume I: Field data collection procedures for phase 2
 626 plots version 4.0. Available online at [http://fia.fs.fed.us/library/field-guides-methods-](http://fia.fs.fed.us/library/field-guides-methods-proc/docs/2006/core_ver_3-0_10_2005.pdf)
 627 [proc/docs/2006/core_ver_3-0_10_2005.pdf](http://fia.fs.fed.us/library/field-guides-methods-proc/docs/2006/core_ver_3-0_10_2005.pdf). Last accessed 28, December 2010.

628 United States Environmental Protection Agency. 1986. Ambient water quality criteria for
 629 bacteria. USEPA, Washington, DC. Available online at [http://www.epa.gov/waterscience](http://www.epa.gov/waterscience/beaches/files/1986crit.pdf)
 630 [/beaches/files/1986crit.pdf](http://www.epa.gov/waterscience/beaches/files/1986crit.pdf). Last accessed 30 December, 2010.

631 United States Environmental Protection Agency. 1999. Action plan for beaches and recreational
 632 waters. ESEPA, Washington, DC. Available online at
 633 <http://www.epa.gov/ord/htm/documents/600r98079.pdf>. Last accessed 30 December,
 634 2010.

635 Vail, J., R. Morgan, C. Merino, F. Gonzales, R. Miller, and J. Ram. 2003. Enumeration of
 636 Waterborne with Petrifilm Plates. Journal of environmental quality 32:368-373.

637 Weseloh, D. C., C. Pekarik, T. Havelka, G. Barrett, and J. Reid. 2002. Population trends and
 638 colony locations of double-crested cormorants in the Canadian Great Lakes and
 639 immediately adjacent areas, 1990–2000: a manager's guide. Journal of Great Lakes
 640 Research 28:125-144.

641 Wires, L. R., and F. J. Cuthbert. 2006. Historic populations of the double-crested cormorant
642 (Phalacrocorax auritus): implications for conservation and management in the 21st
643 century. Waterbirds 29:9-37.

644 Zuur, A., E. N. Ieno, N. Walker, A. A. Saveliev, and G. M. Smith. 2009. Mixed effects models
645 and extensions in ecology with R. Springer Science & Business Media.

646 Zuur, A. F., A. A. Saveliev, and E. N. Ieno. 2012. Zero inflated models and generalized linear
647 mixed models with R. Highland Statistics Limited Newburgh.

648

Table 1. Summary of tree species documented at random tree survey plots on three islands with nesting cormorants (CON, SSS, NS) and reference islands (SEC, WNS, NSS) that lack nesting cormorant on Lake Guntersville, AL, U.S.A., represented by the percentage of the total number of trees recorded in all survey plots per island, rounded to the nearest whole number.

COMMON NAME	Scientific name	COLONY ISLANDS			REFERENCE ISLANDS		
		CON	SSS	NS	SEC	WNS	NSS
AMERICAN ELM	<i>Ulmus americana</i>		4	1		5	1
BALD CYPRESS	<i>Taxodium distichum</i>		4				
BLACK CHERRY	<i>Prunus serotina</i>	8				1	9
BLACK LOCUST	<i>Robinia pseudoacacia</i>	1	1		3		6
BOX ELDER	<i>Acer negundo</i>		3				
CAROLINA BUCKTHORN	<i>Frangula caroliniana</i>	4					
CHINESE PRIVET	<i>Ligustrum sinense</i>		4				
DEVIL'S WALKING STICK	<i>Aralia spinosa</i>	1					
EASTERN RED CEDAR	<i>Juniperus virginiana</i>			1			33
FLOWERING DOGWOOD	<i>Cornus florida</i>	2		4		1	5
HACKBERRY	<i>Celtis occidentalis</i>	2	6		40		8
IRONWOOD	<i>Ostrya virginiana</i>		1			3	
LOBLOLLY PINE	<i>Pinus Taeda</i>	60	29		48		8
MAPLE SPP	<i>Acer spp</i>	1		3		1	
OAK SPP	<i>Quercus spp</i>	1				1	
PERSIMMON	<i>Diospyros virginiana</i>			3	5	12	4
PINE SPP	<i>Pinus spp</i>	1					
RED MAPLE	<i>Acer rubrum</i>	3	4	19		20	8
SASSAFRAS	<i>Sassafras albidum</i>				3		
SHORTLEAF PINE	<i>Pinus echinata</i>						1
SILVER MAPLE	<i>Acer saccharinum</i>	1	22	47		23	1
SOUTHERN RED OAK	<i>Quercus falcata</i>	1				1	1
SWAMP CHESTNUT OAK	<i>Quercus michauxii</i>						1
SWEETGUM	<i>Liquidambar styraciflua</i>	2	4			25	4
SYCAMORE	<i>Plantus occidentalis</i>		1				3
WATER OAK	<i>Quercus nigra</i>	1		6	3	1	4
WHITE OAK	<i>Quercus alba</i>			3			
WILLOW OAK	<i>Quercus phellos</i>			6			
YELLOW POPLAR	<i>Liriodendron tulipifera</i>	7	2	6		2	6
UNKNOWN GENUS		5	10	1			9

Table 2. Study islands on Guntersville Lake, Alabama, U.S.A.

Colony island	Island size (ha)	Cormorant Nests (% with chicks)	Other nests (% with chicks)	Reference island	Island size (ha)	Cormorants nests (% with chicks)	Other nests ^a (% with chicks)
CON	4.69	101 (100)	26 (100)	SEC	1.36	0 (0)	16 (100)
SSS	2.80	272 (100)	22 (91)	NSS	2.21	0 (0)	0 (0)
NS	0.76	7 (100)	28 (96)	WNS	2.20	0 (0)	0 (0)

^a Co-nesting species were exclusively Great blue heron (*Ardea herodias*) with the exception of a single Osprey (*Pandion haliaetus*) nest on South Sauty Island with no chick(s) observed.

658 **Table 3.** Percentage of Double-crested cormorant and great blue heron nesting trees identified as
659 alive or dead during nesting surveys conducted on Guntersville Lake, Alabama, U.S.A.

Island	Cormorant nesting trees			Great blue heron nesting trees		
	N	% alive	% dead	N	% alive	% dead
CON (colony)	34	69	31	20	50	50
SSS (colony)	49	100	0	18	92	8
NS (colony)	4	35	65	12	0	100
SEC (reference)	0	N/A	N/A	9	100	0

660

Figure 1. Study island locations, names and abbreviations (in parentheses) and whether they are reference or colony islands, sampled from mid-May to mid-August, on Guntersville Lake, Alabama, USA. Inset map of the United State of America with a star representing the approximate location of the study area.

666 **Figure 2.** Panel A, median general coliform levels (CFU) relative to distance from shore (m) and compared between mid-May and
667 mid-August, and median *E. coli* levels (CFU) compared between colony islands (nesting cormorants present) and reference
668 islands (nesting cormorants absent) from mid-May to mid-August. Panel B, median pH levels compared between colony
669 islands and references islands from mid-May to mid-August and differences in Ammonia (NH₃) from mid-May to mid-August.
670 For both Panel A and B, only significant results are displayed.
671

672 **Figure 3.** Median differences in soil chemistry values among treatments: (A) phosphorous, (B) Potassium, (C) Magnesium and (D)
673 pH. Differences between treatments are represented by different letters above the quantile values. Treatment is based on
674 location relative to cormorant nesting colonies ranging from 1 to 3: 1 = colony island within a nesting colony, 2 = colony
675 island outside a nesting colony and 3 = reference island. Only significant results are displayed.

Figure 4. Panel A: Tree vigor classes across colony and reference islands relative to treatment; represented as the percentage of total trees surveyed in each treatment by island. Panel B: Tree vigor classes across five categories of tree diameter at breast height (DBH) in centimeters (< 25cm, >25<50cm, >50<75cm, >75<100cm, >100cm) in each treatment; represented as the percentage of total trees surveyed of each size class by treatment. Tree vigor classes range from 1=healthy to 5 = standing dead (see methods for full description). Treatment is based on location relative to cormorant nesting colonies ranging from 1 to 3: 1 = colony island within a nesting colony, 2 = colony island outside a nesting colony and 3 = reference island.