Forest-landscape structure mediates effects of a spruce bark beetle (*Dendroctonus rufipennis*) outbreak on subsequent likelihood of burning in Alaskan boreal forest

Winslow D. Hansen^{1,5*}, F. Stuart Chapin III², Helen T. Naughton³, T. Scott Rupp¹, & David Verbyla⁴

1. Scenarios Network for Alaska and Arctic Planning, University of Alaska Fairbanks, Fairbanks, AK, 99775-6100, USA
2. Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, AK 99775-6100, USA
3. Department of Economics, University of Montana, Missoula, MT 59812-5472, USA
4. Department of Forest Sciences, University of Alaska Fairbanks, Fairbanks, AK 99775-6100, USA
5. Current address, Landscape and Ecosystem Ecology Lab, Department of Zoology, University of Wisconsin, Madison, Madison, WI 53706-1313

*corresponding author: whansen3@wisc.edu

Abstract

Characterizing how variation in forest landscape structure shapes patterns of natural disturbances and mediates interactions between multiple disturbances is critical for anticipating ecological consequences of climate change in high-latitude forest ecosystems. During the 1990s, a massive spruce bark beetle (*Dendroctonus rufipennis*) outbreak took place in boreal spruce forest on the Kenai Peninsula, Alaska allowing us to ask 1) *How did the extent and duration of bark beetle outbreak differ between a homogenous landscape dominated by white spruce (*Picea glauca*), and a landscape in which white spruce and black spruce (*Picea mariana*) were intermixed?* (2) *How has the occurrence and duration of bark beetle outbreak influenced the likelihood of subsequent burning in these two landscapes?* Forest landscape structure had a substantial effect on disturbance patterns and interactions between disturbances in this study. The spruce bark beetle outbreak was smaller in extent and duration where white spruce, the beetle's primary host tree, was intermixed with more beetle-resistant black spruce. However, likelihood of subsequent burning increased where outbreak did occur. Surface fuel loads increased substantially in this landscape following the outbreak, potentially increasing the flammability of white spruce where they once served as fire breaks. In contrast, the outbreak was larger and lasted longer in the landscape with homogeneous stands of white spruce, but was not related to likelihood of subsequent burning, which is consistent with the fire history. Our results suggest that bark beetle outbreaks may have different effects on subsequent patterns of burning than in other systems, such as the Rocky Mountains. These results could inform more effective and targeted management strategies to ameliorate fire risk in beetle-killed stands of Alaska and may help us anticipate the dynamics and consequences of future boreal bark beetle outbreaks as climate warms at high latitudes.

44 **Keywords** conifer forests, climate change, landscape heterogeneity, linked disturbance
45 interactions, natural disturbance, North American boreal forest
46

Introduction

Evaluating how variation in forest-landscape structure shapes natural disturbance patterns and mediates interactions between multiple disturbances is critical for understanding ecological consequences of climate change (Turner 2010, Johnstone et al. 2011). Warming and drying are causing pronounced increases in the frequency, size, and severity of natural disturbances (Millar and Stephenson 2015, Trumbore et al. 2015). In western North America, annual area burned by wildfire is now nearly 6.5 times the 1970-1986 average (Westerling et al. 2006), and bark beetle outbreaks have become substantially more prevalent in recent decades (Bentz et al. 2010). Concern persists that bark-beetle-driven tree mortality could further increase the risk of subsequent fire frequency, size, and severity (Hicke et al. 2012), a concept known as linked disturbance interactions (Simard et al. 2011). However, landscape structure may mediate climate-change effects on disturbances (Turner and Romme 1994, Duffy et al. 2007). Variation in tree-species distributions across landscapes can influence patterns of burning (Schoennagel et al. 2004, Kelly et al. 2013) and determines forest vulnerability to bark beetle outbreak (Hart et al. 2015a, Temperli et al. 2015). Accounting for the role of forest-landscape structure could yield powerful insights into changing disturbance regimes in the 21st century and the nature and magnitude of their interactions. This could help us anticipate when and where forests may be vulnerable to bark beetle outbreaks and whether such outbreaks may influence subsequent patterns of burning.

Few places on Earth are experiencing as rapid climate change as the Alaskan boreal forest, warming at twice the rate of the global average during the 20th century (Wolken et al. 2011). Annual area burned in boreal Canada and Alaska is expected to increase 74% to 118% by 2100 (Balshi et al. 2008, Flannigan et al. 2005). In white spruce (*Picea glauca*) stands of the

western Kenai Peninsula in south-central Alaska, recent climate warming also led to a massive spruce bark beetle (SBB) (*Dendroctonus rufipennis*) outbreak that extended over almost 1.2 million ha during the 1990s (Berg et al. 2006). An estimated 30 million trees were killed per year between 1990 and 1996, when the outbreak peaked (Werner et al. 2006). The distribution of tree species varied markedly in SBB-affected landscapes. This variation may have mediated outbreak dynamics (Seidl et al. 2015, Temperli et al. 2015) and subsequent ecological consequences. White spruce, the beetle's primary host tree in Alaska, dominated the southern portion of the outbreak extent, forming homogenous stands of susceptible mature forest (Berg and Anderson 2006). In the northern portion of the outbreak, white spruce trees were intermixed with black spruce (*Picea mariana*) (DeVolder 1999), which are not vulnerable to SBB (Werner et al. 2006).

The Kenai Peninsula is one of the most densely populated areas in Alaska, and following the outbreak, there has been substantial concern about increases in the subsequent risk of wildfire. A number of wildfires have burned on the Kenai Peninsula since the outbreak. Warming trends suggest that future outbreaks and increased interaction with fire may occur in Alaska (ACIA 2005, Berg et al. 2006, Raffa et al. 2008, Sherriff et al. 2011, Werner and Holsten 1985, Werner et al. 2006). However, it remains poorly resolved when and where Alaskan boreal forests may be vulnerable to future outbreaks and whether future outbreaks could influence subsequent patterns of burning.

In montane to subalpine Rocky-mountain conifer forests, bark beetle outbreaks can alter fire behavior (Hicke et al. 2012) and compromise fire-fighter safety (Jenkins et al. 2012). For example, active crown fire potential may increase 1-4 years post outbreak, before declining, as needles are shed from beetle-killed trees (Hicke et al. 2012). However, outbreaks appear to have modest or little effect on post-outbreak fuel profiles (Simard et al. 2011, Donato et al. 2013,

Schoennagel et al. 2012, Harvey et al. 2014a), wildfire probability (Kulakowski et al. 2003, Lynch et al. 2006, Kulakowski and Veblen 2007, Kulakowski and Jarvis 2011), area burned (Hart et al. 2015b), and fire severity (Bigler et al. 2005, Harvey et al. 2013, 2014b, Andrus et al. 2015). On the Kenai Peninsula, surface-fuel loads of nearly all size classes increased following the outbreak. Duff and moss depth, an indicator of fuel moisture, also decreased (Schulz 1995, 2003). This suggests that post-outbreak changes in forest structure may differ between boreal and Rocky-Mountain forests in ways that could lead to divergent consequences for subsequent patterns of burning.

The SBB outbreak on the Kenai Peninsula offers a tremendous opportunity to evaluate linked disturbance interactions in the boreal forest and the mediating role of forest-landscape structure. In this paper, we ask: (1) *How did the extent and duration of the 1990s SBB outbreak differ between a homogenous landscape dominated by white spruce and a landscape in which white spruce and black spruce were intermixed?* (2) *How has the occurrence and duration of SBB outbreak influenced the likelihood of subsequent burning in these two landscapes, as compared to other factors?* We hypothesized that the SBB outbreak was not as spatially extensive and of shorter duration in the north where white spruce and black spruce were intermixed. Conversely, we expected the outbreak to be substantially more extensive in the south where white spruce dominated. Despite different outbreak extents and durations, we hypothesized that increased surface fuels would lead to higher probabilities of subsequent burning in both regions.

Materials and Methods

Study area

The Kenai Peninsula of south-central Alaska sits south of Anchorage (Fig. 1) and is an ecological transition zone. West of the Kenai Mountains, much of the peninsula is a relatively flat, topographically homogenous plain that lies at the southwestern extent of the physical and ecological conditions characterizing Alaskan boreal forest (Morton et al. 2006). This is where the majority of the SBB outbreak occurred in the 1990s (Fig. 2). On the western Kenai Peninsula, mean annual precipitation varies from 369 mm in the north to 650 mm in the south (1971-2000) (Western Regional Climate Center 2012). Average annual temperature is approximately 1°C. The Kenai Peninsula is also one of the most densely populated areas of Alaska, and people increase numbers of ignitions and decrease area burned by suppressing fires in proximity to roads (Calef et al. 2008). To try to minimize confounding influences of people, we limited our analyses of bark-beetle effects on fire to include only forested areas more than 20 km away from the nearest major road. Tree-species composition varies markedly. Interior stands on the Kenai Peninsula are comprised of white spruce with some resin birch (*Betula neoalaskana*) in the south. Black spruce trees dominate the north of the peninsula, with a transition zone in the middle, where black spruce and white spruce are intermixed (Berg and Anderson 2006). Landscape structure of black spruce and white spruce strongly influence the fire regime. Black-spruce are highly flammable with a historical fire return interval of 60 to 80 years (DeVolder 1999). White spruce forests have burned much less frequently (fire return intervals of 400-600 years) (Berg et al. 2006). Thus, we stratified our study area by tree species distributions and identified two separate study regions; one north of Tustumena Lake (97,000 ha), where black spruce and white spruce are intermixed, and one south of the lake (133,500 ha), dominated by white spruce (Fig. 1) (Berg et al. 2006).

Approach and Data

To determine how forest-landscape structure mediated SBB outbreak extent and duration and to evaluate effects of SBB outbreak on the likelihood of subsequent burning, we assembled gridded geo-spatial datasets of fire perimeters (2001-2014), occurrence/duration of SBB outbreak (1989-2000), previous fire history (1946-2000), mean decadal (2001-2010) fire-season potential aridity (2001-2009), and information on forest structure (Table 1). All data were resampled using nearest neighbor to $\sim 1\text{km}^2$ resolution (chosen because it was the resolution of our coarsest dataset; climate). This yielded 1276 pixels and 1890 pixels in the northern and southern study regions, respectively, that served as observations in analyses.

Fire perimeters between 2001 and 2014 and previous fire history were derived from Alaska Fire Service's Fire History Database (Alaska Fire Service 2012). Perimeter maps are developed using field and aerial surveys and satellite imagery (Kasischke et al. 2002). The database includes all fires >405 ha since 1946 and all fires > 40.5 ha since 1988 (Calef et al. 2015). SBB outbreak and duration between 1989 and 2000 were measured in our study regions using aerial detection surveys from The U.S. Forest Service and Alaska Department of Natural Resource's Alaska Forest Health Protection Program (Fig. 2) (United States Forest Service and Alaska Department of Natural Resources 2012). The number of years that outbreak was recorded in any given pixel varied substantially. To explore whether outbreak duration was important, we conducted all analyses using several different duration thresholds to define whether or not outbreak occurred in a given pixel (e.g., varying whether we consider outbreak to have occurred by if it lasted ≥ 1 year, ≥ 2 years, ≥ 3 years, or ≥ 4 years).

We used the 2001 National Land Cover Database (NLCD) to map vegetation type and % forest canopy cover (Homer et al. 2004). In our study area, these data categorize pixels based on the dominant canopy tree type (e.g, conifer, mixed conifer-deciduous, deciduous) taller than five

meters in forested areas. Other categories included shrub, human development, barren land, water, etc. In the boreal forest, conifers can grow quite slowly. Thus, early to mid-succession forests may be misclassified as shrub in the NLCD. Eighty-two percent of pixels were classified as coniferous forest or shrub and only five percent as deciduous or mixed forest in our study regions. Because fire is most prevalent in conifer stands and to ensure that we had sufficient observations in each forest-type category to conduct robust statistical analyses, we only included pixels classified as conifer or shrub. We defined vegetation type as either late-successional conifer forest (conifer in NLCD) or early-successional conifer/shrub (shrub in NLCD).

Decadal (2001-2010) fire season (May-August) potential aridity, measured as precipitation minus potential evapotranspiration (P-PET), was calculated using gridded historical CRU TS 3.1 0.5° x 0.5° temperature data and CRU TS 3.1.01 0.5° x 0.5° precipitation data (Jones and Harris 2008, Mitchell and Jones 2005). Scenarios Network for Alaska and Arctic Planning downscaled climate data to a ~1km² resolution following the delta methodology and calculated PET using the Hamon equation (SNAP 2012). P-PET incorporates precipitation with evaporative demand, driven by temperature, to provide an estimate of landscape aridity. A lower P-PET value (increased potential aridity) means dryer conditions.

Analysis

We developed regressions to predict likelihood of burning between 2001 and 2014 as a function of SBB outbreak occurrence and duration, while controlling for other factors. Aspect and past fires between 1941 and 2000, which occurred in 1.6% of pixels, were not correlated with fire occurrence in earlier univariate analyses and thus were excluded from final regressions. Collinearity was evaluated using a variance inflation factor cutoff of four. Analyses were conducted separately for north and south study regions, and multiple regressions were run in

each region using different duration thresholds (outbreak occurred for ≥ 1 year, ≥ 2 years, ≥ 3 years, or ≥ 4 years) to define SBB outbreak occurrence. Our response variable, whether a pixel burned or not, is binary (a value of one if fire burned; zero if not), which is conducive to logistic regression. However, we identified spatial autocorrelation in logistic-regression residuals. Spatial autocorrelation often leads to biased standard error estimates. Thus, we used General Linear Mixed Models using Penalized Quasi Likelihood (GLMMPQL), which are widely applied to account for spatial dependence in binary regressions (Dormann et al. 2007). GLMMs combine linear mixed models, which account for correlated residual variance, with generalized linear models that can handle non-normal data (i.e. binary dependent variables) (Bolker et al. 2009).

Results

Bark beetle outbreak extent and duration

SBB outbreak was recorded by aerial detection surveys between 1989 and 2000 in approximately 35% of the northern study region, where black spruce and white spruce were intermixed (Table 1). Recorded outbreak duration ranged from one to seven years (Fig. 3). In 40% of pixels, outbreak was recorded as lasting a single year, and an outbreak duration of $\geq$ four years was recorded in only 5% of pixels. In marked contrast, the outbreak was substantially more extensive and of longer duration in the southern study region, where forests were dominated by white spruce (Fig. 3). Approximately 80% of the area was affected by the outbreak and recorded outbreak duration ranged from one to 11 years. In only 12% of pixels, outbreak was recorded as lasting a single year, and outbreak duration of $\geq$ four years was recorded in 32% of pixels (Fig. 3).

Likelihood of burning

In the north, three fires occurred between 2001 and 2014, burning nearly 30% (30,600 ha) of the region (Table 1). Fire was less prevalent in the south. Five fires burned 15% (19,500 ha) of the study region. Likelihood of burning in the northern study region increased with occurrence of SBB outbreak, if outbreak duration in a pixel was ≥ 2 or ≥ 3 years (Fig. 4). Likelihood of burning also increased in pixels classified as late-succession conifer forest and where percent canopy cover was higher. In the southern study region, likelihood of burning was unrelated to occurrence and duration of SBB outbreak as well as whether or not an observation was classified as late-successional conifer forest (Table 2, Fig. 4). However, likelihood of burning did increase with percent canopy cover. In both study regions, likelihood of burning was unrelated to potential aridity (Table 2).

Discussion

Our study illustrates how variation in forest landscape structure can shape SBB outbreak extent and duration and influences whether SBB outbreaks affect probability of subsequent burning in Alaska. On the Kenai Peninsula, SBB outbreak in the northern study area, where white and black spruce were intermixed, was not nearly as extensive and was often of shorter duration compared to the white-spruce dominated southern study area. Yet, likelihood of subsequent burning was higher when SBB outbreak occurred in the north, likely as a result of increased surface-fuel loads. In contrast, likelihood of burning was not altered by outbreak occurrence in the south, where fires have been historically rare. These results could inform more effective and targeted management strategies to ameliorate fire risk in beetle-killed stands of Alaska and may help us anticipate the dynamics and consequences of future boreal bark beetle outbreaks as climate warms at high latitudes.

SBBs generally do not target black spruce, instead preferring white spruce (Werner et al. 2006). Thus, suitable stands of host trees in the northern study region were less connected in their distribution, with black spruce intermixed, likely explaining smaller outbreak extent, shorter durations, and increased probability of subsequent burning, if outbreak duration was ≥ 2 or ≥ 3 years. Late-successional black spruce forests are highly flammable and have historically burned orders of magnitude more frequently than white-spruce-dominated stands on the Kenai Peninsula (DeVolder 1999, Berg et al. 2006). Forest-structure changes related to the outbreak (e.g., larger and drier fuel loads) may have increased the flammability of white spruce stands when black spruce are intermixed, leading to increased burning in areas that once likely served as fire breaks (Werner et al. 2006, Wolken et al. 2011).

Widespread post-outbreak increases in surface-fuel loads and heights on the western Kenai Peninsula were documented for 1-, 10-, 100-, and non-rotting 1000-hour fuel classes (Schulz 1995, 2003). This is in contrast with montane and subalpine Rocky-Mountain conifer forests where bark beetles appear to have modest effects on fuels (Simard et al. 2011, Donato et al. 2013, Schoennagel et al. 2012, Harvey et al. 2014a) and minimal influence on subsequent likelihood of burning or fire severity (Bigler et al. 2005, Kulakowski and Veblen 2007, Kulakowski and Jarvis 2011, Harvey et al. 2013, 2014b, Meigs et al. 2015). Surface fuels play a key role in the spread of boreal wildfires (Weber and Flannigan 1997). Surface-fuel accumulation following outbreak on the Kenai Peninsula may explain why we saw increased likelihood of burning in boreal forests where black spruce were present, when similar effects of bark-beetle outbreaks on fire have not been documented in other systems.

Increased likelihood of burning in the northern study region may also have resulted from drying surface fuels following outbreak of longer duration. This is supported by observations of

reduced moss and duff depth in post-outbreak stands (Schulz 2003). Substantial uncertainty remains in how bark beetle outbreaks affect hydrologic and radiative fluxes in forest ecosystems (Pugh and Gordon 2013, Brown et al. 2014, Bearup et al. 2014). However, bark beetle outbreaks can cause reduced evapotranspiration and increased sensible heat fluxes (Bearup et al. 2014). For example, summertime evapotranspiration was reduced 19% and sensible heat flux increased 8%, warming surface temperatures 1 °C, five years after a large bark beetle outbreak in British Columbia (Maness et al. 2013). Field-based measurements of air temperature in lodgepole pine stands of Yellowstone National Park varied with time since bark beetle outbreak. Initially, (0-4 years post outbreak) mean daily air temperature remained unchanged. By 30 years post outbreak, temperature was 1.3°C warmer (Griffin et al. 2011).

It is not surprising that outbreaks were more extensive and generally lasted longer in the southern study area. Many pre-outbreak stands in the south were dominated by even-aged, mature white spruce (Berg and Anderson 2006), leading to more than 90% of trees killed and a 87% reduction in stand basal area (Werner et al. 2006, Wolken et al. 2011). Subsequently, calamagrostis, an early successional dominant grass, proliferated extensively throughout SBB-affected stands in the southern region (Boucher and Mead 2006, Holsten et al. 1995, Lieffers et al. 1993, Schulz 1995). The grass can form a thick mat of dead litter that can constrain spruce regeneration by reducing soil temperatures and outcompeting tree seedlings for light and nutrients (Werner et al. 2006, Wolken et al. 2011). Though, inventories suggest sufficient regeneration has occurred to eventually restock most stands (Boucher and Mead 2006).

We found that SBB outbreak did not influence likelihood of subsequent burning in the southern study region. White spruce in this area have burned rarely during the Holocene, with a fire return interval of 400-600 years and some stands not burning for more than 1,000 years

(Berg and Anderson 2006). It appears bark beetle outbreak has done little to alter the fire history of the region (Berg and Anderson 2006), with one caveat. Our study was designed to minimize confounding effects of people on patterns of burning. Yet, some fire suppression still did occur, particularly in the southern study area. Although our finding that bark beetle outbreak has no effect of subsequent likelihood of burning in the south is consistent with past studies and the fire history (Berg and Anderson 2006), it is impossible to completely rule out the potential influence of suppression on patterns of burning. Sufficient data were not available to statistically account for suppression activities that did occur.

Potential aridity was not related to likelihood of burning in either study region. Generally, climate is considered a key driver of boreal fire regimes at regional scales and strong linkages exist between annual area burned and climate trends over long time periods (e.g., Kelly et al. 2013) and broad spatial scales (e.g., Duffy et al. 2005). At finer scales, fire spread is primarily driven by fuels in combination with extreme weather conditions that occur for just a few days or even hours. Gridded spatially explicit data of weather patterns at fine temporal resolutions was unavailable for use in this analysis.

Management applications.

Population density on the Kenai Peninsula is approximately three times greater than the average in Alaska (United States Census Bureau 2010) and there is a pronounced wildland urban interface where homes are interspersed among natural forests (Hansen and Naughton 2013). There has been substantial concern that SBB outbreak would increase the frequency, size, and severity of fires (Flint et al. 2006). Many fires have been suppressed since the 1990s outbreak as a result (Morton et al. 2006). While fire suppression can lead to short-term protection of people

and property, excluding natural-caused wildfire will likely increase the extent of late-succession forest and could heighten the risk of large catastrophic wildfires (Chapin et al. 2003).

Significant federal and state resources have also been expended to develop community wildfire protection plans and reduce hazardous fuels in high-risk areas along the road network (Hansen 2014). These efforts have had some success, and the approach is likely more promising than suppression for effectively protecting people and property over the long term. Results of this study could be used to direct application of fuel treatments on the western Kenai Peninsula and other areas of Alaska, if SBB outbreaks occur in the future. For example, managers could prioritize treatment in areas where white spruce are interspersed with more flammable black spruce.

Conclusion

We characterized how a massive SBB outbreak has influenced subsequent likelihood of burning in boreal forest of the Kenai Peninsula. Forest-landscape structure was a critical determining factor. Likelihood of burning increased in areas where flammable black spruce trees were intermixed with beetle-affected white spruce. We suspect that post-outbreak surface-fuel accumulation may increase the flammability of white spruce stands that would have once served as fire breaks and may also explain why our findings differ from many studies conducted in other systems (e.g., Rocky Mountains). Conversely, SBB outbreak had no effect on subsequent likelihood of burning in stands of uniform white spruce where fires have been historically rare (fire return interval of 400-600 years). Our results may help to anticipate the dynamics and consequences of future boreal bark beetle outbreaks, as climate warms at high latitudes. White spruce is widely distributed throughout interior Alaska (Calef et al. 2005) and interior white spruce burn substantially more frequently than white spruce on the Kenai Peninsula (Yarie

1981). If beetle outbreaks begin to occur in interior Alaska, increased fuel loads may lead to a higher likelihood of subsequent burning, particularly when white spruce are intermixed with or adjacent to more flammable black spruce.

Acknowledgements

This paper is dedicated to the memory of Kipling Rand. A true sage of the mountains, his passion for conservation of wild places will never be forgotten. We thank J. Morton and E. Berg for valuable insight into social and ecological dynamics on the Kenai Peninsula, Alaska. We thank M. Turner, W. Romme, D. Donato, and B. Harvey for insightful feedback on earlier versions of this manuscript. This paper is based upon work supported by the National Science Foundation Graduate Research Fellowship under Grant No. (DGE-1242789). Further funding was provided by the Alaska Climate Science Center, Scenarios Network for Alaska and Arctic Planning, the University of Alaska Fairbanks Resilience and Adaptation Program and Alaska EPSCoR Grant Number EPS-0701898 from the National Science Foundation. Helen Naughton also acknowledges funding from the Joint Fire Science Program under Projects JFSP 12-1-07-1 and JFSP 14-5-01-25.

Literature Cited

ACIA. 2005. Arctic Climate Impact Assessment. Cambridge University Press, New York, NY.
Alaska Fire Service. 2014. Wildfire Historical Database. Alaska Fire Service.
Online:<<http://fire.ak.blm.gov/predsvcs/maps.php>>.
Andrus, R. A., T. T. Veblen, B. J. Harvey, and S. J. Hart. 2015. Fire severity unaffected by spruce beetle outbreak in spruce-fir forests in southwestern Colorado. Ecological Applications In Press.

341 Balshi, M.S., A.D. McGuire, P. Duffy, M. Flannigan, J. Walsh, and J. Melillo. 2008. Assessing
 342 the response of area burned to changing climate in western boreal North America using a
 343 multivariate adaptive regression splines (MARS) approach. *Global Change Biology*
 344 15:578-600.

345 Barber, V. A., G. P. Juday, and B. P. Finney. 2000. Reduced growth of Alaskan white spruce in
 346 the twentieth century from temperature-induced drought stress. *Nature* 405: 668–673.

347 Bearup, L. A., R. M. Maxwell, D. W. Clow, and J. E. McCray. 2014. Hydrological effects of
 348 forest transpiration loss in bark beetle-impacted watersheds. *Nature Climate Change* 4:481–
 349 486.

350 Bentz, B. J., J. Regniere, C. J. Fettig, E. M. Hansen, J. L. Hayes, J. A. Hicke, R. G. Kelsey, J. F.
 351 Negron, and S. J. Seybold. 2010. Climate change and bark beetles of the western United
 352 States and Canada: Direct and indirect effects. *Bioscience* 60:602–613.

353 Berg, E.E., and R.S. Anderson. 2006. Fire history of white and Lutz spruce forests on the Kenai
 354 Peninsula, Alaska, over the last two millennia as determined from soil charcoal. *Forest*
 355 *Ecology and Management* 227:275-283.

356 Berg, E.E., J.D. Henry, C.L. Fastie, A.D.D. Volder, and S.M. Matsuoka. 2006. Spruce beetle
 357 outbreaks on the Kenai Peninsula, Alaska, and Kluane National Park and Preserve,
 358 Yukon Territory: Relationship to summer temperatures and regional differences in
 359 disturbance regimes. *Forest Ecology and Management* 227:219-232.

360 Bigler, C., D. Kulakowski, and T.V. Thomas. 2005. Multiple disturbance interactions and
 361 drought influence fire severity in Rocky Mountain subalpine forests. *Ecology*, 86:3018-
 362 3029.

363 Bolker, B. M., M. E. Brooks, C. J. Clark, S. W. Geange, J. R. Poulsen, M. H. H. Stevens, and J.-
 364 S. S. White. 2009. Generalized linear mixed models: a practical guide for ecology and
 365 evolution. *Trends in Ecology & Evolution* 24:127–35.

366 Boucher, T.V. and B.R. Mead. 2006. Vegetation change and forest regeneration on the Kenai
 367 Peninsula, Alaska following a spruce beetle outbreak, 1987-2000. *Forest Ecology and*
 368 *Management* 227:233-246.

369 Brown, M. G., T. A. Black, Z. Nesic, V. N. Foord, D. L. Spittlehouse, A. L. Fredeen, R. Bowler,
 370 N. J. Grant, P. J. Burton, J. A. Trofymow, D. Lessard, and G. Meyer. 2014.
 371 Evapotranspiration and canopy characteristics of two lodgepole pine stands following
 372 mountain pine beetle attack. *Hydrological Processes* 28:3326–3340.

373 Calef, M.P., A.D. McGuire, F.S. Chapin III. 2008. Human influence on wildfire in Alaska from
 374 1988 through 2005: An analysis of the spatial patterns of human impacts. *Earth*
 375 *Interactions* 12:1-17.

376 Calef, M. P., A. Varvak, A.D. McGuire, F.S. Chapin, & K.B. Reinhold. 2015. Recent changes in
 377 annual area burned in interior Alaska: The impact of fire management. *Earth Interactions*,
 378 19: 1–17.

379 Calef, M. P., A.D. McGuire, H.E. Epstein, T.S. Rupp, and H.H. Shugart. 2005. Analysis of
 380 vegetation distribution in Interior Alaska and sensitivity to climate change using a logistic
 381 regression approach. *Journal of Biogeography* 32: 863–878.

382 Chapin, F.S., T.S. Rupp, A.M. Starfield, L. DeWilde, E.S. Zavaleta, N. Fresco, J. Henkelman,
 383 and A.D. McGuire. 2003. Planning for resilience: modeling change in human-fire
 384 interactions in the Alaskan boreal forest. *Frontiers in Ecology and the Environment*
 385 1:255-261.

386 DeVolder, A. 1999. Fire and Climate History of Lowland Black Spruce Forests, Kenai National
387 Wildlife Refuge, Alaska. Master's Thesis, Northern Arizona University.

388 Donato, D.C., B.J. Harvey, W.H. Romme, M. Simard, and M.G. Turner. 2013. Bark beetle
389 effects on fuel profiles across a range of stand structures in Douglas-fir forests of Greater
390 Yellowstone. *Ecological Applications* 23:3-20.

391 Dormann, F. , C., J. M. McPherson, M. B. Araújo, R. Bivand, J. Bolliger, G. Carl, R. G. Davies,
392 A. Hirzel, W. Jetz, W. Daniel Kissling, I. Kühn, R. Ohlemüller, P. R. Peres-Neto, B.
393 Reineking, B. Schröder, F. M. Schurr, and R. Wilson. 2007. Methods to account for
394 spatial autocorrelation in the analysis of species distributional data: a review. *Ecography*
395 30:609–628.

396 Duffy, P. A., J. Eptin, J. M. Graham, T. S. Rupp, and A. D. McGuire. 2007. Analysis of Alaskan
397 burn severity patterns using remotely sensed data. *International Journal of Wildland Fire*
398 16:277–284.

399 Flannigan, M., K. Logan, B. Amiro, W. Skinner, and B. Stocks. 2005. Future area burned in
400 Canada. *Climatic Change* 72:1-16.

401 Flint, C.G. 2006. Community perspectives on spruce beetle impacts on the Kenai Peninsula,
402 Alaska. *Forest Ecology and Management* 227:207-218.

403 Griffin, J. M., M. G. Turner, and M. Simard. 2011. Nitrogen cycling following mountain pine
404 beetle disturbance in lodgepole pine forests of Greater Yellowstone. *Forest Ecology and*
405 *Management* 261:1077–1089.

406 Hansen, W. D. 2014. Generalizable principles for ecosystem stewardship-based management of
407 social-ecological systems: Lessons learned from Alaska. *Ecology and Society* 19: 13.

408 Hansen, W. D., and H. T. Naughton. 2013. The effects of a spruce bark beetle outbreak and
 409 wildfires on property values in the wildland-urban interface of south-central, Alaska,
 410 USA. *Ecological Economics* 96:141–154.

411 Harvey, B. J., D. C. Donato, W. H. Romme, and M. G. Turner. 2014a. Fire severity and tree
 412 regeneration following bark beetle outbreaks : the role of outbreak stage and burning
 413 conditions. *Ecological Applications* 24:1608–1625.

414 Harvey, B. J., D. C. Donato, W. H. Romme, and M. G. Turner. 2013. Influence of recent bark
 415 beetle outbreak on fire severity and post-fire tree regeneration in montane Douglas-fir
 416 forests. *Ecology* 94:2475–2486.

417 Harvey, B. J., D. C. Donato, and M. G. Turner. 2014b. Recent mountain pine beetle outbreaks,
 418 wildfire severity, and postfire tree regeneration in the US Northern Rockies. *Proceedings*
 419 *of the National Academy of Sciences* 111:15120–15125.

420 Hart, S. J., T. T. Veblen, N. Mietkiewicz, and D. Kulakowski. 2015a. Negative feedbacks on
 421 bark beetle outbreaks: Widespread and severe spruce beetle infestation restricts
 422 subsequent infestation. *Plos One* 10.

423 Hart, S. J., T. Schoennagel, T. T. Veblen, and T. B. Chapman. 2015b. Area burned in the western
 424 United States is unaffected by recent mountain pine beetle outbreaks. *Proceedings of the*
 425 *National Academy of Sciences* 112: 4375–4380.

426 Hicke, J.A., M.C. Johnson, J.L. Hayes, and H.K. Preisler. 2012. Effects of bark beetle-caused
 427 tree mortality on wildfire. *Forest Ecology and Management* 271:81–90.

428 Holsten, E.H., R.A. Werner, and R.L. Develice. 1995. Effects of a spruce beetle (*Coleoptera:*
 429 *Scolytidae*) outbreak and fire on Lutz spruce in Alaska. *Environmental Entomology*
 430 24:1539-1547.

431 Homer, C., C. Huang, L. Lang, B. Wylie, and M. Coan. 2004. Development of a 2001 national
 432 landcover database for the United States. *Photogrammetric Engineering and Remote*
 433 *Sensing* 70:829-840.

434 Jenkins, M. J., W. G. Page, E. G. Hebertson, and M. E. Alexander. 2012. Fuels and fire behavior
 435 dynamics in bark beetle-attacked forests in Western North America and implications for
 436 fire management. *Forest Ecology and Management* 275:23–34.

437 Johnstone, J. F., T. S. Rupp, M. Olson, and D. Verbyla. 2011. Modeling impacts of fire severity
 438 on successional trajectories and future fire behavior in Alaskan boreal forests. *Landscape*
 439 *Ecology*, 26: 487–500.

440 Jones, P.D., and I. Harris. 2008. CRU time series (TS) high resolution gridded datasets.
 441 Online:<http://badc.nerc.ac.uk/view/badc.nerc.ac.uk__ATOM__dataent_1256223773328
 442 276>.

443 Kasischke, E.S., D. Williams, and D. Barry. 2002. Analysis of the patterns of large fires in the
 444 boreal forest region of Alaska. *International Journal of Wildland Fire* 11:131-144.

445 Kelly, R., M.L. Chipman, P.E. Higuera, I. Stefanova, L.B. Brubaker, and F.S. Hu. 2013. Recent
 446 burning of boreal forests exceeds fire regime limits of the past 10,000 years. *Proceedings of*
 447 *the National Academy of Sciences of the United States of America*, 110: 13055–60.

448 Kulakowski, D., T. T. Veblen, and P. Bebi. 2003. Effects of fire and spruce beetle outbreak
 449 legacies on the disturbance regime of a subalpine forest in Colorado. *Journal of*
 450 *Biogeography* 30:1445–1456.

451 Kulakowski, D., and T.T. Veblen. 2007. Effect of prior disturbances on the extent and severity of
 452 wildfire in Colorado subalpine forests. *Ecology* 88:759-769.

453 Kulakowski, D., and D. Jarvis. 2011. The influence of mountain pine beetle outbreaks and
 454 drought on severe wildfires in northwestern Colorado and southern Wyoming: A look at
 455 the past century. *Forest Ecology and Management* 262:1686-1696.

456 Lieffers, V.J., S.E. Macdonald, and E.H. Hogg. 1993. Ecology of and control strategies for
 457 *Calamagrostis canadensis* in boreal forest sites. *Canadian Journal of Forest Research*
 458 23:2070-2077.

459 Lynch, H., R. Renkin, R. Crabtree, and P. Moorcroft. 2006. The influence of previous mountain
 460 pine beetle (*Dendroctonus ponderosae*) activity on the 1988 Yellowstone fires.
 461 *Ecosystems* 9:1318–1327.

462 Maness, H., P.J. Kushner, and I. Fung. 2013. Summertime climate response to mountain pine
 463 beetle disturbance in British Columbia. *Nature Geoscience* 6:65-70.

464 Meigs, G. W., J. L. Campbell, H. S. J. Zald, J. D. Bailey, D. C. Shaw, and R. E. Kennedy. 2015.
 465 Does wildfire likelihood increase following insect outbreaks in conifer forests?
 466 *Ecosphere* 6.

467 Millar, C. I., and N. L. Stephenson. 2015. Temperate forest health in an era of emerging
 468 megadisturbance. *Science* 349: 823–826.

469 Mitchell, T. D., and P. D. Jones. 2005. An improved method of constructing a database of
 470 monthly climate observations and associated high-resolution grids. *International Journal*
 471 *of Climatology* 25:693-712.

472 Morton, J. M., E. Berg, D. Newbould, D. MacLean, and L. O'Brien. 2006. Wilderness fire
 473 stewardship on the Kenai National Wildlife Refuge, Alaska. *International Journal of*
 474 *Wilderness* 12:14-17.

475 Pugh, E., and E. Gordon. 2013. A conceptual model of water yield effects from beetle-induced
 476 tree death in snow-dominated lodgepole pine forests. *Hydrological Processes* 27:2048–
 477 2060.

478 Raffa, K. F., B. H. Aukema, B. J. Bentz, A.L. Carroll, J. A. Hicke, M.G . Turner, and W. H.
 479 Romme. 2008. Cross-scale drivers of natural disturbances prone to anthropogenic
 480 amplification: The dynamics of bark beetle eruptions. *Bioscience* 58:501-517.

481 Schoennagel, T., T. T. Veblen, and W. H. Romme. 2004. The interaction of fire, fuels, and
 482 climate across Rocky Mountain forests. *BioScience* 54: 393–402.

483 Schoennagel, T., T.T. Veblen, J.F. Negron, and J.M. Smith. 2012. Effects of mountain pine
 484 beetle on fuels and expected fire behavior in lodgepole pine forests, Colorado, USA.
 485 *PLoS ONE* 7:1-14.

486 Schulz, B. 1995. Changes Over Time in Fuel-loading Associated with Spruce Beetle-impacted
 487 Stands of the Kenai Peninsula, Alaska. Forest Health Management Technical Report
 488 R10-TP-53, U.S. Forest Service.

489 Schulz, B. 2003. Changes in Downed and Dead Woody Material Following A Spruce Beetle
 490 Outbreak on the Kenai Peninsula, Alaska. Technical Report PNW_RP_559, U.S. Forest
 491 Service.

492 Seidl, R., J. Müller, T. Hothorn, C. Bässler, M. Heurich, and M. Kautz. 2015. Small beetle,
 493 large-scale drivers: how regional and landscape factors affect outbreaks of the European
 494 spruce bark beetle. *Journal of Applied Ecology* In Press.

495 Sherrif, R.L., E.E. Berg, and A.E. Miller. 2011. Climate variability and spruce beetle
 496 (*Dendroctonus rufipennis*) outbreaks in south-central and southwest Alaska. *Ecology*
 497 92:1459-1470.

498 Simard, M., W.H. Romme, J.M. Griffin, and M.G. Turner. 2011. Do mountain pine beetle
 499 outbreaks change the probability of active crown fire in lodgepole pine forests?
 500 Ecological Monographs 81:3-24.

501 SNAP. 2012. Historical Monthly and Temperature and Precipitation- 771m CRU TS 3.1/3.1.01.
 502 Online:< <http://www.snap.uaf.edu/data.php>>.

503 Temperli, C., T. T. Veblen, S. J., Hart, D. Kulakowski, and A. J. Tepley. 2015. Interactions
 504 among spruce beetle disturbance, climate change and forest dynamics captured by a forest
 505 landscape model. Ecosphere 6: 1–20.

506 Trumbore, S., P. Brando, and H. Hartmann. 2015. Forest health and global change. Science 349:
 507 814–818.

508 Turner, M. G., and W. H. Romme. 1994. Landscape dynamics in crown fire ecosystems.
 509 Landscape Ecology 9: 59–77.

510 Turner, M.G. 2010. Disturbance and landscape dynamics in a changing world. Ecology 91:2833-
 511 2849.

512 United States Forest Service and Alaska Department of Natural Resources. 2012. Alaska Forest
 513 Health Survey. Online:<<http://agdc.usgs.gov/data/projects/fhm/>>.

514 United States Census Bureau. 2010. United States Census, 2010. Online:
 515 <<http://quickfacts.census.gov/qfd/index.html>>.

516 Weber, M.G., and M.D. Flannigan. 1997. Canadian boreal forest ecosystem structure and
 517 function in a changing climate: impact on fire regimes. Environmental Reviews 5:145-
 518 166.

519 Werner, R.A., and E.H. Holsten. 1985. Factors influencing generation times of spruce beetles in
 520 Alaska. Canadian Journal of Forest Research 15:438.

521 Werner, R.A., E.H. Holsten, S.M. Matsuoka, and R.E. Burnside. 2006. Spruce beetles and forest
 522 ecosystems in south-central Alaska: A review of 30 years of research. *Forest Ecology and*
 523 *Management* 227:195-206.

524 Westerling, A.L., H.G. Hidalgo, D.R. Cayan, and T.W. Swetnam. 2006. Warming and earlier
 525 spring increase western US forest wildfire activity. *Science* 313:940-943.

526 Western Regional Climate Center. 2012. Western Region Cooperative Climate Science
 527 Summaries. Online:< <http://www.wrcc.dri.edu/summary/Climsmak.html>>.

528 Wolken, J. M., T. N. Hollingsworth, T. S. Rupp, F. S. Chapin, S. F. Trainor, T. M. Barrett, P. F.
 529 Sullivan, a. D. McGuire, E. S. Euskirchen, P. E. Hennon, E. a. Beever, J. S. Conn, L. K.
 530 Crone, D. V. D'Amore, N. Fresco, T. a. Hanley, K. Kielland, J. J. Kruse, T. Patterson, E.
 531 A. G. Schuur, D. L. Verbyla, and J. Yarie. 2011. Evidence and implications of recent and
 532 projected climate change in Alaska's forest ecosystems. *Ecosphere* 2.

533 Yarie, J. 1981. Forest fire cycles and life tables: a case study from interior Alaska. *Canadian*
 534 *Journal of Forest Research* 11:554-562.

535

536

537 **Tables**

538 Table 1. Descriptive statistics of dependent and predictor variables used in analyses of likelihood

539 of burning between 2001 and 2014. Values are separated by region in our study area.

	Total Area	Mean	SE	Min	Max
North Region (ha)	97,334	-	-	-	-
SBB Outbreak (ha)	31,136	-	-	-	-
SBB Outbreak Duration (yrs)	-	2.3	0.10	1	7
Fire (ha)	30,600	10,206	7,195	471	27,797
Forested (ha)	39,906	-	-	-	-
Potential Aridity (mm)	-	-7.6	0.81	-46.4	106.8
Canopy Cover (%)	-	34.6	1.0	0	94
South Region (ha)	133,355	-	-	-	-
SBB Outbreak (ha)	106,477	-	-	-	-
SBB Outbreak Duration (yrs)	-	3.6	0.10	1	11
Fire (ha)	19,500	3,885	2,204	10	12,339
Forested (ha)	58,676	-	-	-	-
Potential Aridity (mm)	-	-19.2	0.39	-49.6	77.2
Canopy Cover (%)	-	38.2	0.81	0	94

540

541 Table 2. Results of GLMM PQL regressions for A. the northern study region and B. the southern
542 study region predicting likelihood of burning between 2001 and 2014. Coefficients (SE) are
543 presented.

Bark Beetle Outbreak Duration				
	≥ 1 year	≥ 2 years	≥ 3 years	≥ 4 years
<i>A. North Region</i>				
SBB Outbreak	0.11 (0.07)	0.15 (0.08)**	0.27 (0.09)***	0.1 (0.11)
Forested	0.2 (0.06)***	0.22 (0.06)***	0.21 (0.06) ***	0.23 (0.06)***
Potential Aridity	-0.002 (0.005)	-0.003 (0.005)	-0.003 (0.005)	-0.002 (0.005)
% Canopy Cover	0.002 (0.001)*	0.002 (0.001)*	0.002 (0.001)*	0.23 (0.06)***
<i>B. South Region</i>				
SBB Outbreak	-0.02 (0.1)	-0.06 (0.09)	0.05 (0.08)	-0.04 (0.07)
Forested	0.04 (0.05)	0.04 (0.05)	0.04 (0.05)	0.04 (0.05)
Potential Aridity	-0.01 (0.01)	-0.01 (0.01)	-0.01 (0.01)	-0.01 (0.01)
% Canopy Cover	0.002 (0.001)**	0.002 (0.001)**	0.002 (0.001)**	0.002 (0.001)**

544 *P<0.1, **P<0.05, ***P<0.01

545

546 Figure 1. Map of the study regions on the Kenai Peninsula in south-central Alaska depicting the
547 major road system (in black).

548 Figure 2. Map of the Kenai Peninsula depicting A. the extent of the 1990s spruce bark beetle
549 outbreak and B. wildfires that burned between 2001 and 2014.

550 Figure 3. Top panel: Log transformed extent of SBB outbreak (ha) in each study region by year.
551 Bottom panel: Proportions of SBB outbreak in each study region binned by outbreak duration
552 (Year).

553 Figure 4. Influence of SBB outbreak occurrence and duration (regression coefficients, β) on
554 likelihood of burning between 2001 and 2014. Astrices denote significant relationships.

Figure 1

Figure 2

Figure 3

Figure 4