**Mechanical vulnerability and resistance to snapping and
uprooting for Central Amazon tree species**

G.H.P.M. RIBEIRO^{1,*}, J.Q. CHAMBERS^{1,2}, C.J. PETERSON³, S.E.
TRUMBORE⁴, D. MAGNABOSCO MARRA^{1,2,5}, C. WIRTH⁵, J.B.
CANNON⁶, R.I. NÉGRON-JUÁREZ⁷, A.J.N. LIMA¹, E.V.C.M de PAULA¹,
J. SANTOS¹, N. HIGUCHI¹

¹National Institute for Amazonian Research (INPA), Av. André Araújo, 2.936, Manaus,
AM, Brazil

²University of California, Department of Geography, Berkeley, CA 94720, USA

³Dept. of Plant Biology, University of Georgia, 2502 Miller Plant Sciences, GA 30602,
Athens, United States

⁴Dept. of Biogeochemical Processes, Max Planck Institute for Biogeochemistry, Hans-
Knöll-Str. 10, 07745, Jena, Germany

⁵AG Spezielle Botanik und Funktionelle Biodiversität, Universität Leipzig, Johannisallee
21, 04103, Leipzig, Germany

⁶Dept. of Forest and Rangeland Stewardship, Colorado State University, Fort Collins, CO,
80523, USA

⁷Lawrence Berkeley National Laboratory, Earth and Environmental Sciences Area, 1
Cyclotron Rd., MS74R316C, CA 94720, Berkeley, United States

* Corresponding author: RIBEIRO, G.H.P.M. E-mail: gabrielgiga@gmail.com

Abstract

High descending winds generated by convective storms are a frequent and a major source of tree mortality disturbance events in the Amazon, affecting forest structure and diversity across a variety of scales, and more frequently observed in western and central portions of the basin. Soil texture in the Central Amazon also varies significantly with elevation along a topographic gradient, with decreasing clay content on plateaus, slopes and valleys respectively. In this study we investigated the critical turning moments (M_{crit} - rotational force at the moment of tree failure, an indicator of tree stability or wind resistance) of 60 trees, ranging from 19.0 to 41.1 cm in diameter at breast height (DBH) and located in different topographic positions, and for different species, using a cable-winch load-cell system. Our approach used torque as a measure of tree failure to the point of snapping or uprooting. This approach provides a better understanding of the mechanical forces required to topple trees in tropical forests, and will inform models of wind throw disturbance. Across the topographic positions, size controlled variation in M_{crit} was quantified for *cardeiro* (*Scleronema mincranthum* (Ducke) Ducke), *mata-matá* (*Eschweilera spp.*), and a random selection of trees from 19 other species. Our analysis of M_{crit} revealed that tree resistance to failure increased with size (DBH and ABG) and differed among species. No effects of topography or failure mode were found for the species either separately or pooled. For the random species, total variance in M_{crit} explained by tree size metrics increased from an R^2 of 0.49 for DBH alone, to 0.68 when both DBH and stem fresh wood density (SWD) were included in a multiple regression model. This mechanistic approach allows the comparison of tree vulnerability induced by wind damage across ecosystems, and facilitates the use of forest structural information in ecosystem models that include variable resistance of trees to mortality inducing factors. Our results indicate that observed topographic differences in windthrow vulnerability are likely due to elevational differences in wind velocities, rather than by differences in soil-related factors that might effect M_{crit} .

Key-words: tree static winching; critical turning moment; tree allometry; functional traits; blowdown; wind-disturbance

1. Introduction

The stability of the erect habit of plant stems is primarily compromised by bending movements in response to wind and gravity (Alméras and Fournier, 2009). Wind is indicated as the primary abiotic agent damaging forests (Sellier and Fourcaud, 2009). Strong winds may cause deflection that may exceed critical thresholds of tree stability, and thus wind interaction with forests causes a large fraction of tree mortality worldwide (Chambers et al., 2007; Dale et al., 2001; Espirito-Santo et al., 2010; Negrón-Juárez et al., 2010; Schelhaas et al., 2003). Wind disturbance also affects forest structure and functioning (Peltola et al., 2010), influencing tree species assembly processes (Denslow, 1980; Marra et al., 2014; Turner et al., 1998), regeneration patterns (Putz et al., 1983) and shifts in forest composition (Trumbore et al., 2015).

High wind events create tree fall gaps at varying spatial scales from stands to landscapes (Asner, 2013; Chambers et al., 2013; Espirito-Santo et al., 2014; Mitchell, 2013). Although a portion of the carbon from decomposing, dead, and damaged trees can be incorporated into the soil (dos Santos et al., 2015) rather than respired to the atmosphere, downed trees represent committed future CO₂ emissions (Chambers et al., 2004). Recruitment and enhanced growth of surviving trees following disturbance can offset some of those emissions (Chambers et al., 2004). Forest carbon stocks are also subject to temporal fluctuations, and decline with increased disturbance event frequency (Chambers et al., 2013). Thus, increasing global temperature can reduce forest carbon balance if more frequent storms or higher peak wind speeds occur under a warming climate (IPCC, 2013).

Variation in vulnerability to tree mortality generated by wind in Amazon forests is a complex phenomenon, involving a suite of biotic and abiotic variables (Chambers et al., 2009;

Garstang et al., 1998; Nelson et al., 1994). Explanations for why some trees uproot (Garstang et al., 1998), while others do not, include differences in the forces exerted on tree crowns (from above and below), and the influence of rooting or soil depth. A recent study (Marra et al., 2014) reported that tree uprooting and snapping were more common on slopes and plateaus in the Central Amazon, with fewer uprooted or snapped trees in valleys. A previous study using wind tunnel measurements reported that wind speeds on hilltops were nearly twice that as observed for valley formations, with a larger variation in wind speed for valleys depending on the direction of the gusts (Ruel et al., 2001). It is also known that small increments in wind speed can result in substantial differences in mode and scale of forest wind disturbance. Wind speed frequently increases with elevation and the pattern of disturbance may be strongly influenced by topography (Quine and Gardiner, 2007).

Soil characteristics might have a strong influence on a tree's ability to withstand high wind events, yet there are few quantitative data linking soils and wind throw vulnerability for Amazon forests (Marra et al., 2014; Toledo et al., 2012). Soil and topography, in the context of tropical forests, has been studied to determine their influence on tree diversity (Baldeck et al., 2012; Laurance et al., 2010; Quesada et al., 2010). Soil texture, organic matter content, soil moisture, soil pH and soil C and N concentrations vary significantly with elevation along topographic gradient in the Central Amazon (Luizão et al., 2004). At regional scales, soil conditions such as fertility (Quesada et al., 2009), and the capacity of trees to develop effective anchorage (Mitchell, 2013) have been linked to variation in tree turnover rates.

Soil depth, soil physical properties, structure and other factors that vary with topography have a strong relationship with tree turnover rates (Chao et al., 2008; Quesada et al., 2009). And have been suggested as factors that control variability in wind-throw rates (Chao et al., 2009; Quesada et al., 2009). The present study was conducted in a region consisting of plateaus with clay-rich soils (with 65-75% of clay in the first 30 cm, increasing to 80-90% clay at 2-4 m of depth) and valleys with sand content approaching 100% near perennial streams (Chauvel et al., 1987).

Slopes connecting plateaus to valleys exhibit a gradual decrease in clay content, and increase in sand content, with elevation (Chauvel et al., 1987). Clay soils are more likely to reduce root growth (Gerard et al 1982), and this may explain previous results of lower uprooting rates in valley positions (Lenart et al., 2010; Marra et al., 2014).

In addition to abiotic factors including topography and soil, biotic factors such as individual or species characteristics may also influence tree stability (Negrón-Juárez et al., 2014; Niklas and Spatz, 2010). A decision support system approach to assess hazard due wind driven mortality for single trees and stands emphasizes the importance of larger tree taper, strength and depth of tap roots, and interlocking root systems as good predictors of low failure hazard (Mickovski et al., 2005). It is expected that trees with a longer life cycle and slower growth rates produce stiffer, stronger stems with greater strength and longevity (Niklas and Spatz, 2010; Williamson and Wiemann, 2010). Inferences of tree stability based on wood density alone are potentially misleading for species with complex life histories (Nock et al., 2009), where the ability of wood to resist mechanical forces may decline with increasing moisture content (Niklas and Spatz, 2010). Large trees vary more than saplings in their wood densities, and are also likely to vary more in their mechanical properties, with potentially important consequences for architecture, growth and survival (van Gelder et al., 2006).

Trees resist their own mass and wind forces by an adequate mechanical design (Sterck and Bongers, 1998), where plant architecture itself can alter the magnitude and spatial distribution of wind loading (Sellier and Fourcaud, 2009). The dynamic behavior of trees and their response to mechanical loads in particular in response to wind loading varies with time (Brüchert and Gardiner, 2006). Critical turning moment (M_{crit}) of a tree is the rotational force (Newton meters) at the moment of failure (snap or uprooting), and indicates tree stability or wind resistance (Cannon et al., 2015). M_{crit} generally increases with tree size and often does not differ significantly across species for similar sized trees, (Cannon et al., 2015; Nicoll et al., 2006; Peltola et al., 2000; Peterson and

Claassen, 2013). The best predictors of M_{crit} vary among studies but include stem volume or above ground biomass (AGB) (Cucchi et al., 2004; Fredericksen et al., 1993; Lundström et al., 2007; Peterson and Claassen, 2013), DBH (Kamimura et al., 2012), and stem mass (Cannon et al., 2015).

To our knowledge, tree pulling experiments based on static load tests have not previously been carried out in Amazon forests. Here we examine relationships among tree stability, topography, mode of failure, species, and tree biometric parameters (i.e. DBH, stem mass, crown mass, AGB, stem volume, stem fresh wood density, and total height). We hypothesized that (1) trees in valleys are more resistant than trees on plateaus (due to lower observed rates of wind mortality in this topographic position, found by Marra *et al* 2014); (2) M_{crit} increases with tree size (e.g. DBH, AGB tree taper or Slenderness, (TH:DBH ratio, dimensionless), Stem, Crown and Center), for all species groups; (3) tree resistance to failure increases with increasing SWD, given that higher wood density implies stiffer stems and stronger or greater strength and longevity (and tropical forests has a very large range in wood density among species); and (4) the total force required to snap or uproot a tree do not differ in relation to M_{crit} (similar number of dead trees by failure mode, found by Marra *et al* 2014).

2. Methods

2.1. Study site

We conducted the experiment at the EEST (Experimental Station of Forest Management) operated by Brazil's National Institute of Amazonian Research (Instituto Nacional de Pesquisas da Amazônia-INPA). The EEST is located at (2.45°– 2.66°S, 60.02°– 60.32°W) 90 km north of Manaus, the capital of Amazonas State, Brazil (Silva et al., 2002) and encompasses an area of 21,000 ha (Andrade and Higuchi, 2009). Annual rainfall for the period 1980–2000 was 2610 ± 124 mm, using data from an EMBRAPA experimental station, located 50 km east from EEST (Silva et al., 2003). The study region has a dry season (≤ 100 mm precipitation) from July to September

(Chambers et al., 2004). Topography at EEST is composed mostly of flat plateaus (90–105 m ASL) incised by a dense drainage network within broad and often swampy valleys (45–55 m ASL) (Rennó et al., 2008). In this region, forests are composed of a high diversity of tree species (Carneiro, 2004; Guillaumet, 1987; Marra et al., 2014; Rankin de Mérona et al., 1992; Vieira et al., 2004). About 280 species can be found in a single hectare (Oliveira A.A. and Mori, 1999), with low frequency (around 4 per ha) of emergent trees that exceed 30 m (Prance et al., 1976) and dominant height of trees $\leq$ 29 m (Lima, 2010).

2.2. Species selection

To test our hypotheses, species were chosen based on the species' abundance along the typical local topographic gradient found at the EEST (Carneiro, 2004). Cardeiro (*Scleronema mincranthum* (Ducke) Ducke [Malvaceae]) and mata-matá (*Eschweilera* spp. [Lecythidaceae]) were selected. *Scleronema* and *Eschweilera* are among the most frequent genera in our study area (Guillaumet, 1987) and both are listed as “hyperdominant” in the entire Amazon basin, with *Eschweilera* represented by 52 species (ter Steege et al., 2013). *Scleronema* and *Eschweilera* have also been described as genera favored (higher abundance) in intermediate and heavily damaged patches, respectively, of a nearby forest disturbed by the 2005 convective storm (Marra et al., 2014). In addition to these two genera, a random selection of trees across 19 species (see supplementary material, Table S1) representing the local species assembly was also sampled to seek trends not specific to the above-mentioned dominant genera.

2.3. Field methods

Trees that were selected for winching followed procedures consistent with previous studies (Cannon et al., 2015; Nicoll et al., 2006; Peterson and Claassen, 2013). We used essentially the same equipment employed in previous studies (Cannon et al., 2015; Peterson and Claassen, 2013) with minor modifications. First, all trees were hand winched without the use of a snatch block, and

the load cell used in this study (Straightpoint model Radiolink Plus) had a smaller rated capacity of 11.8 metric tons that was not exceeded, see Supplementary video. To find the center of mass of each trees, we recorded trunk diameter at 1 m intervals (and DBH) after each tree fall and weighed every 1 m section of the tree with a 300 kg load capacity balance. Details are presented in supplementary data (see section 2).

2.4. Biometric parameters influencing tree stability

For each tree we measured the stump diameter (D_{stump}), diameter at breast height (DBH), stem volume (Vol), stem mass (Stem), crown mass (Crown), above ground biomass (AGB), stem wood density (SWD), tree center of mass (Center) and tree height (TH). Descriptive analyses for each species group versus topographic position are shown in the supplementary material (Table S2-S4, for *Eschweilera* spp., a random set of tree species and *Scleronema mincranthum*, respectively). The measure for linear association between the variables using Pearson product-moment correlation and Bonferroni probabilities are show respectively for *Eschweilera* spp. (Tables S5 and S6), for a random set of tree species (Tables S7 and S8), for *Scleronema mincranthum* (Tables S9 and S10) and for all species groups pooled together (Tables S11 and S12). To compare our results with previous studies, we ran simple (linear and non-linear) and multiple regressions between M_{crit} versus tree biometric parameters, such as DBH, total tree height (TH), above ground biomass (AGB), stem volume (Vol), fresh stem wood density (SWD), for *Eschweilera* spp., a random set of tree species, *Scleronema mincranthum*, and species groups pooled, respectively (Tables S13-S16).

M_{crit} was calculated at the base of each pulled tree (Figure S2). We winched 20 trees for each group (*Scleronema mincranthum*, *Eschweilera* spp., and a random selection of tree species), 10 in each topographic position (plateau and valley), totaling 60 trees ranging from 19 cm to 41.1 cm DBH. Trees larger than 41.1 cm DBH were not winched in this study as a safety precaution. This is a reasonable maximum size since 93% of trees in a nearby 18ha plot had DBH values between 10 – 40 cm (Higuchi et al., 2012), and the mean DBH of trees with $\text{DBH} \geq 10$ cm in

Central Amazon undisturbed forest in the region is about 22 cm (Lima 2010). The mean DBH of dead trees (i.e. snapped and uprooted) found in a contiguous wind-disturbed forest was approximately 27.5 ± 2.3 cm (mean $\pm$ 95%CI) (Marra et al., 2014). Thus the tree size range sampled in this study is representative of forests of the Central Amazon.

2.5. Tree stability

To check the first hypothesis (trees in valleys are more resistant than trees on plateaus), we compared M_{crit} of species groups versus topographic position (as a factor) by means of one-way ANOVA. Subsequently, we ran an ANOVA including topographic position as a predictor of M_{crit} with DBH as an independent variable. After checking for differences among each species group, individuals from all species were pooled for the same analysis. To check the second hypothesis (evaluate whether M_{crit} increases with tree size), the effects of species group (as a factor), DBH, AGB, tree taper or Slenderness (TH:DBH ratio, dimensionless), Stem, Crown and Center as independent variables, we used analysis of covariance (ANCOVAs), one for each independent variable. Tree height was not a good predictor of M_{crit} in previous studies (Cannon et al., 2015; Peterson and Claassen, 2013). To check the third hypothesis (evaluate if M_{crit} increases with SWD), the effects of species group, tree mode of failure (both as factors), and fresh stem wood density (as independent variable) on M_{crit} were tested using analysis of covariance (ANCOVA). To check the fourth hypothesis (M_{crit} varies among tree mode of failure), individuals of all species groups were pooled, and the effects of tree mode of failure (categorical variable) and DBH (independent variable) on M_{crit} were tested using simple ANCOVA. To test which tree biometric parameter better predict M_{crit} , simple and multiple linear and nonlinear regressions were carried out separately for each species groups and for species groups pooled together.

3. Results

A total of 60 trees ranging from 19 cm to 41.1cm in DBH were winched from August 2014 to April 2015 totaling 76,229 kg of above ground biomass. The mode of failure (snapping and uprooting) differed significantly between the species groups. All *Scleronema micranthum* trees snapped in both topographic positions. For *Eschweilera* spp., equal number of trees in the valleys were snapped and uprooted, while in the plateau 60% failed by snapping and 40% by uprooting. The random set of trees had equal numbers of failures in both modes in the valley, but 40% snapped versus 60% uprooted on the plateau. Topographic position was not a significant predictor of Critical turning moment (M_{crit}) of trees in our study for any species groups tested, even when all species were pooled. According to One-way ANOVAs, topography had no effect on M_{crit} for random species ($p = 0.45$), for *S. micranthum* ($p = 0.70$), for *Eschweilera* spp. ($p = 0.73$), or for all species groups pooled ($p = 0.72$). Results from subsequent ANOVAs with DBH are presented in figure 1.

Figure 1. Analysis of variance (ANOVA) using topographic position as factor and DBH as independent variable on critical turning moments for species groups and for species pooled together with regression lines and 95% confidence envelopes.

M_{crit} varied among species groups when measures of tree size (e.g. DBH, AGB and Slenderness) were used as independent variables in an ANCOVA analysis. The interaction terms between species group with DBH, AGB and Slenderness were not significant ($p = 0.72, 0.12, 0.07$ respectively) with no evidence of different slopes. Nonetheless, the intercepts differ among groups, DBH, AGB and Slenderness showed significant increases in M_{crit} ($p < 0.005$). For a given DBH *Eschweilera* spp. was the most resistant group followed by *S. micranthum* and random species. When increasing AGB, *Eschweilera* spp. became more prone to failure (figure 2). The AGB distribution behavior for *Eschweilera* spp. together with tree Slenderness can be explained by differences in allometry among the other tested groups in relation to tree center of mass, stem and crown biomass. The interaction terms between species group with Stem and Center were non-significant ($p = 0.871, 0.2511$ respectively) with no evidence of different slopes. Nonetheless, the

intercepts differ among groups, Stem, Crown and Center showed significant increases in M_{crit} ($p < 0.005$) (figure 3). This difference in allometry, changes the tree center of mass of *Eschweilera* spp. to a higher point from the ground mainly by differences in crown mass, and wind loading is expected to occur at a higher effective height (Hale et al., 2012)

Figure 2. Analysis of covariance (ANCOVA) using species group as factor and DBH (left panel), AGB (right panel) and slenderness (lower panel) as independent variable on critical turning moments with regression lines and 95% confidence envelopes.

Figure 3. Analysis of covariance (ANCOVA) using species group as factor and Stem (left panel), Crown (right panel) and Center (lower panel) as independent variable on critical turning moments with regression lines and 95% confidence envelopes.

Pos-hoc Tukey tests were performed to evaluate differences among species groups (Table 1). Variance inflation factors (VIF) were assessed to check multicollinearity among the previous mentioned size attributes with species groups. VIF values were 4.83, 2.44, 4.24, 2.88, 1.70 and 2.37 for DBH, AGB, Slenderness, Stem, Crown and Center. Note that DBH values were fixed around values between 20 cm to 40 cm for all sampled species groups as mentioned in section 2.4.

Table 1. Pos-hoc Tukey tests for species groups using DBH, AGB, Slenderness, Stem, Crown and Center.

The interaction terms between species group and mode of failure (both categorical variables) with fresh stem wood density were non-significant ($p = 0.63$ and 0.77 , respectively) but SWD showed effects on critical turning moment ($p = 0.036$) for species groups (mainly by large differences in the random set of trees, see Table S2) but no effect for mode of failure ($p = 0.08$). Multiple regression combining DBH and SWD showed an increase of about 40% in explained variance (R^2) related to the simple regression for the random set of trees (Table S15), but no increases were found for *Eschweilera* spp. (Table S13), nor were increases found for *Scleronema micranthum* (Table S14). Because the higher variance in SWD for the random set of tree species,

this group was analyzed separately from the others so the effect of SWD was not confounded with species groups. The graphical relationship among M_{crit} , SWD and the random set of tree species did not show a linear trend, but show increases in M_{crit} with increasing SWD values (figure 4).

Figure 4. Scatter plot of M_{crit} and SWD for the random species.

The presence of decay and rot in tree parts can change SWD values. For two similar sized trees in DBH of *Eperua glabriflora*, presented SWD values of 1.02 g cm^{-3} (38 cm in DBH) and 0.594 g cm^{-3} (35.3 cm in DBH). For the second tree, decay and rot could be noted over 6 m from the stem base. Among the architectural attributes related to tree resistance to wind, the measure of slenderness; quotient of height and diameter (total tree height divided by DBH, both variables in meters) have been used as a indices of tree stability for Scots pine (*Pinus sylvestris*) and Norway spruce (*Picea abies*) stands (Jelonek et al., 2013; Mickovski et al., 2005; Slodicak and Novak, 2006). Information's about Stem, Crown and Slenderness, to further explore other tree architecture attributes, were included in graphical relationships among M_{crit} , SWD and the random species (figure 5). Trees can have different above ground biomass allocation patterns. Thus the Slenderness factor can be influenced by the relationship among crown biomass and stem biomass and consequently affect tree stability to with stand strong winds.

Figure 5. Scatter plot of M_{crit} and SWD for the random species with correspondent slenderness factor (top panel), stem mass in kg (left panel) and crown mass in kg (right panel).

Critical turning moments were influenced by tree mode of failure with DBH as independent variable. The interaction term between species group with DBH was not significant ($p = 0.823$) with no evidence of different slopes. The intercepts differ among mode of failure, and DBH was

positively related to M_{crit} ($p < 0.001$). For different trees with the same DBH, more force is required to cause stem breakage (figure 6).

Figure 6. Analysis of covariance (ANCOVA) using tree mode of failure as factor, DBH as independent variable on critical turning moments, with regression lines and 95% confidence envelopes.

Values of M_{crit} were best predicted by different biometric parameters according to the species groups. For *Scleronema micranthum*, DBH was the best individual predictor (table S14). For *Eschweilera* spp., the random set of species and species groups pooled, the best individual predictor was AGB (tables S13, S15 and S16 respectively).

4. Discussion

Although the tree wind-damage was assessed in different studies, the correlation between topographic exposure and intensity of damage is unclear (Zhu et al., 2004). Direction of slope aspect and orientation of valleys with respect to the prevailing wind direction can have a considerable influence on the magnitude of wind speed experienced at a location, and in many storms, the highest wind speeds were found on ridge tops and on gentle slopes facing the wind (Quine and Gardiner, 2007). In our study, there was no overall difference in M_{crit} of trees on plateaus and valleys, independent of whether species groups were analyzed separately or pooled. We have evidence to reject our first hypothesis, as we found no differences in tree stability between valleys and plateaus. Other factors such as higher wind-exposure and differences in wind speed and direction in plateaus (upland) areas (Negrón-Juárez et al., 2014; Quine and Gardiner, 2007; Ruel et al., 2001) may explain the higher wind-related mortality reported for plateaus and top of slopes (Marra et al., 2014). Regressions of M_{crit} with tree size descriptors (i.e. DBH and AGB) showed that the topographic position did not influence tree resistance to failure, and therefore differences in soil texture between valleys and plateaus (Chauvel et al., 1987; Luizão et al., 2004) do not influence tree resistance to failure. In same soil condition, when the influence of slope in anchorage was analyzed

for 40-year-old Sitka spruce (*Picea sitchensis*), more force was need to pull trees upslope then downslope. However no overall difference in anchorage was found between trees grown on the horizontal and sloping parts of the site (Nicoll et al., 2005). Studies that used crown architecture improve the understanding of dynamic response of standing trees to wind (Hale et al., 2012), but in Central Amazon forests calculating the average crown width based on measurements of maximum crown radius in eight directions was not possible due to difficulties defining neighbors for each sampled tree. Our winching experiment confirmed the effect of tree size on M_{crit} suggested by previous studies carried out in different forest types (i.e. structure and species composition), where tree stability increases substantially with increases in tree size (Cannon et al., 2015; Nicoll et al., 2006; Peltola, 2006). The same pattern was not confirmed for slenderness, for which values bellow 80 indicated higher stability against wind damage (Slodicak and Novak, 2006).

The different species groups investigated in this study suggest that there is a relation between mode of failure and tree biometric parameters. All individuals of *S. micranthum* snapped independent of topographic position and corresponding soil texture. This pattern suggests that tap root systems, typical of *S. micranthum*, partly influence tree mode of failure (see figure S1, supplementary material). In contrast, this pattern was not observed for *Eschweilera* spp. The lack of pattern in *Eschweilera* spp. can be attributed to the higher variation on biometric attributes from this genus, including the presence or absence of buttress depending on the species (Ribeiro *et al.*, 1999). Failure mode was closely linked to soil type for *Pinus radiata* in a range of New Zealand soils were 92% of trees uprooted in non-cohesive soils (Moore, 2000). In our study the same pattern was not observed, 40% of Random trees and 60% *Eschweilera* spp. uprooted on cohesive plateaus soil. In some cases, uprooting one tree can additionally uproot four surrounding trees when roots are interlocked (Coutts, 1983). Despite recent advances in the understanding of the relative roles of root components (lateral, superficial or tap roots) in tree anchorage strength (Fourcaud et al., 2008), general knowledge about roots architecture in tropical species is still scarce. Although this is the

first study to describe M_{crit} for Amazonian tree species, our results (tables S2 to S4) are in accord with previous studies (Peterson and Claassen 2013, Cannon *et al* 2015) for similar-sized hardwood and coniferous species of the United States. As in other static winching studies, M_{crit} of trees from this study also increased with tree size, supporting our second hypothesis. M_{crit} differs among Amazonian tree species, depending on the tree size (i.e., DBH, AGB, Slenderness, Stem, Crown, Center), and some species are simply more resistant than others (see figure 2 and figure 3). For instance, for a given DBH, *Eschweilera* spp. are more resistant than the other tested species. However, *Eschweilera* spp. trees with an AGB $> \sim 2,000$ kg are more susceptible to failure than similar AGB *S. micranthum*, probably due to the large contribution of crown biomass observed for this genus (see figure 2 left panel and figure 3 left panel). This result does not corroborate previous studies (Cannon et al., 2015; Nicoll et al., 2006; Peltola, 2006; Peterson and Claassen, 2013), in which no differences in M_{crit} between species were reported. One hectare in Amazon forest can hold more tree species than in all of Canada and the continental US (Coley and Kursar, 2014), thus there is a much larger range in interspecific differences.

In contrast with wood engineering studies, cell walls in living trees are fully water-saturated (Fournier et al., 2013), reducing the ability of wood to resist mechanical forces (Niklas and Spatz, 2010). Due to low variance in wood density within species, there was no significant difference in M_{crit} with wood density for *S. micranthum* and *Eschweilera* spp. (see tables S2 and S4). However, among the random set of tree species explained variance with single entry models, one linear and one non-linear, explained 0.28 and 0.31, respectively. But total explained variance increased from an R^2 of 0.49 for DBH alone, to 0.68 when including both DBH and SWD in a multiple regression model (model 12 and model 17, respectively, from table S15). As previously suggested (Niklas and Spatz, 2010, 2012), SWD has a strong effect on M_{crit} . Further work is needed to explore a larger range of wood density and its effect on the variation of M_{crit} together with tree architectural attributes. In this study, considering the random set of tree species, a non-linear relationship was

found, and species with higher SWD tends to exhibit higher values of M_{crit} . Other studies indicate that species with high wood density have more slender stems, but larger crowns than similar sized trees of low wood density species (Iida et al., 2012). Theoretical studies, in which the experiments have been conducted from the wood technological or timber use perspective (focusing on properties of dry wood), limit's the understanding of the biomechanics of living plants (Niklas and Spatz, 2010) and the understanding of wood characteristics from an ecological or evolutionary perspective (Larjavaara and Muller-Landau, 2012).

We found differences in M_{crit} related with tree mode of failure as a factor and DBH as independent variable. Based on these findings, we have strong evidence to reject our fourth hypothesis that M_{crit} does not vary with mode of failure. For a given DBH, more force is required to snap the stem than to uproot the tree. This result is in contrast with Peltola et al. (2000), but similar to Cannon et al. (2015). In the later study, tree modes of failure were reported with stem mass as independent variable of M_{crit} . This result suggests that tree mode of death in wind throw areas in Amazon forests depends on tree biometric parameters (e.g. root architecture, root: shoot ratio, crown architecture) and should be analyzed at the species level in future studies.

5. Conclusion

This mechanistic approach allows the comparison of tree vulnerability and resistance to snapping and uprooting across tropical and temperate forests and facilitates the use of current findings in the context of ecosystem models. Higher wind-induced tree mortality observed on plateaus and top of slopes may be explained by different wind speeds and gusts direction (valleys have different aspects and the wind can blow parallel or perpendicular), rather than by differences in soil-related factors that might effect M_{crit} . Further studies on root architecture can enhance our understanding about the vulnerability of Amazonian tree species to wind disturbance and consequent changes in tree species communities with greater storm intensity with a warming

climate. In this study, values of M_{crit} were governed by species-specific size attributes and vary among Amazonian tree species.

Acknowledgements

The authors gratefully thank the staff of EEST. This study was financed by the Brazilian Council for Scientific and Technological Development (CNPq) within the projects Succession After Windthrows (SAWI) (Chamada Universal MCTI/No14/2012, Proc. 473357/2012-7), INCT - Madeiras da Amazônia, and the Next Generation Ecosystem Experiment (NGEE) Tropics at Lawrence Berkeley National Laboratory, and also supported by the Max-Planck-Institute for Biogeochemistry within the Tree Assimilation and Carbon Allocation Physiology Experiment (TACAPE). Robinson Negrón-Juárez was supported by the Director, Office of Science, Office of Biological and Environmental Research of the US Department of Energy under Contract No. DE-AC02-05CH11231 as part of Next-Generation Ecosystems Experiments (NGEE Tropics) and the Regional and Global Climate Modeling (RGCM) and Programs. The authors declare no conflict of interest.

References

- Alméras, T., Fournier, M., 2009. Biomechanical design and long-term stability of trees: morphological and wood traits involved in the balance between weight increase and the gravitropic reaction. *J. Theor. Biol.* 256, 370–381. doi:10.1016/j.jtbi.2008.10.011
- Andrade, E.A., Higuchi, N., 2009. Produtividade de quatro espécies arbóreas de Terra Firme da Amazônia Central. *Acta Amaz.* 39, 105–112. doi:10.1590/S0044-59672009000100011
- Asner, G.P., 2013. Geography of forest disturbance. *Proc. Natl. Acad. Sci. U. S. A.* 110, 3711–2. doi:10.1073/pnas.1300396110
- Baldeck, C. A, Harms, K.E., Yavitt, J.B., John, R., Turner, B.L., Navarrete, H., Davies, S.J., Chuyong, G.B., Kenfack, D., Duncan, W., Madawala, S., Gunatilleke, N., Gunatilleke, S., Bunyavejchewin, S., Kiratiprayoon, S., Yaacob, A., Supardi, M.N.N., Dalling, J.W., Valencia, R., Thomas, D.W., 2012. Soil resources and topography shape local tree community structure in tropical forests. *Proc. R. Soc.* 280, 20122532. doi:10.1098/rspb.2012.2532
- Brüchert, F., Gardiner, B., 2006. THE EFFECT OF WIND EXPOSURE ON THE TREE

424 AERIAL ARCHITECTURE AND BIOMECHANICS OF SITKA SPRUCE (*PICEA*
425 *SITCHENSIS*, PINACEAE). *Am. J. Bot.* 93, 1512–1521.

426 Cannon, J.B., Barrett, M.E., Peterson, C.J., 2015. The effect of species, size, failure mode,
427 and fire-scarring on tree stability. *For. Ecol. Manage.* doi:10.1016/j.foreco.2015.07.014

428 Carneiro, V.M.C., 2004. COMPOSIÇÃO FLORÍSTICA E ANÁLISE ESTRUTURAL DA
429 FLORESTA PRIMÁRIA DE TERRA FIRME NA BACIA DO RIO CUIEIRAS,
430 MANAUS, AM. Instituto Nacional de Pesquisas da Amazônia.

431 Chambers, J.Q., Fisher, J.I., Zeng, H., Chapman, E.L., Baker, D.B., Hurtt, G.C., 2007.
432 Hurricane Katrina's carbon footprint on U.S. Gulf Coast forests. *Science* 318, 1107.
433 doi:10.1126/science.1148913

434 Chambers, J.Q., Higuchi, N., Teixeira, L.M., Dos Santos, J., Laurance, S.G., Trumbore,
435 S.E., 2004. Response of tree biomass and wood litter to disturbance in a Central
436 Amazon forest. *Oecologia* 141, 596–614. doi:10.1007/s00442-004-1676-2

437 Chambers, J.Q., Negrón-Juárez, R.I., Hurtt, G.C., Marra, D.M., Higuchi, N., 2009. Lack of
438 intermediate-scale disturbance data prevents robust extrapolation of plot-level tree
439 mortality rates for old-growth tropical forests. *Ecol. Lett.* 12, E22–E25.
440 doi:10.1111/j.1461-0248.2009.01398.x

441 Chambers, J.Q., Negrón-Juarez, R.I., Marra, D.M., Di Vittorio, A., Tews, J., Roberts, D.,
442 Ribeiro, G.H.P.M., Trumbore, S.E., Higuchi, N., 2013. The steady-state mosaic of
443 disturbance and succession across an old-growth Central Amazon forest landscape.
444 *Proc. Natl. Acad. Sci. U. S. A.* 110, 3949–54. doi:10.1073/pnas.1202894110

445 Chao, K.-J., Phillips, O.L., Gloor, E., Monteagudo, A., Torres-Lezama, A., Martínez, R.V., 2008.
446 Growth and wood density predict tree mortality in Amazon forests. *J. Ecol.* 96, 281–292.
447 doi:10.1111/j.1365-2745.2007.01343.x

448 Chao, K.-J., Phillips, O.L., Monteagudo, A., Torres-Lezama, A., Vásquez Martínez, R., 2009. How
449 do trees die? Mode of death in northern Amazonia. *J. Veg. Sci.* 20, 260–268.
450 doi:10.1111/j.1654-1103.2009.05755.x

451 Chauvel, A., Lucas, Y., Boulet, R., 1987. On the genesis of the soil mantle of the region of
452 Manaus, Central Amazonia, Brazil. *Experientia* 43, 234–241.
453 doi:10.1007/BF01945546

454 Coley, P.D., Kursar, T. a., 2014. On Tropical Forests and Their Pests. *Science* (80-.). 343,
455 35–36. doi:10.1126/science.1248110

456 Coutts, M.P., 1983. Root architecture and tree stability. *Plant Soil* 71, 171–188.

457 Cucchi, V., Meredieu, C., Stokes, A., Berthier, S., Bert, D., Najar, M., Denis, A., Lastennet,
458 R., 2004. Root anchorage of inner and edge trees in stands of Maritime pine (*Pinus*
459 *pinaster* Ait.) growing in different podzolic soil conditions. *Trees* 18, 460–466.
460 doi:10.1007/s00468-004-0330-2

461 Dale, V.H., Joyce, L.A., McNulty, S., Neilson, R.P., Ayres, M.P., Flannigan, M.D., Hanson,
462 P.J., Irland, L.C., Lugo, A.E., Peterson, C.J., Simberloff, D., Swanson, F.J., Stocks,
463 B.J., Wotton, B.M., 2001. Climate change and forest disturbances. *Bioscience* 51,
464 723–734 ST – Climate change and forest disturbanc. doi:10.1641/0006-
465 3568(2001)051[0723:CCAFD]2.0.CO;2

466 Denslow, J.S., 1980. Patterns of plant species diversity during succession under different
467 disturbance regimes. *Oecologia* 46, 18–21. doi:10.1007/BF00346960

468 dos Santos, L.T., Magnabosco Marra, D., Trumbore, S., Camargo, P.B., Chambers, J.Q.,
469 Negrón-Juárez, R.I., Lima, A. J.N., Ribeiro, G.H.P.M., dos Santos, J., Higuchi, N.,
470 2015. Windthrows increase soil carbon stocks in a Central Amazon forest.
471 *Biogeosciences Discuss.* 12, 19351–19372. doi:10.5194/bgd-12-19351-2015

472 Espírito-Santo, F.D.B., Gloor, M., Keller, M., Malhi, Y., Saatchi, S., Nelson, B., Junior,
473 R.C.O., Pereira, C., Lloyd, J., Frolking, S., Palace, M., Shimabukuro, Y.E., Duarte, V.,
474 Mendoza, A.M., López-González, G., Baker, T.R., Feldpausch, T.R., Brien, R.J.W.,
475 Asner, G.P., Boyd, D.S., Phillips, O.L., 2014. Size and frequency of natural forest
476 disturbances and the Amazon forest carbon balance. *Nat. Commun.* 5, 3434.
477 doi:10.1038/ncomms4434

478 Espírito-Santo, F.D.B., Keller, M., Braswell, B., Nelson, B.W., Frolking, S., Vicente, G.,
479 2010. Storm intensity and old-growth forest disturbances in the Amazon region.
480 *Geophys. Res. Lett.* 37. doi:10.1029/2010GL043146

481 Fourcaud, T., Ji, J.N., Zhang, Z.Q., Stokes, A., 2008. Understanding the impact of root
482 morphology on overturning mechanisms: A modelling approach. *Ann. Bot.* 101,
483 1267–1280. doi:10.1093/aob/mcm245

484 Fournier, M., Dlouhá, J., Jaouen, G., Alméras, T., 2013. Integrative biomechanics for tree
485 ecology: beyond wood density and strength. *J. Exp. Bot.* 60, 4397–4410.
486 doi:10.1093/jxb/ert279

487 Fredericksen, T.S., Hedden, R.L., Williams, S.A., 1993. Testing loblolly pine wind firmness
488 with simulated wind stress. *Can. J. For. Res.* 23, 1760–1765.

489 Garstang, M., White, S., Shugart, H.H., Halverson, J., 1998. Convective cloud downdrafts
490 as the cause of large blowdowns in the Amazon rainforest. *Meteorol. Atmos. Phys.* 67,
491 199–212. doi:10.1007/BF01277510

492 Guillaumet, J.L., 1987. Some structural and floristic aspects of the forest. *Experientia* 43,
493 241–251. doi:10.1007/BF01945547

494 Hale, S.E., Gardiner, B. a., Wellpott, A., Nicoll, B.C., Achim, A., 2012. Wind loading of
495 trees: Influence of tree size and competition. *Eur. J. For. Res.* 131, 203–217.
496 doi:10.1007/s10342-010-0448-2

497 Higuchi, F.G., Siqueira, J.D.P., Lima, A.J.N., Filho, A.F., Higuchi, N., 2012. Influência do
498 tamanho da parcela na precisão da função de distribuição diamétrica de weibull na
499 floresta primária da amazônia central. *Floresta* 42, 599–606.

500 Iida, Y., Poorter, L., Sterck, F.J., Kassim, A.R., Kubo, T., Potts, M.D., Kohyama, T.S.,
501 2012. Wood density explains architectural differentiation across 145 co-occurring
502 tropical tree species. *Funct. Ecol.* 26, 274–282. doi:10.1111/j.1365-2435.2011.01921.x

503 IPCC, 2013. IPCC - Climate Change 2013: The Physical Science Basis. Contribution of
504 Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on
505 Climate Change ed Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J.
506 Boschung, A. Nauels,.

507 Jelonek, T., Walkowiak, R., Jakubowski, M., Tomczak, A., 2013. Indices of tree stability in

508 Scots pine stands damaged by wind. *Sylwan* 157, 323–329.

509 Kamimura, K., Kitagawa, K., Saito, S., Mizunaga, H., 2012. Root anchorage of hinoki
510 *Chamaecyparis obtuse* (Sieb . Et Zucc .) Endl .) under the combined loading of wind
511 and rapidly supplied water on soil□: analyses based on tree-pulling experiments. *Eur.*
512 *J. For. Res.* 131, 219–227. doi:10.1007/s10342-011-0508-2

513 Larjavaara, M., Muller-Landau, H.C., 2012. Still rethinking the value of high wood density.
514 *Am. J. Bot.* 99, 165–8. doi:10.3732/ajb.1100324

515 Laurance, S.G.W., Laurance, W.F., Andrade, A., Fearnside, P.M., Harms, K.E., Vicentini,
516 A., Luizão, R.C.C., 2010. Influence of soils and topography on Amazonian tree
517 diversity: A landscape-scale study. *J. Veg. Sci.* 21, 96–106. doi:10.1111/j.1654-
518 1103.2009.01122.x

519 Lenart, M.T., Falk, D.A., Scatena, F.N., Osterkamp, W.R., 2010. Estimating soil turnover
520 rate from tree uprooting during hurricanes in Puerto Rico. *For. Ecol. Manage.* 259,
521 1076–1084. doi:10.1016/j.foreco.2009.12.014

522 Lima, A.J.N., 2010. Avaliação de um sistema de inventário florestal contínuo em áreas
523 manejadas e não manejadas do estado do Amazonas (AM). Ph.D. Diss. Univ. Fed. do
524 Amaz. Manaus, Brazil, p. 183 (in Port. with English Abstr. Instituto Nacional de
525 Pesquisas da Amazônia.

526 Luizão, R.C.C., Luizão, F.J., Paiva, R.Q., Monteiro, T.F., Sousa, L.S., Kruijt, B., 2004.
527 Variation of carbon and nitrogen cycling processes along a topographic gradient in a
528 central Amazonian forest. *Glob. Chang. Biol.* 10, 592–600. doi:10.1111/j.1529-
529 8817.2003.00757.x

530 Lundström, T., Jonas, T., Stöckli, V., Ammann, W., 2007. Anchorage of mature conifers:
531 resistive turning moment, root-soil plate geometry and root growth orientation. *Tree*
532 *Physiol.* 27, 1217–1227. doi:10.1093/treephys/27.9.1217

533 Marra, D.M., Chambers, J.Q., Higuchi, N., Trumbore, S.E., Ribeiro, G.H.P.M., Dos Santos,
534 J., Negrón-Juárez, R.I., Reu, B., Wirth, C., 2014. Large-scale wind disturbances
535 promote tree diversity in a Central Amazon forest. *PLoS One* 9, 1–17.
536 doi:10.1371/journal.pone.0103711

537 Mickovski, S.B., Stokes, A., Van Beek, L.P.. H., 2005. A decision support tool for
538 windthrow hazard assessment and prevention. *For. Ecol. Manage.* 216, 64–76.
539 doi:10.1016/j.foreco.2005.05.043

540 Mitchell, S.J., 2013. Wind as a natural disturbance agent in forests: A synthesis. *Forestry*
541 00, 1–11. doi:10.1093/forestry/cps058

542 Moore, J.R., 2000. Differences in maximum resistive bending moments of *Pinus radiata*
543 trees grown on a range of soil types. *For. Ecol. Manage.* 135, 63–71.

544 Negrón-Juárez, R., Baker, D.B., Chambers, J.Q., Hurtt, G.C., Goosem, S., 2014. Multi-
545 scale sensitivity of Landsat and MODIS to forest disturbance associated with tropical
546 cyclones. *Remote Sens. Environ.* 140, 679–689. doi:10.1016/j.rse.2013.09.028

547 Negrón-Juárez, R.I., Chambers, J.Q., Guimaraes, G., Zeng, H., Raupp, C.F.M., Marra,
548 D.M., Ribeiro, G.H.P.M., Saatchi, S.S., Nelson, B.W., Higuchi, N., 2010. Widespread
549 Amazon forest tree mortality from a single cross-basin squall line event. *Geophys.*

550 Res. Lett. 37, L16701. doi:10.1029/2010GL043733

551 Nelson, B.W., Kapos, V., Adams, J.B., Oliveira, W.J., Oscar, P.G., 1994. Forest
552 Disturbance by Large Blowdowns in the Brazilian Amazon. *Ecology* 75, 853–858.

553 Nicoll, B.C., Achim, A., Mochan, S., Gardiner, B.A., 2005. Does steep terrain influence
554 tree stability? A field investigation. *Can. J. For. Res.* 35, 2360–2367.
555 doi:10.1139/X05-157

556 Nicoll, B.C., Gardiner, B.A., Rayner, B., Peace, A.J., 2006. Anchorage of coniferous trees
557 relation to species, soil type and rooting depth. *Can. J. For. Res.* 36, 1871–1883.
558 doi:10.1139/x06-072

559 Niklas, K.J., Spatz, H.-C., 2012. Mechanical properties of wood disproportionately increase
560 with increasing density. *Am. J. Bot.* 99, 169–170. doi:10.3732/ajb.1100567

561 Niklas, K.J., Spatz, H.-C., 2010. Worldwide correlations of mechanical properties and
562 green wood density. *Am. J. Bot.* 97, 1587–1594. doi:10.3732/ajb.1000150

563 Nock, C.A., Geihofer, D., Grabner, M., Baker, P.J., Bunyavejchewin, S., Hietz, P., 2009.
564 Wood density and its radial variation in six canopy tree species differing in shade-
565 tolerance in western Thailand. *Ann. Bot.* 104, 297–306. doi:10.1093/aob/mcp118

566 Oliveira A.A., D., Mori, S. a, 1999. A central Amazonian terra firme forest.I. High species
567 richness on poor soils. *Biodivers. Conserv.* 8, 1219–1244.

568 Peltola, H., Ikonen, V.P., Gregow, H., Strandman, H., Kilpeläinen, a., Venäläinen, A.,
569 Kellomäki, S., 2010. Impacts of climate change on timber production and regional
570 risks of wind-induced damage to forests in Finland. *For. Ecol. Manage.* 260, 833–845.
571 doi:10.1016/j.foreco.2010.06.001

572 Peltola, H., Kellomäki, S., Hassinen, A., Granander, M., 2000. Mechanical stability of
573 Scots pine , Norway spruce and birch: an analysis of tree-pulling experiments in
574 Finland. *For. Ecol. Manage.* 135, 143–153.

575 Peltola, H.M., 2006. Mechanical stability of trees under static loads. *Am. J. Bot.* 93, 1501–
576 11. doi:10.3732/ajb.93.10.1501

577 Peterson, C.J., Claassen, V., 2013. An evaluation of the stability of *Quercus lobata* and
578 *Populus fremontii* on river levees assessed using static winching tests. *Forestry* 86,
579 201–209. doi:10.1093/forestry/cps080

580 Prance, G.T., Rodrigues, W.A., da Silva, M.F., 1976. Inventário florestal de um hectare de
581 mata de terra firme km 30 da estrada Manaus - Itacoatiara. *Acta Amaz.* 6, 9–35.

582 Putz, F.E., Coley, P.D., Lu, K., Montalvo, A., Aiello, A., 1983. Uprooting and snapping of
583 trees: structural determinants and ecological consequences. *Can. J. For. Res.*

584 Quesada, C. A., Lloyd, J., Schwarz, M., Patiño, S., Baker, T.R., Czimczik, C., Fyllas, N.M.,
585 Martinelli, L., Nardoto, G.B., Schmerler, J., Santos, a. J.B., Hodnett, M.G., Herrera,
586 R., Luizão, F.J., Arneith, A., Lloyd, G., Dezzio, N., Hilke, I., Kuhlmann, I., Raessler,
587 M., Brand, W. a., Geilmann, H., Filho, J.O.M., Carvalho, F.P., Filho, R.N.A., Chaves,
588 J.E., Cruz, O.F., Pimentel, T.P., Paiva, R., 2010. Variations in chemical and physical
589 properties of Amazon forest soils in relation to their genesis. *Biogeosciences* 7, 1515–
590 1541. doi:10.5194/bg-7-1515-2010

- Quesada, C.A., Lloyd, J., Schwarz, M., Baker, T.R., Phillips, O.L., Patino, S., Czimczik, C., Hodnett, M.G., Herrera, R., Arneeth, A., Lloyd, G., Malhi, Y., Dezzeo, N., Luizao, F.J., Santos, A.J.B., Schmerler, J., Arroyo, L., Silveira, M., Priante Filho, N., Jimenez, E.M., Paiva, R., Vieira, I., Neill, D.A., Silva, N., Penuela, M.C., Monteagudo, A., Vasquez, R., Prieto, A., Rudas, A., Almeida, S., Higuchi, N., Lezama, A.T., Lopez-Gonzalez, G., Peacock, J., Fyllas, N.M., Alvarez Davila, E., Erwin, T., di Fiore, A., Chao, K.J., Honorio, E., Killeen, T., Pena Cruz, A., Pitman, N., Nunez Vargas, P., Salomao, R., Terborgh, J., Ramirez, H., 2009. Regional and large-scale patterns in Amazon forest structure and function are mediated by variations in soil physical and chemical properties. *Biogeosciences Discuss* 6, 3993–4057.
- Quine, C.P., Gardiner, B. a, 2007. Understanding How the Interaction of Wind and Trees Results in Windthrow, Stem Breakage, and Canopy Gap Formation, in: *Plant Disturbance Ecology the Process and the Response*. pp. 103–156. doi:10.1016/B978-0-12-088778-1.50006-6
- Rankin de Mérona, J.M., Prance, G.T., Hutchings, R.W., Silva, M.F. da, Rodrigues, W.A., Uehling, M.E., 1992. Preliminary results of a large-scale tree inventory of the upland rain forest in the Central Amazon. *Acta Amaz.* 22, 493–534.
- Rennó, C.D., Nobre, C.A., Cuartas, L.A., Soares, J. V, Hodnett, M.G., Tomasella, J., Waterloo, M.J., 2008. Remote Sensing of Environment HAND , a new terrain descriptor using SRTM-DEM: Mapping terra- firme rainforest environments in Amazonia. *Remote Sens. Environ.* 112, 3469–3481. doi:10.1016/j.rse.2008.03.018
- Ribeiro, J. E. L. S., Hopkins, M. J. G., Vicentini, A., Sothers, C. A., Costa, M. A. da S., Brito, J.M., de, Souza, M. A. D. de, Martins, L. H. P., Lohmann, L. G., Assunção, P. A. C. L., Pereira, E., da C., Silva, C. F. da, Mesquita, M. R., and Procópio, L.C., 1999. *Flora da Reserva Ducke: Guia de Identificação das Plantas Vasculares de uma Floresta de Terra-firme na Amazônia Central*. INPA, Manaus.
- Ruel, J.C., Pin, D., Cooper, K., 2001. Windthrow in riparian buffer strips: Effect of wind exposure, thinning and strip width. *For. Ecol. Manage.* 143, 105–113. doi:10.1016/S0378-1127(00)00510-7
- Schelhaas, M.-J., Nabuurs, G.-J., Schuck, A., 2003. Natural disturbances in the European forests in the 19th and 20th centuries. *Glob. Chang. Biol.* 9, 1620–1633. doi:10.1046/j.1529-8817.2003.00684.x
- Sellier, D., Fourcaud, T., 2009. Crown structure and wood properties: Influence on tree sway and response to high winds. *Am. J. Bot.* 96, 885–896. doi:10.3732/ajb.0800226
- Silva, R.P. da, dos Santos, J., Tribuzy, E.S., Chambers, J.Q., Nakamura, S., Higuchi, N., 2002. Diameter increment and growth patterns for individual tree growing in Central Amazon, Brazil. *For. Ecol. Manage.* 166, 295–301. doi:10.1016/S0378-1127(01)00678-8
- Silva, R.P. da, Nakamura, S., Azevedo, C.P. de, Rocha, R.D.M., Carlos, A., Pinto, M., 2003. *Uso De Banda Dendrométrica Na Definição Diâmetro De Árvores Da Bacia Do Rio Cuieiras Use of Metallic Dendrometers for Individual Diameter Growth Patterns of Trees At Cuieiras River Basin*. *Acta Amaz.* 33, 67–84.
- Slodicak, M., Novak, J., 2006. Silvicultural measures to increase the mechanical stability of

634 pure secondary Norway spruce stands before conversion. *For. Ecol. Manage.* 224,
635 252–257. doi:10.1016/j.foreco.2005.12.037

636 Sterck, F.J., Bongers, F., 1998. ONTOGENETIC CHANGES IN SIZE , ALLOMETRY ,
637 AND MECHANICAL DESIGN OF TROPICAL RAIN FOREST TREES. *Am. J. Bot.*
638 85, 266–272.

639 ter Steege, H., Pitman, N.C.A., Sabatier, D., Baraloto, C., Salomão, R.P., Guevara, J.E.,
640 Phillips, O.L., Castilho, C. V, Magnusson, W.E., Molino, J.-F., Monteagudo, A.,
641 Núñez Vargas, P., Montero, J.C., Feldpausch, T.R., Coronado, E.N.H., Killeen, T.J.,
642 Mostacedo, B., Vasquez, R., Assis, R.L., Terborgh, J., Wittmann, F., Andrade, A.,
643 Laurance, W.F., Laurance, S.G.W., Marimon, B.S., Marimon, B.-H., Guimarães
644 Vieira, I.C., Amaral, I.L. do, Brien, R., Castellanos, H., Cárdenas López, D.,
645 Duivenvoorden, J.F., Mogollón, H.F., Matos, F.D.D.A., Dávila, N., García-Villacorta,
646 R., Stevenson Diaz, P.R., Costa, F., Emilio, T., Levis, C., Schietti, J., Souza, P.,
647 Alonso, A., Dallmeier, F., Montoya, A.J.D., Fernandez Piedade, M.T., Araujo-
648 Murakami, A., Arroyo, L., Gribel, R., Fine, P.V.A., Peres, C.A., Toledo, M., C, A., A,
649 G., Baker, T.R., Cerón, C., Engel, J., Henkel, T.W., Maas, P., Petronelli, P., Stropp, J.,
650 Zartman, C.E., Daly, D., Neill, D., Silveira, M., Paredes, M.R., Chave, J., Lima Filho,
651 D.D.A., Jørgensen, P.M., Fuentes, A., Schöngart, J., Cornejo Valverde, F., Di Fiore,
652 A., Jimenez, E.M., Peñuela-Mora, M.C., Phillips, J.F., Rivas, G., van Andel, T.R., von
653 Hildebrand, P., Hoffman, B., Zent, E.L., Malhi, Y., Prieto, A., Rudas, A., Ruschell,
654 A.R., Silva, N., Vos, V., Zent, S., Oliveira, A. a, Schutz, A.C., Gonzales, T., Trindade
655 Nascimento, M., Ramirez-Angulo, H., Sierra, R., Tirado, M., Umaña Medina, M.N.,
656 van der Heijden, G., Vela, C.I. a, Vilanova Torre, E., Vriesendorp, C., Wang, O.,
657 Young, K.R., Baider, C., Balslev, H., Ferreira, C., Mesones, I., Torres-Lezama, A.,
658 Urrego Giraldo, L.E., Zagt, R., Alexiades, M.N., Hernandez, L., Huamantupa-
659 Chuquimaco, I., Milliken, W., Palacios Cuenca, W., Pauletto, D., Valderrama
660 Sandoval, E., Valenzuela Gamarra, L., Dexter, K.G., Feeley, K., Lopez-Gonzalez, G.,
661 Silman, M.R., 2013. Hyperdominance in the Amazonian tree flora. *Science* 342,
662 1243092. doi:10.1126/science.1243092

663 Toledo, J.J., Magnusson, W.E., Castilho, C. V., Nascimento, H.E.M., 2012. Tree mode of
664 death in Central Amazonia: Effects of soil and topography on tree mortality associated
665 with storm disturbances. *For. Ecol. Manage.* 263, 253–261.
666 doi:10.1016/j.foreco.2011.09.017

667 Trumbore, S., Brando, P., Hartmann, H., 2015. Forest health and global change. *Science*
668 (80-.). 349, 814–818.

669 Turner, M.G., Baker, W.L., Peterson, C.J., Peet, R.K., 1998. Factors Influencing
670 Succession: Lessons from Large, Infrequent Natural Disturbances. *Ecosystems* 1, 511–
671 523. doi:10.1007/s100219900047

672 van Gelder, H.A., Poorter, L., Sterck, F.J., 2006. Wood mechanics, allometry, and life-
673 history variation in a tropical rain forest tree community. *New Phytol.* 171, 367–378.
674 doi:10.1111/j.1469-8137.2006.01757.x

675 Vieira, S., de Camargo, P.B., Selhorst, D., da Silva, R., Hutya, L., Chambers, J.Q., Brown,
676 I.F., Higuchi, N., dos Santos, J., Wofsy, S.C., Trumbore, S.E., Martinelli, L.A., 2004.
677 Forest structure and carbon dynamics in Amazonian tropical rain forests. *Oecologia*

678 140, 468–479. doi:10.1007/s00442-004-1598-z
679 Williamson, G.B., Wiemann, M.C., 2010. Measuring wood specific gravity...correctly. *Am.*
680 *J. Bot.* 97, 519–524. doi:10.3732/ajb.0900243
681 Zhu, J., Liu, Z., Li, X., Matsuzaki, T., Gonda, Y., 2004. Review: effects of wind on trees. *J.*
682 *For. Res.* 15, 153–160. doi:10.1007/BF02856753
683

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Table 1

Tukey multiple comparisons of means	DBH	AGB	Slenderness	Stem	Crown	Center
	p	p	p	p	p	p
Random and <i>Eschweilera</i> spp.	0.0034	0.3758	0.1923	0.0200	0.0865	0.1998
<i>S. micranthum</i> and <i>Eschweilera</i> spp.	0.7251	0.1951	0.9840	0.9611	0.9760	0.9844
<i>S. micranthum</i> and Random	0.0286	0.0084	0.1394	0.0101	0.0541	0.1459