New Quantum Chemistry-based Descriptors for Better Prediction of Melting point and Viscosity of Ionic Liquids

Amirhossein Mehrkesh* and Arunprakash T. Karunanithi

Center for Sustainable Infrastructure Systems, University of Colorado Denver, Denver, Colorado, 80217

Abstract

Ionic liquids (ILs) are a group of chemicals that possess negligible vapor pressure and have the potential to replace volatile organic compounds in industrial applications. A large number of ILs, through the combination of different cations and anions, can potentially be synthesized. In this context, it will be useful to intelligently design customized ILs through computer-aided methods. Practical limitations dictate that any successful attempt to design new ILs for industrial applications requires the ability to accurately predict their melting point and viscosity as experimental data will not be available for designed structures. In this paper, we present two new correlation equations towards the more precise prediction of melting point and viscosity of ILs solely based on the inputs from quantum chemistry calculations (no experimental data or simulation results are needed). To develop these correlations we utilized data related to size, shape, and electrostatic properties of cations and anions that constitutes ILs. In this work, new descriptors such as dielectric energy of cations and anions as well as the values predicted by an ‘ad-hoc’ model for the radii of cations and anions (instead of their van der waals radii) were used. An enormous form of correlation equations constituent of all different combinations of descriptors (as the inputs to the model) were tested. The average relative errors were measured to be 3.16% and 6.45% for the melting point, T_m , and $\ln(vis)$, respectively.

1. Introduction

Ionic liquids (ILs) are an emerging group of chemicals having interesting properties from negligible volatility, and high thermal stability to wide electrochemical window, and high solvency power. The

aforementioned characteristics make ionic liquids a promising candidate for multitude of applications. Ionic liquids are mainly composed of large organic cations and organic or inorganic anions. Physical properties of ionic liquids can significantly vary from their sister compounds, ionic salts. The properties of ionic salts can be completely and solely attributed to their ionic nature since strong ionic bonds hold their particles together. Ionic salts are mostly made of small monoatomic ions in the close vicinity of one and other in their lattice crystal network. The lattice energy of crystalline compounds is proportional to the inverse of the distance between their lattice points. The ionic bonds in salts are very strong due to the relatively short distance between the small ions contributing to high values of melting point. On the contrary, ionic liquids are made up of larger multi-atomic cations and anions resulting in weaker ionic bonds. This explains the considerably low melting point and viscosity of ionic liquids compared to those of ionic salts. The multi-atomic nature of cations and anions in ionic liquids is a great opportunity for researchers to fine tune their properties with customizing them for different applications. In ILs, cations, and occasionally anions, are composed of alkyl side chain groups (CH_2 , CH_3 , etc.) which can be accompanied with a number of functional groups (OH , NH_2 , COOH , etc.) to functionalize ionic liquids for different applications. A vast number of ILs (estimated to be about 10^{14})¹ can be potentially synthesized through distinct combinations of different cation-cores, alkyl groups, functional groups, and anions. Careful evaluation of experimental data from literature on the physical and thermodynamic properties of ILs shows that substituting functional groups can drastically alter a property of interest in ionic liquids.

Despite the fact that the ionic bonds in ILs are relatively weak, their properties can still be attributed to their ionic nature as even a weak ionic bond is still stronger than any other type of intermolecular forces. Crystal network of ionic materials (e.g. ionic liquids) are made of cations and anions held together by electrostatic attraction. The ionic force between charged particles is directly proportional to the charge of each particle and inversely to the distance between them. The larger the cations and anions are, the weaker the ionic bonds between them will be. This is due to the fact that by increasing the distance between two ions the electrostatic attraction, which holds them together, is reduced.

Computer-aided optimization methods can help in designing optimal ionic liquids suitable for different applications from solvent extraction to thermal energy storage.^{2,3} ILs are generally salts that are in liquid phase below an arbitrary temperature, commonly 100°C. Therefore, several ILs are in solid state at room temperature. When it comes to the melting point of ILs, those with significantly lower melting points or “room temperature ionic liquids [RTILs]” ($T_m < 25^\circ\text{C}$), are of great interest to researchers seeking new industrial applications. One main reason for the desirability of low melting point of ILs is the fact that they are being considered as candidates (solvents or absorbents) for separation tasks involving selective dissolution of solutes in gas (e.g. CO_2), liquid (e.g. toluene), or solid (e.g. cellulose) states. They are also widely considered as liquid solvents to promote chemical reactions.^{4,5} From a practical point of view, for an IL to be used as an industrial solvent it needs to be transported (pumped) across multiple unit operations and therefore it must be in the liquid phase.

Another significant barrier towards commercialization of IL-based applications is their high viscous characteristics due to their ionic nature (existence of relatively strong ionic bonds) resulting in transportation difficulties. This necessitates the use of powerful pumping equipment and efficient process equipment to handle viscous fluids. Therefore, identification and customization of ILs that have relatively low viscosity and appropriate melting point will greatly aid in their accelerated discovery and commercialization. Several models exist for prediction of melting point and viscosity of ILs. In general, these models are either Quantitative Structure Property Relationship (QSPR) correlations⁶⁻⁸, group contribution (GC) type⁹⁻¹⁶, or molecular dynamics (MD) simulations.¹⁷⁻²³ The main drawback of the QSPR and GC models is the narrow restrictions in choice of applicable compounds while the main limitation of MD simulations is the high computational demands and the need for starting geometry.²⁴ In this paper, we present two empirical correlations to predict melting point and viscosity of ILs in a way that does not need experimental input or tedious simulations, but rely on inputs from simple quantum chemical calculations.

2. Methods

Widely available data and research findings related to melting point and viscosity of salts explicitly shows that there is a meaningful relationship between the magnitude of the two properties and the lattice energy of ionic bonds. Generally, in an ionic compound, size (volume and area), shape (sphericity), molecular weight, and dielectric energy of ions, play important roles in determining the strength of an ionic bond. Normally, larger size of cations and anions, results in longer distances between these ions in the crystal lattice, making the ionic bond weaker. On the other hand, the shape of the ions is also important as they are better packed together when they are symmetrical in nature. It has been suggested that asymmetry of ions in an ionic compound, most likely, will decrease the melting point since the ions are more loosely connected and can be easily separated from each other (by applying lower amount of energy). When we studied the size of cations and anions of a wide variety of ionic liquids and compared with their melting points and viscosities, we found that there were several ILs that violated these general trends, i.e. ILs with relatively large cations and anions did not necessarily have low melting point or viscosity. One reason could be that in ionic liquids, ionic bonds are not the only intermolecular forces responsible for holding the particles together and other types of forces such as hydrogen bonding and polar-polar forces also come into play. Therefore we focused on using size, shape, and electrostatic properties to develop new correlations for the prediction of ILs' melting points and viscosities. In order to better account for deviations discussed above we included, in addition to the three descriptors, several other quantum chemical descriptors to refine the correlations to cover a wide variety of ionic liquids. Another challenge was that for several ILs, there were multiple experimental values reported in literature for these two physical properties. This inconsistency was especially observed in the case of melting points which could be due to the existence of impurities or different crystal network systems. In this study, we did not include ILs that had inconsistent experimental values, during the development of the correlations. Ionic liquids selected for this study were all 1:1 (one cation and one anion) with delocalized charges. These types of ILs are normally able to avoid crystallization and form glasses compounds far below room temperature.²⁵ Currently, the most widely used approach to predict the melting point of ILs is quantitative structure-property relationship (QSPR) models, mostly combined with artificial neural networks (ANNs).²⁵ In this

approach, there is a reasonably good correlation between actual and predicted melting points within a standard deviation of less than 10°C.²⁵ The limited availability of experimental data on physical properties of ILs is the main drawback of constructing good QSPR models. In recent years, simulations with molecular dynamics (MD) have evolved to study the behavior of ILs. The quality of these simulations strongly depends on the employed force fields. Several groups have tuned them specifically for ILs, while others have modified previously existing force fields. For example, Alavi and Thompson²⁵ have used MD simulations to predict the melting temperature of [C₂MIm][PF₆]. The demanding simulation indicated a melting point that was approximately 43°C too high.²⁵

Quantum chemistry data related to the constituents of the ILs, i.e. cations and anions, were obtained using TURBOMOLE software which is a powerful, general-purpose Quantum Chemistry (QC) program, for ab-initio electronic structure calculations.²⁶ This software allows for accurate prediction of cluster structures, conformational energies, excited states, and dipoles that can be utilized for different purposes. When a chemical compound, in our case a cation or an anion, is simulated using TURBOMOLE, it can be exported as a Cosmo file, which can be used in COSMOtherm software.²⁶ COSMOtherm is a universal tool, which combines quantum chemistry (QC) and thermodynamic laws to calculate different properties of chemicals and solvents.²⁷ This method can estimate the chemical potential of different molecules (in pure or mixed forms) at different temperatures. In contrast to other methods, COSMOtherm can predict thermodynamic properties of compounds as a function of concentration and temperature by equations, which are thermodynamically consistent.²⁷ In the this study, for all the cations and anions, their corresponding COSMO files were imported from COSMO*baseIL* database (Version 1501), provided by the Cosmologic Company, in which the compounds geometries are optimized, the ground state energies are calculated, and the corresponding COSMO and gasphase files are generated all using TZVP (BP_TZVP_C30_1501) basis set. In the next step, the computational chemistry information of the selected cations and anions, required for the development of the correlative equations, were collected from their COSMO files. These data were either used directly (without any changes) or as a combination

with other data (as the secondary descriptors) as the inputs to the correlative equations utilized to predict the melting point and viscosity of ILs.

3. Results and Discussion

Molar mass of cation (MW_C), molar mass of anion (MW_A), volume of anion $[A^\circ]^3$ (Vol_A), volume of cation $[A^\circ]^3$ (Vol_C), Area of anion $[A^\circ]^2$ ($Area_A$), Area of cation $[A^\circ]^2$ ($Area_C$), dielectric energy of cation (Di_C), dielectric energy of anion (Di_A), symmetrical values of ions (σ), density $[\frac{kg}{m^3}]$ (ρ) and the average distance between a cation and an anion in the network $[A^\circ]^3$ (R_t) were selected as descriptors to develop the proposed empirical correlations. Accurate measurements of volume, surface area and ionic radii of cations and anions are only possible through X-ray diffraction method. When X-ray data is unavailable correlative or approximation methods (e.g. van der Waals model) need to be used to estimate the characteristics that are related to the size and shape of ions. X-ray diffraction data is very sparse for the case of ILs and since we are interested in development of a universal correlation covering a wide range of IL structures, we have to rely on predictive data. Typical cations and anions of ILs, do not usually have spherical shapes, so it is necessary to estimate their ionic radii using correlative approaches. We selected ions for which the value of the experimental ionic radii were available. Next, for those selected ions, we also estimated the ionic radii using van der Waals model where all cations and anions are assumed to have a spherical shape. The volumes of corresponding ions were estimated using COSMOtherm software and were converted to ionic radii using Eqn.1.

$$R_{Calc} = \frac{3}{4} \times \frac{1}{\pi} \times Vol^{1/3} \quad \text{Eqn.1}$$

A linear correlation between the experimental values of the cations and anions radii and the corresponding values of their van der Waals radii is shown in Figure 1. As it can be seen from Figure 1, the actual and predicted ionic radii are perfectly correlated through a linear relationship ($R^2=0.98622$). The corresponding linear fit is shown in Eqn.2.

$$R_{Pred} = 0.48601 \times R_{Calc} - 0.01725$$

Eqn.2

Figure 1. A correlation between the van der Waals (VdW) and actual radii of ions in ILs

Although, there were only few experimental data points available for use in this correlation, we see that this data covers a wide range of VdW ionic radii from 3.8 to 7.5 Å⁰. Therefore, we were comfortable in using this linear relationship to estimate the unknown values of ionic radii of other cations and anions considered in this study. First, the ionic radius of each cation or anion was calculated using Eqn.1 from the volume calculated by COSMOtherm software. This value was corrected using Eqn.2 to arrive at the predicted ionic radii. The distance between anions and cations, R_t , were estimated as the sum of ionic radii of cations and anions present in the crystal network of ILs. The symmetrical value of ionic liquids, σ , were calculated using the sphericity of cations and anions. Sphericity is a measure of how spherical an object is and can be estimated using Eqn.3.

$$Sph = \frac{\pi^{\frac{1}{3}}(6V_p)^{\frac{1}{3}}}{A_p}$$

Eqn.3

where V_p and A_p are the volume and surface area, respectively.

Further, the symmetrical value of ionic liquids were calculated using Eqn.4.

$$\sigma = \sqrt{Sph_C * Sph_A} \quad \text{Eqn.4.}$$

where Sph_C and Sph_A are the sphericity of cations and anions in the IL of interest, respectively.

3.1 Melting point

Experimental melting point data of several ILs covering a wide range of categories (different type of cation cores, anions, functional groups etc.), were gathered from the literature.²⁸⁻⁴⁶ A multivariate correlation with inputs based on quantum chemistry descriptors, and the output of the experimental values of melting point, T_m , was performed utilizing Eureka software, a data analysis tool developed by Nutonian, Inc. Out of the 37 experimental melting point data, 17 (i.e. 45% of the data points) were chosen as part of the validation set (depicted in green in Figure 2), which were not used in the process of the model development.

The experimental data along with the correlated multivariate trend line, that was developed to predict the melting point of ILs as a function of their QC parameters, is shown in Figure 2. It can be seen from Figure 2 that the trend line strongly captures the fluctuations in the melting point data.

Figure 2. Experimental vs. model predicted melting point data (black-Training set; Green-Validation set)

Figure 3 presents a comparison between the experimental (observed) melting point data and the corresponding predictions based on the developed correlation.

Figure 3. Goodness of the model predictions for the melting point of ILs

The best multivariate correlation (having highest R^2) for predicting the melting point of ILs, shown in Eqn.5, uses the QC descriptors of the cation and anion radii, density of ionic liquids, symmetrical value, molecular weight of cation, and the dielectric energy of anion.

$$T_m = a(R_A) + b(\rho) + c(R_C)(\sigma) - d - e(MW_C) - f(Di_A) \quad (5)$$

$$a=11.38, b=0.05413, c=196.6, d=434.3, e=0.649, f=1661$$

Table 1 lists the characteristics related to the selected predictive empirical correlation.

Table 1. Model Accuracy for prediction of melting point

Parameter	Value-All data	Value-Test data	Value-Train data
R ² (Goodness of Fit)	0.91	---	---
Maximum Relative Error (%)	9.87	9.87	9.41
Avg. Relative Error (%)	3.16	2.77	3.48

3.2 Viscosity

Experimental data on the viscosity of several ILs at different temperatures, were gathered from the literature.⁴⁶⁻⁶⁵ Once more, a multivariate correlation with several quantum chemistry descriptors along with temperature as input parameters, and $\ln(\text{viscosity})$, as the output parameter, was developed using Eureqa software.

In this case, we used 131 viscosity data points, out of which, 83 were chosen as part of the validation set (depicted in green in Figure 4), which again were not used for model development purposes. The experimental viscosity data points and the correlated multivariate trend line that was developed to predict the viscosity of ILs as a function of QC descriptors, is shown in Figure 4. It can be seen from Figure 4 that, the trend line strongly captures the fluctuations in the viscosity of selected ILs.

Figure 4. Experimental vs. model predicted viscosity data (black-Training set; Green-Validation set)

Figure 5 presents a comparison between the experimental viscosity data and the corresponding predictions based on the developed correlation.

Figure 5. Goodness of the model for prediction of viscosity of ILs-test data only

The best multivariate correlation (having highest R^2) for viscosity, shown in Eqn.6, uses the descriptors of the symmetrical value, distance between anions and cations, anion volume and surface area, temperature and the dielectric energy of anion and cation.

$$\ln(\text{vis}) = a (\sigma) + b (R_t) + c (\text{Vol}_A) - d - e (\text{Area}_A) - f (T) - g (\text{Di}_A) - h (\text{Di}_C) \quad \text{Eqn.6}$$

$$a=16.513, b=2.2179, c=0.00892, d=15.0073, e=0.02686, f=0.02975, g=15.8297, h=48.1367$$

Table 2 lists the characteristics related to the empirical model used to predict the viscosity of ILs as a function of QC parameters and temperature.

It is noteworthy that the average error in the prediction of viscosity of different classes of ionic liquids (i.e. ionic liquids with different cation head groups at a wide range of temperature) using the semi-empirical model of this study was calculated to be 6.45%. This shows a significant improvement in the accuracy of the prediction compared to the model used in Preiss et al.²⁵ (which has ~18% absolute error for the same group of ionic liquids) study which is currently incorporated in COSMOTherm software.

Table 2. Model Accuracy for prediction of viscosity

Parameter	Value-All data	Value-Test data	Value-Train data
R^2 (Goodness of Fit)	0.91	---	---
Maximum Relative Error (%)	20.78	20.51	20.78
Avg. Relative Error (%)	6.45	7.40	4.81

4. Comparison

There have been several studies focusing⁹⁻¹⁶ on prediction of the physical properties of ionic liquids using group contribution (GC) approaches. In these methods, a value is attributed to each structural group (a cation head group, a side chain group attached to the cation and an anion). Optimization frameworks are used to regress experimental data points on the physical properties of chemicals, in this case ionic liquids, to equations comprising the weighted summation of the contribution of structural groups (i.e. number of each structural group multiply by its corresponding contribution). GC methods are limited to the

structural groups for which experimental data are available. Therefore, these methods, even if accurate, can only be used for the structural groups which were available in the initial set of data used to develop the model. This study is not to invalidate the group contribution models used predict viscosity and melting point of ionic liquids since these methods can be very useful in trend analysis in which the effect of different structural groups on physical properties can be studied. Only it is noteworthy that models errors in this study for prediction of two of the most important physical properties of ionic liquids (melting point and viscosity) are comparable to those achieved by group contribution models⁹⁻¹⁶ (i.e. less than 10% absolute error). The main difference is that the models in this study are solely *ab initio* meaning that there is no need for experimental data and they can be used to predict properties of all classes of ionic liquids even those for which no experimental data is available thereby GC models cannot be applied.

5. Conclusion

In this study, we present two multivariate empirical correlations to predict the melting point and viscosity of ILs with relatively high accuracy. Prediction of these two properties of ILs is critical as they have a significant influence on the practical applicability of ILs for industrial applications. The input parameters in the case of melting point (T_m) prediction were all quantum chemistry (QC) descriptors and in the case of viscosity were QC descriptors and temperature. These correlations can be utilized as long as the structure of a new ionic liquid is known and the quantum chemistry calculations can be performed to simulate the IL and estimate the descriptors related to size, shape, electrostatic energies, and the polarity which can be used to predict the macroscopic properties, melting point and viscosity.

Acknowledgement

This material is based upon work supported by the United States National Science Foundation (CAREER Program) under Grant No. 1151182.

Author Information

Corresponding Author

* E-mail: Amirhossein.mehrkehsh@ucdenver.edu

References

- (1) Rogers, R. D.; Seddon, K. R. Ionic liquids--solvents of the future? *Science* **2003**, 302(5646), 792-793.
- (2) Karunanithi, A. T.; Mehrkesh, A. Computer-aided design of tailor-made ionic liquids. *AIChE Journal* **2013**, 59, 4627-4640.
- (3) Mehrkesh, A.; Karunanithi, A. T. Optimal Design of Ionic Liquids for Thermal Energy Storage. *Under Review by the Journal of Computers & Chemical Engineering* **2015**.
- (4) Cole, A. C.; Jensen, J. L.; Ntai, I.; Tran, K. L. T.; Weaver, K. J.; Forbes, D. C.; Davis, J. H. Novel Brønsted acidic ionic liquids and their use as dual solvent-catalysts. *Journal of the American Chemical Society* **2002**, 124, 5962-5963.
- (5) Wilkes, J. S. Properties of ionic liquid solvents for catalysis. *Journal of Molecular Catalysis A: Chemical* **2004**, 214, 11-17.
- (6) Yan, F.; Xia, S.; Wang, Q.; Yang, Z.; Ma, P. Predicting the melting points of ionic liquids by the Quantitative Structure Property Relationship method using a topological index. *The Journal of Chemical Thermodynamics* **2013**, 62, 196-200.
- (7) Trohalaki, S.; Pachter, R.; Drake, G. W.; Hawkins, T. Quantitative structure-property relationships for melting points and densities of ionic liquids. *Energy & Fuels* **2005**, 19(1), 279-284.
- (8) Gharagheizi, F.; Mirkhani, S. A.; Keshavarz, M. H.; Farahani, N.; Tumba, K. A molecular-based model for prediction of liquid viscosity of pure organic compounds: A quantitative structure property relationship (QSPR) approach. *Journal of the Taiwan Institute of Chemical Engineers* **2013**, 44(3), 359-364.
- (9) Gardas, R. L.; Coutinho, J. A. A group contribution method for viscosity estimation of ionic liquids. *Fluid Phase Equilibria* **2008**, 266(1), 195-201.
- (10) Matsuda, H.; Yamamoto, H.; Kurihara, K.; Tochigi, K. Computer-aided reverse design for ionic liquids by QSPR using descriptors of group contribution type for ionic conductivities and viscosities. *Fluid Phase Equilibria* **2007**, 261(1), 434-443.
- (11) Gharagheizi, F.; Ilani-Kashkouli, P.; Mohammadi, A. H.; Ramjugernath, D.; Richon, D. Development of a group contribution method for determination of viscosity of ionic liquids at atmospheric pressure. *Chemical engineering science* **2012**, 80, 326-333.
- (12) Gharagheizi, F.; Ilani-Kashkouli, P.; Mohammadi, A. H. Computation of normal melting temperature of ionic liquids using a group contribution method. *Fluid Phase Equilibria* **2012**, 329, 1-7.
- (13) Lazzús, J. A. A group contribution method to predict the melting point of ionic liquids. *Fluid Phase Equilibria* **2012**, 313, 1-6.

- (14) Huo, Y.; Xia, S.; Zhang, Y.; Ma, P. Group contribution method for predicting melting points of imidazolium and benzimidazolium ionic liquids. *Industrial & engineering chemistry research* **2009**, 48(4), 2212-2217.
- (15) Aguirre, C. L.; Cisternas, L. A.; Valderrama, J. O. Melting-point estimation of ionic liquids by a group contribution method. *International Journal of Thermophysics* **2012**, 33(1), 34-46.
- (16) Gardas, R. L.; Coutinho, J. A. Group contribution methods for the prediction of thermophysical and transport properties of ionic liquids. *AIChE Journal* **2009**, 55(5), 1274-1290.
- (17) Alavi, S.; Thompson, D. L. Molecular dynamics studies of melting and some liquid-state properties of 1-ethyl-3-methylimidazolium hexafluorophosphate [emim][PF₆]. *The Journal of chemical physics* **2005**, 122(15), 154704.
- (18) Jayaraman, S.; Maginn, E. J. Computing the melting point and thermodynamic stability of the orthorhombic and monoclinic crystalline polymorphs of the ionic liquid 1-n-butyl-3-methylimidazolium chloride. *The Journal of chemical physics* **2007**, 127(21), 214504.
- (19) Batista, M. L.; Coutinho, J. A.; Gomes, J. R. Prediction of Ionic Liquids Properties through Molecular Dynamics Simulations. *Curr. Phys. Chem.* **2014**, 4, 151-172.
- (20) Ghatee, M. H.; Zare, M.; Moosavi, F.; Zolghadr, A. R. Temperature-dependent density and viscosity of the ionic liquids 1-alkyl-3-methylimidazolium iodides: experiment and molecular dynamics simulation. *Journal of Chemical & Engineering Data* **2010**, 55(9), 3084-3088.
- (21) Weingärtner, H. Understanding ionic liquids at the molecular level: facts, problems, and controversies. *Angewandte Chemie International Edition* **2008**, 47(4), 654-670.
- (22) Yan, T.; Li, S.; Jiang, W.; Gao, X.; Xiang, B.; Voth, G. A. Structure of the liquid-vacuum interface of room-temperature ionic liquids: a molecular dynamics study. *The Journal of Physical Chemistry B* **2006**, 110(4), 1800-1806.
- (23) Köddermann, T.; Paschek, D.; Ludwig, R. Molecular dynamic simulations of ionic liquids: A reliable description of structure, thermodynamics and dynamics. *Chem. Phys. Chem.* **2007**, 8(17), 2464-2470.
- (24) Preis, T.; Virnau, P.; Paul, W.; Schneider, J. J. GPU accelerated Monte Carlo simulation of the 2D and 3D Ising model. *Journal of Computational Physics* **2009**, 228(12), 4468-4477.
- (25) Preiss, U.; Bulut, S.; Krossing, I. In silico prediction of the melting points of ionic liquids from thermodynamic considerations: a case study on 67 salts with a melting point range of 337° C. *The Journal of Physical Chemistry B* **2010**, 114, 11133-11140.
- (26) TURBOMOLE V7.0 2015, a development of University of Karlsruhe and Forschungszentrum Karlsruhe GmbH, 1989-2007, TURBOMOLE GmbH, since 2007; available from <http://www.turbomole.com>

- (27) Eckert, F., and A. Klamt. "COSMOtherm Version C30, Release 14.01. COSMOlogic GmbH & Co. KG, Leverkusen, Germany." Available at www.cosmologic.de (2014).
- (28) Fredlake, C. P.; Crosthwaite, J. M.; Hert, D. G.; Aki, S. N.; Brennecke, J. F. Thermophysical properties of imidazolium-based ionic liquids. *Journal of Chemical & Engineering Data* **2004**, *49*, 954-964.
- (29) Hu, H. C.; Soriano, A. N.; Leron, R. B.; Li, M. H. Molar heat capacity of four aqueous ionic liquid mixtures. *Thermochimica Acta* **2011**, *519*, 44-49.
- (30) Vila, J.; Fernández-Castro, B.; Rilo, E.; Carrete, J.; Domínguez-Pérez, M.; Rodríguez, J. R.;... & Cabeza, O. Liquid–solid–liquid phase transition hysteresis loops in the ionic conductivity of ten imidazolium-based ionic liquids. *Fluid Phase Equilibria* **2012**, *320*, 1-10.
- (31) Bradley, J-C. **2011**, Alfa Aesar melting point data now openly available.
- (32) Hughes, T. J.; Syed, T.; Graham, B. F.; Marsh, K. N.; May, E. F. Heat capacities and low temperature thermal transitions of 1-hexyl and 1-octyl-3-methylimidazolium bis (trifluoromethylsulfonyl) amide. *Journal of Chemical & Engineering Data* **2011**, *56*, 2153-2159.
- (33) Lourenço, C.; Melo, C. I.; Bogel-Lukasik, R.; Bogel-Lukasik, E. Solubility advantage of pyrazine-2-carboxamide: Application of alternative solvents on the way to the future pharmaceutical development. *Journal of Chemical & Engineering Data* **2012**, *57*, 1525-1533.
- (34) Bradaric, C. J.; Downard, A.; Kennedy, C.; Robertson, A. J.; Zhou, Y. Industrial preparation of phosphonium ionic liquids. *Green Chemistry* **2003**, *5*, 143-152.
- (35) Coker, T. G.; Ambrose, J.; Janz, G. J. Fusion properties of some ionic quaternary ammonium compounds. *Journal of the American Chemical Society* **1970**, *92*, 5293-5297.
- (36) ChemFiles, Enabling Technologies-Ionic Liquids, Vol.5 No.6. www.sigmaaldrich.com/content/dam/sigma-aldrich/docs/Aldrich/Brochure/al_chemfile_v5_n6.pdf
- (37) Li, H.; Zhao, G.; Liu, F.; Zhang, S. Physicochemical Characterization of MF m–Based Ammonium Ionic Liquids. *Journal of Chemical & Engineering Data* **2013**, *58*, 1505-1515.
- (38) Verevkin, S. P.; Zaitsau, D. H.; Emel'yanenko, V. N.; Ralys, R. V.; Yermalayeu, A. V.; Schick, C. Does alkyl chain length really matter? Structure–property relationships in thermochemistry of ionic liquids. *Thermochimica Acta* **2013**, *562*, 84-95.
- (39) Nishida, T.; Tashiro, Y.; Yamamoto, M. Physical and electrochemical properties of 1-alkyl-3-methylimidazolium tetrafluoroborate for electrolyte. *Journal of Fluorine Chemistry* **2003**, *120*, 135-141.
- (40) Fletcher, S. I.; Sillars, F. B.; Hudson, N. E.; Hall, P. J. Physical properties of selected ionic liquids for use as electrolytes and other industrial applications. *Journal of Chemical & Engineering Data* **2009**, *55*, 778-782.

- (41) Wachter, P.; Schweiger, H. G.; Wudy, F.; Gores, H. J. Efficient determination of crystallization and melting points at low cooling and heating rates with novel computer controlled equipment. *The Journal of Chemical Thermodynamics* **2008**, *40*, 1542-1547.
- (42) Zhang, Z. H.; Tan, Z. C.; Li, Y. S.; Sun, L. X. Thermodynamic investigation of room temperature ionic liquid. *Journal of thermal analysis and calorimetry* **2006**, *85*, 551-557.
- (43) MacFarlane, D. R.; Meakin, P.; Amini, N.; Forsyth, M. Structural studies of ambient temperature plastic crystal ion conductors. *Journal of physics: condensed matter* **2001**, *13*, 8257.
- (44) Lourenço, C.; Melo, C. I.; Bogel-Lukasik, R.; Bogel-Lukasik, E. Solubility advantage of pyrazine-2-carboxamide: Application of alternative solvents on the way to the future pharmaceutical development. *Journal of Chemical & Engineering Data* **2012**, *57*, 1525-1533.
- (45) Liu, Q. S.; Yang, M.; Li, P. P.; Sun, S. S.; Welz-Biermann, U.; Tan, Z. C.; Zhang, Q. G. Physicochemical properties of ionic liquids [C₃py][NTf₂] and [C₆py][NTf₂]. *Journal of Chemical & Engineering Data* **2011**, *56*, 4094-4101.
- (46) Zhang, S.; Sun, N.; He, X.; Lu, X.; Zhang, X. Physical properties of ionic liquids: database and evaluation. *Journal of physical and chemical reference data* **2006**, *35*(4), 1475-1517.
- (47) Carvalho, P. J.; Regueira, T.; Santos, L. M.; Fernandez, J.; Coutinho, J. A. Effect of Water on the Viscosities and Densities of 1-Butyl-3-methylimidazolium Dicyanamide and 1-Butyl-3-methylimidazolium Tricyanomethane at Atmospheric Pressure†. *Journal of Chemical & Engineering Data* **2009**, *55*, 645-652.
- (48) Harris, K. R.; Kanakubo, M.; Woolf, L. A. Temperature and pressure dependence of the viscosity of the ionic liquid 1-butyl-3-methylimidazolium tetrafluoroborate: viscosity and density relationships in ionic liquids. *Journal of Chemical & Engineering Data* **2007**, *52*, 2425-2430.
- (49) Seddon, K. R.; Stark, A.; Torres, M. J. Influence of chloride, water, and organic solvents on the physical properties of ionic liquids. *Pure and Applied Chemistry* **2000**, *72*, 2275-2287.
- (50) Baker, S. N.; Baker, G. A.; Kane, M. A.; Bright, F. V. The cybotactic region surrounding fluorescent probes dissolved in 1-butyl-3-methylimidazolium hexafluorophosphate: effects of temperature and added carbon dioxide. *The Journal of Physical Chemistry B* **2001**, *105*, 9663-9668.
- (51) Tokuda, H.; Tsuzuki, S.; Susan, M. A. B. H.; Hayamizu, K.; Watanabe, M. How ionic are room-temperature ionic liquids? An indicator of the physicochemical properties. *The Journal of Physical Chemistry B* **2006**, *110*, 19593-19600.
- (52) Jacquemin, J.; Husson, P.; Padua, A. A.; Majer, V. Density and viscosity of several pure and water-saturated ionic liquids. *Green Chemistry* **2006**, *8*, 172-180.
- (53) Almeida, H. F.; Passos, H.; Lopes-da-Silva, J. A.; Fernandes, A. M.; Freire, M. G.; Coutinho, J. A. Thermophysical properties of five acetate-based ionic liquids. *Journal of Chemical & Engineering Data* **2012**, *57*, 3005-3013.

- (54) Ghatee, M. H.; Zare, M.; Moosavi, F.; Zolghadr, A. R. Temperature-dependent density and viscosity of the ionic liquids 1-alkyl-3-methylimidazolium iodides: experiment and molecular dynamics simulation. *Journal of Chemical & Engineering Data* **2010**, *55*, 3084-3088.
- (55) Freire, M. G.; Teles, A. R. R.; Rocha, M. A.; Schröder, B.; Neves, C. M.; Carvalho, P. J.; ... & Coutinho, J. A. Thermophysical characterization of ionic liquids able to dissolve biomass. *Journal of Chemical & Engineering Data* **2011**, *56*, 4813-4822.
- (56) Schreiner, C.; Zugmann, S.; Hartl, R.; Gores, H. J. Fractional walden rule for ionic liquids: examples from recent measurements and a critique of the so-called ideal KCl line for the Walden Plot†. *Journal of Chemical & Engineering Data* **2009**, *55*, 1784-1788.
- (57) Deive, F. J.; Rivas, M. A.; Rodríguez, A. Thermophysical properties of two ionic liquids based on benzyl imidazolium cation. *The Journal of Chemical Thermodynamics* **2011**, *43*, 487-491.
- (58) Restolho, J.; Serro, A. P.; Mata, J. L.; Saramago, B. Viscosity and surface tension of 1-ethanol-3-methylimidazolium tetrafluoroborate and 1-methyl-3-octylimidazolium tetrafluoroborate over a wide temperature range. *Journal of Chemical & Engineering Data* **2009**, *54*, 950-955.
- (59) Neves, C. M.; Carvalho, P. J.; Freire, M. G.; Coutinho, J. A. Thermophysical properties of pure and water-saturated tetradecyltrihexylphosphonium-based ionic liquids. *The Journal of Chemical Thermodynamics* **2011**, *43*, 948-957.
- (60) Crosthwaite, J. M.; Muldoon, M. J.; Dixon, J. K.; Anderson, J. L.; Brennecke, J. F. Phase transition and decomposition temperatures, heat capacities and viscosities of pyridinium ionic liquids. *The Journal of Chemical Thermodynamics* **2005**, *37*, 559-568.
- (61) Sanchez, L. G.; Espel, J. R.; Onink, F.; Meindersma, G. W.; Haan, A. B. D. Density, viscosity, and surface tension of synthesis grade imidazolium, pyridinium, and pyrrolidinium based room temperature ionic liquids. *Journal of Chemical & Engineering Data* **2009**, *54*, 2803-2812.
- (62) Gaciño, F. M.; Regueira, T.; Lugo, L.; Comuñas, M. J.; Fernández, J. Influence of molecular structure on densities and viscosities of several ionic liquids. *Journal of Chemical & Engineering Data* **2011**, *56*, 4984-4999.
- (63) Bhattacharjee, A.; Carvalho, P. J.; Coutinho, J. A. The effect of the cation aromaticity upon the thermophysical properties of piperidinium-and pyridinium-based ionic liquids. *Fluid Phase Equilibria* **2014**, *375*, 80-88.
- (64) Królikowska, M.; Zawadzki, M.; Królikowski, M. Physicochemical and thermodynamic study on aqueous solutions of dicyanamide-based ionic liquids. *The Journal of Chemical Thermodynamics* **2014**, *70*, 127-137.
- (65) Gaciño, F. M., Regueira, T., Lugo, L., Comuñas, M. J., & Fernández, J. Influence of molecular structure on densities and viscosities of several ionic liquids. *Journal of Chemical & Engineering Data* **2011** *56*(12), 4984-4999.