1
2
3 **Revised manuscript**
4
5
6
7

8 Distinct but milder phenotypes with choreiform movements in siblings with
9 compound heterozygous mutations in the transcription preinitiation
10 mediator complex subunit 17 (*MED17*)
11
12

13
14
15 Shinichi Hirabayashi^{a,*}, MD, Hirotomo Saitsu^b, MD, Naomichi Matsumoto^b,
16 MD
17
18

19
20 ^aDivision of Neurology, Nagano Children's Hospital, Azumino, Japan
21
22 ^bDepartment of Human Genetics, Yokohama City University Graduate
23 School of Medicine, Yokohama, Japan
24
25

26
27 *Corresponding author. Division of Neurology, Nagano Children's Hospital,
28 3100 Toyoshina, Azumino-shi, Nagano 399-8288, Japan. Tel.: +81 263 73
29 6700, Fax: +81 263 73 5432
30
31 e-mail address: shirarin@naganoch.gr.jp
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61

Abstract

Two siblings born to non-consanguineous parents showed nystagmus and sudden opitotonic posturing from the early infancy, and subsequent developmental delay and marked choreiform movements with hypotonia in the childhood. The brother had a mild postnatal microcephaly. Brain MRI of the sister showed mild delay of myelination, dilated anterior horn and mild cerebellar atrophy. Whole exome sequencing (WES) revealed compound heterozygous mutations in *MED17* gene in both siblings: c.1013-5A>G and c.1484T>G mutations transmitted from their father and mother, respectively. The c.1013-5A>G mutation caused insertion of 4 bases of intron 6 in the transcript, resulting in frameshift (p. Ser338Asnfs*15), and mutant transcript underwent nonsense-mediated mRNA decay in lymphoblastoid cells derived from two siblings. The c.1484T>G mutation substituted a leucine residue, which is highly conserved among the vertebrates, and was predicted to be damaging by *in silico* analysis programs. Both mutations were not registered in dbSNP data and in our 575 control exomes. These results suggest that the siblings' mutations are likely to be pathogenic.

This is the second case report concerning *MED17* mutations. Compared with the first reported cases of Caucasian Jewish origin, the clinical symptoms and courses are much milder and slower, respectively, in our cases. Genotype difference (a homozygous mutation versus compound heterozygous mutations) might explain these clinical differences between two cases, though early-onset nystagmus and later choreiform movements were unique in our cases. Clinical spectrum and phenotype-genotype correlations in this rare mutation should be further elucidated.

Key word: whole exome sequencing, mediator complex subunit 17, compound heterozygous mutation, choreiform movement, nystagmus

Introduction

Whole exome sequencing (WES) is now established as a useful diagnostic method for the identification of molecular defects in patients with genetic disorders. It has been reported that the diagnostic yields of patients, mainly children with neurological disorders, with potential genetic causes [1] and of those with severe intellectual disability [2] were 25% and 16% respectively.

We experienced two siblings with very similar neurological deficits suggestive of autosomal recessive inheritance, but without hallmarks of any clinically recognizable diseases. The WES revealed compound heterozygous mutations in the transcription preinitiation mediator complex subunit 17 gene (*MED17*) in both siblings. One homozygous mutation in *MED17* had been consistently found in five infants from four unrelated families of Caucasian Jewish origin [3]. We describe here the clinical and genetic features of the siblings and compare them with those of the reported cases.

Sibling 1 (II-1 in Figure 2A)

She was born at term without complications to non-consanguineous and healthy parents. Her head circumference (HC) was normal. Horizontal nystagmus and poor visual pursuit were noticed since the age of 3 months, which led her to our hospital at the age of 6 months. She was hypotonic with the decreased voluntary movements, but showed a sudden opisthotonic posturing. The deep tendon reflexes could not be elicited. The organomegaly was not detected. The ophthalmologic examinations and brain MRI, ABR, ERG, VEP were normal. The analysis of amino acids and organic acids were also normal.

At the age of 1 year-and-7 month, she could sit for a short time. The Gallant reflex remained positive. Her visual pursuit was improved with decreased nystagmus. The brain MRI at that time showed a mild delay of myelination at the frontal lobe, with the dilated bilateral anterior horn suggestive of the caudate atrophy, and with the mild atrophy of the anterior vermis. Soon after, she could not sit alone possibly because of the opisthotonic postures. Though she seemed to recognize her parents, she could not speak any words nor use her hands purposefully.

When she was over the age of 2 years, choreiform movements were noticed,

1
2
3 especially on her face, tongue and wrists, whereas opisthotonic postures were
4 subsided. The neurotransmitter metabolites and pteridine analysis in the
5 CSF were normal. The effect of L-dopa was equivocal, though her emotional
6 response might be slightly improved.
7
8

9
10 At the age of 9 years, marked choreiform movements persisted on her face,
11 tongue and the distal portions of the upper extremities at the resting state,
12 and were exaggerated by emotion or lack of adequate sleep. Her HC
13 remained within normal limits. The nystagmus was not so evident, but
14 sometimes seen. Her voluntary movement was only rolling from supine to
15 prone, and there were no meaningful words. Epileptic seizures had not been
16 observed, though her EEG was mildly abnormal with left posterior small
17 spikes. Her brain MRI showed normal myelination with no obvious changes
18 in the anterior horn enlargement and the cerebellar atrophy (Fig.1). No
19 definite deterioration had been observed during her clinical course.
20
21
22
23
24
25
26
27
28

29 Sibling 2 (II-2 in Figure 2A)

30
31 He was born at term without complications, and his HC was normal. The
32 left renal enlargement was detected in utero by ultrasonography. Urological
33 examinations revealed the bilateral hydronephrosis due to the pelvoureteral
34 stenosis and also the bilateral vesicoureteral reflux. Poor visual fixation
35 and the abnormal eye movements were detected since the age of 2-3 months
36 in spite of the normal ophthalmological examinations.
37
38
39
40

41 At the age of 5 months, he showed an asynchronous pendular and
42 sometimes downbeat nystagmus with the right-sided predominance.
43 Dystonic posturing predominantly in the right upper extremity was
44 constantly seen, and opisthotonic postures were easily elicited when he was
45 held. The deep tendon reflexes were normal. The MRI revealed a mild delay
46 of myelination without obvious brain atrophy. L-dopa seemed mildly
47 effective to his eye pursuit and opisthotonic postures.
48
49
50
51
52
53

54 He could roll over, but not sit alone at the age of 6 years and 3 months.
55 His HC was just below -2SD. The nystagmus and inner strabismus were
56 observed in his right eye. His muscle tonus was decreased, with the absent
57 deep tendon reflexes, and he also showed the clubfoot. Choreiform
58
59
60
61
62
63
64
65

movements comparable to his sister were seen in his extremities. Any meaningful words could not be acquired. Epileptic seizures have not been noticed yet. His renal symptoms were stable.

Whole exome sequencing (WES)

The experimental protocols were approved by the Institutional Review Board of Yokohama City University Graduate School of Medicine. Written informed consents were obtained from the parents in accordance with the requirements of Japanese regulations. Genomic DNA was isolated from peripheral blood leukocytes from two siblings and their parents. Genomic DNA of two siblings were captured using a SureSelect Human All Exon v5 Kit (51 Mb; Agilent Technologies, Santa Clara, CA, USA) and sequenced using an Illumina HiSeq2000 (Illumina, San Diego, CA, USA), with 101 bp paired-end reads. Exome data processing, variant calls, and variant annotation were performed as previously described [4].

Coverage data and variant filtering are shown in Table S1. Among two candidate genes possessing two heterozygous variants (possible compound heterozygous mutations) or a homozygous mutation based on the autosomal recessive model (Table S2), *MED17* was of our interest because a homozygous mutation had been reported in five infants with brain atrophy [3]. Compound heterozygous mutations were confirmed by Sanger sequencing (Fig. 2A, B). The c.1484T>G (p.Leu495Trp) mutation transmitted from their mother substituted a leucine residue which is highly conserved among the vertebrates, and was predicted to be damaging by SIFT, PolyPhen-2, and Mutation Taster (Fig. 2B and Table S2). The c. 1013-5A>G mutation transmitted from their father caused insertion of 4 bases of intron 6 in the transcript, resulting in a frameshift (p. Ser338Asnfs*15) in lymphoblastoid cells derived from two siblings (Fig. 2C, data not shown in sibling 2). The increased electropherogram peaks of mutant allele by cycloheximide treatment, which inhibits nonsense-mediated RNA decay, indicated that the mutant transcript with a premature stop codon underwent nonsense-mediated mRNA decay. Both mutations were not found in dbSNP137 data or our in-house 575 control exomes. These results suggest

that the *MED17* mutations are likely to be pathogenic.

Discussion

These two siblings shared common clinical course and symptoms: the abnormal eye movements with poor visual fixation and dystonic postures were seen from the early infancy, and as these features were subsided, choreiform involuntary movements appeared from the early childhood. These strongly indicate that they had a same disease. Frontal lobe-dominant brain atrophy observed in the MRI of sibling 1 suggests that abnormal eye movements with poor visual fixation may be related to the dysfunctions of the frontal visual fields, which have an integrative role in voluntary eye movements and visual fixation, and also dystonic postures and choreiform movements may be related to the basal ganglia dysfunctions.

Among the diseases showing choreiform movement, Huntington disease is a prototypical disorder with progressive striatal degeneration. The abnormal eye movement is also sometimes seen in this disease. But, autosomal dominant inheritance of this disease is not compatible with our cases, though it can't be excluded that the father is still in a pre-symptomatic stage. In addition, infantile onset has not been reported in this disease, though approximately 10% of the affected patients show symptoms before 20 years of age and 5% before 14 years of age. Furthermore, rigidity, which is more common than choreiform movement in such juvenile-onset type, was not observed in our cases. Other childhood-onset genetic disorders mainly with chorea, including abetalipoproteinemia, ataxia-telangiectasia, Fahr disease, glutaric aciduria, Wilson disease and neuroacanthocytosis can be ruled out by the clinical features and laboratory findings in our cases.

The fact that their symptoms were not compatible with any known diseases led us to the whole exome sequencing analysis of the patients. Surprisingly, both of them showed compound heterozygous mutations in *MED 17*. It is difficult to expect *MED17* abnormality from the clinical standpoint due to the scarcity of the patients, and WES is the only way to confirm this rare abnormality. It is suggested that WES is very powerful and highly recommended in order to confirm the genetic defects in clinically

unidentifiable diseases with suspected genetic background as such in our cases.

MED17 encodes one of 25 subunits constituting a huge transcription preinitiation mediator complex (MC). MC itself does not bind DNA, but works as a bridge to convey information from the transcription factors that recognize the enhancer site in DNA to the basic transcription apparatus on the promoter site, thus directly regulating the transcription initiation by RNA polymerase II (Pol II). MC is consisted of 4 modules: head, middle, tail and CDK/Cyclin modules. The head module forms 3 domains and interacts with Pol II and other basic transcription apparatus, putting them into an appropriate constellation [5]. MED17 is one of 7 subunits that constitute the head module. It has been shown that the direct interaction of MED17 with Rpb3 subunit of Pol II is essential for global Pol II recruitment in vivo [6]. Although the precise role of MED17 has not yet been established, it is reasonable to speculate that defective MED17 functions influence delicate mechanism of transcription. As the gene transcription is controlled subtly and strictly especially in the CNS, the defective MED17 is likely to disturb the CNS functions seriously.

The biological significance of MC in controlling signals involved in cell growth, development and differentiation has been stressed by an expanding list of human diseases with MC defects, including not only neurodevelopmental disorders but also cancers, cardiovascular and behavioral disorders [7]. As far as neurodevelopmental disorders are concerned, several human diseases have been known to be associated with MC defects. Charcot-Marie-Tooth disease type 2B2 (MIM 605589), an autosomal recessive axonal neuropathy, is caused by a missense mutation in *MED25* (MIM 610197) [8]. Two X-linked dysmorphic mental retardation syndromes, Opitz-Kaveggia syndrome (MIM 305450) and Fryns-Lujan syndrome (MIM 309520) are associated with mutations in *MED12* (MIM 300188) [9,10]. A distinct clinical syndrome with congenital retinal folds, microcephaly and mental retardation is associated with haploinsufficiency of *CDK19* (MIM 614720)[11].

Recently, Kaufmann et al reported a missense mutation in *MED17* (MIM

603810) in five infants from unrelated families with postnatal progressive microcephaly, cerebral and cerebellar atrophy (MIM 613668) [3]. All of them were Caucasian Jews, and had the same homozygous mutation (p.L371P), indicating a founder effect in the Caucasus Jewish community. Though the head circumferences were normal at birth, swallowing difficulties, jitteriness, poor visual fixation, opisthotonic posturing and overt seizures with microcephaly, increased muscle tone were seen from the early infancy. No acquisition of developmental milestones, marked spasticity and progressive microcephaly were evident at the age of 9 months. EEGs were highly abnormal with multifocal spikes and waves or hypsarrhythmia. Brain MRI revealed severe and diffuse cerebral and cerebellar atrophy already evident at 3 months of age, subsequently with poor myelination, small thalami and thin brainstems.

Our siblings are the second reported cases with *MED17* mutations. Compared with the previous cases, our cases show much milder clinical symptoms and slower progression with more restricted MRI lesions (Table 1). The homozygous mutation in their cases and the compound heterozygous mutations in our cases might explain these differences. In our cases, it is assumed that an allele harboring the c.1484T>G mutation can be translated to some functionally active protein, though another allele harboring c.1013-5A>G is likely to be functionally null. Alternatively, the allele harboring c.1013-5A>G mutation might produce a small amount of normally spliced transcripts, which would be translated to fully functional protein, resulting in milder phenotype. It seems difficult to find common features between their cases and our cases other than developmental delay with poor visual fixation and opisthotonic posturing from the infancy. Though the authors postulate that *MED17* mutation interferes with the expression of the gene(s) involved with oligodendrocyte development, our cases do not show any severe myelination defects. Choreiform movements with hypotonia in our cases and spasticity with increased muscle tones in the reported cases may reflect pathophysiological differences of the involving brain networks. It is likely that the defects in *MED17* protein influence many functions in the CNS, resulting in the quite diverse clinical features. To accumulate more

cases is needed in order to elucidate the clinical diversities and phenotype-genotype correlations in *MED17* mutations.

Acknowledgments

We thank the patients and their families for their participation in this study. We also thank Nobuko Watanabe for her excellent technical assistance. This work was supported by a Grant-in-Aid for the Ministry of Health, Labour and Welfare of Japan; Grants-in-Aid for Scientific Research (B) (25293085) and (A) (13313587), and challenging Exploratory Research (26670505) from the Japan Society for the Promotion of Science; the Takeda Science Foundation; the fund for Creation of Innovation Centers for Advanced Interdisciplinary Research Areas Program in the Project for Developing Innovation Systems from the Japan Science and Technology Agency; the Strategic Research Program for Brain Sciences (11105137); and a Grant-in-Aid for Scientific Research on Innovative Areas (Transcription Cycle) from the Ministry of Education, Culture, Sports, Science and Technology of Japan (12024421).

References

- [1] Yang Y, Muzny DM, Reid JG, Brainbridge MN, Willis A, Ward PA, et al. Clinical whole-exome sequencing for the diagnosis of Mendelian disorders. *N Engl J Med* 2013;369:1502-11.
- [2] de Ligt J, Willemsen MH, van Bon BWM, Kleefstra T, Yntema HG, Kroes T, et al. Diagnostic exome sequencing in persons with severe intellectual disability. *N Eng J Med* 2012;367:1921-9.
- [3] Kaufmann R, Straussberg R, Mandel H, Fattal-Valevski A, Ben-Zeev B, Naamati A, et al. Infantile cerebral and cerebellar atrophy is associated with a mutation in the MED17 subunit of the transcription preinitiation mediator complex. *Am J Hum Genet* 2010;87:667-70.
- [4] Ohba C, Osaka H, Iai M, Yamashita S, Suzuki Y, Aida N, et al. Diagnostic utility of whole exome sequencing in patients showing cerebellar and/or vermis atrophy in childhood. *Neurogenetics* 2013;14:225-32.
- [5] Imasaki T, Calero G, Cai G, Tsai K-L, Yamada K, Cardelli F, et al. Architecture of the Mediator Head module. *Nature* 2011;475:240-3.
- [6] Soutourina J, Wydau S, Ambroise Y, Boschiero C, Werner M. Direct interaction of RNA polymerase II and mediator required for transcription in vivo. *Science* 2011;331:1451-4.
- [7] Spaeth JM, Kim NH, Boyer TG. Mediator and human disease. *Semin Cell Dev Biol* 2011;22:776-87.
- [8] Leal A, Huehne K, Bauer F, Sticht H, Berger P, Suter U, et al. Identification of the variant Ala335Val of MED25 as responsible for CMT2B2: molecular data, functional studies of the SH3 recognition motif and correlation between wild-type MED25 and PMP22 RNA levels in CMT1A animal models. *Neurogenetics* 2009;10:275-87.
- [9] Risheg H, Graham JM Jr, Clark RD, Rogers RC, Opitz JM, Moeschler JB, et al. A recurrent mutation in MED12 leading to R961W causes Opitz-Kaveggia syndrome. *Nat Genet* 2007;39:451-3.
- [10] Schwartz CE, Tarpey PS, Lubs HA, Verloes A, May MM, Risheg H, et al. The original Lujan syndrome family has a novel missense mutation (p.N1007S) in the MED12 gene. *J Med Genet* 2007;44:472-7.
- [11] Mukhopadhyay A, Kramer JM, Merks G, Lugtenberg D, Smeets DF, Oortveld MA, et al. CDK19 is disrupted in a female patient with bilateral

congenital retinal folds, microcephaly and mild mental retardation. Hum
Genet 2010;128:281-91.

Figure & Table legends

Figure 1. Brain MRI of sibling 1

(A) Axial T2WI at the age of 1y.7m. (B) Saggital T1WI at the age of 1y.7m. (C) Axial T2WI at the age of 10y.7m. (D) Saggital T1WI at the age of 10y.7m.

Figure 2. *MED17* mutations in two siblings

(A) Familial pedigree of the two affected siblings and the *MED17* mutations. (B) Electropherogram of two mutations is shown. The missense mutation (p.Leu495Trp) occurred at leucine residue which is evolutionarily conserved among vertebrates. (C) Schematic representation of location of the c. 1013-5A>G mutation and primers used for mRNA analysis. Sequence of reverse transcriptase PCR amplicons using cDNA with cycloheximide (CHX) treatment clearly showed presence of mutant transcript with insertion of four base pair sequences of intron 6. Peak of mutant transcript was much weaker in the PCR amplicons using cDNA with dimethyl sulfoxide (DMSO) treatment as vehicle control.

Table 1. Clinical symptoms and MRI findings at the age of 10y and 6y in sibling 1 and 2, respectively, and those of the reported cases.

Table(s)

	sibling 1 (II-1)	sibling 2 (II-2)	Kaufmann’s Cases
Head circumference at birth	normal	normal	normal
Failure to thrive	absent	absent	present
Eye symptoms	horizontal nystagmus, poor visual fixation	asynchronous nystagmus	Poor visual fixation, lack of tracking
Opisthotonic posture in infancy	present	present	present
Muscle tone	decreased	decreased	increased
Developmental delay	moderate	moderate	profound
Seizures	absent	absent	present
Microcephaly	absent	present (-2SD at 6y)	present (-6SD at 9 mo)
Choreiform movements	present	present	absent
MRI: cerebral atrophy	mild (frontal lobe)	absent?	diffuse
MRI: cerebellar atrophy	mild (anterior vermis)	absent	diffuse
MRI: poor myelination	mild in infancy	mild in infancy	severe
MRI: small thalami	absent	absent	present
MRI: thin brainstem	absent	absent	present

Figure(s)

A

C

B

<i>H. sapiens</i> MED17	SIQLQLNIGVE
<i>B. Taurus</i> MED17	SIQLQLNIGVE
<i>M. musculus</i> Med17	SIQLQLNIGVE
<i>G. gallus</i> MED17	SIQLQLNIGVE
<i>X. tropicalis</i> med17	SIQLQLNIGVD
<i>D. rerio</i> med17	SIQLQLNICVE