Title: Using Quantitative Image Analysis to Classify Axillary Lymph Nodes on Breast MRI: A New Application for the Z 0011 Era

Authors: David V. Schacht MD MPH, Karen Drukker PhD MBA, Iris Pak, Hiroyuki Abe MD PhD, Maryellen L Giger PhD

Institution of Origin: The University of Chicago, Department of Radiology, Section of Breast Imaging, 5841 S. Maryland Avenue, MC 2026, Chicago, Illinois 60637

Corresponding Author:

David V. Schacht, M.D. M.P.H.

Department of Radiology

The University of Chicago

5841 S. Maryland Avenue, MC 2026, Chicago, Illinois 60637

Phone: +1(773) 702-2781, Fax: +1(773) 702-1161

E-mail address: dschacht@radiology.bsd.uchicago.edu

Co-Author contact information:

Karen Drukker, PhD MBA
5841 S Maryland Ave
Department of Radiology, MC2026
University of Chicago, IL 60637

E-mail: kdrukker@uchicago.edu

Iris Pak
5841 S Maryland Ave
Department of Radiology, MC2026
University of Chicago, IL 60637

E-mail: irisgpak@gmail.com

Hiroyuki Abe, MD PhD
5841 S Maryland Ave
Department of Radiology, MC2026
University of Chicago, IL 60637

E-mail: habe@radiology.bsd.uchicago.edu

Maryellen Giger, PhD
5841 S Maryland Ave
Department of Radiology, MC2026
University of Chicago, IL 60637

E-mail: m-giger@uchicago.edu

ABSTRACT:

Purpose: To assess the performance of computer extracted feature analysis of dynamic contrast enhanced (DCE) magnetic resonance images (MRI) of axillary lymph nodes. To determine which quantitative features best predict nodal metastasis.

Methods: This institutional board-approved HIPAA compliant study, in which informed patient consent was waived, collected enhanced T1 images of the axilla from patients with breast cancer. Lesion segmentation and feature analysis were performed on 192 nodes using a laboratory-developed quantitative image analysis (QIA) workstation. The importances of 28 features were assessed. Classification used the features as input to a neural net classifier in a leave-one-case-out cross-validation and evaluated with receiver operating characteristic (ROC) analysis.

Results: The area under the ROC curve (AUC) values for features in the task of distinguishing between positive and negative nodes ranged from just over 0.50 to 0.70. Five features yielded AUCs greater than 0.65: two morphological and three textural features. In cross-validation, the neural net classifier obtained an AUC of 0.88 (SE 0.03) for the task of distinguishing between positive and negative nodes.

Conclusion: QIA of DCE MRI demonstrated promising performance in discriminating between positive and negative axillary nodes.

INTRODUCTION

Breast MRI is often used in the clinical staging of patients with newly diagnosed breast cancer for defining extent of disease in the breast, detecting contralateral cancers [1], and detecting adenopathy. Axillary and internal mammary lymph nodes are readily detectable on MRI, and T2 weighted sequences and post-contrast dynamic sequences can both demonstrate the size and morphology of axillary lymph nodes. With these high-resolution sequences, the axillae can be viewed three dimensionally and a high level of anatomic detail is discernable. Such images are especially useful for determining architectural details of lymph nodes such as cortical size, morphology and the presence or absence of a fatty hilum (Figure 1).

Quantitative image analysis (QIA) is an area of active research and includes rather well-established applications, such as computer-aided detection (CAdE), and applications not yet available for everyday clinical use, such as computer-aided prognosis. Within radiology, and especially within the subspecialty of breast imaging, CAdE has become mainstream for some imaging modalities and is often integrated within clinical workstations. On mammograms, CAdE serves as a “second reader” and is used to detect masses and calcifications that could indicate the presence of invasive or in-situ carcinoma [2].

In this paper, we investigate the potential of computer-aided prognosis through axillary lymph node assessment in breast MRI. Currently, most commercially available software is more limited in its abilities and performs volumetric assessment of defined lesions, which can aid in surgical planning. Similarly, in cases where the patient will receive neoadjuvant chemotherapy, comparison measurements performed before and after therapy can be used as an imaging biomarker for response [3]. The use of more sophisticated QIA for breast MRI, however, remains an area of active research both for tumor classification [4], and for staging and prognosis [5]. In previous research studies, promising performance was obtained using image-based

1
2
3
4 biomarkers for computer analysis of breast lesions in MRI, whereby the computer performed
5
6 segmentation, extraction of morphologic and kinetic characteristics (features), and subsequent
7
8 classification [6-9]. In this study, we investigated whether a QIA scheme utilizing a digital
9
10 analysis of lymph nodes imaged on breast MRI is able to distinguish between lymph nodes that
11
12 were positive for metastasis ('positive' nodes) and those that were negative for metastasis
13
14 ('negative' nodes). In the future, such a scheme, if successful, could potentially help guide
15
16 clinical management in the axilla.
17
18
19
20
21
22
23
24

25 METHODS

26
27
28 This study was an institutional review board-approved, HIPPA compliant study, with
29
30 waiver of informed consent. A retrospective review was performed on 66 cancer patients who
31
32 underwent staging MRI at our institution between 2006 and 2010.
33
34
35
36

37 MR images were obtained by using 1.5 and 3.0 T systems depending on clinical
38
39 availability. MRI was performed with a dedicated breast coil and the patient in the prone
40
41 position (Table 1). Contrast material was injected IV (0.1 mmol/kg of gadodiamide [Omniscan,
42
43 GE Healthcare]) and followed by a 20-mL saline flush at a rate of 2 mL/s. The same contrast
44
45 material/protocol was used for all systems.
46
47
48
49

50 A database from 66 cancer patients was retrospectively collected for the assessment of
51
52 QIA of axillary lymph nodes on MRI (Table 1). Analysis was performed on 154 negative lymph
53
54 nodes and 38 positive lymph nodes, identified *a posteriori* by a board certified expert radiologist
55
56 with 9 years of experience. Review of surgical pathology reports, radiology reports, and
R 1 #3
R1 #8
ultrasound images were used to establish the 'gold standard' of positivity or negativity for these

1
2
3
4 lymph nodes. All axillary lymph nodes of the patients with negative sentinel lymph node biopsy
5
6 were considered negative. All metastatic lymph nodes proven by ultrasound guided core needle
7
8 biopsy were regarded as positive for metastasis. To correlate between the biopsy proven
9
10 metastatic axillary LNs on ultrasound (US) and LNs shown on MRI, the same radiologist
11
12 identified each metastatic axillary LN on MRI by comparing MR images with images from the
13
14 US guided biopsy. When multiple lymph nodes (> 8) were proven positive for metastasis at
15
16 axillary lymph node dissection, highly suspicious lymph nodes on MRI (up to three lymph nodes
17
18 per patient) were presumed positive for metastasis.
19
20
21
22
23

R 1 #9

24 The methodology involved several steps which were all automated except for MR image
25 acquisition (as detailed above) and the identification of the image locations of lymph nodes . The
26
27 steps in the methodology not specific to the current application have been described extensively
28
29 elsewhere [10, 11] and are briefly summarized here for clarity. An expert board-certified
30
31 radiologist identified the locations of axillary lymph nodes visible in the MR images. The lymph
32
33 nodes were then automatically segmented) using a method previously developed for breast
34
35 tumors [10]. Subsequently, computer-extracted MR-based features were calculated to
36
37 characterize the lymph nodes. The feature set describing each node consisted of 28 features,
38
39 including 5 kinetic curve assessment, 4 variance of kinetics, 14 enhancement texture, and 5
40
41 morphological features (Figure 2a and Table 2). For the task of distinguishing between nodes
42
43 positive for metastasis and those negative for metastasis, each feature was assessed individually
44
45 to gain insight into which type of feature was relevant for this task. Subsequently, all features
46
47 were used in combination through the use of a classifier to predict which nodes were positive and
48
49 which were negative for metastasis. We used a Markov-Chain Monte-Carlo Bayesian Neural Net
50
51 classifier [12] (MCMC-BNN) within leave-one-case-out cross-validation. Leave-one-case-out
52
53
54
55
56
57
58
59
60
61
62
63
64
65

cross-validation is an accepted training and testing method with the aim to minimize database bias (overtraining) of a classifier. Here, for N cases, in each cross-validation a single case was assigned as the testing case and the remaining N-1 cases were used for classifier training. This process was repeated N times until all cases had served a test case.

Hence, the novel aspects of the current study included (a) the application of a 3D segmentation method, which was previously developed on breast tumors [10], for the segmentation of axillary lymph nodes (Figure 2b), (b) the use of computer-extracted features (mathematical descriptors) to quantitatively characterize the nodes from MR image data [6-8,13], and (c) a neural network classifier to distinguish between nodes that were positive for metastasis and nodes that were negative for metastasis .

Performance was assessed both qualitatively – using box plots - and quantitatively using receiver operator characteristic (ROC) analysis with the area under the ROC curve (AUC) as the figure of merit [14, 15]. The performance of individual features was assessed to gain insight into their relative importance for the task at hand, and thus, statistical comparisons amongst performance of individual features were not performed. The aim was to determine whether we can learn anything at all from imaging features regarding lymph node status, not whether image-based features were as good as the ‘gold-standard’ of more invasive and not necessarily risk-free biopsy (assumption AUC=1.0). Hence, the null-hypothesis here was that a computer-extracted nodal feature gives no insight into the presence of metastasis, and the AUC value for each individual feature was compared to the baseline of random chance (AUC=0.5) through the estimation of the symmetrical 95% confidence interval of the AUC value as $AUC \pm 1.96$ standard error. The performance of all features used in conjunction with the neural net classifier was similarly assessed.

R 1 #4

R 1 #10

RESULTS

When each of the 28 features was analyzed individually, 13 features achieved performance substantially better than guessing (Figure 3). These 13 traits were 2 kinetic curve assessment, 9 enhancement texture, and 2 morphological features. The five best performing individual features, in order of decreasing AUC were: variance of radial gradient histogram (vRGH) (morphology), correlation (enhancement texture), difference in variance (enhancement texture), energy (enhancement texture), and circularity (morphology). Note that no kinetic curve assessment features were in this group of 5 best performing individual characteristics. Also note that the highest AUC for an individual characteristic of 0.70 (0.05) was obtained for a morphological feature that describes how well enhancing structures within a lesion extend from the center of a lesion outwards.

Feature values for the 5 features with the highest individual merit (as noted above) and for the best performing kinetic feature (washout) are shown with box plots and whiskers for positive and negative nodes (Figure 4). Based on this analysis, positive nodes were more likely to have: increased variance of radial gradient histogram (increase in radially outward enhancement patterns), increased circularity, decreased energy (less homogeneous appearance), decreased difference in variance (less variation in the difference in gray levels between voxel pairs), increased correlation (larger image linearity), and decreased washout (kinetic curve assessment).

The use of all lymph node characteristics combined in leave-one-case-out cross-validation considerably improved performance compared to the performance of any individual feature on its own, yielding AUC=0.88 (standard error 0.03 and estimated 95% confidence interval [0.82;

0.94]) (Figure 5). Note that the best performing individual feature, as discussed above, obtained AUC = 0.70.

DISCUSSION

The results of this study, though preliminary in scope, show how quantitative methods that had previously been applied to breast lesions could also be applied to axillary lymph nodes in newly diagnosed cancer patients. The results showed that several individual features, and also the combination of all studied lymph node characteristics could be used to assess lymph nodes. The majority of features that achieved performance better than guessing were textural features. This could be due to the inherent differences at the cellular level that occur in metastatic versus benign lymph nodes and the resultant MRI signal changes. Not surprisingly, several morphological features also performed well. For example, the morphology descriptor of circularity is generally higher in positive nodes. This result is concordant with the typical imaging appearance of metastatic nodes, which tend to be rounder and less reniform. Kinetic features, in general, showed less performance in distinguishing between the positive and negative lymph nodes. It is well known that even normal axillary lymph nodes can display typically malignant patterns of rapid uptake and washout kinetics. And therefore it is not unexpected that there were many outlier negative lymph nodes that demonstrated significant washout (Figure 4 a). The finding of a significantly higher AUC value with the utilization of all characteristics, by merging with a classifier, is important because it highlights the potential clinical utility of this work. By using the combination of all selected computer-extracted lymph node characteristics, a high AUC was obtained.

1
2
3
4 This study has several limitations. First, the MRI images analyzed in this study represent
5
6 only those from dynamic contrast enhancement (DCE) MRI protocols. QIA on primary breast
7
8 tumors has demonstrated improved performance in distinguishing between malignant and benign
9
10 lesions with the incorporation of QIA on the T2 weighted images [16]. Future studies of axillary
11
12 node QIA could incorporate images from an additional sequence, such as the T2 images.
13
14
15 Similarly, if QIA could also incorporate multi-modality information, the utility might be further
16
17 improved.
18
19
20
21

22 Another limitation is that the results presented here might be affected by the relatively
23
24 small and unbalanced data set. Because of the modest size of the data set, we did not extensively
25
26 investigate different feature selection protocols to improve performance, since such an approach
27
28 would likely result in database bias, i.e., in a too optimistic performance assessment and a lack of
29
30 generalize-ability. The single leave-one-case-out cross-validation presented here provided a more
31
32 realistic evaluation. A similar study with a larger and more balanced set of lymph nodes,
33
34 however, may have some effect on the characteristics that were seen to be most influential in
35
36 determining lymph node status, and thus further studies are planned. In addition, inclusion of
37
38 more lymph-node specific features, such as cortical thickness, may lead to further improvements
39
40 as well. A third limitation is a level of ambiguity in the ‘gold standard’ due to the difficult task of
41
42 identifying which lymph nodes were biopsied or dissected and of matching the pathology to the
43
44 imaged nodes.
45
46
47
48
49
50
51

52 Staging the axilla has recently been a topic of much research, discussion and debate. The
53
54 publication and implementation of the ACOSOG Z 0011 study [17] has led to changes in the
55
56 imaging work up of many breast cancer patients at this institution. Specifically, axillary
57
58 ultrasound is being used much less frequently. Prior to Z 0011, every newly diagnosed cancer
59
60
61
62
63
64
65

1 patient would undergo an axillary ultrasound in the search for a suspicious lymph node. If a
2
3
4
5
6 suspicious node was found and biopsy proven malignant, then surgeons would proceed directly
7
8
9 to complete axillary dissection at the time of breast surgery. However, this paradigm is no
10
11 longer used in light of Z 0011, which showed no survival benefit to more extensive axillary
12
13 surgery, even with positive sentinel nodes (up to two). Now patients with small breast cancers
14
15 (T1 or T2) usually only have a sentinel lymph node biopsy (rather than a complete axillary
16
17 dissection) even when the sentinel node is positive. As a result, the rate of staging axillary
18
19 ultrasound and ultrasound guided axillary biopsy has significantly decreased.
20
21
22
23

24 In institutions where axillary ultrasound is being used less frequently, MRI may become
25
26 more important for providing imaging staging of the axilla. While the results presented here are
27
preliminary in nature, it is possible that a new paradigm could depend on MRI with quantitative
analysis to provide the initial staging view of the axilla, and if there are multiple suspicious
lymph nodes (suggesting at least 3 positive nodes, and therefore suggesting exclusion from Z
0011 criteria) then further work-up with axillary ultrasound and axillary ultrasound guided
biopsy could be performed (Figure 6). Given this potential new framework, gaining the most
information from the MRI images of the axilla is critical, and further development of QIA
methods of evaluating the axilla could increase the diagnostic utility and prognostic value of
staging MRI.

R 1 #2

References:

- [1] Lehman C, Gatsonis C, Kuhl C, Hendrick RE, Pisano ED, Hanna L, et al. MRI evaluation of the contralateral breast in women with recently diagnosed breast cancer. *N Engl J Med* 2007; 356(13):1295-1303.
- [2] Doi K, Giger ML, Nishikawa RM, Schmidt RA. Computer-aided diagnosis of breast cancer on mammograms. *Breast Cancer* 1997 Dec 25; 4(4): 228-233.
- [3] Hylton NM, Blume JD, Bernreuter WK, Pisano ED, Rosen MA, Morris EA, et al. Locally advanced breast cancer: MR imaging for prediction of response to neoadjuvant chemotherapy—results from ACRIN 6657/I-SPY trial. *Radiology* 2012 Jun; 263(3): 663-72.
- [4] Bhooshan N, Giger M, Edwards D, Yuan Y, Jansen S, Li H, et al. Computerized three-class classifications of MRI-based prognostic markers for breast cancer. *Phys Med Biol* 2011 Sep 21; 56 (18): 5995-6008.
- [5] Bhooshan N, Giger M, Jansen S, Li H, Lan L, Newstead G. Cancerous breast lesions on DCE-MRI: computerized characterization for image-based prognostic markers. *Radiology* 2010 March; 254(3): 680-690.
- [6] Chen W, Giger ML, Li H, Bick U, Newstead G. Volumetric texture analysis of breast lesions on contrast-enhanced magnetic resonance images. *Magn Reson Med* 2007; 58:562-571.

- 1
2
3
4 [7] Chen W, Giger ML, Li H, Bick U and Newstead G. Automatic identification and
5
6 classification of characteristic kinetic curves of breast lesions on DCE-MRI. Med Phys
7
8 2006 Aug; 33(8): 2878-2887.
9
- 10
11 [8] Chen W, Giger ML, Lan L , Bick U. Computerized interpretation of breast MRI:
12
13 investigation of enhancement –variance dynamics. Med Phys 2004 May; 31(5): 1076-
14
15 1082.
16
17
- 18
19 [9] Shimauchi A, Giger ML, Bhooshan N, Lan L, Pesce LL, Lee JK, et al. Evaluation of
20
21 clinical breast MR imaging performed with prototype computer-aided diagnosis breast
22
23 MR imaging workstation: reader study. Radiology 2011 Mar; 258(3): 696-704.
24
25
- 26 [10] Bhooshan N, Giger ML, Edwards D, Yuan Y, Jansen S, Li H, et al. Computerized
27
28 three-class classification of MRI-based prognostic markers for breast cancer. Phys Med
29
30 Biol 2011 Sep 21; 56(18): 5995-6008.
31
32
- 33 [11] Chen W, Giger ML, Bick U. A fuzzy c-means (FCM)-based approach for
34
35 computerized segmentation of breast lesions in dynamic contrast-enhanced MRI
36
37 images. Acad Radiol 2006; 13(1): 63–72.
38
39
- 40 [12] Cronin B, Stevenson IH, Sur M, Kording KP. Hierarchical Bayesian modeling
41
42 and Markov chain Monte Carlo sampling for tuning-curve analysis. J Neurophysiol 2010;
43
44 103 (1):591-602.
45
46
- 47 [13] Chen W, Giger ML, Newstead G, Bick U, Jansen S, Li H, Li L. Computerized
48
49 assessment of breast lesion malignancy using DCE-MRI: robustness study on two
50
51 independent clinical datasets from two manufacturers. Acad Radiol 2010 July; 17(7):
52
53 822-829.
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4 [14] Pesce LL, Metz CE. Reliable and computationally efficient maximum-likelihood
5
6 estimation of "proper" binormal ROC curves. Acad Radiol 2007 July; 14(7): 814-829.
7
8
9 [15] Metz, CE. Basic principles of ROC analysis. Semin Nucl Med 1978 Oct; 8(4):
10
11 283-298.
12
13
14 [16] Bhooshan N, Giger ML, Lan L, Li H, Marquez A, Shimauchi, Newstead G.
15
16 Combined use of T2-weighted MRI and T1-weighted dynamic contrast-enhanced MRI in
17
18 the automated analysis of breast lesions. Magn Reson Med 2011; 66: 555-563.
19
20
21 [17] Guiliano AE, Hunt KK, Ballman KV, Beitsch PD, Whitworth PE, Blumencranz
22
23 PW, et al. Axillary dissection vs no axillary dissection in women with invasive breast
24
25 cancer and sentinel node metastasis: a randomized clinical trial. JAMA 2011 Feb 9;
26
27 305(6): 569-575.
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1: (a) Normal morphology right axillary lymph node (arrow) on an axial post-contrast T1 fat saturated subtracted sequence. Note the normal appearing enhancement of the lymph node and normal appearance of the fatty hilum with density similar to the background fat.

(b) Abnormal right axillary lymph node (arrow) on a post- contrast T1 fat saturated subtracted axial image: Enlarged nodal size and near complete loss of the fatty hilum are seen.

Figure 2a: Pictorial representation of the characteristics analyzed

Figure 2b: Example of a positive axillary lymph node with automated segmentation overlay (red outline)

Figure 3: AUC values for individual features in the task of distinguishing between nodes that were positive for metastasis and those that were negative (with error bars denoting $\pm$ standard error). The numbers indicate the order of the best performing 5 features, and features that individually achieved significantly better performance than a coin-toss are marked with an asterisk. The dashed line is a visual guide to separate those features from the ones that failed to outperform a coin-toss.

Figure 4: Box plots of feature values for positive and negative nodes (after standardization to zero mean and unit standard deviation for the entire dataset): a) the most discriminatory kinetic

feature (washout), and the 5 best performing features, i.e., 3 textural (b-d) and 2 morphological (e-f). On each box, the central mark is the median, the edges of the box are the 25th and 75th percentiles, the whiskers extend to the most extreme data points not considered outliers, and outliers are plotted individually.

Figure 5: ROC curve for the MCMC-BNN classifier in the task of distinguishing between nodes that were positive for metastasis and those that were negative for metastasis (leave-one-case-out cross-validation).

Figure 6: Imaging work-up of the axilla of newly diagnosed T1 or T2 breast cancer in the Z0011 era

Table 1. Acquisition Protocols and Lymph Node Status of the MRI database of 66 cancer patients

		1.5 T Achieva Philips	3 T Achieva Tx. Philips
Lymph Nodes	Number of Positive Lymph Nodes	32	6
	Number of Negative Lymph Nodes	123	31
Dynamic Studies	Study type	3D gradient echo sequences One pre- and six post-contrast enhancement (axial)	3D gradient echo sequences One pre- and five post-contrast enhancement (axial)
	TR/TE	5.5/2.7	5.5/2.7
	Flip angle	10°	10°
	Slice thickness	2 mm	2 mm
	Matrix size	480x480	448x448
	FOV	360 mm	340 mm
	Acquisition time	75 second	65 second

Table 2 revised

Feature category	Image feature	Description
Kinetic Curve Assessment	Maximum enhancement	Maximum contrast enhancement
	Time to peak (s)	Time at which the maximum enhancement occurs
	Uptake rate (1/s)	Uptake speed of the contrast enhancement
	Washout rate (1/s)	Washout speed of the contrast enhancement
	Curve shape index	Difference between late and early enhancement
Enhancement-Variance Kinetics	Maximum variance of enhancement	Maximum spatial variance of contrast enhancement over time
	Time to peak at maximum variance (s)	Time at which the maximum variance occurs
	Enhancement variance increasing rate (1/s)	Rate of increase of the enhancement-variance during uptake
	Enhancement variance decreasing rate (1/s)	Rate of decrease of the enhancement-variance during washout
Enhancement Texture	Angular second moment (Energy)	Image homogeneity
	Contrast	Local image variations
	Correlation	Image linearity
	Entropy	Randomness of the gray-levels
	Sum of squares (Variance)	Spread in the gray-level distribution
	Difference entropy	Randomness of the difference of neighboring voxels' gray-levels
	Difference variance	Variations of difference of gray-levels between voxel-pairs
	Inverse difference moment	Image homogeneity
	Sum average	Overall brightness
	Sum entropy	Randomness of the sum of gray-levels of neighboring voxels
	Sum variance	Spread in the sum of the gray-levels of voxel-pairs distribution
	Information measure of correlation (IMC) 1	Nonlinear gray-level dependence
	Information measure of correlation (IMC) 2	Nonlinear gray-level dependence
	Maximum correlation coefficient	Nonlinear gray-level dependence
Morphology	Sphericity/circularity	Similarity of the lesion shape to a sphere

Irregularity	Deviation of the lesion surface from the surface of a sphere
Margin sharpness	Mean of the image gradient at the lesion margin
Variance of margin sharpness	Variance of the image gradient at the lesion margin
Variance of radial gradient histogram	Degree to which the enhancement structure extends in a radial pattern originating from the center of the lesion

Figure(s)

Figure(s)

Figure2a revised

Computer-extracted image features

	Kinetic curve assessment	such as time to peak, washout rate
Enhancement		Enhancement variance kinetics	such as spatial variance of contrast enhancement over time
	Enhancement texture	such as entropy, homogeneity
Morphology		Shape	such as circularity, irregularity
	Other morphology	such as margin sharpness, radial gradient

Figure(s)

Figure(s)

Figure(s)

Figure(s)

1
2
3
4 Acknowledgements:
5
6
7

8 This work was supported in part by Department of Energy grant DE-FG02-08ER6478, National
9

10
11 Institute of Health grants NIH P50-CA125183 and NIH S10 RR021039.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65