1
2
3 **Changes in acyl and total ghrelin concentrations and their association with dry matter**
4 **intake, average daily gain, and feed efficiency of finishing beef steers and heifers**

5 A. P. Foote¹, K. E. Hales, and H. C. Freetly

6 ^aUSDA, Agriculture Research Service, U.S. Meat Animal Research Center, Clay Center, NE

7 68933-0166²

8

¹ Corresponding Author: Tel.: 402-762-4205, E-mail: Andrew.Foote@ars.usda.gov

² Mention of trade names or commercial products in this article is solely for the purpose of providing specific information and does not imply recommendations or endorsement by the USDA. USDA is an equal opportunity provider and employer.

9 **ABSTRACT**

10 Ghrelin is a peptide hormone produced in the gut that is implicated in signaling appetite and
11 regulating dry matter intake (DMI). The objective of this experiment was to determine the
12 change in acyl ghrelin, total ghrelin, and the ghrelin ratio (acyl ghrelin/total ghrelin) over an 84-d
13 DMI and average daily BW gain (ADG) measurement period and to determine the association of
14 those ghrelin measurements with DMI, ADG, ADG:DMI ratio (G:F), and residual feed intake
15 (RFI) in finishing beef steers and heifers. Blood samples were collected on day 0 and day 83
16 prior to feeding and between 0730 h and 1130 h. Samples were analyzed for acyl and total
17 ghrelin using commercially available RIA. Dry matter intake in steers was greater during the last
18 35-d period of the experiment compared to the first 35-d ($P < 0.01$), and was greater than heifers
19 regardless of period ($P < 0.01$). Steers had greater acyl ghrelin concentrations on day 0 than
20 heifers, but concentrations decreased by day 83 to equal concentrations in heifers ($P < 0.01$).
21 Total ghrelin concentrations were lower on day 0 in heifers but increased by day 83 and did not
22 differ from steers on day 83 ($P < 0.01$). A mixed model analysis was utilized to determine the
23 association of ghrelin concentrations and ratio with production traits, independent of breed and
24 sire effects. There was an interaction of day 0 acyl ghrelin concentrations with time of sample
25 collection for 84-d DMI ($P < 0.01$), ADG ($P < 0.01$), and G:F ($P = 0.09$), indicating a general
26 positive association of acyl ghrelin with production traits, but the association weakened as time
27 of sample collection increased. The mean ghrelin ratio tended ($P = 0.08$) to be positively
28 associated with DMI in the last 35-d period. The ghrelin ratio on day 0 interacted with time of
29 sample collection for ADG and G:F ($P < 0.05$), indicating an overall positive association of the
30 ghrelin ratio with ADG and G:F. Results indicate that ghrelin is associated with DMI, ADG, and

feed efficiency of finishing beef cattle, and data lend more evidence that ghrelin is involved in appetite regulation of ad libitum fed cattle.

Keywords: beef cattle; feed efficiency; feed intake regulation; gut peptide

1. Introduction

The gut peptide ghrelin is thought to be a stimulator of appetite in many species [1-3], including cattle [4]. Ghrelin was first discovered because of its ability to bind the growth hormone secretagogue receptor when it is acylated on the serine-3 residue, which leads to the release of growth hormone [5]. In addition to the growth-hormone releasing activity of acyl ghrelin, it is also thought to increase appetite by stimulating the release of neuropeptide Y and agouti-related protein in the hypothalamus [6,7]. Foote, et al. [8] reported that a single measurement of acyl ghrelin was positively associated with dry matter intake (DMI) and average daily body weight gain (ADG) in ad libitum fed finishing steers and heifers, indicating that differences in circulating ghrelin concentrations could be partially responsible for the observed differences in DMI of ad libitum fed cattle. It was also shown that the ratio of acyl:total ghrelin (total ghrelin includes both acyl and desacyl ghrelin) was more associated with DMI and ADG than acyl ghrelin alone [8]. The single measurement of acyl and total ghrelin used by Foote et al. [8] was collected after the completion of DMI and ADG collection period, which indicates that ghrelin concentrations could be more representative of previous DMI and not predictive of future DMI. Given the known biological function of ghrelin in other species [1-3], it is likely that greater ghrelin concentrations would be indicative of a greater appetite, and therefore greater DMI in ad libitum fed animals. Given the potential actions of ghrelin on growth and DMI regulation, it is possible that ghrelin concentrations could be used as a biomarker for feed

efficiency evaluation; however, it isn't clear if the association of ghrelin concentrations with production traits will remain significant in other groups of cattle. If ghrelin is shown to have a resilient association with production traits, the data will provide evidence that the ghrelin biological system should be evaluated further for having direct effects on observed variation in DMI of ad libitum fed cattle.

The objective of this experiment was to determine the association of acyl and total ghrelin measured at the beginning and end of an 84-d feeding period with production traits. Additionally, the changes in acyl and total ghrelin concentrations were determined. It was hypothesized that ghrelin concentrations will be associated with feed intake.

2. Materials and methods

This experiment was reviewed and approved by the U.S. Meat Animal Research Center (USMARC) Animal Care and Use Committee (IACUC approval number 5438-31000-0092-03) under IACUC standards meeting the requirements outlined by the USDA.

2.1 Animals

Details of the animals and experimental procedures have been previously described [9]. Calves ($n = 236$ total; $n = 127$ steers and $n = 109$ heifers) used in this experiment were the progeny from composite breed cows that were mated to Angus, Charolais, Gelbvieh, Limousin, Red Angus, and Simmental bulls that were in current use on industry ranches. Both steers (Angus- [$n = 35$], Charolais- [$n = 8$], Gelbvieh- [$n = 21$], Limousin- [$n = 21$], Red Angus- [$n = 17$], and Simmental-sired [$n = 25$]) and heifers (Angus- [$n = 25$], Charolais- [$n = 6$], Gelbvieh- [$n = 30$], Limousin- [$n = 18$], Red Angus- [$n = 14$], and Simmental-sired [$n = 16$]) were used in the study. At the beginning of the study, heifers weighed 395 ± 3.5 kg and steers weighed 416 ± 3.5

kg. Mean age of cattle at the beginning of the study was 298 ± 0.4 d. Calves were housed in a facility with Calan-Broadbent electronic headgates (American Calan, Inc., Northwood, NH) to measure individual feed intake. The calves were sorted and penned by BW prior to beginning the study to minimize aggressive behavior of larger cattle toward smaller cattle. Steers and heifers were penned separately. Thirteen pens were used in this experiment and housed either 19 or 20 animals each. The cattle were trained to use individual Calan headgates during the adaptation period (approximately 45 d). The steers were implanted (200 mg trenbolone acetate and 40 mg estradiol 17 β ; Revalor XS, Merck Animal Health, Kenilworth, NJ) 86 d before the start of the experiment. Heifers received a Revalor H implant (140 mg trenbolone acetate and 14 mg estradiol; Merck Animal Health) 86 d before the beginning of the study and a Revalor 200 implant (200 mg trenbolone acetate and 20 mg estradiol; Merck Animal Health) 57 d after the study began.

Cattle were placed on a diet that on a dry matter (DM) basis consisted of 67.75% dry-rolled corn, 20% wet distillers grains with solubles, 8% chopped alfalfa hay, and 4.25% commercial vitamin and mineral supplement 7 d before the feed intake and growth trial began. The supplement contained monensin to supply 300 mg/head daily. Feed bunks were evaluated visually each day of the experiment at approximately 0730 h to determine the quantity of feed to offer each animal. The bunk management approach was designed to allow for 0.25 to 0.50 kg of feed remaining in the feed bunk at the time of evaluation. After the quantity of feed to be provided to each bunk was determined, a portion of the ration sufficient to supply the feed for all pens was mixed in the feed truck (Roto-Mix IV 274-12B, scale readability ± 0.09 kg) for approximately 5 min. Cattle were fed once daily throughout the experiment, starting at approximately 0800 h. Feed was subsampled daily, and a weekly composite sample was made to

determine feed DM. Orts were determined once per week. Total DMI was the sum of total DM fed minus total Orts. Values reported and used in the models are average daily DMI (referred to as DMI for simplicity). The feeding study lasted 84 d and cattle were weighed on days 0, 1, 21, 42, 63, 83, and 84. A quadratic equation was used to regress BW on day of study and total BW gain was determined by solving the equation for 84 d on study. Average daily gain was calculated as total BW gain divided by days on study. The ratio of ADG:DMI (G:F) was calculated as the quotient of ADG divided by DMI. Residual feed intake (RFI) was calculated by regressing observed DMI against ADG and metabolic BW at the midpoint of the experiment, and the residual was RFI. Steers and heifers were treated as separate cohorts for the RFI calculation.

2.2 Sample collection, processing, and analyses

Blood (9 mL) was collected on day 0 and day 83 of the DMI and ADG measurement period prior to feeding, via jugular venipuncture into tubes containing EDTA (1.7 µg/mL of blood) and placed on ice immediately. Samples were then centrifuged at $3,000 \times g$ for 25 min at 4°C within 1.5 h of collection to obtain plasma. Two 1-mL aliquots for acyl and total ghrelin analysis were treated with 50 µL of 1 M HCl and 10 µL of phenylmethanesulfonyl fluoride (10 mg/mL in 100% methanol; Sigma Aldrich, St. Louis, MO). All plasma samples were stored at -20°C until analysis. Cattle were processed prior to feeding and samples were collected between 0730 h and 1130 h on day 0 and day 83. Pens were processed in the same order each day. An electronic time stamp was recorded for each animal when they were weighed, and this time stamp was used in the analysis as the time of sample collection within day. Feeding was delayed until cattle were processed.

Acyl and total ghrelin were analyzed as previously described [8], using commercially available RIA kits (GHRA-88HK and GHRT-89HK for acyl and total ghrelin, respectively; EMD Millipore, Billerica, MA). Intra- and inter-assay CV for the acyl ghrelin assay were 3.7 and 4.0%, respectively. Intra- and inter-assay CV for the total ghrelin assay were 2.4 and 2.6 %, respectively.

2.3 Statistical analysis

All analyses used animal as the experimental unit and were performed in SAS 9.3 (SAS Inst. Inc., Cary, NC). Dry matter intake data were broken up into the first 35 d, the last 35 d, and the entire 84 d, and differences in DMI between the periods were tested using a mixed model analysis with breed, sex, period, and the interaction as main effects and sire as a random variable. Average daily gain and G:F differences were tested using a mixed model with breed and sex as main effects and sire as a random variable. Differences in concentrations of acyl and total ghrelin, and the ratio of acyl:total ghrelin (referred to as ghrelin ratio) between sexes and day of measurement were tested using a mixed model with breed, sex, day, and the interaction as main effects, and sire as a random variable. All variables were tested for normality using the Shapiro-Wilk test in the univariate procedure. Acyl ghrelin and the ghrelin ratio were determined to be non-normal with a Shapiro-Wilk test statistic (W) < 0.95 and were log- (base 10) transformed to achieve near normality (W > 0.98).

To evaluate the association of acyl ghrelin, total ghrelin, and the ghrelin ratio with DMI, ADG, G:F, and RFI, a mixed model analysis was utilized. All models included breed and sex as fixed effects and sire as a random variable. The dependent variables of interest were analyzed in separate models and included acyl ghrelin (log transformed) concentration on day 0 or day 83, total ghrelin concentration on day 0 or day 83, and the ghrelin ratio (log transformed) on day 0 or

day 83, the mean of day 0 and day 83 values, and the delta (difference in day 83 and day 0 ghrelin concentrations or ratio) of the 3 measures of ghrelin (acyl ghrelin, total ghrelin, and the ratio of acyl:total ghrelin). Blood sample collection time was included as an interaction with the ghrelin variable of interest; however, the interaction was removed from the model if $P > 0.10$ and the blood sample collection time was removed completely from the model if $P > 0.10$. The interaction of ghrelin with time of sample collection was included to determine if the association of ghrelin with the production traits changed over time within sampling day. Solutions $\pm$ SEM of the mixed model are presented to illustrate the directionality and magnitude of the associations between ghrelin and the production traits, and for future modeling studies based on the data. To determine the amount of variation explained by the models, a pseudo- R^2 and the variance explained by the dependent variables was calculated as described previously [10]. Variables were considered significant when $P < 0.05$.

3. Results

Dry matter intake was greater in the final 35-d period than the first 35-d period of the trial for steers but not heifers ($P < 0.01$; Table 1). Average DMI was approximately 0.56 kg/d greater in the final 35-d period compared to the first 35-d of the trial for steers. The 84-d average DMI was intermediate and essentially the mean of the first and last 35-d DMI for steers. Regardless of period, DMI was greater for steers than heifers ($P < 0.01$). Heifer DMI did not change from the first 35-d period to the last 35-d period. Steers also had greater ADG ($P < 0.01$) and G:F ($P < 0.01$) than heifers (Table 1) over the 84-d trial.

There was a sex $\times$ day interaction ($P < 0.01$; Table 2) for acyl ghrelin concentrations, indicating that acyl ghrelin concentrations decreased from day 0 to day 83 in steers, but

concentrations did not change over time in heifers. Total ghrelin concentrations did not change from day 0 to day 83 for steers, but increased over time for heifers ($P < 0.01$). Total ghrelin concentrations for heifers were lower on day 0 than day 0 concentrations in steers, and increased about 55% by day 83. There was no sex effect on the ghrelin ratio ($P = 0.17$), but there was a day effect ($P < 0.01$), indicating that the ratio of acyl:total ghrelin decreased from day 0 to day 83 in both steers and heifers.

3.1 Association of acyl and total ghrelin concentrations with DMI

Associations of day 0 ghrelin with production traits were dependent on the time of day the sample was collected, indicated by a significant ghrelin $\times$ time of day interaction (Table 3). This was observed for day 0 acyl and total ghrelin association with DMI (84-d, first 35-d, and last 35-d). In all instances, the interaction solution was negative, indicating that as the time of sample collection was later in the day, the association of acyl ghrelin with 84-d DMI was decreased. While it is necessary to consider the significant interactions when interpreting the data, the focus will remain on the main association of ghrelin concentrations with production traits to simplify the presentation of the data.

There was an overall positive association of day 0 acyl ghrelin with 84-d DMI ($P = 0.01$), and the last 35-d DMI ($P = 0.02$), and a tendency for the first 35-d DMI ($P = 0.06$), indicating that cattle that consumed more feed had greater acyl ghrelin concentrations on day 0. Acyl ghrelin concentrations on day 0 accounted for 1.09% of the variation (Table 4) for 84-d DMI, 0.50 % of the variation explained by the first 35-d DMI model, and 0.88% of the variation explained by the last 35-d DMI model. Neither day 83 acyl ghrelin, the mean acyl ghrelin, nor the delta acyl ghrelin were significantly associated with DMI ($P > 0.26$).

As with acyl ghrelin, the day 0 total ghrelin parameter was positive in all 3 DMI models, indicating that cattle with greater DMI also had greater total ghrelin concentrations. Total ghrelin concentrations on day 0 accounted for 1.39% of the variation explained by the model for 84-d DMI, 0.22% of the variation explained by the first 35-d DMI model, and 1.03% of the variation explained by the last 35-d DMI model. Total ghrelin concentrations on day 83 were not associated with DMI in any model ($P > 0.10$). The mean total ghrelin concentrations tended to be negatively associated with 84-d DMI and the last 35-d DMI ($P = 0.08$) and accounted for 0.75% of the variation explained by each model.

The ratio of acyl:total ghrelin concentration on day 0 and day 83 were not associated with DMI in any of the models ($P > 0.10$; Table 3). The mean acyl:total ghrelin ratio tended to be positively associated with the last 35-d DMI ($P = 0.08$) and accounted for 0.75% of the variation explained by the model.

3.2 Association of acyl and total ghrelin concentrations with ADG, G:F, and RFI

As previously stated, day 0 acyl ghrelin concentration associations with ADG and G:F, and the ghrelin ratio association with ADG were dependent on the time of day the sample was collected, which was indicated by a significant ghrelin $\times$ time of day interaction. Additionally, the solution for the interaction was negative, indicating that as the time of sample collection was later in the day, the association of ghrelin with ADG and G:F was decreased. The focus of the presentation of the data and discussion will be on the association of ghrelin with production traits to simplify the interpretation.

Day 0 acyl ghrelin concentration was positively associated with ADG ($P < 0.01$; Table 5) and accounted for 1.76% of the variation explained by the ADG model (Table 6). Acyl ghrelin

measured on day 83, the mean of day 0 and day 83 acyl ghrelin concentrations, and the delta from day 0 to day 83 were not associated with ADG ($P > 0.10$). Total ghrelin concentrations on day 0 or day 83, the mean, and the delta were not associated with ADG ($P > 0.10$). There was a positive association ($P = 0.02$) between day 0 ghrelin ratio and ADG. Day 0 ghrelin ratio accounted for 1.66% of the variation explained by the ADG model. The ghrelin ratio on day 83, the mean ghrelin ratio, and the delta ghrelin ratio were not associated with ADG ($P > 0.10$).

There was a tendency for a positive association ($P = 0.09$) of day 0 acyl ghrelin with G:F and day 0 acyl ghrelin concentrations accounted for 0.26% of the variation of the G:F model. Total ghrelin measured on day 0 tended to be positively associated with G:F ($P = 0.08$) and accounted for 1.03% of the variation of the G:F model. Ghrelin ratio on day 0 had a positive association with G:F ($P < 0.05$ for the interaction). Day 0 ghrelin ratio accounted for 2.03% of the variation.

All of the models for RFI had a pseudo $R^2 < 0.01$ (Table 6), indicating almost none of the variation of RFI was explained by the parameters included in the models. However, there was a tendency for the delta acyl and total ghrelin to have a positive association with RFI ($P = 0.10$).

4. Discussion

Acyl ghrelin is thought to stimulate appetite in animals by binding the growth hormone secretagogue receptor 1A [11], which is expressed in neurons in the arcuate nucleus of the hypothalamus [12] and alters the release of neuropeptide Y and agouti-related peptide [6,7,13]. It has previously been shown that chronic injection of bovine acyl ghrelin increases feed intake in cattle [4] and circulating concentrations are elevated in chronic nutrient restriction [14,15] as well as short-term feed restriction [4]. In a previous study, acyl ghrelin measured prior to

feeding was shown to have a positive association with DMI in finishing beef cattle, yet the ghrelin ratio was more associated with DMI [8]. In the Foote, et al. [8] study, ghrelin was measured after the completion of the DMI measurement period, indicating the association of ghrelin with DMI could be reflective of previous feed intake; however it is unclear how ghrelin concentrations change over time. Jennings, et al. [16] showed that acyl-ghrelin concentrations increase with age and/or body fatness in beef steers, but the change was dependent on composition of growth, where cattle accreting more fat had greater acyl-ghrelin concentrations. Another study showed that ghrelin concentrations (presumably acyl-ghrelin) were increased in mature cows compared to calves [17]; however, this could be partially due to energy balance differences [18]. Data presented here indicate that sex is an important factor in the change of ghrelin concentration over a finishing period. Acyl ghrelin decreased in steers, while it did not change in heifers, and total ghrelin increased in heifers only. The decrease in acyl ghrelin for steers from day 0 to day 83 did not reflect the change in DMI from the first 35-d to the last 35-d, where DMI was greater in the last 35-d than the first 35-d. The proportion of ghrelin that was acylated was not different between the sexes, but decreased from day 0 to day 83. For steers, the decrease of the ghrelin ratio was mainly attributed to a decrease in acyl ghrelin, while in heifers the change in the ghrelin ratio was attributed to an increase in total ghrelin. It appears there could be some sex differences in the expression of the enzymes that acylate (GOAT) and deacylate (LYPLA1) ghrelin [19,20], although it is unclear mechanistically how this would occur.

Even though cattle in this experiment were only sampled once each collection day, it appears there were also some trends for ghrelin to change within day. Associations of ghrelin with production variables were dependent on the time of sample collection early in the study. This observation was likely not due to feed restriction; however, there is insufficient data in the

literature regarding peri-prandial ghrelin concentration changes in cattle fed a finishing ration as in this experiment. Additionally, this observation was not due to differences in processing order.

In cattle, ghrelin increases prior to feeding [4,21], but the increase could be smaller in ruminants fed more often [22]. In a study similar to the current experiment, Foote, et al. [8] showed that acyl ghrelin concentration after the feed intake measurement period had a positive association with DMI and accounted for about 6% of the variance explained by the DMI model. In the model for the 84-d DMI presented here, acyl ghrelin had a smaller relative contribution to the variance explained by the model (1%), but a greater amount of the total variance was explained by the model ($R^2 = 0.41$), which was likely due to utilizing a mixed model with breed and sire data. By including breed and sire in the model, the contribution of acyl ghrelin to explain DMI is actually more meaningful as it is independent of breed effects. It should be noted that neither this current experiment or the previous experiment [8] have enough power to truly determine breed effects on ghrelin concentrations.

Acyl ghrelin concentrations in this study were more associated with the 84-d DMI trait than the first 35-d DMI and contributed more to the variance explained by the model. It is interesting that day 83 acyl ghrelin was not associated with DMI. This is in slight contrast to Foote, et al. [8], which showed that acyl ghrelin measured 29 days after the DMI measurement period was associated with DMI.

The contribution of total ghrelin to the variance explained by the model for 84-d DMI, first 35-d DMI, and last 35-d DMI was similar to the contribution of acyl ghrelin. This is in contrast to a previous study that showed no association of total ghrelin concentrations with DMI [8]. It is thought that ghrelin's appetite stimulating effect is mediated through the growth hormone secretagogue receptor 1A, which requires the presence of the acyl group on serine-3 to

bind and activate the receptor [5]. Given this, it is interesting that total ghrelin, which includes acyl and desacyl ghrelin, associates with DMI as one would assume that acyl ghrelin is a more important measure for DMI.

In addition to the effect of ghrelin on feed intake in many species [2,4,6], it appears that ghrelin has effects on BW gain and body composition [3,6,23]. Intraperitoneal and intracerebroventricular administration of ghrelin to rats and mice has been shown to cause increased BW gain, and is mostly attributed to fat mass gain [3,6,23]. Data presented here and in Foote, et al. [8] indicates that acyl ghrelin, or the ratio of acyl:total ghrelin, could play a role in regulating BW gain in beef cattle, as seen in the association of ghrelin concentrations with ADG. In contrast to Foote, et al. [8], the data presented here show no association of total ghrelin with ADG. It is still unclear, however, if acyl ghrelin is associated with body composition, or composition of BW gain. Foote, et al. [8] showed no correlation of acyl ghrelin with carcass composition; however, Jennings, et al. [16] showed that cattle accreting fat at a greater rate also had greater acyl ghrelin concentrations, which would concur with rodent literature [3,6,23].

Rats chronically treated with acyl ghrelin were shown to be more efficient in conversion of feed to BW gain [3]. The data presented here indicates that acyl ghrelin concentrations may not be as important for feed efficiency as the ratio of acyl:total ghrelin. However, Foote, et al. [8] showed that only total ghrelin was associated with G:F, which was a negative association opposed to the positive association observed in the data for total ghrelin. It is not clear what contributed to the divergence in results, but taken together, acyl ghrelin and the ratio of acyl:total ghrelin should be considered in the evaluation of feed efficiency.

Data presented here indicates that there may not be much value in measuring ghrelin concentrations across multiple days. Using either the mean or delta of ghrelin concentrations did

not greatly increase the variance explained by the models compared to a single day 0 measurement. Given that there were interactions of ghrelin concentrations with time of sample collection and the ultradian nature of ghrelin release, there may be some value in having multiple samples within a single day. However, this is difficult with the large numbers of animals that are required to find these small associations, and concentrations shouldn't fluctuate much in ad libitum fed animals [22].

In conclusion, the gut peptide ghrelin appears to be associated with DMI and growth, independent of breed and sire effects. The direction of the association of ghrelin with DMI lends more evidence that ghrelin is involved in the regulation of feed intake and growth of finishing beef cattle, and differences in circulating forms of ghrelin may be responsible for some of the variation in ad libitum feed intake of cattle. It is possible that ghrelin concentrations could be a component of a group of biomarkers for production traits in beef cattle. However, it would be necessary to consider the form of ghrelin and time of collection, as ghrelin concentrations can fluctuate during the feeding period, but in different patterns for steers and heifers.

Acknowledgments

This project was partially funded by NIFA Grant 2011-68004-30214 through the National Program for Genetic Improvement of Feed Efficiency in Beef Cattle.

References

- [1] Kojima M, Kangawa K. Ghrelin, an orexigenic signaling molecule from the gastrointestinal tract. *Curr Opin Pharmacol* 2002; 2:665-668.
- [2] Cummings DE, Purnell JQ, Frayo RS, Schmidova K, Wisse BE, Weigle DS. A preprandial rise in plasma ghrelin levels suggests a role in meal initiation in humans. *Diabetes* 2001; 50:1714-1719.
- [3] Wren AM, Small CJ, Abbott CR, Dhillon WS, Seal LJ, Cohen MA, Batterham RL, Taheri S, Stanley SA, Ghatei MA, Bloom SR. Ghrelin causes hyperphagia and obesity in rats. *Diabetes* 2001; 50:2540-2547.
- [4] Wertz-Lutz AE, Knight TJ, Pritchard RH, Daniel JA, Clapper JA, Smart AJ, Trenkle A, Beitz DC. Circulating ghrelin concentrations fluctuate relative to nutritional status and influence feeding behavior in cattle. *J Anim Sci* 2006; 84:3285-3300.
- [5] Kojima M, Hosoda H, Date Y, Nakazato M, Matsuo H, Kangawa K. Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature* 1999; 402:656-660.
- [6] Nakazato M, Murakami N, Date Y, Kojima M, Matsuo H, Kangawa K, Matsukura S. A role for ghrelin in the central regulation of feeding. *Nature* 2001; 409:194-198.
- [7] Shintani M, Ogawa Y, Ebihara K, Aizawa-Abe M, Miyanaga F, Takaya K, Hayashi T, Inoue G, Hosoda K, Kojima M, Kangawa K, Nakao K. Ghrelin, an endogenous growth hormone secretagogue, is a novel orexigenic peptide that antagonizes leptin action through the activation of hypothalamic neuropeptide y/y1 receptor pathway. *Diabetes* 2001; 50:227-232.

345 [8] Foote AP, Hales KE, Lents CA, Freetly HC. Association of circulating active and total
346 ghrelin concentrations with dry matter intake, growth, and carcass characteristics of
347 finishing beef cattle. *J Anim Sci* 2014; 92:5651-5658.

348 [9] Foote AP, Hales KE, Tait RGJ, Berry ED, Lents CAW, J. E., Lindholm-Perry AK,
349 Freetly HC. Relationship of glucocorticoids and hematological measures with feed
350 intake, growth, and efficiency of finishing beef cattle. *J Anim Sci* 2016; 94:275-283.

351 [10] Foote AP, Hales KE, Kuehn LA, Keisler DH, King DA, Shackelford SD, Wheeler TL,
352 Freetly HC. Relationship of leptin concentrations with feed intake, growth, and efficiency
353 in finishing beef steers. *J Anim Sci* 2015; 93:4401-4407.

354 [11] Bednarek MA, Feighner SD, Pong S-S, McKee KK, Hreniuk DL, Silva MV, Warren VA,
355 Howard AD, Van der Ploeg LHY, Heck JV. Structure–function studies on the new
356 growth hormone-releasing peptide, ghrelin: Minimal sequence of ghrelin necessary for
357 activation of growth hormone secretagogue receptor 1a. *J Med Chem* 2000; 43:4370-
358 4376.

359 [12] Guan X-M, Yu H, Palyha OC, McKee KK, Feighner SD, Sirinathsinghji DJS, Smith RG,
360 Van der Ploeg LHT, Howard AD. Distribution of mrna encoding the growth hormone
361 secretagogue receptor in brain and peripheral tissues. *Mol Brain Res* 1997; 48:23-29.

362 [13] Druce MR, Small CJ, Bloom SR. Minireview: Gut peptides regulating satiety.
363 *Endocrinology* 2004; 145:2660-2665.

364 [14] Field ME, Deaver SE, Rhoads RP, Collier RJ, Rhoads ML. Effects of prolonged nutrient
365 restriction on baseline and periprandial plasma ghrelin concentrations of postpubertal
366 holstein heifers. *J Dairy Sci* 2013; 96:6473-6479.

- 367 [15] Wertz-Lutz AE, Daniel JA, Clapper JA, Trenkle A, Beitz DC. Prolonged, moderate
368 nutrient restriction in beef cattle results in persistently elevated circulating ghrelin
369 concentrations. J Anim Sci 2008; 86:564-575.
- 370 [16] Jennings JS, Wertz-Lutz AE, Pritchard RH, Weaver AD, Keisler DH, Bruns K.
371 Circulating ghrelin and leptin concentrations and growth hormone secretagogue receptor
372 abundance in liver, muscle, and adipose tissue of beef cattle exhibiting differences in
373 composition of gain. J Anim Sci 2011; 89:3954-3972.
- 374 [17] Miura H, Tsuchiya N, Sasaki I, Kikuchi M, Kojima M, Kangawa K, Hasegawa Y,
375 Ohnami Y. Changes in plasma ghrelin and growth hormone concentrations in mature
376 holstein cows and three-month-old calves. J Anim Sci 2004; 82:1329-1333.
- 377 [18] Bradford BJ, Allen MS. Negative energy balance increases periprandial ghrelin and
378 growth hormone concentrations in lactating dairy cows. Dom Anim Endocrinol 2008;
379 34:196-203.
- 380 [19] Shanado Y, Kometani M, Uchiyama H, Koizumi S, Teno N. Lysophospholipase i
381 identified as a ghrelin deacylation enzyme in rat stomach. Biochem Biophys Res
382 Commun 2004; 325:1487-1494.
- 383 [20] Yang J, Brown MS, Liang G, Grishin NV, Goldstein JL. Identification of the
384 acyltransferase that octanoylates ghrelin, an appetite-stimulating peptide hormone. Cell
385 2008; 132:387-396.
- 386 [21] Wertz-Lutz AE, Jennings JS, Clapper JA. Plasma ghrelin concentrations of beef cattle
387 consuming a similar amount of dietary energy supplied by different ingredients. J Anim
388 Sci 2010; 88:2289-2299.

- 389 [22] Sugino T, Yamaura J, Yamagishi M, Ogura A, Hayashi R, Kurose Y, Kojima M,
390 Kangawa K, Hasegawa Y, Terashima Y. A transient surge of ghrelin secretion before
391 feeding is modified by different feeding regimens in sheep. *Biochem Biophys Res*
392 *Commun* 2002; 298:785-788.
- 393 [23] Tschop M, Smiley DL, Heiman ML. Ghrelin induces adiposity in rodents. *Nature* 2000;
394 407:908-913.

395

396

397 **Table 1.** Production traits of individually fed beef steers and heifers fed a finishing ration during an 84-d

398 evaluation period

Item	Steers			Heifers			SEM ¹	Sex effect	Day effect	Sex × Day effect
	First 35 d	Last 35 d	84 d	First 35 d	Last 35 d	84 d				
DMI (kg/d)	9.36 ^b	9.92 ^a	9.63 ^{ab}	8.36 ^c	8.28 ^c	8.37 ^c	0.14	< 0.01	0.04	< 0.01 ²
ADG (kg/d)	.	.	1.86 ^a	.	.	1.49 ^b	0.027	< 0.01	.	.
G:F (kg/kg)	.	.	0.193 ^a	.	.	0.177 ^b	0.002	< 0.01	.	.

399 Abbreviations: DMI, Dry matter intake; ADG, average daily gain; G:F, gain:feed ratio (average daily

400 gain:dry matter intake)

401 ¹Standard error of the mean (n = 236).402 ^{a,b,c}Values within row with differing superscripts differ $P < 0.05$.

Table 2. Acyl and total ghrelin concentrations and the ghrelin ratio (acyl:total ghrelin) in finishing beef steers and heifers measured on day 0 and 84 of a DMI and ADG evaluation period

Item	Steers		Heifers		SEM ¹	Sex effect	Day effect	Sex × Day effect
	Day 0	Day 83	Day 0	Day 83				
Log acyl ghrelin (log pg/mL)	2.07 ^a	1.94 ^b	1.90 ^b	1.93 ^b	0.021	< 0.01	< 0.01	< 0.01
Acyl ghrelin (pg/mL)	117.5	87.1	79.4	85.1	.	.	.	.
Total ghrelin (pg/mL)	345.3 ^a	363.5 ^{ab}	245.6 ^c	381.1 ^b	8.31	< 0.01	< 0.01	< 0.01
Log ghrelin ratio	-0.45	-0.60	-0.47	-0.64	0.025	0.17	< 0.01	0.58
Ghrelin ratio	0.35	0.25	0.34	0.23	.	.	.	.

¹Standard error of the mean (n = 236).

^{a,b,c}Values within row with differing superscripts differ $P < 0.05$.

Table 3. Mixed model estimates for acyl ghrelin, total ghrelin, the ghrelin ratio (acyl:total ghrelin) concentrations on day 0 or 83, the mean of day 0 and 83, and the delta (change) from day 0 to day 83 associated with dry matter intake (DMI) of 236 individually fed finishing beef steers and heifers.

Dependent Variable	84-d DMI			First 35-d DMI			Last 35-d DMI		
	Solution ^a	SEM	P-value	Solution	SEM	P-value	Solution	SEM	P-value
Acyl ghrelin									
Day 0	7.01 ^b	2.881	0.01 ^d	5.59 ^b	2.910	0.06 ^c	7.49 ^b	3.220	0.02 ^d
Day 83	0.52 ^c	0.459	0.26	0.49	0.461	0.45	0.62 ^c	0.520	0.20
Mean	0.40	0.484	0.41	0.14	0.480	0.77	0.64	0.545	0.24
Delta	0.001	0.001	0.53	0.001	0.001	0.36	0.001	0.001	0.85
Total ghrelin									
Day 0	0.02 ^{b,d}	0.001	0.05	0.014 ^{b,e}	0.001	0.11	0.017 ^{b,e}	0.0094	0.08
Day 83	-0.001 ^c	0.0009	0.33	-0.001	0.0009	0.28	-0.001 ^c	0.0010	0.36
Mean	-0.002	0.0012	0.08	-0.002	0.0012	0.12	-0.002	0.0013	0.07
Delta	-0.001	0.0008	0.70	-0.0001	0.0008	0.91	-0.001	0.0009	0.58
Ghrelin ratio									
Day 0	0.36	0.316	0.26	0.17	0.316	0.59	0.031 ^c	0.4018	0.94
Day 83	0.48 ^c	0.359	0.17	0.46	0.361	0.20	0.54 ^c	0.401	0.18
Mean	0.58	0.386	0.13	0.40	0.385	0.30	0.76	0.434	0.08
Delta	0.11	0.340	0.74	0.25	0.340	0.47	-0.054	0.3840	0.89

Abbreviation: DMI, dry matter intake

^a Ghrelin solution is the estimate from the fixed effect of ghrelin concentration (acyl, total, and acyl:total ghrelin ratio on day 0, day 83, the mean of the 2 d, or the change [delta] from day 0 to day 83) in the respective mixed model analysis. All measures of ghrelin were analyzed in separate models. Other factors included the random effect of sire and fixed effect of breed and sex. Additional fixed effects were included as noted.

^b Model parameters also included the covariate of time of day of blood sample collection and ghrelin × time of day of sample collection interaction.

420 ^c Model parameters also included covariate of time of day of blood sample collection.

421 ^d Ghrelin $\times$ time of day of sample collection interaction significant ($P < 0.05$)

422 ^e Ghrelin $\times$ time of day of sample collection interaction tendency ($0.05 < P < 0.10$)

423

424

425

Table 4. Percentage of the variance explained by ghrelin concentrations and the pseudo R² values for the mixed model analysis of the association of acyl, total, or the ratio of acyl:total ghrelin concentrations measured on day 0 or 83 and the mean of the 2 measures and the delta (change) from day 0 to day 83 with dry matter intake (DMI) of 236 individually fed finishing beef steers and heifers.

Dependent Variable	84-d DMI		First 35-d DMI		Last 35-d DMI	
	% Variance Explained	Pseudo R ²	% Variance Explained	Pseudo R ²	% Variance Explained	Pseudo R ²
Acyl ghrelin						
Day 0	1.09	0.41	0.50	0.34	0.88	0.45
Day 83	0.09	0.39	< 0.01	0.33	0.23	0.44
Mean	< 0.01	0.37	< 0.01	0.33	0.17	0.41
Delta	< 0.01	0.37	< 0.01	0.33	< 0.01	0.41
Total ghrelin						
Day 0	1.39	0.40	0.22	0.33	1.03	0.44
Day 83	< 0.01	0.39	0.38	0.34	< 0.01	0.44
Mean	0.75	0.38	0.36	0.34	0.75	0.42
Delta	< 0.01	0.37	< 0.01	0.33	< 0.01	0.41
Ghrelin ratio						
Day 0	0.22	0.37	< 0.01	0.33	0.03	0.43
Day 83	0.20	0.40	0.13	0.45	0.20	0.44
Mean	0.50	0.38	< 0.01	0.33	0.75	0.41
Delta	< 0.01	0.37	< 0.01	0.33	< 0.01	0.41

Abbreviation: DMI, Dry matter intake.

Table 5. Mixed model estimates for acyl ghrelin, total ghrelin, the ghrelin ratio (acyl:total ghrelin) concentrations on day 0 or 83, the mean of day 0 and 83, and the delta (change) from day 0 to day 83 associated with growth and feed efficiency of 236 individually fed finishing beef steers and heifers.

Dependent Variable	ADG			G:F			RFI		
	Solution ^a	SEM	P-value	Solution	SEM	P-value	Solution	SEM	P-value
Acyl ghrelin									
Day 0	1.97 ^{b,d}	0.654	< 0.01 ^d	0.085 ^{b,e}	0.0500	0.09 ^c	-0.11	0.203	0.59
Day 83	2.15 ^{b,e}	1.120	0.06 ^e	-0.009	0.0078	0.27	0.28	0.261	0.29
Mean	0.070	0.110	0.53	-0.001	0.008	0.96	-0.018	0.260	0.94
Delta	-0.001	0.0003	0.81	-0.001	0.0001	0.29	0.001	0.0007	0.10
Total ghrelin									
Day 0	0.0002 ^c	0.0002	0.52	0.0003 ^c	0.0002	0.08	-0.0004	0.0004	0.44
Day 83	-0.0001 ^c	0.0002	0.67	-0.0001 ^c	0.0001	0.80	-0.0007	0.0005	0.17
Mean	-0.0002	0.0003	0.42	0.00001	0.0002	0.58	0.0019	0.0006	0.76
Delta	-0.0001	0.0003	0.81	-0.0001	0.0001	0.29	0.001	0.0006	0.10
Ghrelin ratio									
Day 0	1.49 ^{b,d}	0.631	0.02 ^d	0.10 ^{b,d}	0.048	0.59 ^d	0.048	0.179	0.77
Day 83	0.012 ^c	0.082	0.88	-0.008 ^c	0.0061	0.22	-0.005	0.204	0.93
Mean	0.079	0.0885	0.38	-0.003	0.0066	0.70	-0.024	0.219	0.91
Delta	0.0032	0.0790	0.97	-0.002	0.0059	0.72	0.145	0.190	0.45

Abbreviations: ADG, average daily gain, G:F, gain:feed ratio (average daily gain:dry matter intake); RFI, residual feed intake.

^a Ghrelin solution is the estimate from the fixed effect of ghrelin concentration(acyl, total, and acyl:total ghrelin ratio on day 0, day 83, the mean of the 2 d, or the change [delta] from day 0 to day 83) in the respective mixed model analysis . All measures of ghrelin were analyzed in separate models. All models included the random effect of sire and fixed effect of breed and sex, except for RFI where sex was excluded because RFI was calculated within sex group. Other fixed effects were included as noted.

445 ^b Model parameters also included the covariate of time of day of blood sample collection and
446 ghrelin $\times$ time of day of sample collection interaction.

447 ^c Model parameters also included covariate of time of day of blood sample collection.

448 ^d Ghrelin $\times$ time of day of sample collection interaction significant ($P < 0.05$)

449 ^e ghrelin $\times$ time of day of sample collection interaction tendency ($0.05 < P < 0.10$)

450

Table 6. Percentage of the variance explained by ghrelin concentrations and the pseudo R² values for the mixed model analysis of the association of acyl, total, or the ratio of acyl:total ghrelin concentrations measured on day 0 or 83 and the mean of the 2 measures and the delta (change) from day 0 to day 83 with growth and feed efficiency of 236 individually fed finishing beef steers and heifers.

Dependent Variable	ADG		G:F		RFI	
	% Variance Explained	Pseudo R ²	% Variance Explained	Pseudo R ²	% Variance Explained	Pseudo R ²
Acyl ghrelin						
Day 0	1.76	0.50	0.26	0.19	< 0.01	< 0.01
Day 83	0.41	0.48	< 0.01	0.16	< 0.01	< 0.01
Mean	< 0.01	0.45	< 0.01	0.16	< 0.01	< 0.01
Delta	< 0.01	0.45	0.26	0.17	0.95	< 0.01
Total ghrelin						
Day 0	0.48	0.48	1.03	0.19	< 0.01	< 0.01
Day 83	< 0.01	0.48	< 0.01	0.17	0.52	< 0.01
Mean	0.35	0.45	< 0.01	0.16	< 0.01	< 0.01
Delta	0.65	0.46	0.77	0.17	0.83	< 0.01
Ghrelin ratio						
Day 0	1.66	0.50	2.03	0.19	< 0.01	< 0.01
Day 83	< 0.01	0.48	< 0.01	0.18	< 0.01	< 0.01
Mean	0.42	0.45	< 0.01	0.16	< 0.01	< 0.01
Delta	< 0.01	0.45	< 0.01	0.16	< 0.01	< 0.01

Abbreviations: ADG, average daily gain; G:F, gain:feed ratio (average daily gain:dry matter intake); RFI, residual feed intake.