Coagulopathy of Trauma

Authors: Mitchell J. Cohen MD¹ and S. Ariane Christie MD²

Affiliation for all authors: ¹ Department of Surgery, San Francisco General Hospital and the University of California, San Francisco; San Francisco, California.

Email: ¹ Mitchell.Cohen@ucsf.edu and ² Sabrinah.Christie@ucsf.edu

Address for all authors:

Mitchell J. Cohen MD,
Department of Surgery, Ward 3A
San Francisco General Hospital
1001 Potrero Avenue, Room 3C-38
San Francisco, CA 94110
Phone: (415) 206-3764
Fax: (415) 206-5484

Corresponding Author: Mitchell J. Cohen

Key Words (3-8) Trauma, hemorrhage, coagulation, coagulopathy, Trauma-induced coagulopathy, Targeted Resuscitation, Bleeding

Abstract (*<=100 words*)

Coagulopathy is common following injury and develops independently from iatrogenic, hypothermic, and dilutional causes. Despite considerable research on the topic over the past decade, trauma induced coagulopathy continues to portend poor outcomes including decreased survival. In this article, we review the current evidence regarding the diagnosis and mechanisms underlying trauma induced coagulopathy and summarize the debates regarding optimal management strategy including product resuscitation, potential pharmacologic adjuncts, and targeted approaches to hemostasis. Throughout, we will identify areas of continued investigation and controversy in the understanding and management of trauma induced coagulopathy (TIC).

Key Points (3-5)

- Trauma-induced coagulopathy is an endogenous hypocoagulable state distinct from iatrogenic causes.
- Activation of Protein C pathway is a key mechanistic mediator of traumatic coagulopathy via multiple downstream effects including thrombin diversion, deactivation of coagulation factors, and de-repression of fibrinolysis.
- Standard coagulation tests and functional viscoelastic assays are commonly utilized in the diagnosis and management of TIC.
- Balanced resuscitation is the mainstay of coagulopathy treatment, but precise ratios for empiric resuscitation and optimal monitoring protocols for transfusion practice remain unknown.
- Patients with traumatic coagulopathy have worse outcomes including increased rates of transfusion, infection, thromboembolism, acute lung injury, multiorgan failure, and death.

Outline

- **Introduction**
- **Patient Evaluation and Overview**
 - **Mechanism and Pathophysiology**
 - **APC and Fibrinolysis**

- Platelet
 - Endothelial Involvement
 - Microparticles
- Diagnosis
 - Standard clinical and laboratory assessment
 - Diagnostic criteria
 - Standard Assays
 - Viscoelastic Assays
 - Scoring Systems
 - Phenotypes of Traumatic Coagulopathy
- Pharmacologic Treatment Options
 - Antifibrinolytics
 - Recombinant Factor Concentrates
 - Recombinant Activated Factor VII
 - Prothrombin Complex Concentrate
 - Fibrinogen Concentrate
- Non- Pharmacologic Treatment Options
 - Balanced Resuscitation and Product Ratios
- Combination Therapies
 - Targeted Resuscitation Practice
- Surgical Treatment Options
- Treatment Resistance/ Complications
 - Compounding Coagulopathy
 - Hypothermia
 - Acidosis
 - Hemodilution

- **Disseminated Intravascular Coagulopathy**
- **Evaluation of Outcomes/ Long-Term Recommendations**
 - **Transfusion Requirements**
 - **Hypercoagulability**
 - **Acute Lung Injury, Multiorgan Failure and Death**
- **Summary**

Introduction

Bleeding remains the leading cause of preventable death following injury¹. Contributing to this problem, coagulopathy develops in approximately one-third of all injured patients²⁻⁴, resulting in worsened outcomes including higher transfusion requirements, increased multi-organ system failure, increased hospital, ICU, and ventilator days, and increased mortality^{2,3,5,6}.

History of Coagulopathy in Trauma

Although coagulopathy was known to occur following injury, until recently coagulation was not viewed as a critical driver of post-injury physiology. Instead, injured patients were thought to be coagulopathic due only to the iatrogenic secondary effects of hemodilution, hypothermia, and acidosis^{7,8}. In 2003, two independent investigators described admission perturbations of prothrombin time (PT) and partial thromboplastin time (PTT) in newly injured patients prior to significant fluid administration^{2,3}. This phenomenon, which correlated with increasing injury severity and mortality, became known as “Acute Traumatic Coagulopathy” (now “Trauma-induced coagulopathy”) and effectively changed the paradigm of modern trauma care^{2,3,9}. The study of coagulation and inflammation derangements after injury now constitutes one of the most active areas of ongoing trauma research.

This review will address the current evidence regarding the diagnosis, mechanisms, and management of trauma-induced coagulopathy, highlighting areas of ongoing debate and controversy. While TIC will be emphasized, it is equally important to recognize that coagulopathy following trauma is often caused or compounded by additional contributors of disordered coagulation including hypothermia, acidosis, dilution with large volume of intravenous fluid or unbalanced blood product all termed iatrogenic coagulopathy. Management of the injured coagulopathic patient must therefore include a high suspicion for and treatment of multiple different potential etiologies of dysfunctional clotting.

Patient Evaluation and Overview

Mechanism and Pathophysiology

Multiple distinct but highly integrated pathways have been implicated as mediators of trauma-induced coagulopathy (see Figure 1). Delineating the exact pathophysiology and interplay between disordered coagulation and inflammation mechanisms remains the subject of ongoing research. Here, we describe the most important known contributors to TIC.

1) *Activated Protein C and Fibrinolysis*

Trauma-induced coagulopathy is an endogenous hypocoagulable state that occurs in the setting of tissue hypo-perfusion (base deficit) and is primarily mediated by activation of Protein C (see Figure 2).¹⁰ Multiple prospective clinical studies have linked early coagulopathy in critically injured trauma patients to Protein C depletion (activated Protein C elevation), and increased risk of acute lung injury (ALI), ventilator-associated pneumonia (VAP), multi- system organ failure (MOSF), and death^{6,10,11}. Protein C is a serine protease with both anticoagulant and inflammomodulatory functions^{12,13}. When severe injury is accompanied by shock (tissue hypoperfusion), increased endothelial and circulating thrombomodulin bind thrombin forming thrombomodulin-thrombin complex which subsequently activates Protein C¹⁰. Activated Protein C (aPC) deactivates factors V and VIII inhibiting clot formation, and depletes plasminogen activator inhibitor-1 (PAI-1) leading to unopposed fibrinolysis with increased levels of tPA and D-dimer¹⁴.

Fibrinolytic activity is further exacerbated by reduced activation of thrombin-activatable fibrinolysis inhibitor (TAFI) as thrombin is diverted to PC activation.¹⁵ Severe fibrinolysis in TIC portends increased mortality,¹⁶⁻¹⁸ and even low degrees of clot lysis have been associated with poor outcomes¹⁹. In a murine model aPC inhibition prevented TIC following trauma and hemorrhagic shock¹².

Conversely, over-inhibition of fibrinolysis, termed “fibrinolysis shut down”, has been demonstrated to be an independent predictor of adverse outcomes after injury including increased mortality.²⁰ Furthermore, recent prospective cohort data suggests that more severely injured patients present with fibrinolysis shutdown than either hyper- or physiologic fibrinolysis.²¹ This data has important implications with regards to the utility of potential fibrinolytic inhibitors, which will

need to be carefully targeted to the physiologic range and avoid over-inhibition leading to fibrinolysis shutdown.

2) *Platelets*

Platelet deficit and dysfunction are also likely to be significant contributors to trauma-induced coagulopathy²².

a) *Relative Thrombocytopenia*

Multiple prospective cohort studies have demonstrated that relatively lower admission platelet counts are associated with increased all-cause, hemorrhagic and CNS mortality, and increased blood utilization after injury, even when the initial platelet count remains well within the normal limits^{23,24}. Additionally, platelet count has been shown to decrease substantially over the course of hospitalization.

b) *Functional Platelet Impairment*

Severe injury is also associated with impaired platelet function. In 2001, Jacoby et al used flow cytometry and light aggregometry first identified decreased admission platelet function in injury non-survivors,²⁵ and, in a separate analysis, in patients with head injury at 24 hours. Thrombelastograph platelet mapping data demonstrated that severely-injured trauma patients had impaired platelet stimulation in response to adenosine diphosphate and arachadonic acid stimulation compared to healthy human volunteers, with impairment proportionate to injury severity²⁶.

Another prospective clinical study demonstrated 46% of severely-injured patients on admission and 91% of patients at 120 hours had some degree of platelet dysfunction by multiplate impedance aggregometry despite normal platelet counts.²⁷ Impaired admission platelet function in response to arachadonic acid, collagen, and thrombin receptor activating peptide were predictive of death. Data from multiple additional animal and clinical studies have corroborated findings of impaired platelet function after severe-injury and traumatic brain injury.^{28 29}

3) *Endothelial Involvement*

Recent investigations have also demonstrated evidence that endothelial dysfunction likely plays a role in development of TIC³⁰. Admission plasma samples from severely injured patients demonstrate elevated circulating levels of Syndecan-1, a protein normally found in the glycocalyx of

the endothelium. Soluble syndecan-1 was associated with increased aPC, prolonged PTT, and elevated adrenaline levels, suggesting that tissue hypoperfusion and catecholamine stimulation after injury may result in degradation of the endothelial glycocalyx and contribute to coagulopathy. To date, however, there is no experimental confirmation of this theory making it possible that the association between catecholamines and coagulopathy is only correlative.

4) Microparticles

Emerging research suggests that microparticles may also play a role in the mechanism of TIC. Some evidence suggests that systemic release of thrombin-rich microparticles, which likely function normally in local hemostasis after tissue injury, may cause a coagulopathic state similar to DIC. Elevated circulating endothelial-, erythrocyte-, and leukocyte-derived microparticles have been identified in the plasma of injured patients compared with non-injured controls whereas coagulopathic patients demonstrated lower levels of platelet-derived and tissue-factor-positive microparticles compared to non-coagulopathic patients³¹. In a single small study elevated circulating microparticles were also found in patients with traumatic brain injury compared with controls³².

Diagnosis

Standard clinical and laboratory assessment after severe injury

As with the management of all severely injured patients, diagnosis of TIC should be performed within the context of the Advanced Trauma Life Support evaluation. Obvious sources of active bleeding should be promptly temporized with compression or other hemostatic measures, hypotension should be addressed with administration of PRBC, and the patient should be warmed with heated blankets as the patient is exposed. Severe injury and hypotension should prompt a high degree of suspicion for traumatic coagulopathy. Standard trauma labs should be obtained as soon as intravenous access is secured and can provide indicators of the presence of tissue hypoperfusion and coagulopathy. Table 1 describes standard admission trauma labs and their role in the diagnosis of TIC.

Table 1: Standard Admission Trauma Laboratory Measures in the Evaluation of Trauma-induced Coagulopathy

Lab	Level	Utility in the diagnosis of TIC
pH	Low	Significant hypoperfusion probable

Base Deficit/Excess	Negative	Significant hypoperfusion probable
Hemoglobin/ Hematocrit	Low	Likely significant blood loss
Platelet Count	Low	Absolute/relative thrombocytopenia
Partial Thromboplastin Time	Prolonged	Diagnostic of TIC
Prothrombin Time/International Normalized Ratio	Prolonged	Diagnostic of TIC
<p>We also recommend the standard collection of baseline fibrinogen and d-dimer, which can provide a surrogate estimation of factor consumption and fibrinolysis respectively.</p>		

Diagnostic criteria

Multiple assays currently play a role in the diagnosis of trauma-induced coagulopathy.

1) Standard Assays

Trauma-induced coagulopathy was initially defined by prolongation of the standard coagulation assays partial thromboplastin time (PTT) and prothrombin time/international normalized ratio (PT/INR), and these remain the most widely utilized method for the diagnosis of TIC. Different cutoffs for these assays have been described in the literature (See Table 2).

Table 2: Commonly Utilized Cutoffs for Conventional Coagulation Assays in Trauma-induced Coagulopathy

Assay	Cutoff		
Partial Thromboplastin Time (PTT)	> 34-60 seconds ^{3, 30}	or	Any value > 1.5x institutional reference range³³
Prothrombin Time (PT)	> 18 seconds		
International Normalized Ratio (INR)	> 1.2-1.5 ^{3,6,34,35}		

Macleod et. al. reported that alterations in the cutoffs for PTT and PT did not alter the predictive value of these variables for predicting death in an adjusted regression model³. Lower thresholds for INR cutoffs demonstrated improved sensitivity to discriminate patients with higher transfusion requirements in shock in severely-injured population (ISS>15) in a multicenter retrospective study but at least one prospective study has refuted this³⁶.

However, concerns have been raised regarding the use of standard coagulation assays as the benchmark for trauma-induced coagulopathy. PTT and INR were initially designed to test heritable coagulopathy, and standard reference ranges were generated using data from healthy volunteers.

Additionally, concerns have been raised regarding the length of time required to run standard coagulation tests when rapid and ongoing diagnosis and treatment of the coagulopathic patient are essential to reverse pathophysiology and improve outcomes.

2) *Viscoelastic Assays*

There has been increased interest and utilization of point-of-care functional tests such as thromboelastography for the diagnosis and management of patients with TIC. Thromboelastography is a modality which assesses multiple real-time viscoelastic properties of coagulation including:

- Time to clot initiation
- Clot propagation
- Clot strength
- Clot breakdown (fibrinolysis).

The addition of specific clotting activators and inhibitors can be used to assay different contributors to clot formation or breakdown. Multiple studies have demonstrated the capacity of thromboelastography to diagnose hypocoagulability and predict transfusion and mortality in the trauma population³⁷⁻⁴⁰. Thromboelastography has been validated against standard coagulation tests⁴⁰, and assays of both thrombin generation⁴¹ and fibrinolysis⁴². However, there remains ongoing debate regarding the superiority of thromboelastography compared to other assays, in particular with regards to fibrinolysis as plasmin-antiplasmin levels have demonstrated high sensitivity⁴³.

3) *Scoring systems*

Multiple scoring systems have been generated to predict need for massive transfusion, including:

- The Trauma-Associated Severe Hemorrhage (TASH) score⁴⁴
- The McLaughlin score⁴⁵
- The Assessment of Blood Consumption (ABC) score⁴⁶.

A retrospective review comparing the three scoring systems failed to demonstrate a difference between their capacity to predict massive transfusion⁴⁶. Importantly, the scoring systems do not take into account coagulation parameters and likely reflect significant hemorrhage due to injury rather than TIC proper. None of the scoring systems have been widely utilized in the diagnosis of TIC.

Phenotypes of Trauma-Induced Coagulopathy (TIC)

There is evidence to suggest that traumatic coagulopathy is not one single entity, but rather consists of multiple distinct but related pathophysiologic subtypes. A principle component analysis of a large prospective cohort of injured patients demonstrated two such phenotypes:

1. Coagulation Factor Deficiency TIC

- characterized by abnormality of standard coagulation tests and increased mortality

2. Fibrinolytic TIC

- characterized by aPC elevation and is associated with increased end organ failure, infectious complications and mortality⁹

There are likely even more TIC subtypes that have yet to be identified. By diagnosing and treating specific deficits underlying different coagulopathic phenotypes, it may be possible to streamline individualized care and improved outcomes.

Pharmacologic Treatment Options

Although balanced product transfusion currently remains the mainstay of treatment for TIC, there are several pharmacologic agents which have the potential to be efficacious based on our mechanistic understanding of traumatic coagulopathy.

Antifibrinolytics

Evidence of hyperfibrinolysis as a critical mechanism underlying traumatic coagulopathy has led to interest antifibrinolytic agents as potential adjuncts for the treatment of TIC. These agents include Tranexamic acid (TXA), Aminocaproic acid and Aprotinin, of which only TXA has been widely studied in trauma patients.

The Clinical Randomization of an Antifibrinolytic in Significant Hemorrhage (CRASH-2) trial was a large international randomized trial which reported a 1.5% absolute mortality reduction in patients administered TXA compared to placebo, although there was no difference in blood product transfusion between groups. Furthermore, the subset of patients who received TXA three or more hours after injury had increased mortality⁴⁷. Of note, enrollment criteria for this study failed to incorporate coagulation data and included any patients with or at risk for significant hemorrhage limiting the generalizability of these findings.

The Military Application of Tranexamic-acid in Trauma Emergency Resuscitation (MATTERs) study was a retrospective observational study which identified reduced mortality in patients who received TXA, with larger differences identified in patients requiring massive transfusion⁴⁸. Despite achieving considerable acceptance as an adjunctive treatment for TIC, there has not yet been sufficient evidence to support the routine use of TXA in the trauma setting⁴⁹.

Recombinant factor concentrates

Additional pharmaceutical hemostatic agents with potential utility in the treatment of TIC include recombinant factor concentrates including recombinant factor VIIa, prothrombin complex concentrate (PCC), and fibrinogen.

Recombinant factor VII

Fresh frozen plasma (FFP) is often utilized to reverse coagulopathy, but has limitations including the requirement for thawing and crossmatching, incomplete and variable factor level repletion⁵⁰⁻⁵² and coagulation test reversal^{53,54} and complications such as transfusion related acute lung injury (TRALI) and circulatory overload (TACO)^{55,56}.

Physiologically targeted resuscitation could theoretically arrest coagulopathy in a rapid fashion and avoid many of the pitfalls of traditional plasma. There was significant interest in utilizing recombinant factor VIIa (rFVIIa)

to do this, however, key trials failed to show benefit and suggested increased thrombotic complications⁵⁷. It has been pointed out that these trials generally utilized rFVIIa late in the resuscitation process, when poor outcomes were relatively certain. It is not known whether rFVIIa could potentially have a role in trauma resuscitation as part of a more targeted empiric therapy for specific subpopulations of patients with TIC, but at present factor VIIa has not gained widespread support for use in the trauma setting.

Prothrombin Complex Concentrate

Prothrombin complex concentrate (PCC) comes in several varieties including 3 and 4-factor formulations. The most common formulation is four-factor PPC (4F-PCC), a human-plasma derived concentrate of Vitamin K dependent clotting factors II, VII, IX, X which received FDA approval in April, 2013 as an alternative to urgent warfarin reversal in the setting of acute bleeding or need for urgent surgery. Since that time it has developed usage for reversal of non-warfarin coagulopathy including coagulopathy induced by new oral anticoagulants and in coagulopathy of non-medication etiologies⁵⁸⁻⁶⁰.

In a porcine trauma models, PCC demonstrated more rapid and effective hemostasis than FFP in correction of acquired coagulopathy⁶¹. Multiple retrospective studies have reported decreased time to reversal, decreased product usage, and decreased mortality in patients treated with PCC in the setting of traumatic coagulopathy in general and TBI population^{62,63}. However, there are ongoing concerns regarding the potential for increased thromboembolic complications and cost with 4F-PCC compared to plasma. At present, the paucity of prospective data limits our ability to draw conclusions regarding the safety and efficacy of 4F-PCC in severe injury.

Fibrinogen Concentrate

Fibrinogen deficit has been shown to predict trauma-induced coagulopathy¹¹, and as described above fibrinolysis constitutes an important component of the pathophysiology of TIC. In a prospective cohort study of 517 trauma patients, low fibrinogen levels were independent predictor of mortality at 24 hours and 28 days ($p < 0.001$), and administration of cryoprecipitate was associated with improved survival. Although the threshold of sufficient fibrinogen to support normal clotting has not been rigorously studied in trauma patients, routine testing and repletion of low admission levels or function may be a reasonable adjunct to the treatment of TIC. Fresh frozen

plasma does not contain sufficient amounts of fibrinogen for adequate replacement⁶⁴ and in the United States cryoprecipitate is commonly utilized for this purpose. In Europe, retrospective studies in a trauma population have reported good efficacy of fibrinogen concentrate in correcting functional deficits⁶⁵, however this product is not currently approved in the United States. Current European guidelines for hemorrhage management in trauma patients with fibrinogen levels < 1.5-2.0 g/L or viscoelastic signs of a functional fibrinogen deficit recommend an initial fibrinogen concentrate dose of 3-4g (equivalent to 15-20 single donor units of cryoprecipitate), with further dosing guided by laboratory or viscoelastic testing (Figure 4).

Non-Pharmacologic Treatment Options

Treatment of coagulopathy in the injured patient includes early diagnosis, prompt hemostasis and early hemorrhage control, prevention of complicating causes of coagulopathy(hypothermia, acidosis, hemodilution), and blood product transfusion with ffp, platelets, and cryoprecipitate. There is ongoing debate regarding the optimal protocol for delivery of blood product resuscitation.

Balanced resuscitation and resuscitation ratios

Although early transfusion was conducted with whole blood, in the last quarter of the 21st century standard practice favored resuscitation of the injured patient with large volumes of crystalloid and packed red blood cells(PRBC)⁶⁶. Then, in the early 2000's, retrospective military data from Afghanistan and Iraq suggested a mortality benefit to trauma resuscitation with a balanced ratio of packed red blood cells⁶⁷. Civilian retrospective data echoed these findings using a 1:1:1 ratio of PRBCs, FFP and platelets^{68,69} and the trauma community subsequently began to shift toward balanced blood product resuscitation (see Figure 3).

The Prospective, Observational, Multicenter, Major Trauma Transfusion (PROMMTT) study demonstrated in a large multicenter cohort that patients who received increased plasma to RBC ratios had reduced 6-hour mortality compared to those who received less plasma⁷⁰. In an attempt to delineate the ideal empiric transfusion ratio, The Pragmatic, Randomized Optimal Platelet and Plasma Ratios (PROPPR) trial randomized severely-injured patients to 1:1:1 vs 1:1:2 PRBC to FFP to platelet resuscitation but ultimately failed to demonstrate a difference between resuscitation groups for 24 hour or 28 day mortality⁷¹, which was likely at least in part due to poor

separation between treatment groups. At present, the precise “ideal” PRBC to plasma to platelet ratio remains undetermined, however, it remains clear that limitation of crystalloid and balanced product transfusion approximating whole blood improve outcomes and should be priorities of trauma resuscitation.

Combination Therapies

Targeted Resuscitation Practice

A different paradigm for treatment of the patient with TIC proposes a targeted approach to resuscitation. In this approach, laboratory assays, including point-of-care thromboelastography are obtained in a serial fashion and used to guide product transfusion and administration of pharmaceutical adjuncts. This allows resuscitation to be tailored to the individual patient in real time, and coordinates the different modalities available for treatment. An additional advantage to this approach is the ability to provide dynamic management as the patient’s condition changes. Objections to this approach include the need for infrastructure to support serial and rapid thromboelastography assays and the fact that these tests are dependent on and vary by user skill and interpretation.

A recent Cochrane review suggested that there was insufficient evidence to recommend thromboelastography-based transfusion guidelines as superior to established transfusion practice⁷². However, recently published prospective randomized data from the Denver group shows a mortality benefit when viscoelastic functional testing (thromboelastography, TEG) was used to guide massive transfusion protocols compared to conventional coagulation assays (CCA)⁷³. Survival in the TEG group was significantly higher than the CCA group (28-day mortality 36.4% CCA vs. 19.6% TEG) and 6-hour mortality was significantly lower in the TEG group (21.7% CCA vs. 7.1% TEG, $p=0.032$). Importantly, not all centers have access to TEG, and these centers should perform frequent serial measurements of PT/ INR, PTT, platelets, Hgb/Hct, fibrinogen and D-dimer to help guide resuscitation practice.

Surgical Treatment Options

Surgical management of trauma coagulopathy should be directed toward prompt cessation of any anatomic causes of hemorrhage, and thereby avoidance of what has been termed the “lethal triad” of coagulopathy, hypothermia, and acidosis.

Treatment Resistance/Complications

Compounding coagulopathy

If prompt diagnosis and tailored resuscitation fails to improve TIC, compounding etiologies of coagulopathy should be strongly suspected. These can include hypothermia, acidosis, hemodilution, DIC, and heritable coagulopathies. Treatment will include the approaches described above, as well as the following considerations.

1) Hypothermia

Hypothermia ($T < 36^{\circ}\text{C}$) is present in approximately 2/3 of injured patients on admission, and 9% of patients present with severe hypothermia ($T < 33^{\circ}\text{C}$) due to combination of exposure in the field and during transport and administration of cold intravenous fluids⁷⁴. Patients are at further risk for worsening of hypothermia in the emergency room and operating room, if necessary. Mild to moderate hypothermia (T 33-36°C) results in impaired platelet aggregation and adhesion and decreased tissue factor activity leading to coagulopathy that is typically not detectable using standard or functional coagulations assays as samples are routinely warmed prior to testing⁷⁵. Therefore, it is essential to maintain a high level of vigilance against hypothermia with continuous temperature monitoring. All trauma patients should receive passive rewarming with removal of clothing and application warmed blankets, as well as warming of administered fluids. Central warming should be considered in cases of severe or resistant hypothermia.

2) Acidosis

Severely- injured patients frequently present with or develop acidosis during the course of their resuscitation. Assembly of functional coagulation factor complexes are inhibited in acidotic environments ($\text{pH} < 7.2$) with increasing dysfunction as acidosis worsens^{76,77}. Arterial blood gas should be obtained at

admission and repeated at serial time points throughout resuscitation with correction of acidosis as necessary.

3) *Hemodilution*

Dilutional coagulopathy, also known as “iatrogenic” or “resuscitation–associated coagulopathy” occurs when coagulation factor proteins are diluted by large volumes of crystalloid, colloid, or PRBCs. Coagulation factor dilution with large volume administration has been demonstrated in multiple laboratory, modeling and healthy control studies⁷⁸⁻⁸⁰. Retrospective data has shown admission coagulopathy to be significantly more prevalent among injured patients who received >3L of pre-hospital fluids compared to those who received little (<500cc) or no volume⁴. To avoid dilutional coagulopathy, resuscitation should primarily consist of balanced product transfusion with frequent monitoring of ongoing coagulation status. Crystalloid should no longer be considered a resuscitation fluid and only used to facilitate administration of medications and blood products.

4) *Disseminated intravascular coagulation*

Disseminated intravascular coagulation (DIC) occurs as a result of systemic microvascular thrombosis causing severe consumptive coagulopathy. Although now understood to be a distinct pathology, some features of DIC overlap with traumatic coagulopathy making diagnosis challenging. Trauma patients are at increased risk for DIC due to the potential for embolization of tissue-specific thromboplastin following long bone fractures, amniotic disruption, or brain injury, or later in the clinical course as a result of sepsis. DIC should be considered as an alternate or concomitant etiology of late, recurrent, or treatment resistant coagulopathy in the severely injured patient but is not a cause of acute traumatic coagulopathy.

Evaluation of Outcome and Long-Term Recommendations

Outcomes following treatment of TIC should be assessed by:

- 1) ***Improvement in the clinical condition of the injured patient including a trend toward global hemostasis and hemodynamic stability and***
- 2) ***Reversal of standard and functional laboratory coagulation abnormalities.***

Despite advances in the understanding, diagnosis and resuscitation of injured patients with TIC, patients with traumatic coagulopathy go on to have worse outcomes than non-coagulopathic patients.

1) Transfusion Requirements

Compared to non-coagulopathic patients, patients with TIC receive substantially increased blood product transfusion¹⁰ which independently confers higher risk of complications such as Acute Respiratory Distress Syndrome⁸¹, Systemic Inflammatory Response Syndrome⁸², and mortality following injury⁸³. While, as above, there is no consensus regarding ideal ratios of resuscitation, ongoing observational experience from institutions with massive transfusion protocols have described decreased crystalloid and overall product usage, and lower PRBC:FFP ratios with concomitant survival benefit⁸⁴.

2) Hypercoagulability

Multiple investigators have reported increased incidence of thromboembolic complications in patients with early TIC.^{36,85} Given the potential risks of anticoagulation in this patient population, multiple scoring systems including the Trauma Embolic Scoring System (TESS) and the Risk Assessment Profile(RAP) have attempted to stratify patients at high risk for venous thromboembolism (VTE), but recent retrospective data suggests scoring systems often fail to discriminate between patients who go on to develop VTE.⁸⁶ It also remains unclear whether standard or functional assays can accurately predict clinically significant hypercoagulability in critically ill trauma patients.⁸⁷ Many studies investigating VTE in trauma patients rely on duplex screening, and the significance of these incidentally discovered events remains unknown. Finally, there has not been sufficient evidence to definitively establish whether standard chemoprophylaxis is efficacious in mitigating thromboembolic risk in this population⁸⁸. Further prospective clinical studies are necessary to better delineate the mechanism of VTE in previously coagulopathic trauma patients and to identify potential targets for prevention and treatment.

3) Acute Lung Injury, Multiorgan Failure, and Death

Despite substantive progress in delineation of the mechanism of TIC and improved outcomes resulting from changes in resuscitation practice, patients with traumatic coagulopathy have vastly increased rates of acute lung injury, multiorgan failure and death. Unsurprisingly, these patients also have worse hospital metrics including more days spent on the ventilator, in the Intensive Care Unit, and in the hospital than non-coagulopathic patients. This highlights the need for continued research efforts and clinical innovation to combat this considerable clinical challenge and improve outcomes. One potential approach is through focus on individualization of diagnosis and resuscitation. As discussed above, there is evidence to suggest that traumatic coagulopathy is not one single entity, but rather consists of multiple phenotypes with unique implications for outcomes and management.

Summary (*repeat of key points*)

- Trauma-induced coagulopathy (TIC) is an endogenous hypocoagulable state distinct from iatrogenic, dilution, or hypothermic causes.
- Activation of Protein C pathway is a key mechanistic mediator of TIC via multiple downstream effects including thrombin diversion, deactivation of coagulation factors, and de-repression of fibrinolysis.
- Standard coagulation tests and functional viscoelastic assays are commonly utilized in the diagnosis and management of TIC.
- Balanced resuscitation is the mainstay of TIC treatment, but precise ratios for empiric resuscitation and optimal monitoring protocols for transfusion practice remain hotly debated.
- Patients with TIC have worse outcomes including increased rates of transfusion, infection, thromboembolism, acute lung injury, multiorgan failure, and death.

Figure Legends:

Figure 1: Pathophysiology of Traumatic Coagulopathy

Multiple distinct but highly integrated pathways have been implicated as mediators of trauma-induced coagulopathy. Delineation and integration of these pathways remains an area of ongoing research.

Figure 2: Critical Role of Activated Protein C in the Pathophysiology of Trauma-induced Coagulopathy.

Severe injury accompanied by tissue hypoperfusion leads to increased endothelial and circulating thrombomodulin which subsequently binds thrombin. The resultant thrombomodulin-thrombin complex converts Protein C into its activated form (aPC). APC decreases clot formation via deactivation of factors V and VIII while simultaneously consuming plasminogen activator inhibitor-1 (PAI-1), disinhibiting tPA and

leading to unopposed fibrinolysis. APC, activated Protein C; PAI, Plasminogen Activator Inhibitor; TPA, Tissue Plasminogen Activator; VIIIa, Activated Factor VIII; Va, Activated Factor V.

Figure 3: Evolution of Paradigms and Resuscitation of Injured Patients over Time

Early transfusion was conducted with whole blood. In the 1970's resuscitation practice shifted toward an emphasis on perfusion with heavy utilization of large volumes of crystalloid and packed red blood cells (PRBC). Coagulopathy was understood to occur in severely injured patients but was not recognized as a distinct entity. Rather, it was attributed to iatrogenic dilution, hypothermia, or acidosis. In 2003, trauma-induced coagulopathy was identified as a distinct entity and treatment shifted away from crystalloid and toward balanced hemostatic resuscitation.

Figure 4. Recommendations of European Guideline on management of major bleeding and coagulopathy following trauma, 2016.

From Rossaint R, Bouillon B, Cerny V et al. From The European guideline on management of major bleeding and coagulopathy following trauma: fourth edition. Crit Care. 2016; 20: 100; with permission.

References

1. Sauaia A, Moore FA, Moore EE, et al. Epidemiology of trauma deaths: a reassessment. *J Trauma*. 1995;38(2):185-193.
2. Brohi K, Singh J, Heron M, Coats T. Acute traumatic coagulopathy. *J Trauma*. 2003;54(6):1127-1130.
3. MacLeod JB, Lynn M, McKenney MG, Cohn SM, Murtha M. Early coagulopathy predicts mortality in trauma. *J Trauma*. 2003;55(1):39-44.
4. Maegele M, Lefering R, Yucel N, et al. Early coagulopathy in multiple injury: an analysis from the German Trauma Registry on 8724 patients. *Injury*. 2007;38(3):298-304.
5. Cohen MJ, Bir N, Rahn P, et al. Protein C depletion early after trauma increases the risk of ventilator-associated pneumonia. *J Trauma*. 2009;67(6):1176-1181.
6. Cohen MJ, Call M, Nelson M, et al. Critical role of activated protein C in early coagulopathy and later organ failure, infection and death in trauma patients. *Ann Surg*. 2012;255(2):379-385.
7. Hess JR, Brohi K, Dutton RP, et al. The coagulopathy of trauma: a review of mechanisms. *J Trauma*. 2008;65(4):748-754.
8. Kirkpatrick AW, Chun R, Brown R, Simons RK. Hypothermia and the trauma patient. *Can J Surg*. 1999;42(5):333-343.
9. Kutcher ME, Ferguson AR, Cohen MJ. A principal component analysis of coagulation after trauma. *J Trauma Acute Care Surg*. 2013;74(5):1223-1229; discussion 1229-1230.
10. Brohi K, Cohen MJ, Ganter MT, Matthay MA, Mackersie RC, Pittet JF. Acute traumatic coagulopathy: initiated by hypoperfusion: modulated through the protein C pathway? *Ann Surg*. 2007;245(5):812-818.
11. Cohen MJ, Kutcher M, Redick B, et al. Clinical and mechanistic drivers of acute traumatic coagulopathy. *J Trauma Acute Care Surg*. 2013;75(1 Suppl 1):S40-47.
12. Chesebro BB, Rahn P, Carles M, et al. Increase in activated protein C mediates acute traumatic coagulopathy in mice. *Shock*. 2009;32(6):659-665.
13. Esmon CT. Protein C pathway in sepsis. *Ann Med*. 2002;34(7-8):598-605.

14. Rezaie AR. Vitronectin functions as a cofactor for rapid inhibition of activated protein C by plasminogen activator inhibitor-1. Implications for the mechanism of profibrinolytic action of activated protein C. *J Biol Chem*. 2001;276(19):15567-15570.
15. Bajzar L, Jain N, Wang P, Walker JB. Thrombin activatable fibrinolysis inhibitor: not just an inhibitor of fibrinolysis. *Crit Care Med*. 2004;32(5 Suppl):S320-324.
16. Schochl H, Frietsch T, Pavelka M, Jambor C. Hyperfibrinolysis after major trauma: differential diagnosis of lysis patterns and prognostic value of thrombelastometry. *J Trauma*. 2009;67(1):125-131.
17. Kashuk JL, Moore EE, Sawyer M, et al. Primary fibrinolysis is integral in the pathogenesis of the acute coagulopathy of trauma. *Ann Surg*. 2010;252(3):434-442; discussion 443-434.
18. Cotton BA, Harvin JA, Kostousouv V, et al. Hyperfibrinolysis at admission is an uncommon but highly lethal event associated with shock and prehospital fluid administration. *J Trauma Acute Care Surg*. 2012;73(2):365-370; discussion 370.
19. Chapman MP, Moore EE, Ramos CR, et al. Fibrinolysis greater than 3% is the critical value for initiation of antifibrinolytic therapy. *J Trauma Acute Care Surg*. 2013;75(6):961-967; discussion 967.
20. Moore HB, Moore EE, Gonzalez E, et al. Hyperfibrinolysis, physiologic fibrinolysis, and fibrinolysis shutdown: the spectrum of postinjury fibrinolysis and relevance to antifibrinolytic therapy. *J Trauma Acute Care Surg*. 2014;77(6):811-817; discussion 817.
21. Moore HB, Moore EE, Liras IN, et al. Acute Fibrinolysis Shutdown after Injury Occurs Frequently and Increases Mortality: A Multicenter Evaluation of 2,540 Severely Injured Patients. *J Am Coll Surg*. 2016;222(4):347-355.
22. Davenport RA, Brohi K. Coagulopathy in trauma patients: importance of thrombocyte function? *Curr Opin Anaesthesiol*. 2009;22(2):261-266.
23. Brown LM, Call MS, Margaret Knudson M, et al. A normal platelet count may not be enough: the impact of admission platelet count on mortality and transfusion in severely injured trauma patients. *J Trauma*. 2011;71(2 Suppl 3):S337-342.
24. Stansbury LG, Hess AS, Thompson K, Kramer B, Scalea TM, Hess JR. The clinical significance of platelet counts in the first 24 hours after severe injury. *Transfusion*. 2013;53(4):783-789.

25. Jacoby RC, Owings JT, Holmes J, Battistella FD, Gosselin RC, Paglieroni TG. Platelet activation and function after trauma. *J Trauma*. 2001;51(4):639-647.
26. Wohlauser MV, Moore EE, Thomas S, et al. Early platelet dysfunction: an unrecognized role in the acute coagulopathy of trauma. *J Am Coll Surg*. 2012;214(5):739-746.
27. Kutcher ME, Redick BJ, McCreery RC, et al. Characterization of platelet dysfunction after trauma. *J Trauma Acute Care Surg*. 2012;73(1):13-19.
28. Sillesen M, Johansson PI, Rasmussen LS, et al. Platelet activation and dysfunction in a large-animal model of traumatic brain injury and hemorrhage. *J Trauma Acute Care Surg*. 2013;74(5):1252-1259.
29. Solomon C, Traintinger S, Ziegler B, et al. Platelet function following trauma. A multiple electrode aggregometry study. *Thromb Haemost*. 2011;106(2):322-330.
30. Johansson PI, Stensballe J, Rasmussen LS, Ostrowski SR. A high admission syndecan-1 level, a marker of endothelial glycocalyx degradation, is associated with inflammation, protein C depletion, fibrinolysis, and increased mortality in trauma patients. *Ann Surg*. 2011;254(2):194-200.
31. Matijevic N, Wang YW, Wade CE, et al. Cellular microparticle and thrombogram phenotypes in the Prospective Observational Multicenter Major Trauma Transfusion (PROMMTT) study: correlation with coagulopathy. *Thromb Res*. 2014;134(3):652-658.
32. Nekludov M, Mobarrez F, Gryth D, Bellander BM, Wallen H. Formation of microparticles in the injured brain of patients with severe isolated traumatic brain injury. *J Neurotrauma*. 2014;31(23):1927-1933.
33. Hoyt DB, Dutton RP, Hauser CJ, et al. Management of coagulopathy in the patients with multiple injuries: results from an international survey of clinical practice. *J Trauma*. 2008;65(4):755-764; discussion 764-755.
34. Brohi K, Cohen MJ, Davenport RA. Acute coagulopathy of trauma: mechanism, identification and effect. *Curr Opin Crit Care*. 2007;13(6):680-685.
35. Kashuk JL, Moore EE, Johnson JL, et al. Postinjury life threatening coagulopathy: is 1:1 fresh frozen plasma:packed red blood cells the answer? *J Trauma*. 2008;65(2):261-270; discussion 270-261.

36. Peltan ID, Vande Vusse LK, Maier RV, Watkins TR. An International Normalized Ratio-Based Definition of Acute Traumatic Coagulopathy Is Associated With Mortality, Venous Thromboembolism, and Multiple Organ Failure After Injury. *Crit Care Med*. 2015;43(7):1429-1438.
37. Kaufmann CR, Dwyer KM, Crews JD, Dols SJ, Trask AL. Usefulness of thrombelastography in assessment of trauma patient coagulation. *J Trauma*. 1997;42(4):716-720; discussion 720-712.
38. Carroll RC, Craft RM, Langdon RJ, et al. Early evaluation of acute traumatic coagulopathy by thrombelastography. *Transl Res*. 2009;154(1):34-39.
39. Rugeri L, Levrat A, David JS, et al. Diagnosis of early coagulation abnormalities in trauma patients by rotation thrombelastography. *J Thromb Haemost*. 2007;5(2):289-295.
40. Cotton BA, Faz G, Hatch QM, et al. Rapid thrombelastography delivers real-time results that predict transfusion within 1 hour of admission. *J Trauma*. 2011;71(2):407-414; discussion 414-407.
41. Rivard GE, Brummel-Ziedins KE, Mann KG, Fan L, Hofer A, Cohen E. Evaluation of the profile of thrombin generation during the process of whole blood clotting as assessed by thrombelastography. *J Thromb Haemost*. 2005;3(9):2039-2043.
42. Levrat A, Gros A, Rugeri L, et al. Evaluation of rotation thrombelastography for the diagnosis of hyperfibrinolysis in trauma patients. *Br J Anaesth*. 2008;100(6):792-797.
43. Raza I, Davenport R, Rourke C, et al. The incidence and magnitude of fibrinolytic activation in trauma patients. *J Thromb Haemost*. 2013;11(2):307-314.
44. Yucel N, Lefering R, Maegele M, et al. Trauma Associated Severe Hemorrhage (TASH)-Score: probability of mass transfusion as surrogate for life threatening hemorrhage after multiple trauma. *J Trauma*. 2006;60(6):1228-1236; discussion 1236-1227.
45. McLaughlin DF, Niles SE, Salinas J, et al. A predictive model for massive transfusion in combat casualty patients. *J Trauma*. 2008;64(2 Suppl):S57-63; discussion S63.
46. Nunez TC, Voskresensky IV, Dossett LA, Shinall R, Dutton WD, Cotton BA. Early prediction of massive transfusion in trauma: simple as ABC (assessment of blood consumption)? *J Trauma*. 2009;66(2):346-352.

47. collaborators C-t, Shakur H, Roberts I, et al. Effects of tranexamic acid on death, vascular occlusive events, and blood transfusion in trauma patients with significant haemorrhage (CRASH-2): a randomised, placebo-controlled trial. *Lancet*. 2010;376(9734):23-32.
48. Morrison JJ, Dubose JJ, Rasmussen TE, Midwinter MJ. Military Application of Tranexamic Acid in Trauma Emergency Resuscitation (MATTERs) Study. *Arch Surg*. 2012;147(2):113-119.
49. Napolitano LM, Cohen MJ, Cotton BA, Schreiber MA, Moore EE. Tranexamic acid in trauma: how should we use it? *J Trauma Acute Care Surg*. 2013;74(6):1575-1586.
50. Chowdary P, Saayman AG, Paulus U, Findlay GP, Collins PW. Efficacy of standard dose and 30 ml/kg fresh frozen plasma in correcting laboratory parameters of haemostasis in critically ill patients. *Br J Haematol*. 2004;125(1):69-73.
51. Cardigan R, Lawrie AS, Mackie IJ, Williamson LM. The quality of fresh-frozen plasma produced from whole blood stored at 4 degrees C overnight. *Transfusion*. 2005;45(8):1342-1348.
52. Stanworth SJ. The evidence-based use of FFP and cryoprecipitate for abnormalities of coagulation tests and clinical coagulopathy. *Hematology Am Soc Hematol Educ Program*. 2007:179-186.
53. Abdel-Wahab OI, Healy B, Dzik WH. Effect of fresh-frozen plasma transfusion on prothrombin time and bleeding in patients with mild coagulation abnormalities. *Transfusion*. 2006;46(8):1279-1285.
54. Holland LL, Brooks JP. Toward rational fresh frozen plasma transfusion: The effect of plasma transfusion on coagulation test results. *Am J Clin Pathol*. 2006;126(1):133-139.
55. Kor DJ, Stubbs JR, Gajic O. Perioperative coagulation management--fresh frozen plasma. *Best Pract Res Clin Anaesthesiol*. 2010;24(1):51-64.
56. Toy P, Gajic O, Bacchetti P, et al. Transfusion-related acute lung injury: incidence and risk factors. *Blood*. 2012;119(7):1757-1767.
57. Levi M, Levy JH, Andersen HF, Truloff D. Safety of recombinant activated factor VII in randomized clinical trials. *N Engl J Med*. 2010;363(19):1791-1800.
58. Marlu R, Hodaj E, Paris A, Albaladejo P, Cracowski JL, Pernod G. Effect of non-specific reversal agents on anticoagulant activity of dabigatran and rivaroxaban: a randomised crossover ex vivo study in healthy volunteers. *Thromb Haemost*. 2012;108(2):217-224.

59. Gorlinger K, Dirkmann D, Hanke AA, et al. First-line therapy with coagulation factor concentrates combined with point-of-care coagulation testing is associated with decreased allogeneic blood transfusion in cardiovascular surgery: a retrospective, single-center cohort study. *Anesthesiology*. 2011;115(6):1179-1191.
60. Tanaka KA, Mazzeffi M, Durila M. Role of prothrombin complex concentrate in perioperative coagulation therapy. *J Intensive Care*. 2014;2(1):60.
61. Dickneite G, Pragst I. Prothrombin complex concentrate vs fresh frozen plasma for reversal of dilutional coagulopathy in a porcine trauma model. *Br J Anaesth*. 2009;102(3):345-354.
62. Schochl H, Nienaber U, Hofer G, et al. Goal-directed coagulation management of major trauma patients using thromboelastometry (ROTEM)-guided administration of fibrinogen concentrate and prothrombin complex concentrate. *Crit Care*. 2010;14(2):R55.
63. Joseph B, Pandit V, Khalil M, et al. Use of prothrombin complex concentrate as an adjunct to fresh frozen plasma shortens time to craniotomy in traumatic brain injury patients. *Neurosurgery*. 2015;76(5):601-607; discussion 607.
64. Theusinger OM, Baulig W, Seifert B, Emmert MY, Spahn DR, Asmis LM. Relative concentrations of haemostatic factors and cytokines in solvent/detergent-treated and fresh-frozen plasma. *Br J Anaesth*. 2011;106(4):505-511.
65. Schlimp CJ, Voelckel W, Inaba K, Maegele M, Schochl H. Impact of fibrinogen concentrate alone or with prothrombin complex concentrate (+/- fresh frozen plasma) on plasma fibrinogen level and fibrin-based clot strength (FIBTEM) in major trauma: a retrospective study. *Scand J Trauma Resusc Emerg Med*. 2013;21:74.
66. Duchesne JC, Kimonis K, Marr AB, et al. Damage control resuscitation in combination with damage control laparotomy: a survival advantage. *J Trauma*. 2010;69(1):46-52.
67. Borgman MA, Spinella PC, Perkins JG, et al. The ratio of blood products transfused affects mortality in patients receiving massive transfusions at a combat support hospital. *J Trauma*. 2007;63(4):805-813.

68. de Biasi AR, Stansbury LG, Dutton RP, Stein DM, Scalea TM, Hess JR. Blood product use in trauma resuscitation: plasma deficit versus plasma ratio as predictors of mortality in trauma (CME). *Transfusion*. 2011;51(9):1925-1932.
69. Brown LM, Aro SO, Cohen MJ, et al. A high fresh frozen plasma: packed red blood cell transfusion ratio decreases mortality in all massively transfused trauma patients regardless of admission international normalized ratio. *J Trauma*. 2011;71(2 Suppl 3):S358-363.
70. Holcomb JB, del Junco DJ, Fox EE, et al. The prospective, observational, multicenter, major trauma transfusion (PROMTT) study: comparative effectiveness of a time-varying treatment with competing risks. *JAMA Surg*. 2013;148(2):127-136.
71. Holcomb JB, Tilley BC, Baraniuk S, et al. Transfusion of plasma, platelets, and red blood cells in a 1:1:1 vs a 1:1:2 ratio and mortality in patients with severe trauma: the PROPPR randomized clinical trial. *JAMA*. 2015;313(5):471-482.
72. Hunt H, Stanworth S, Curry N, et al. Thromboelastography (TEG) and rotational thromboelastometry (ROTEM) for trauma induced coagulopathy in adult trauma patients with bleeding. *Cochrane Database Syst Rev*. 2015;2:CD010438.
73. Gonzalez E, Moore EE, Moore HB, et al. Goal-directed Hemostatic Resuscitation of Trauma-induced Coagulopathy: A Pragmatic Randomized Clinical Trial Comparing a Viscoelastic Assay to Conventional Coagulation Assays. *Ann Surg*. 2016;263(6):1051-1059.
74. Tsuei BJ, Kearney PA. Hypothermia in the trauma patient. *Injury*. 2004;35(1):7-15.
75. Wolberg AS, Meng ZH, Monroe DM, 3rd, Hoffman M. A systematic evaluation of the effect of temperature on coagulation enzyme activity and platelet function. *J Trauma*. 2004;56(6):1221-1228.
76. Martini WZ. Coagulopathy by hypothermia and acidosis: mechanisms of thrombin generation and fibrinogen availability. *J Trauma*. 2009;67(1):202-208; discussion 208-209.
77. Engstrom M, Schott U, Romner B, Reinstrup P. Acidosis impairs the coagulation: A thromboelastographic study. *J Trauma*. 2006;61(3):624-628.
78. Hirshberg A, Dugas M, Banez EI, Scott BG, Wall MJ, Jr., Mattox KL. Minimizing dilutional coagulopathy in exsanguinating hemorrhage: a computer simulation. *J Trauma*. 2003;54(3):454-463.

79. Coats TJ, Brazil E, Heron M. The effects of commonly used resuscitation fluids on whole blood coagulation. *Emerg Med J.* 2006;23(7):546-549.
80. Brazil EV, Coats TJ. Sonoclot coagulation analysis of in-vitro haemodilution with resuscitation solutions. *J R Soc Med.* 2000;93(10):507-510.
81. Silverboard H, Aisiku I, Martin GS, Adams M, Rozycki G, Moss M. The role of acute blood transfusion in the development of acute respiratory distress syndrome in patients with severe trauma. *J Trauma.* 2005;59(3):717-723.
82. Dunne JR, Malone DL, Tracy JK, Napolitano LM. Allogenic blood transfusion in the first 24 hours after trauma is associated with increased systemic inflammatory response syndrome (SIRS) and death. *Surg Infect (Larchmt).* 2004;5(4):395-404.
83. Charles A, Shaikh AA, Walters M, Huehl S, Pomerantz R. Blood transfusion is an independent predictor of mortality after blunt trauma. *Am Surg.* 2007;73(1):1-5.
84. Kutcher ME, Kornblith LZ, Narayan R, et al. A paradigm shift in trauma resuscitation: evaluation of evolving massive transfusion practices. *JAMA Surg.* 2013;148(9):834-840.
85. Knudson MM, Collins JA, Goodman SB, McCrory DW. Thromboembolism following multiple trauma. *J Trauma.* 1992;32(1):2-11.
86. Zander AL, Van Gent JM, Olson EJ, et al. Venous thromboembolic risk assessment models should not solely guide prophylaxis and surveillance in trauma patients. *J Trauma Acute Care Surg.* 2015;79(2):194-198.
87. Van Haren RM, Valle EJ, Thorson CM, et al. Hypercoagulability and other risk factors in trauma intensive care unit patients with venous thromboembolism. *J Trauma Acute Care Surg.* 2014;76(2):443-449.
88. Allen CJ, Murray CR, Meizoso JP, et al. Coagulation Profile Changes Due to Thromboprophylaxis and Platelets in Trauma Patients at High-Risk for Venous Thromboembolism. *Am Surg.* 2015;81(7):663-668.