Review Article for 11th US-Japan Scientific Seminar

Title: Citrus as a molecular contact point for co-evolution of *Alternaria* pathogens

Kazuya Akimitsu, Kouhei Ohtani, Takuya Shimagami, Mai Katsumoto, Chika Igarashi,
Sawa Tanaka, Syu Matsuoka, Susumu Mochizuki, Takashi Tsuge¹, Mikihiro Yamamoto²,
Motoichiro Kodama³, Kazuya Ichimura, and Kenji Gomi

Faculty of Agriculture, Kagawa University, Kagawa 761-079; ¹Graduate School of
Bioagricultural Sciences, Nagoya University, Nagoya 464-8601; ²Faculty of Agriculture,
Okayama University, Okayama 700-8530, Japan; ⁴Faculty of Agriculture, Tottori
University, Tottori 680-8553, Japan

Correspondence: Kazuya Akimitsu (K. Akimitsu) Tel.: +81 87 891 3131; Fax: +81 87
891 3021; e-mail: kazuya@ag.kagawa-u.ac.jp

ABSTRACT:

Alternaria black rot, Alternaria leaf spot of rough lemon, and Alternaria brown spot of tangerines are three major citrus Alternaria pathogens. Citrus could be considered as a molecular contact point for host-selective toxin (HST)-mediated co-evolution of these Alternaria pathogens and susceptibility in the field. ACR-toxin is an HST produced by the rough lemon pathotype, and the target site of the toxin was identified as rough lemon mitochondria. The biosynthetic gene cluster for ACR-toxin production is on the 1.5Mb-chromosome of the rough lemon pathotype. Another gene cluster for ACT-toxin production is located on the 1.9Mb-chromosome of the tangerine pathotype. These TOX genes shown to have a role in ACR- or ACT-toxin biosynthesis by using gene disruption and silencing.

Keywords: Host-selective toxin; *Alternaria alternata*; pathogenicity; conditionally dispensable chromosome; co-evolution

1. Introduction

Alternaria species cause three major diseases of citrus including Alternaria black rot of many citrus fruits, Alternaria leaf spot of rough lemon, and Alternaria brown spot of tangerines [1]. Alternaria black rot causes internal rotting in fruits of all citrus species and is one of the major citrus post-harvest diseases worldwide. The Alternaria black rot pathogen is hardly distinguishable in morphological characteristics from the Alternaria brown spot or Alternaria leaf spot pathogens, and they are considered to be the same species, *A. alternata* [2]. Unlike the black rot pathogen, the brown spot and leaf spot pathogens produce host-selective toxins (HST); ACT-toxin for the brown spot pathogen and ACR-toxin for the leaf spot pathogen, respectively [3, 4]. The host ranges of these two pathogens are exactly the same as the sensitivity of citrus cultivars to these toxins [3, 4]. The ACT-toxin producing brown spot pathogen affects many ACT-toxin-sensitive tangerines and mandarins causing necrotic spots on leaves, stems, and fruits, while the ACR-toxin producing leaf spot pathogen causes similar symptoms on ACR-toxin-sensitive rough lemon and rangpur lime [1, 3].

The Alternaria black rot pathogen causes black-colored rotting at the central columella usually without showing any external symptoms in fruits [5]. Endopolygalacturonase (endoPG) appears to be an essential virulence factor for this pathogen [6]. A series of studies demonstrated that EndoPG is essential for the penetration of citrus peel and development of black rot symptoms [6, 7]. The gene encoding endoPG is regulated by carbon catabolite repression mediated by the catabolite repressive element A (CreA) regulation factor, and endoPG does not have a central role for hyphal growth and colonization in the juice sac area in fruits, which contains sugars such as D-fructose and D-glucose [6-9]. The endoPG gene is conserved

among *Alternaria* species [2], and endoPG sequences from black rot and leaf spot pathogens are highly similar [6]. However, necrotic symptoms induced by an endoPG-minus mutant of the rough lemon pathotype were unchanged, while an endoPG-minus mutant of the black rot pathogen was significantly reduced in its ability to cause symptoms [6]. It is probably a reasonable result for the *Alternaria* black rot pathogen may require endoPG because pectin degradation is likely essential to cause the typical maceration symptom as well as for successful infection by this pathogen, which penetrates fruit tissues that are very rich in pectin. Contrary to this case, host-selective ACR-toxin production by necrotrophic *Alternaria* leaf spot pathogen induces necrosis without extensive penetration and colonization as well as maceration of the cell wall, and the role of endoPG might be masked by the strong toxicity of HST which may allow necrotic symptoms to appear without endoPG functions [6]. If this hypothesis is correct, HST is a pathogenicity factor controlling host specificity with a strong virulence effect, which masks the effects of other virulence factors, e.g. endoPG.

2. The ACRT cluster for ACR-toxin biosynthesis in the *A. alternata* rough lemon pathotype

The genus *Alternaria* belongs to the Hyphomycetes in the Deuteromycotina [10]. Phylogenetic studies using multiple genes among hundreds of *Alternaria* isolates from citrus fields indicated that all citrus-associated *Alternaria* species are *A. alternata* [2]. All isolates of *A. alternata* possess a potential general aggressiveness, which is the fundamental ability to penetrate plant tissues [11, 12]. Nishimura [11] proposed that HST-producing pathogens should be considered as a distinct pathotype of *A. alternata*, and one biotype among two citrus-associated *A. alternata* producing HST was identified

as the rough lemon pathotype of *A. alternata*. The major toxin, ACR-toxin I, of the rough lemon pathotype has structural features typical of polyketides consisting of a C19 polyalcohol with an α -dihydropyrone ring [13-15]. Masunaka *et al.* [16] found that the rough lemon pathotype has pathotype-specific chromosomes with sizes of 1.2 to 1.5 Mb. Draft sequence analysis of the 1.5-Mb chromosome of the standard isolate HC-1 identified multiple *ACRT* genes, responsible for ACR-toxin biosynthesis. *ACRTS1*, encoding a hydroxylase, is essential for ACR-toxin biosynthesis in the rough lemon pathotype and is required for pathogenicity [17]. *ACRTS2* encodes a polyketide synthase of 2,513 amino acids and is also essential for ACR-toxin biosynthesis in the rough lemon pathotype [18]. A third gene *ACRTS3* encoding a putative cyclase, was recently found to be essential for ACR-toxin biosynthesis (Nikaïdo *et al.*, unpublished data). All three of these genes have multiple paralogs and were studied using targeted gene disruption and RNA silencing to knock-out and/or -down the functional copies for examination of their roles in ACR-toxin biosynthesis. Disruption of a few copies of the respective genes reduced toxin production, and the transformants in which these genes were silenced did not produce ACR-toxin and lost pathogenicity [17, 18]. These genes are unique to ACR-toxin producers of the rough lemon pathotype, and they are clustered in the 1.5 Mb-chromosome. This chromosome exists only in the rough lemon pathotype and is known to be dispensable. Gain of these clustered genes on the 1.5 Mb-chromosome during evolution may have led to ACR-toxin production and hence the pathogenicity to rough lemon and rangpur lime. However, the origin of these genes, the gene cluster, or the chromosome has not been identified yet.

3. Target site of ACR-toxin in rough lemon

ACR-toxin has high specificity and toxicity and thus is generally recognized as a necrosis-inducing agent. However, ACR-toxin also causes delay or suppression of defense-related gene expression in rough lemon leaves inoculated with the rough lemon pathotype producing ACR-toxin [19-21].

The target site of ACR-toxin was found to be the mitochondria by electron microscopic examination [22] and monitoring of the toxin effects on oxidative phosphorylation using isolated physiologically active mitochondria [23]. ACR-toxin causes uncoupling of mitochondrial oxidative-phosphorylation with a loss of membrane potential and leakage of the co-factor NAD⁺ from the TCA cycle [23]. The *ACRS* gene that confers sensitivity to ACR-toxin has been identified from the mitochondrial genome of rough lemon [24]. The gene confers ACR-toxin sensitivity to *E. coli*, and the mechanism of specificity in plants is alternative processing of transcripts of the gene conferring ACR-toxin sensitivity [24]. *ACRS* is 171 bp and encodes a putative 7 kDa oligomeric pore-forming transmembrane protein controlling ACR-toxin sensitivity [24]. *ACRS* antibodies detected three proteins with molecular weights of 14, 21 and 28 kDa in extracts from rough lemon mitochondria, which could be the dimer, trimer and tetramer that are typical features of pore-forming receptors and not fully dissociated during SDS-PAGE [24]. Identification and evaluation of a protein complex which regulates alternative processing of the *ACRS* transcript is currently underway (Shimagami *et al.*, unpublished data).

4. *ACTT* cluster for ACT-toxin biosynthesis in *A. alternata* tangerine pathotype

Alternaria brown spot disease on Emperor mandarin was first reported in Australia in 1903 [25], and the pathogen was initially identified as *A. citri* Ellis & Pierce

due to the morphological similarity to the black rot pathogen [26, 27]. Later the pathogen was classified as *A. alternata* [4], as one of two biotypes causing leaf spot of rough lemon [13, 14, 28] and brown spot of tangerine [26, 29].

The tangerine pathotype affects many species of tangerines and mandarins worldwide but there is no occurrence in Japan, because a major citrus cultivar grown in Japanese citrus fields Unshu is resistant to this pathogen [1]. In laboratory tests, this pathotype is also pathogenic to Japanese pear cultivars susceptible to the Japanese pear pathotype. However, the pathogen has never encountered susceptible pear cultivars, which are grown only in Japanese fields, and thus there is no field occurrence of this disease in Japan either [29]. The tangerine pathotype produces ACT-toxins I and II; ACT-toxin I is toxic to both citrus and pear, and toxin II is highly toxic to pear and slightly toxic to citrus [29]. ACT-toxin I causes necrosis on leaves of susceptible cultivars at concentrations as low as 10^{-8} to 10^{-9} M but no necrosis on leaves of resistant cultivars even at higher concentrations. ACT-toxins consist of three moieties, 9,10-epoxy-8-hydroxy-9-methyl-decatrienoic acid (EDA), valine and a polyketide [29]. Toxins of the tangerine, Japanese pear and strawberry pathotypes are likely structurally analogous metabolites each containing a EDA moiety of (2E,4Z,6E)EDA for AK- and ACT-toxins or (2E,4E,6Z)EDA for AF-toxin [29-31].

ACTT genes responsible for the biosynthesis of the EDA moiety include pathotype-specific genes as well as genes common to the three pathotypes, which have multiple copies of functional or nonfunctional homologs sharing >90% nucleotide identity in the genomes from the tangerine, Japanese pear, and strawberry pathotypes [16, 33-38]. These biosynthetic genes appear to be clustered on a chromosome with a size of 2.0-Mb in the standard isolate SH-20 of the tangerine pathotype [16, 32, 34-37].

Functional analyses of *ACTT1*, *ACTT2*, *ACTT3*, *ACTTR*, *ACTT5*, and *ACTT6* in ACT-toxin biosynthesis have been carried out using a combination of target gene disruption and gene silencing [16, 32, 34-37] (Tanaka *et al.*, unpublished data). Four additional genes, *ACTTS1* to *ACTTS4*, were also identified as the tangerine pathotype-specific genes responsible for ACT-toxin biosynthesis; *ACTTS1* and *ACTTS4* encode a putative non-ribosomal peptide synthase, *ACTTS2* encodes an enoyl-reductase, and *ACTTS3* encodes a polyketide synthase [16, 37] (Katsumoto *et al.*, unpublished data). Targeted gene disruption or RNA silencing of any one of *ACTTS1* to *ACTTS4* resulted in the complete loss of ACT-toxin production and pathogenicity [16, 37] (Katsumoto *et al.*, unpublished data).

5. Concluding remarks: A role of citrus HST in the field as a molecular contact point of co-evolution.

Rough lemon defense-related genes including lipoxygenase (*RlemLOX*) [20], chalcone synthase [39, 40], polygalacturonase-inhibiting protein (*RlemPGIP1*, 2, A and B) [39-41], chitinases (*RlemAchiI* and *RlemAchiII*) [19], hydroperoxide lyase (*RlemHPL*) [21], epoxide hydrolase (*RlemEH*) [42], miraculin-like protein (*RlemMLP1* and *RlemMLP2*) [43], thaumatin-like protein (*RlemTLP*) [44] and nonspecific lipid transfer protein (*RlemLTP*) [45], are highly inducible in rough lemon leaves by inoculation of the tangerine pathotype producing ACT-toxin. Rough lemon is insensitive to ACT-toxin. The timing and intensity of the induction of these genes by the nonpathogenic strain O-94 of *A. alternata* were not as fast or strong as induction by the tangerine pathotype. There are no apparent morphological or biochemical differences between the tangerine pathotype and the nonpathogenic strain except

ACT-toxin production. The intensity and timing of the induction pattern were similar to those induced by nonpathogenic strain O-94, when an ACT-toxin-minus, nonpathogenic mutant of the tangerine pathotype, which was made by silencing the ACT-toxin biosynthetic gene *ACTT2* [35], was inoculated [46]. These results indicated that ACT-toxin is a pathogenicity factor with functions of both toxicity and suppression in toxin-sensitive plants and that it also has a role as a defense elicitation factor in toxin-insensitive plants.

Interestingly, Masunaka *et al.* [16] discovered an *A. alternata* isolate from a rough lemon field in Florida that produces two citrus-associated HSTs, ACT-toxin and ACR-toxin, and is pathogenic to both rough lemon and tangerines. Although inoculation of the isolate in lab tests resulted in necrosis on leaves of both ACT-toxin-sensitive and ACR-toxin-sensitive citrus cultivars, the dual pathotype producing both toxins is not dominant among hundreds of isolates examined to date and is not commonly found in populations of *A. alternata* on citrus. This isolate carries two chromosomes with sizes of 1.05- and 2.0-Mb, which encode the respective clusters of *ACTT* and *ACRT* genes [16]. Dual production of two distinct HSTs is expected to be a benefit due to the expansion of host range and may increase the fitness of pathogens by increasing the capacity for infection of and reproduction on a larger number of host plant species. However, one of the HSTs produced by this dual pathotype could act as an elicitor of plant defense responses in the toxin insensitive plants. For example, when the dual pathotype infects rough lemon leaves, ACR-toxin functions work as a pathogenicity factor, while at the same time, ACT-toxin might act as an elicitor of defense responses in the same leaves. Based on these assumptions, an isolate that produces a single HST is probably more dominant in infection as well as colonization

compared with isolates of the dual pathotype, and the population of the dual pathotype might be limited to a minimum level in co-evolution processes on citrus.

Acknowledgement

The authors are deeply indebted to the late Dr. S. Nishimura, Dr. K. Kohmoto, Dr. H. Otani, and Dr. T. L. Peever for their vision, efforts, and collaboration for citrus *Alternaria* HST studies. This work was supported by Grants-in-Aids for Scientific Research (S) (21228001 for KA and 19108001 for TT), Scientific Research (A) (26252007 for KA and 23248007 for TT) and Scientific Research (B) (20380028 and 23380025 for MK) from the Japanese Society for Promotion of Sciences, Special Coordination Funds for Promoting Sciences from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (TT), and Precursory Research for Embryonic Science and Technology from Japan Science and Technology Agency (KA).

References

- [1]. K. Akimitsu, T.L. Peever, L.W. Timmer, Molecular, ecological and evolutionary approaches to understanding *Alternaria* diseases of citrus, *Mol. Plant Pathol.* 4 (2003) 435-446.
- [2]. T.L. Peever, G. Su, L. Carpenter-Boggs, L.W. Timmer, Molecular systematics of citrus-associated *Alternaria* species, *Mycologia* 96 (2004) 119-134.
- [3]. K. Kohmoto, K. Akimitsu, H. Otani, Correlation of resistance and susceptibility of citrus to *Alternaria alternata* with sensitivity to host-specific toxins, *Phytopathology* 81 (1991) 719-722.
- [4]. K. Kohmoto, R.P. Scheffer, J.O. Whiteside, Host-selective toxins from *Alternaria*

citri, Phytopathology 69 (1979) 667-671.

[5]. G.E. Brown, A.A. McCornack, Decay caused by *Alternaria citri* in Florida citrus fruit, Plant Dis. Rep. 56 (1972) 909–912.

[6]. A. Isshiki, K. Akimitsu, M. Yamamoto, H. Yamamoto, Endopolygalacturonase is essential for citrus black rot caused by *Alternaria citri* but not brown spot caused by *Alternaria alternata*, Mol. Plant-Microbe Interact. 14 (2001) 749–757.

[7]. A. Isshiki, K. Ohtani, M. Kyo, H. Yamamoto, K. Akimitsu, Green fluorescent detection of fungal colonization and endopolygalacturonase gene expression in the interaction of *Alternaria citri* with citrus, Phytopathology 93 (2003) 768–773.

[8]. H. Katoh, K. Ohtani, H. Yamamoto, K. Akimitsu, Overexpression of a gene encoding a catabolite repression element in *Alternaria citri* causes severe symptoms of black rot in citrus fruit, Phytopathology 97 (2007) 557-563.

[9]. K. Ohtani, A. Isshiki, H. Katoh, H. Yamamoto, K. Akimitsu, Involvement of carbon catabolite repression on regulation of endopolygalacturonase gene expression in citrus fruit, J. Gen. Plant Pathol. 69 (2003) 120–125.

[10]. J. Rotem, The Genus *Alternaria*: Biology, Epidemiology, and Pathogenicity. The American Phytopathological Society Press, St. Paul, MN. (1994).

[11]. S. Nishimura, Host-specific toxins from *Alternaria alternata*: problems and prospects, Proc. Jpn. Acad. 56B (1980) 362-366.

[12]. S. Nishimura, K. Kohmoto, Host-specific toxins and chemical structures from *Alternaria* species, Annu. Rev. Phytopathol. 21 (1983) 87-116.

[13]. J.M. Gardner, Y. Kono, J.H. Tatum, Y. Suzuki, S. Takeuchi, Structure of major component of ACRL toxins, host-specific phytotoxic compound produced by *Alternaria citri*, Agri. Biol. Chem. 49 (1985a) 1235-1238.

- [14]. J.M. Gardner, Y. Kono, J.H. Tatum, Y. Suzuki, S. Takeuchi, Plant pathotoxins from *Alternaria citri*: the major toxin specific for rough lemon plants, *Phytochemistry* 24 (1985b) 2861-2868.
- [15]. S. Nakatsuka, T. Goto, K. Kohmoto, S. Nishimura, Host-specific phytotoxins. *Natural Products and Biological Activities* (Imura H, Goto T, Murachi T & Nakajima T, eds), pp 11-18. University of Tokyo Press, Tokyo (1986a).
- [16]. A. Masunaka, K. Ohtani, T.L. Peever, L.W. Timmer, T. Tsuge, M. Yamamoto, H. Yamamoto, K. Akimitsu, An isolate of *Alternaria alternata* that is pathogenic to both tangerines and rough lemon and produces two host-selective toxins, ACT- and ACR-toxins, *Phytopathology* 95 (2005) 241-247.
- [17]. Y. Izumi, E. Kamei, Y. Miyamoto, K. Ohtani, A. Masunaka, T. Fukumoto, K. Gomi, Y. Tada, K. Ichimura, T.L. Peever, K. Akimitsu, Role of pathotype-specific *ACRTS1* gene encoding a hydroxylase involved in the biosynthesis of host-selective ACR-toxin in the rough lemon pathotype of *Alternaria alternata*, *Phytopathology* 102 (2012a) 741–748.
- [18]. Y. Izumi, K. Ohtani, Y. Miyamoto, A. Masunaka, T. Fukumoto, K. Gomi, Y. Tada, K. Ichimura, T.L. Peever, K. Akimitsu, A polyketide synthase gene, *ACRTS2*, is responsible for biosynthesis of host-selective ACR-toxin in the rough lemon pathotype of *Alternaria alternata*, *Mol. Plant Microbe Interact.* 25 (2012b) 1419–1429.
- [19]. K. Gomi, N. Itoh, H. Yamamoto, K. Akimitsu, Characterization and functional analysis of class I and II acidic chitinase cDNA from rough lemon, *J. Gen. Plant Pathol.* 68 (2002a) 191-199.
- [20]. K. Gomi, H. Yamamoto, K. Akimitsu, Characterization of lipoxygenase gene in

- rough lemon induced by *Alternaria alternata*, J. Gen. Plant Pathol. 68 (2002b) 21-30.
- [21]. K. Gomi, Y. Yamasaki, H. Yamamoto, K. Akimitsu, Characterization of hydroperoxide lyase gene and effect of C₆-volatiles on expression of genes of the oxylipin metabolism in *Citrus*, J. Gen. Plant Pathol. 160 (2003b) 1219-1231.
- [22]. K. Kohmoto, Y. Kondoh, T. Kohguchi, H. Otani, S. Nishimura, R.P. Scheffer, Ultrastructural changes in host cells caused by host-selective toxin of *Alternaria alternata* from rough lemon, Can J Bot 62 (1984) 2485-2492.
- [23]. K. Akimitsu, K. Kohmoto, H. Otani, S. Nishimura, Host-specific effect of toxin from the rough lemon pathotype of *Alternaria alternata* on mitochondria, Plant Physiol. 89 (1989) 925-931.
- [24]. K. Ohtani, H. Yamamoto, K. Akimitsu, Sensitivity to *Alternaria alternata* toxin in citrus because of altered mitochondrial RNA processing, PNAS 99 (2002) 2439-2444.
- [25]. N.A. Cobb, Letters on the diseases of plants — *Alternaria* of the citrus tribe, Agri. Gaz. NSW. 14 (1903) 955–986.
- [26]. K.G. Pegg, Studies of a strain of *Alternaria citri* Pierce, the causal organism of brown spot of emperor mandarin (*Citrus reticulata*), Queensland J. Agric. Anim. Sci. 23 (1966) 15-28.
- [27]. J.O. Whiteside, A newly recorded *Alternaria*-induced brown spot disease on Dancy tangerines in Florida, Plant Dis. Rep. 60 (1976) 326–329.
- [28]. E.M. Doidge, A study of some *Alternarias* infecting citrus in South Africa, Union South Africa Dept. Agric. Sci. Bull. 69 (1929) 1-29.
- [29]. K. Kohmoto, Y. Itoh, N. Shimomura, Y. Kondoh, H. Otani, M. Kodama, S.

Nishimura, S. Nakatsuka, Isolation and biological activities of two host-specific toxins from the tangerine pathotype of *Alternaria alternata*, *Phytopathology* 83 (1993) 495-502.

[30]. T. Nakashima, T. Ueno, H. Fukami, T. Taga, H. Masuda, K. Osaki, H. Otani, H. Kohmoto, S. Nishimura, Isolation and structures of AK-toxin I and II, host-specific phytotoxic metabolites produced by *Alternaria alternata* Japanese pear pathotype, *Agric. Biol. Chem.* 49 (1985) 807-815.

[31]. S. Nakatsuka, K. Ueda, T. Goto, M. Yamamoto, S. Nishimura, K. Kohmoto, Structure of AF-toxin II, one of the host-specific toxins produced by *Alternaria alternata* strawberry pathotype, *Tetrahedron Lett.* 27 (1986b) 2753-2756.

[32]. N. Ajiro, Y. Miyamoto, A. Masunaka, T. Tsuge, M. Yamamoto, K. Ohtani, T. Fukumoto, K. Gomi, T.L. Peever, Y. Izumi, Y. Tada, K. Akimitsu, Role of the host-selective ACT-toxin synthesis gene *ACTTS2* encoding an enoyl-reductase in pathogenicity of the tangerine pathotype of *Alternaria alternata*, *Phytopathology* 100 (2010) 120–126.

[33]. R. Hatta, K. Ito, Y. Hosaki, T. Tanaka, A. Tanaka, M. Yamamoto, K. Akimitsu, T. Tsuge, A conditionally dispensable chromosome controls host-specific pathogenicity in the fungal plant pathogen *Alternaria alternata*, *Genetics* 161 (2002) 59-70.

[34]. A. Masunaka, A. Tanaka, T. Tsuge, T.L. Peever, L.W. Timmer, H. Yamamoto, M. Yamamoto, K. Akimitsu, Distribution and characterization of *AKT* homologs in the tangerine pathotype of *Alternaria alternata*, *Phytopathology* 90 (2000) 762-768.

[35]. Y. Miyamoto, A. Masunaka, T. Tsuge, M. Yamamoto, K. Ohtani, T. Fukumoto, K.

Gomi, T.L. Peever, K. Akimitsu, Functional analysis of a multi-copy host-selective ACT-toxin biosynthesis gene in the tangerine pathotype of *Alternaria alternata* using RNA silencing, Mol. Plant-Microbe In. 21 (2008) 1591-1599.

[36]. Y. Miyamoto, Y. Ishii, A. Masunaka, T. Tsuge, M. Yamamoto, K. Ohtani, T. Fukumoto, K. Gomi, T.L. Peever, K. Akimitsu, Function of genes encoding acyl-CoA synthetase and enoyl-CoA hydratase for host-selective ACT-toxin biosynthesis in the tangerine pathotype of *Alternaria alternata*, Phytopathology 99 (2009) 369-377.

[37]. Y. Miyamoto, A. Masunaka, T. Tsuge, M. Yamamoto, K. Ohtani, T. Fukumoto, K. Gomi, T.L. Peever, Tada Y, Ichimura K, K. Akimitsu, *ACTTS3* encoding a polyketide synthase is essential for the biosynthesis of ACT-toxin and pathogenicity in the tangerine pathotype of *Alternaria alternata*, Mol. Plant-Microbe In. 23 (2010) 406–414.

[38]. S.R. Ruswandi, K. Kitani, K. Akimitsu, T. Tsuge, T. Shiraishi, M. Yamamoto, Structural analysis of cosmid clone pcAFT-2 carrying *AFT10-1* encoding an acyl-CoA dehydrogenase involved in AF-toxin production in the strawberry pathotype of *Alternaria alternata*, J. Gen. Plant Pathol. 71 (2005) 107-116.

[39]. Y. Gotoh, S. Nalumpang, A. Isshiki, T. Utsumi, K. Gomi, H. Yamamoto, K. Akimitsu, A cDNA encoding polygalacturonase-inhibiting protein induced in citrus leaves by polygalacturonase of *Alternaria citri*, J. Gen. Plant Pathol. 68 (2002) 57-61.

[40]. S. Nalumpang, Y. Gotoh, H. Tsuboi, K. Gomi, H. Yamamoto, K. Akimitsu, Functional characterization of citrus polygalacturonase-inhibiting protein, J. Gen.

Plant Pathol. 68 (2002a) 118-127.

[41]. S. Nalumpang, Y. Gotoh, Y. Yamasaki, K. Gomi, H. Yamamoto, K. Akimitsu, Comparison and characterization of polygalacturonase-inhibiting protein genes from the genus of *Citrus* and its close related genera, Thai J. Agr. Sci. 35 (2002b) 147-164.

[42]. K. Gomi, H. Yamamoto, K. Akimitsu, Epoxide hydrolase: a mRNA induced by fungal pathogen *Alternaria alternata* on rough lemon (*Citrus jambhiri* Lush), Plant Mol. Biol. 53 (2003a) 189-199.

[43]. S. Tsukuda, K. Gomi, H. Yamamoto, K. Akimitsu, Characterization of cDNA encoding two distinct miraculin-like proteins and stress-related modulation of the corresponding mRNAs in *Citrus jambhiri* Lush, Plant Mol. Biol. 60 (2006) 125-136.

[44]. B.G. Kim, T. Fukumoto, S. Tatano, K. Gomi, K. Ohtani, Y. Tada, K. Akimitsu, Molecular cloning and characterization of a thaumatin-like protein-encoding cDNA from rough lemon, Physiol. Mol. Plant Pathol. 74 (2009) 3-10.

[45]. S. Nishimura, S. Tatano, K. Gomi, K. Ohtani, T. Fukumoto, K. Akimitsu, Chloroplast-localized nonspecific lipid transfer protein with anti-fungal activity from rough lemon, Physiol. Mol. Plant Pathol. 72 (2008) 134-140.

[46]. H. Shishido, Y. Miyamoto, R. Ozawab, S. Taniguchia, J. Takabayashib, K. Akimitsua, K. Gomi, Geraniol synthase whose mRNA is induced by host-selective ACT-toxin in the ACT-toxin-insensitive rough lemon (*Citrus jambhiri*), J. Plant Physiol. 169 (2012) 1401–1407.