False idolatry of the mythical growth versus immunity tradeoff in molecular systems plant pathology

Daniel J. Kliebenstein^{1,2}

¹Department of Plant Sciences

University of California, Davis

One Shields Avenue

Davis, CA 95616 USA

²DynaMo Center of Excellence, University of Copenhagen, Thorvaldsensvej 40, DK-1871,

Frederiksberg C, Denmark

DJK is the corresponding author. Email is kliebenstein@ucdavis.edu

Abstract

Systems studies of signal transduction pathways that modulate plant growth and immunity are rapidly identifying a large number of interactions within these pathways. These interactions are frequently presented as mechanisms allowing a plant to make proper decisions with regards to how to partition energy and resources in a proposed growth versus immunity tradeoff. This is a reinterpretation of the classical costs of resistance theory that has a long history in the ecology research community. While the ecology community is reinterpreting this theory, the reinterpretation has not been introduced into the molecular systems biology community that is studying the intersection of regulatory pathways. In this article, I describe evidence against a simple growth versus immunity tradeoff concept and propose an alternative wherein the intersection of these regulatory pathways is instead designed to coordinate these pathways, and not simply link them in mutual antagonism, to optimize fitness in complex environments where resistance/immunity and growth do not have simple linear relationships with fitness.

Introduction

Recent years have seen the re-introduction of ecological theories into studies focused on molecular plant pathology. This includes a burgeoning interest in the concept of costs of resistance and how this may drive an interplay between growth and immunity [1, 2]. A cost of resistance is broadly defined as any negative effect that expressing a resistance trait has on plant fitness [2-5]. In plant pathology, this is being increasingly linked to an interest in tradeoffs between growth and immunity [6-14]. This is being driven by molecular studies that are finding interactions between regulatory genes involved in the control of growth and defense. These links are being found in related signaling pathways that interlink the brassinosteroid and effector triggered immunity pathways [7, 11, 12, 15]. Similar links between development and plant defense signaling have been found for gibberellins, salicylic acid, auxin and jasmonic acid [16-22]. Interestingly, just as the molecular plant pathology community is beginning to investigate their systems using a cost of resistance framework, the ecology community is reassessing if there are costs of resistance and what this concept fundamentally means [3-5]. However, this reconsideration in the ecological community has not yet been translated into studies of molecular plant pathology. This article is intended to communicate misperceptions about costs of resistance that are permeating the molecular oriented literature and to provide alternate concepts of what is truly meant by the cost of immunity.

What do flux costs mean in a natural context?

The most common interpretation of resistance costs makes the implicit assumption that energy and elements are universally limiting for plant growth and/or fitness in the wild [13, 14]. This is often pictorialized as a teeter-totter wherein the plant has to decide if any specific element or energy is placed into growth or defense. This simplistic representation also makes the implicit statement that there are no other options or avenues open to the plant in this process. This teeter-totter model can be classified as the idea that any flux of a nutrient into a defense process necessarily removes that nutrient from what otherwise would have been growth or biomass accumulation, i.e. a flux cost model (Figure 1). However, the way the flux cost model is

presented assumes that all nutrients are equally limiting which is an oversimplification of how nutrients limit plant fitness in the wild [23]. In natural ecosystems, growth is limited by imbalances in the availability of nutrients and micronutrients rather than singular nutrient limitations. This means that potassium or phosphorus may be growth limiting in an environment where carbon and nitrogen may be freely available and non-limiting [23]. As such, it is the balance of nutrients that is more critical than any single nutrient when considering how nutrients may limit growth.

The observation that the plant is more responsive to the specific balance of nutrients and micronutrients raises the possibility that nutrient limitations on growth may be different than nutrient limitations on defense. Most plant defense mechanisms are based on proteins and metabolites that are predominantly Carbon and Nitrogen based with almost no associated Potassium and Phosphorus [24-26]. Thus, it is possible that in environments where Potassium, Phosphorus or other micronutrients are growth limiting that the plant has spare Carbon and Nitrogen available that cannot be used for growth. This spare Carbon and Nitrogen could then be used to create defenses at essentially no cost to growth. Thus, flux costs are not universal and instead are highly conditional based on the available nutrient and energy status of the plants environment. This suggests that the specific defense metabolism and mechanisms in a plant may be co-adapted to the plants typical nutrient profile to maximize the utilization of nutrients that are not rate limiting in that plants ecosystem niche. For example, a plant that typically grows in nitrogen limiting environments may shift from nitrogen-heavy defenses like alkaloids towards carboniferous defenses like terpenoids.

But constitutive defense mutants grow smaller?

A common argument in support of the idea that growth and defense must be in a tradeoff relationship is the observation that constitutive defense mutants are frequently small [27-32]. This argument posits that the elevated defense in these mutants is removing nutrients that would otherwise be utilized for growth. However, there is little evidence about the exact molecular reason for the diminutive growth in these mutants. A similar growth defect argument was proposed to explain why the *reduced epidermal fluorescence (ref)* mutants deficient in

phenolic metabolism displayed growth defects [33-36]. It was posited that the deficiency in lignin production and structure created a simple growth defect due to altered lignin physics [33-36]. However, this was later found to be an incorrect interpretation as shown by a suppressor screen that identified mutants that alleviated the growth defects in the *ref* mutants [35, 37-39]. These suppressors had wild-type growth rates while still containing the deficient lignin indicating that the growth effect was not caused by physical issues associated with the lignin deficit. Instead all of the suppressor mutations were in genes encoding components of the mediator transcriptional complex showing that the growth defect was caused by altered signal transduction and regulation of development and not by the physical or flux based costs of the *ref* mutant [35, 37-39]. Thus, it is possible that mutants displaying constitutive defense responses may show a growth defect for regulatory or developmental reasons that have nothing to do with the associated flux cost of producing those defense responses. For example, constitutive expression of jasmonate mediated defenses didn't specifically alter growth but instead altered flowering time and other phenological processes suggesting that this constitutive jasmonate mutant displayed regulatory costs more than elemental costs [40]. Thus, more work needs to be done to understand why constitutive defense mutants display growth defects.

Modern biochemical genetics is providing a more explicit assessment of the flux costs associated with specific defenses by providing the ability to genetically delete individual defense genes or pathways and then assess any resulting growth effect. The simple growth vs immunity tradeoff model would suggest that the energy no longer used for defense in these mutants should be redirected from into growth. Analysis of genotypes altered in glucosinolate accumulation was unable to find a significant link with absolute growth [41-43]. Instead, it was only possible to see a small increase in relative growth rate in these mutants and only at early developmental stages and not in larger mature plants [41, 42]. Importantly, this slight growth rate boost only benefited these plants when they are competing with other plant genotypes, i.e. a low glucosinolate plant would outgrow a high glucosinolate plant when they are in direct competition. Similar results had been found in the wild using Brassica [44-46]. Thus, if there is a direct growth benefit obtained by redirecting defenses to growth, these benefits appear to be

specific to conditions when two genotypes are in direct competition. Thus, studying any growth vs immunity tradeoff hypothesis must involve competition studies.

What are other costs of immunity/resistance?

The experiment showing potentially the most overt cost of a resistance mechanism was a set of studies that linked the presence of an R gene to a fitness cost in the absence of the pathogen [47-49]. This study was originally controversial because the estimated flux cost, the energy required to produce a single R gene encoded protein that accumulates to low levels, did not agree with the measured 9% cost in fitness in plants expressing the functional R protein [48]. However, flux costs are not the only potential cost for a resistance mechanism. There are also ecological or opportunity costs wherein a resistance mechanism that may provide a fitness benefit by aiding the plants interaction with one organism may actually create a cost by decreasing the plants ability to interact with another organism [44-46, 50, 51]. In this particular case, an ecological cost hypothesis could be as simple as the presence of the R gene may have altered the microbiome of these plants leading to the decreased fitness. Resistance signaling genes have been found to specifically affect the plants associated microbiome supporting this potential ecological cost [52, 53]. Another possibility is that this specific R gene actually generates sensitivity to an unknown pathogen as has been found for the LOV1 locus [54-58]. In this case, the cost or benefit of the R gene would depend upon the presence or absence of the respective species within the plants environment. Supporting the potential for ecological costs in a broader context is work on defense metabolites linking them to a number of detrimental side effects depending upon the specific insect community in which the plant exists [59-62].

Supporting the concept that ecological costs may be more important than flux costs, the authors utilized a set of R gene isogenic lines to assess how plant fitness responded to infection with the pathogen in the presence or absence of competition with uninfected plants [49]. In the presence of competition by uninfected plants, the ability to resist the pathogen increased a plants fitness. Interestingly, when the plants were grown in the absence of other plants (no competition) there was no fitness benefit provided by the R gene and the associated pathogen

resistance. If anything, the R gene led to decreased fitness in the presence of the pathogen [49]. In this case, the susceptible plants were able to compensate for the infection by increasing their seed production on secondary inflorescences. Thus, depending upon the organisms present within an environment, a specific resistance mechanism can invert from being costly to being beneficial. In this case, susceptibility to a pathogen was beneficial to plant fitness under specific conditions [49]. The ability of fluctuations in the environment to shift the fitness effect of specific resistance mechanisms has also been seen for defense metabolites [63]. Thus, the environment can impart ecological fitness costs that vastly outweigh any potential flux cost. Importantly, this also shows that it is not possible to classify all defenses as a single group when developing a model linking development to immunity.

For what other reasons could development and immunity be linked?

If flux costs of resistance are not the driving force behind the observed link between growth and immunity, this raises the question of what would be other reasons for these two processes to be linked. The growth vs immunity tradeoff concept fundamentally imagines growth and immunity as opposing contretemps in a zero sum game where for one to increase, the other must necessarily decrease. This is however, a misreading of the relationship between development and resistance. An alternative interpretation is that properly structured development is a key component of resistance. For example, shifting flowering time is a potentially effective way for plants to resist specific foliar herbivores by allowing the plant to evade these herbivores as seeds. Similarly as shown above, plants can alleviate the effects of a plant/pathogen interaction by a developmental reprogramming that increases the production of secondary inflorescences to compensate [49]. Developmental compensation is also a major mechanism that relieves deleterious consequences of plant herbivore interactions [64-66]. As such, major morphological shifts in development can be key aspects of how a plant interacts with other organisms.

Another facet of the interplay between growth and defense that is often overlooked by the growth vs immunity tradeoff concept is that defense mechanisms are not ubiquitously distributed across a whole plant but are instead placed in specific locations for potentially

specific reasons. In these instances, an explicit intersection of immunity and developmental pathways is central to placing defenses in their proper whole organism context to maximize their utility. This includes placing defense metabolites in specific structures like laticifers and glandular trichomes or enhancing pathogen defenses by elevating their expression in stomata [67-69]. While the specific developmental patterning of defenses can be thought to place them in the proper location for optimal function, there are other potential reasons for this patterning. The developmental patterning of anti-herbivory defenses in *Arabidopsis* actually affects the feeding behavior of the lepidopteran herbivores [70, 71]. This raises the possibility that the specific pattern could be a part of the defense activity by potentially leading the herbivore to a detrimental feeding behavior that may maximize its exposure to predators or parasites [72]. Another potential benefit from distributing defenses unequally across a specific tissue is that this patterning may decrease the ability of the attacking organisms to evolve a counter-defense strategy [73, 74]. This suggests that it is better to view development and immunity as two processes that are coordinated to optimize a plants interaction with its environment and not as opponents in a zero sum game.

Summary:

Growth(development) and immunity(resistance) are highly interlinked processes as exemplified by the increasing evidence for cross-talk in the different pathways. While this has begun to be described as a tradeoff relationship between growth and immunity locked in a zero sum game, the above studies and observations illustrate that this is likely a misreading of the growth to immunity relationship within a plant. Instead, we need to develop a deeper conceptualization of the link between growth and immunity wherein the two are integrating, sometimes as a tradeoff and other times synergistically, to maximize the plants ensuing fitness depending upon the specific environment in which that plant exists (Figure 1B). It is only by developing this more conditional and realistic framework in which to study the connections between growth and immunity that we will make full progress in understanding how plant fitness is maximized in the wild.

Acknowledgements

Funding for this work was provided by the NSF awards IOS 1339125 and MCB 1330337 to DJK, the USDA National Institute of Food and Agriculture, Hatch project number CA-D-PLS-7033-H to DJK, and by the Danish National Research Foundation (DNRF99) grant to DJK.

Figure Legends

Figure 1. Alternative models of growth and defense

- A. Standard illustration of the growth vs immunity tradeoff where the two are in opposition.
- B. Alternative conceptualization whereby growth and defense are in a continual conversation to coordinate.

Literature Cited

- [1] Strauss SY, Agrawal AA. The ecology and evolution of plant tolerance to herbivory. *Trends in Ecology & Evolution*. 1999;14:179-85.
- [2] Karban R, Baldwin IT. *Induced Responses to Herbivory*. Chicago, IL, USA: University of Chicago Press; 1997.
- [3] Agrawal AA, Strauss SY, Stout MJ. Costs of induced responses and tolerance to herbivory in male and female fitness components of wild radish. *Evolution*. 1999;53:1093-104.
- [4] Agrawal AA. Current trends in the evolutionary ecology of plant defence. *Functional Ecology*. 2011;25:420-32.
- [5] Agrawal AA. New synthesis-trade-offs in Chemical Ecology. *Journal of Chemical Ecology*. 2011;37:230-1.
- [6] Lin WW, Lu DP, Gao XQ, Jiang S, Ma XY, Wang ZH, et al. Inverse modulation of plant immune and brassinosteroid signaling pathways by the receptor-like cytoplasmic kinase BIK1. *Proceedings of the National Academy of Sciences of the United States of America*. 2013;110:12114-9.
- [7] Lozano-Duran R, Macho AP, Boutrot F, Segonzac C, Somssich IE, Zipfel C. The transcriptional regulator BZR1 mediates trade-off between plant innate immunity and growth. *Elife*. 2013;2.
- [8] Chandran D, Rickert J, Huang YX, Steinwand MA, Marr SK, Wildermuth MC. Atypical E2F Transcriptional Repressor DEL1 Acts at the Intersection of Plant Growth and Immunity by Controlling the Hormone Salicylic Acid. *Cell Host & Microbe*. 2014;15:506-13.
- [9] Fan M, Bai MY, Kim JG, Wang TN, Oh E, Chen L, et al. The bHLH Transcription Factor HBI1 Mediates the Trade-Off between Growth and Pathogen-Associated Molecular Pattern-Triggered Immunity in Arabidopsis. *Plant Cell*. 2014;26:828-41.
- [10] Malinovsky FG, Batoux M, Schwessinger B, Youn JH, Stransfeld L, Win J, et al. Antagonistic Regulation of Growth and Immunity by the Arabidopsis Basic Helix-Loop-Helix Transcription Factor HOMOLOG OF BRASSINOSTEROID ENHANCED EXPRESSION2 INTERACTING WITH INCREASED LEAF INCLINATION1 BINDING bHLH1. *Plant physiology*. 2014;164:1443-55.

- [11] Wang WF, Wang ZY. At the Intersection of Plant Growth and Immunity. *Cell Host & Microbe*. 2014;15:401-3.
- [12] Lozano-Duran R, Zipfel C. Trade-off between growth and immunity: role of brassinosteroids. *Trends in plant science*. 2015;20:12-9.
- [13] Herms DA, Mattson WJ. THE DILEMMA OF PLANTS - TO GROW OR DEFEND. *Quarterly Review of Biology*. 1992;67:283-335.
- [14] Pieterse CMJ, Van der Does D, Zamioudis C, Leon-Reyes A, Van Wees SCM. Hormonal Modulation of Plant Immunity. In: Schekman R, editor. *Annual Review of Cell and Developmental Biology*, Vol 28 2012. p. 489-521.
- [15] Wierzbza MP, Tax FE. An Allelic Series of bak1 Mutations Differentially Alter bir1 Cell-Death, Immune Response, Growth, and Root Development Phenotypes in *Arabidopsis thaliana*. *Genetics*. 2015.
- [16] Navarro L, Dunoyer P, Jay F, Arnold B, Dharmasiri N, Estelle M, et al. A plant miRNA contributes to antibacterial resistance by repressing auxin signaling. *Science*. 2006;312:436-9.
- [17] Navarro L, Bari R, Achard P, Lison P, Nemri A, Harberd NP, et al. DELLAs control plant immune responses by modulating the balance and salicylic acid signaling. *Current Biology*. 2008;18:650-5.
- [18] Yang DL, Yao J, Mei CS, Tong XH, Zeng LJ, Li Q, et al. Plant hormone jasmonate prioritizes defense over growth by interfering with gibberellin signaling cascade. *Proceedings of the National Academy of Sciences of the United States of America*. 2012;109:E1192-E200.
- [19] Attaran E, Major IT, Cruz JA, Rosa BA, Koo AJK, Chen J, et al. Temporal Dynamics of Growth and Photosynthesis Suppression in Response to Jasmonate Signaling. *Plant physiology*. 2014;165:1302-14.
- [20] Noir S, Bomer M, Takahashi N, Ishida T, Tsui TL, Balbi V, et al. Jasmonate Controls Leaf Growth by Repressing Cell Proliferation and the Onset of Endoreduplication while Maintaining a Potential Stand-By Mode. *Plant physiology*. 2013;161:1930-51.
- [21] Zhang C, Xie Q, Anderson RG, Ng G, Seitz NC, Peterson T, et al. Crosstalk between the Circadian Clock and Innate Immunity in *Arabidopsis*. *Plos Pathogens*. 2013;9.
- [22] Wang W, Barnaby JY, Tada Y, Li H, Toer M, Caldelari D, et al. Timing of plant immune responses by a central circadian regulator. *Nature*. 2011;470:110-4.
- [23] Fay PA, Prober SM, Harpole WS, Knops JMH, Bakker JD, Borer ET, et al. Grassland productivity limited by multiple nutrients. *Nature Plants*. 2015;1:15080.
- [24] D'Auria JC, Gershenzon J. The secondary metabolism of *Arabidopsis thaliana*: growing like a weed. *Current Opinion in Plant Biology*. 2005;8:308-16.
- [25] Halkier BA, Gershenzon J. Biology and biochemistry of glucosinolates. *Annual Review of Plant Biology*. 2006;57:303-33.
- [26] Glazebrook J. Contrasting mechanisms of defense against biotrophic and necrotrophic pathogens. *Annual Review of Phytopathology*. 2005;43:205-27.
- [27] Rate DN, Cuenca JV, Bowman GR, Guttman DS, Greenberg JT. The Gain-of-Function *Arabidopsis* *acd6* Mutant Reveals Novel Regulation and Function of the Salicylic Acid Signaling Pathway in Controlling Cell Death, Defenses, and Cell Growth. *Plant Cell*. 1999;11:1695-708.
- [28] Song JT, Lu H, McDowell JM, Greenberg JT. A key role for ALD1 in activation of local and systemic defenses in *Arabidopsis*. *Plant Journal*. 2004;40:200-12.
- [29] Ellis C, Turner JG. The *Arabidopsis* mutant *cev1* has constitutively active jasmonate and ethylene signal pathways and enhanced resistance to pathogens. *Plant Cell*. 2001;13:1025-33.
- [30] Lu H, Salimian S, Gamelin E, Wang GY, Fedorowski J, LaCourse W, et al. Genetic analysis of *acd6-1* reveals complex defense networks and leads to identification of novel defense genes in *Arabidopsis*. *Plant Journal*. 2009;58:401-12.
- [31] Todesco M, Balasubramanian S, Hu TT, Traw MB, Horton M, Epple P, et al. Natural allelic variation underlying a major fitness trade-off in *Arabidopsis thaliana*. *Nature*. 2010;465:632-U129.

- [32] Ng GN, Seabolt S, Zhang C, Salimian S, Watkins TA, Lu H. Genetic Dissection of Salicylic Acid-Mediated Defense Signaling Networks in *Arabidopsis*. *Genetics*. 2011;189:851-U214.
- [33] Hemm MR, Ruegger MO, Chapple C. The *Arabidopsis* *ref2* mutant is defective in the gene encoding CYP83A1 and shows both phenylpropanoid and glucosinolate phenotypes. *Plant Cell*. 2003;15:179-94.
- [34] Humphreys JM, Chapple C. Rewriting the lignin roadmap. *Current Opinion in Plant Biology*. 2002;5:224-9.
- [35] Kim JI, Ciesielski PN, Donohoe BS, Chapple C, Li X. Chemically Induced Conditional Rescue of the Reduced Epidermal Fluorescence8 Mutant of *Arabidopsis* Reveals Rapid Restoration of Growth and Selective Turnover of Secondary Metabolite Pools. *Plant physiology*. 2014;164:584-95.
- [36] Landry LG, Chapple CCS, Last RL. *Arabidopsis* mutants lacking phenolic sunscreens exhibit enhanced Ultraviolet-B injury and oxidative damage. *Plant physiology*. 1995;109:1159-66.
- [37] Bonawitz ND, Soltau WL, Blatchley MR, Powers BL, Hurlock AK, Seals LA, et al. REF4 and RFR1, Subunits of the Transcriptional Coregulatory Complex Mediator, Are Required for Phenylpropanoid Homeostasis in *Arabidopsis*. *Journal of Biological Chemistry*. 2012;287:5434-45.
- [38] Bonawitz ND, Kim JI, Tobimatsu Y, Ciesielski PN, Anderson NA, Ximenes E, et al. Disruption of Mediator rescues the stunted growth of a lignin-deficient *Arabidopsis* mutant. *Nature*. 2014;509:376-+.
- [39] Kim JI, Dolan WL, Anderson NA, Chapple C. Indole Glucosinolate Biosynthesis Limits Phenylpropanoid Accumulation in *Arabidopsis thaliana*. *Plant Cell*. 2015;27:1529-46.
- [40] Cipollini D. Constitutive expression of methyl jasmonate-inducible responses delays reproduction and constrains fitness responses to nutrients in *Arabidopsis thaliana*. *Evolutionary Ecology*. 2010;24:59-68.
- [41] Paul-Victor C, Züst T, Rees M, Kliebenstein DJ, Turnbull LA. A new method for measuring relative growth rate can uncover the costs of defensive compounds in *Arabidopsis thaliana*. *New Phytologist*. 2010;187:1102-11.
- [42] Züst T, Joseph B, Shimizu KK, Kliebenstein DJ, Turnbull LA. Using knockout mutants to reveal the growth costs of defensive traits. *Proceedings of the Royal Society B-Biological Sciences*. 2011;278:2598-603.
- [43] Joseph B, Corwin JA, Zuest T, Li B, Iravani M, Schaepman-Strub G, et al. Hierarchical nuclear and cytoplasmic genetic architectures for plant growth and defense within *Arabidopsis*. *Plant Cell*. 2013;25:1929-45.
- [44] Lankau RA, Kliebenstein DJ. Competition, herbivory and genetics interact to determine the accumulation and fitness consequences of a defence metabolite. *Journal of Ecology*. 2009;97:78-88.
- [45] Lankau RA, Strauss SY. Mutual feedbacks maintain both genetic and species diversity in a plant community. *Science*. 2007;317:1561-63.
- [46] Lankau RA, Strauss SY. Community complexity drives patterns of natural selection on a chemical defense of *Brassica nigra*. *American Naturalist*. 2008;171:150-61.
- [47] Mauricio R, Stahl EA, Korves T, Tian DC, Kreitman M, Bergelson J. Natural selection for polymorphism in the disease resistance gene *Rps2* of *Arabidopsis thaliana*. *Genetics*. 2003;163:735-46.
- [48] Tian D, Traw MB, Chen JQ, Kreitman M, Bergelson J. Fitness costs of R-gene-mediated resistance in *Arabidopsis thaliana*. *Nature*. 2003;423:74-7.
- [49] Korves T, Bergelson J. A novel cost of R gene resistance in the presence of disease. *American Naturalist*. 2004;163:489-504.
- [50] Kliebenstein DJ, Rowe HC. Ecological costs of biotrophic versus necrotrophic pathogen resistance, the hypersensitive response and signal transduction. *Plant Science*. 2008;174:551-6.
- [51] Lankau RA. Specialist and generalist herbivores exert opposing selection on a chemical defense. *New Phytologist*. 2007;175:176-84.

- [52] Kniskern JM, Traw MB, Bergelson J. Salicylic acid and jasmonic acid signaling defense pathways reduce natural bacterial diversity on *Arabidopsis thaliana*. *Molecular Plant-Microbe Interactions*. 2007;20:1512-22.
- [53] Lebeis SL, Paredes SH, Lundberg DS, Breakfield N, Gehring J, McDonald M, et al. Salicylic acid modulates colonization of the root microbiome by specific bacterial taxa. *Science*. 2015;349:860-4.
- [54] Lorang JM, Carkaci-Salli N, Wolpert TJ. Identification and characterization of victorin sensitivity in *Arabidopsis thaliana*. *Molecular Plant-Microbe Interactions*. 2004;17:577-82.
- [55] Lorang JM, Sweat TA, Wolpert TJ. Plant disease susceptibility conferred by a "resistance" gene. *Proceedings of the National Academy of Sciences of the United States of America*. 2007;104:14861-6.
- [56] Sweat TA, Wolpert TJ. Thioredoxin h5 is required for victorin sensitivity mediated by a CC-NBS-LRR gene in *Arabidopsis*. *Plant Cell*. 2007;19:673-87.
- [57] Sweat TA, Lorang JM, Bakker EG, Wolpert TJ. Characterization of natural and induced variation in the LOV1 gene, a CC-NB-LRR gene conferring victorin sensitivity and disease susceptibility in *Arabidopsis*. *Molecular Plant-Microbe Interactions*. 2008;21:7-19.
- [58] Lorang J, Kidarsa T, Bradford CS, Gilbert B, Curtis M, Tzeng SC, et al. Tricking the Guard: Exploiting Plant Defense for Disease Susceptibility. *Science*. 2012;338:659-62.
- [59] Mauricio R, Rausher MD. Experimental manipulation of putative selective agents provides evidence for the role of natural enemies in the evolution of plant defense. *Evolution*. 1997;51:1435-44.
- [60] Bidart-Bouzat MG, Kliebenstein DJ. Differential levels of insect herbivory in the field associated with genotypic variation in glucosinolates in *Arabidopsis thaliana*. *Journal of Chemical Ecology*. 2008;34:1026-37.
- [61] Siemens DH, Haugen R. Plant chemical defense allocation constrains evolution of tolerance to community change across a range boundary. *Ecology and Evolution*. 2013;3:4339-47.
- [62] Kessler A, Halitschke R, Poveda K. Herbivory-mediated pollinator limitation: negative impacts of induced volatiles on plant-pollinator interactions. *Ecology*. 2011;92:1769-80.
- [63] Kerwin R, Feusier J, Corwin J, Rubin M, Lin C, Muok A, et al. Natural genetic variation in *Arabidopsis thaliana* defense metabolism genes modulates field fitness. *Elife*. 2015;4.
- [64] Mauricio R, Rausher MD, Burdick DS. Variation in the defense strategies of plants: Are resistance and tolerance mutually exclusive? *Ecology*. 1997;78:1301-11.
- [65] Agrawal AA. Overcompensation of plants in response to herbivory and the by-product benefits of mutualism. *Trends in plant science*. 2000;5:309-13.
- [66] Paige KN. Overcompensation in response to mammalian herbivory - From mutualistic to antagonistic interactions. *Ecology*. 1992;73:2076-85.
- [67] Kliebenstein DJ. Making new molecules - evolution of structures for novel metabolites in plants. *Curr Opin Plant Biol*. 2013;Online.
- [68] Zhao Z, Zhang W, Stanley BA, Assmann SM. Functional Proteomics of *Arabidopsis thaliana* Guard Cells Uncover New Stomatal Signaling Pathways. *Plant Cell*. 2008;20:3210-26.
- [69] Melotto M, Underwood W, He SY. Role of stomata in plant innate immunity and foliar bacterial diseases. *Annual Review of Phytopathology*. 2008;46:101-22.
- [70] Shroff R, Vergara F, Muck A, Svatos A, Gershenzon J. Nonuniform distribution of glucosinolates in *Arabidopsis thaliana* leaves has important consequences for plant defense. *Proceedings of the National Academy of Sciences of the United States of America*. 2008;105:6196-201.
- [71] Hoelscher D, Shroff R, Knop K, Gottschaldt M, Crecelius A, Schneider B, et al. Matrix-free UV-laser desorption/ionization (LDI) mass spectrometric imaging at the single-cell level: distribution of secondary metabolites of *Arabidopsis thaliana* and *Hypericum* species. *Plant Journal*. 2009;60:907-18.
- [72] Gols R, Wagenaar R, Bukovinszky T, van Dam NM, Dicke M, Bullock JM, et al. Genetic variation in defense chemistry in wild cabbages affects herbivores and their endoparasitoids. *Ecology*. 2008;89:1616-26.

[73] Shelton AL. Variation in chemical defences of plants may improve the effectiveness of defence. *Evolutionary Ecology Research*. 2004;6:709-26.

[74] Shelton AL. Within-plant variation in glucosinolate concentrations of *Raphanus sativus* across multiple scales. *Journal of Chemical Ecology*. 2005;31:1711-32.

Figure 1. Alternative models of growth and defense

A. Standard illustration of the growth vs immunity tradeoff where the two are in opposition.

B. Alternative conceptualization whereby growth and defense are in a continual conversation to coordinate.