Original research article

Mineral micronutrient and prebiotic carbohydrate profiles of USA-grown kale (*Brassica oleracea* L. var. *acephala*)

Dil Thavarajah^{1*}, Pushparajah Thavarajah^{1, 2}, Alex Abare¹, Suranga Basnagala¹, Craig Lacher³, Powell Smith⁴, Gerald F. Combs Jr⁵.

¹Agricultural and Environmental Sciences, 270 Poole Agricultural Center, Clemson University, Clemson, SC 29634, USA.

²International College Beijing, China Agricultural University, No. 17 Qinghua Dong Road, Haidian District, Beijing, Post Code 100083, China.

³Grand Forks Human Nutrition Research Centre, ARS/USDA, 2420 2nd Ave. N, STOP 9034, Grand Forks, ND 58202-9034, USA.

⁴CUCES-Lexington County, 605 W. Main St. Ste. 109, Lexington, SC 29072, USA.

⁵Jean Mayer USDA Human Nutrition Research Center on Aging, Tufts University, Boston, MA 02111, USA.

*Corresponding author: dthavar@clemson.edu; Tel: +18646562607; Fax: +18646564960

Abstract

Kale (*Brassica oleracea* L. var. *acephala*.) is a leafy green brassica vegetable, the production levels of which have increased over the past 10 years due to increased demand by North American consumers. Kale is perceived to be highly nutritious leafy green vegetable even though

its nutrient profile has not been well characterized to date. The objective of this study was to determine the nutritional composition (energy, protein, mineral, and prebiotic carbohydrate concentrations) of 25 different kale genotypes grown in Pelion, South Carolina, USA, and assess its potential as a whole food source of daily essential minerals and dietary fiber. The results of this study show a single 100 g serving of fresh kale can provide a significant percentage of the recommended daily intake of mineral micronutrients (188-873 mg K; 35-300 mg Ca; 20-100 mg Mg) identified by the 2015 Dietary Guidelines Advisory Committee as being underconsumed by Americans. This serving of kale can also provide approximately 0.4-6.7 g of prebiotic carbohydrates, including sugar alcohols (45.4-59.8 mg), simple sugars (0.4-3348 mg), and hemicellulose (245-703 mg). Fresh kale is a low calorie food (36-98 kcal/100 g) with moderate levels of protein (1.6-5.9 g/100 g). Among the kale genotypes studied, 'Frizzy Lizzy', 'Dauro' and 'Fizz' have significantly high levels of essential minerals and prebiotic carbohydrates, moderate protein content, and low caloric value. These data confirm that kale is a whole food that can provide significant quantities of daily essential minerals and prebiotic carbohydrates. By genetic and location sourcing, the nutritional quality of kale could be further enhanced to benefit North American consumers.

Key words: Kale; Nutritional composition; Prebiotic carbohydrates; Mineral; Recommended daily intake; Food analysis; Food composition

1 Introduction

More than 50 nutrients are essential for human physiological functions and general health. However, billions of people worldwide do not receive sufficient daily quantities of micronutrients due to their low bioavailability in commonly eaten foods. Inadequate intake of

these essential minerals leads to “hidden hunger”, also known as micronutrient malnutrition. Among these, potassium (K), calcium (Ca), and magnesium (Mg) deficiencies are common in the USA, while deficiencies in other essential minerals including iron (Fe) and zinc (Zn) are prevalent in Southeast Asia and Africa (USDA, 2015; WHO, 2015). Cruciferous green leafy vegetables including kale (*Brassica oleracea* L. var. *acephala*) are major sources of these essential micronutrients, however, few Americans consume the daily recommended intake of leafy vegetables. In 2015, the Dietary Guidelines Advisory Committee (USDA, 2015) urged Americans to increase their vegetable intake as a strategy to prevent mineral deficiencies, obesity, and overweight. According to the Center for Disease Control (CDC, 2015), no state’s population is meeting the nationally recommended vegetable consumption level of 5-6 servings per day. In fact, the average American consumes only ~59% of the recommended intake of vegetables (USDA, 2015). The result is widespread under consumption of several nutrients (e.g., K, Ca, Mg) and dietary fiber relative to nutritional needs (USDA, 2015).

Kale has received recent attention from the health and nutrition sectors due to its nutrient profile. However, current kale nutritional quality data are based on a very narrow sampling base conducted more than 10 years ago and are lacking with respect to information on prebiotic carbohydrates (USDA, 2015). The available data support the consumer perception that kale places high on lists of healthy foods. On average, kale contains a total of 8% carbohydrates and 3.6% dietary fiber (USDA, 2015); however, the levels of phytonutrients in kale genotypes have not been well characterized.

Several studies show prebiotic-rich, micronutrient-rich, low-calorie diets to play important roles in supporting intestinal health, with potential to prevent obesity and promote gut health (Ley et al., 2005; Dumas et al., 2006; Wu et al., 2011). Prebiotic-rich diets promote intestinal microbial

diversity, stimulate the immune system, promote mineral micronutrient absorption, decrease the risk of developing colon cancer, reduce excessive blood glucose and cholesterol levels, and improve insulin sensitivity (Johnston et al., 2010; Lee and Mazmanian, 2010; Caselato et al., 2011). As such, products enriched with prebiotics are becoming more popular, in the belief that they may also have probiotic effects.

Naturally occurring prebiotic carbohydrates include dietary fiber and sugar alcohols (Roberfroid, 2007). Dietary fiber is comprised of non-digestible starch polysaccharides including resistant starch (RS) and non-starch polysaccharides. Sugar alcohols (e.g., sorbitol, mannitol) are derived from simple sugars (Roberfroid, 2007). Americans 20 years of age and over consume less than half of the recommended amounts of prebiotic carbohydrates (e.g., 10 g/d of fructooligosaccharides; 7 g/d of galactooligosaccharide). Legumes, cereals, and vegetables are rich in these prebiotic carbohydrates. For example, a single 100 g serving of lentils (*Lens culinaris* Medik) provides 13 g of prebiotics, the level of which can as much as double after cooking/processing (Johnson et al., 2013, 2015). However, no prebiotic data are available for commonly consumed leafy green vegetables and none are available for kale. Therefore, this study was undertaken to determine the nutritional composition (energy, protein, minerals, and prebiotic carbohydrate concentration) of 25 different kale genotypes grown in Pelion, South Carolina (SC), USA, to the end of developing an evidence-based understanding of the nutritional value of kale.

2 Materials and methods

2.1 Materials

Standards, reagents, acids, and high-purity solvents used for sample and reagent preparation were purchased from Sigma-Aldrich Co. (St. Louis, MO, USA) and VWR International (Radnor, PA, USA) and used without further purification. Water, distilled and deionized (ddH₂O) to a resistance of ≥ 18.2 M Ω (Milli-Q Water System, Millipore, Milford, MA, USA), was used for sample and reagent preparation.

2.2 Kale samples

Field-grown fresh kale samples were collected from variety trials conducted by Walter P. Rawls and Sons, Inc. in Pelion, SC, USA in the fall of 2014. Twenty-five commonly grown kale genotypes were selected based on market class, consumer demand, disease resistance, biological yield, and preference for future breeding and selection research (**Table 1**). Kale plants were harvested at physiological or horticultural maturity (ready to eat). Fresh leaf samples (250 g) were taken randomly from the entire harvested plant of each of four independent replicated field plots and subjected to energy, protein, mineral, and carbohydrate analyses. A total of 100 replicate kale leaf samples were collected. Moisture content of fresh sub-samples (105 °C for 16 h) and the remaining samples were immediately freeze-dried and stored at -40 °C until analysis. Prior to each analysis, the freeze dried samples were finely ground using a mortar and pestle. Nutrient composition data are reported on a fresh weight basis (85% moisture).

2.3 Energy and protein analysis

Finely ground freeze-dried kale samples (1 g) were compressed into a pellet, and then ignited in an oxygen-rich closed environment using an oxygen bomb calorimeter (Parr Instrument

Company, Moline, Illinois 61265). Quality control and calibration was accomplished using benzoic acid as the standard (Schmidt-Rohr, 2015). Samples were analyzed in duplicate, and a NIST standard (Typical diet SRM 1548A; NIST Gaithersburg, MD 20899, USA) was used as a reference. Total nitrogen was determined on freeze dried kale samples using a LECO FP3000 CNS analyzer; protein content was calculated ($N \times 6.25$).

2.4 Mineral concentrations

Minerals were determined by inductively coupled argon plasma emission spectroscopy (ICP) using a Thermo 6500 Duo instrument (Thermo Fisher Scientific, [no city], PA, USA) after HNO_3 - H_2O_2 digestion (Thavarajah et al., 2009). Aliquots (500 mg) of freeze-dried samples were digested with 6 mL of 70% HNO_3 overnight at room temperature. They were then heated to 90 °C for 1 h, after which 3 mL of 30% H_2O_2 were added and the sample was held at 90 °C for 15-20 min. Finally, 3 mL of 6M hydrochloric acid was added and samples were held at 90 °C for another 5 min. Samples were then filtered and made up to 10 mL in Milli-Q water. Detection limits were 30 µg/L for K, Ca, and Mg and 5 µg/L for Fe, Zn, Mn, Cu, and Se. Analytical quality assurance was accomplished using authentic calibration standards and the NIST standard reference material peach leaves 1547. Results were used to calculate the percentages of recommended daily allowance values (%RDA, National Academy of Sciences, 2004) provided by a 100 g portion size of fresh kale

2.5 Carbohydrate analysis

Freeze-dried kale samples (~500 mg) were extracted as described by Muir et al. (2009). Briefly, this method involved extraction with distilled water at 80 °C for 1 h. The resulting extract was passed through a 13 mm × 0.45 µm nylon syringe filter (Chromatographic Specialties,

Brockville, ON, Canada). Analyses for sugar alcohols, monosaccharides, disaccharides, and oligosaccharides were performed by high-performance liquid-liquid partition chromatography (ICS-5000 Dionex, Sunnyvale, CA, USA) as described by Feinberg (2009) with pulsed amperometric detection (PAD, Dionex) in series with electrochemical detection (Thermo Fisher Scientific, Inc.). Carbohydrates were separated by a CarboPac PA-100 4 × 250 mm column in series with a CarboPac PA-100 4 × 50 mm guard column (Dionex, Sunnyvale, CA, USA). The Mobile phase flow rate was maintained at 1 mL/min. Solvents used for elution were 100 mM sodium hydroxide/600 mM sodium acetate (solvent A), 200 mM sodium hydroxide (solvent B), and ddH₂O (solvent C). Solvents B and C were used at 50% for the initial 16 min. This was followed by a gradient change to 3% A, 48.5% B, and 48.5% C at the 16 min mark. At 18 min, the gradient changed linearly to 16% A, 42% B, and 42% C. Finally, at the 19 min mark the gradient returned to 50% B and 50% C. The silver-silver chloride electrode was set at 2.0 µA. The minimum detectible level (signal:noise ≥3) of each analyte was 0.1 ppm. Peak areas for a reference lentil sample were routinely analyzed with an error of less than 5%, and linear calibration models for standards had errors <5%. Concentrations of carbohydrates in the filtrate (C) were calculated from the calibration model used to calculate concentrations in sample dry matter in the expression $X = (C \times V) / m$, where X is the concentration of the carbohydrate in the sample (corrected for moisture), V is the final diluted volume, and m is the mass of the dry sample aliquot. Hemicellulose was measured as described above method after digesting 500 mg of sample with 5 mL of 7% (w/w) hydrochloric acid (HCl) at 55 °C for 120 minutes. Data are reported as mg/100g fresh weight; data for genotypes ‘Ripbor’, ‘Reflex’, ‘Vates’, ‘Starbor’, and ‘Scarlet’ were below detection limits (and are not included in Table 2).

2.6 Statistical analysis

The experimental design was a complete randomized block design with four replicates for 25 commercial kale genotypes (n=100). Replicates and genotypes were considered as random factors. Class variables include genotype and replication. A mixed model analysis of variance was performed using the PROC GLM procedure of SAS version 9.4 (SAS Institute SAS User's Guide, 2012). Means were separated by Fisher's protected least significant difference at $P < 0.05$.

3 Results

Kale grown in SC was found to be low in energy (36-98 Kcal/100 g) with moderate protein content (1.6-5.9 g/100 g). Contents of essential minerals and prebiotic carbohydrates were variable (Table 2). Some genotypes of kale were rich source of minerals (Table 2, **Fig. 1**). These genotypes also differ in prebiotic carbohydrates including sugar alcohols, simple sugars, and hemicellulose (Table 2, Fig. 2). The total identified prebiotic carbohydrates in kale ranged from 0.4-6.7 g/100 g; however, 5.0-8.0 g/100 g of unidentified soluble carbohydrates were detected. Each of these nutrients showed significant within-variety differences. Kale genotypes varied considerably in nutrient contents. 'Black Magic', 'Fizz', 'Stars & Stripes', and 'Winterbor' had the highest protein levels. 'Black Magic', 'Darkibor', 'Lacinato', 'Maribor', and 'Red Russian' had significantly low energy levels compared to other kale genotypes. Among these genotypes, 'Black Magic' had higher level of protein and low level of energy compared to the other genotypes (**Fig 3**).

Overall, a 100 g serving of fresh kale can provide a significant percentage of the RDA for all analyzed mineral micronutrients, with the exception of Fe for females (**Table 3**). Kale can be

considered a “good source” of mineral micronutrients as one serving (100 g) provides more than 10% of the RDA. Furthermore, substantial genotypic differences in mineral concentration were observed for all elements, and more than 50% of tested genotypes could provide at least 10% of the RDA for the minerals considered (Figure 1).

The greatest contents of K, Ca, Mg, Fe, and Zn were found in the curly kale genotype ‘Frizzy Lizzy’, followed by ‘Lacinato’, ‘Fizz’, and ‘Frizzy Joe’ (Fig 2). Most of these kale genotypes were also high in Mn and Cu; however, Se concentrations were relatively low in all genotypes (data not shown). Prebiotic carbohydrate concentrations also varied by genotype. ‘Frizzy Joe’ (39.2 mg/100 g) and ‘Darkibor’ (33.2 mg/100 g) had the highest concentrations of sorbitol and mannitol, respectively (Fig 3). ‘Italian Kale’ (145 mg/100 g), ‘X’ (cultivar name withheld by grower) (103 mg/100 g), and ‘Frizzy Joe’ (101 mg/100 g) had the highest concentration of sucrose; and ‘Frizzy Joe’ and ‘Dauro’ had the highest concentrations of both fructose (1246-1080 mg/100 g) and glucose (1715-1947 mg/100 g). ‘Frizzy Joe’ had the highest concentrations of mannose (403 mg/100 g) and xylose (201 mg/100 g). ‘Darkibor’ (166 mg/100 g) and ‘X’ (156 mg/100 g) had the highest arabinose contents.

4 Discussion

Kale is one of the oldest brassica vegetables originating from the eastern Mediterranean region, and is consumed either raw or cooked. Recently, kale has received attention from the health and nutrition sectors due to its superior nutrient profile despite the lack of nutritional quality data for cultivars currently in US production. Our data show that US-grown kale has the potential to provide significant quantities of several essential micronutrients and adequate quantities of prebiotic carbohydrates with moderate to low levels of protein and energy. Indeed, a 100 g single

serving of fresh kale can provide significant concentrations of K, Ca, Mg, Mn, Cu, and Se as well as 0.4-6.7 g of total beneficial prebiotic carbohydrates, 1.6-5.9 g of protein, and 36-98 Kcal of energy (Tables 2 and 3). Some genotypes, ‘Lacinato’, ‘Frizzy Lizzy’, and ‘Dauro’ are particularly rich in nutrients that are underconsumed by Americans, making them low-calorie foods that are good sources of essential minerals as well as prebiotic carbohydrates.

A recent study published by the Center for Disease Control ranked 47 “powerhouse” fruits and vegetables (a serving providing $\geq 10\%$ of 17 essential nutrients) that are strongly associated with reducing the risk of heart disease and other non-communicable diseases (Noia, 2014). Kale was ranked 15th based on nutrient concentrations as reported in the USDA National Nutrient Database. The results of the present study show that several genotypes of kale have greater nutrient contents than those listed in the current version of the USDA National Nutrient Database (USDA, 2015), which are based on a very narrow sampling of unspecified cultivars (Table 1; Figure 2). The comprehensive nutrient composition data provided herein on contemporary cultivars represent a significant improvement of the evidence base for decisions about the use of this vegetable in diets.

Despite its gaining popularity, kale remains an understudied vegetable. Only a few studies have compared the nutritional quality of kale with that of other leafy green vegetables (Kawashima and Soares, 2003; Ayaz et al., 2006; Fadigas et al., 2010; Lefsrud et al., 2006). For example, Kawashima and Soares (2003) compared K, Ca, and Mg levels in Brazilian-grown kale to other green vegetables, including Chinese cabbage, cabbage, butter head lettuce, and spinach. They found kale to have the highest concentration of K (712 mg/100 g fresh weight), Ca (283 mg/100 g fresh weight), and Mg (51 mg/100 g of fresh weight). In addition, slow cooking did not change kale’s mineral concentrations. The present results are comparable to their data, with the

differences present attributable to genetic, environmental, and analytical differences. However, other studies report mineral data on a dry weight basis, which hinders comparisons when moisture content data are not reported (Ayaz et al., 2006; Fadigas et al., 2010). The clear variation between genotypes in essential minerals and prebiotic carbohydrates suggests the possibility of further increasing the mineral micronutrients in kale through genetic biofortification (enrichment of micronutrient through plant breeding and genetic selection). Thus, selection of kale genotypes with a greater potential for mineral uptake may be part of a food-based solution to address micronutrient undernutrition.

Glucose and fructose are the major soluble sugars in kale. Our data on soluble sugars compare favorably with those of (Ayaz et al., 2006). Those genotypes with the greatest levels of glucose and fructose, 'Frizzy Lizzy', 'Dauro', 'Blue Ridge', 'Darkibor' and 'Winterbor' are known to require fall temperatures (7-21 °C) to obtain a sweeter taste, crispy leaf quality, and better yield. Kale grown in higher temperatures (>25 °C), such as during southern summers, is bitter compared to the late fall-grown kale. Generally, kale cultivars produce higher levels of water-soluble prebiotics when grown under cooler temperatures, which lead to a sweeter taste and higher nutritional quality.

Consumption of diets rich in fruits and vegetables can reduce the prevalence of chronic diseases, including diabetes, cancer, heart disease, and stroke (Slavin and Lloyd, 2012). Additional health benefits are better control of blood sugar levels, glycemic index, and inflammation as well as reduced risk of obesity due to the high fiber and low calorie content of such a diet (Slavin and Lloyd, 2012). Vegetables only account for about 5-10% of the total calories consumed, yet make a significant contribution to overall human health (Slavin and Lloyd, 2012). A healthy diet rich in vegetables is inversely associated with inflammatory markers C-reactive protein, TNF- α ,

soluble vascular cell adhesion molecules (sVCAM-1), and E-selectin; conversely, a diet rich in refined grains and added sugar is positively associated with systemic inflammation (Cai et al., 2005). In particular, diets rich in prebiotic carbohydrates can decrease systemic inflammation via associations with hindgut microflora (Delzenne et al., 2011). The present results show that kale can be a significant food source of several essential minerals and dietary fiber, while being low in calories and providing moderate amounts of protein. Further, the genetic variation evident in these nutrient contents indicates the potential for breeding to enhance further the nutritional value of kale.

5 Conclusion

In conclusion, data from this study provide evidence that kale is a whole food source of essential minerals K, Ca, Mg, Mn, and Cu as well as dietary fiber including sugar alcohols, simple sugars, and hemicellulose. Kale is also low in calories and provides moderate levels of protein.

Therefore, kale is a potentially good source of minerals and prebiotic carbohydrates and could be promoted for consumption in American diets as per the recommendations of the 2015 Dietary Guidelines Advisory Committee (USDA, 2015).

Acknowledgments

This publication [or project] was supported by the Specialty Crop Block Grant Program at the U.S. Department of Agriculture through grant 15-SCBGP-SC-0013. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the USDA. In addition, funding support was provided by New Faculty Startup funds from Clemson University, SC, USA. We also acknowledge Mr. Charles Wingard, WP Rawl & Sons, Inc. for providing kale samples and technical assistance (Josh Hall, Matt Warren) with field harvest.

References

- Ayaz, F. A., Glew, R. H., Millson, M., Huang, H. S., Chuang, L. T., Sanz, C., Hayırlıoglu-Ayaz, S. (2006). Nutrient contents of kale (*Brassica oleraceae* L. var. *acephala* DC.). *Food Chemistry*, 96(4), 572-579.
- Cai, D., Yuan, M., Frantz, D. F., Melendez, P. A., Hansen, L., Lee, J., et al. (2005). Local and systemic insulin resistance resulting from hepatic activation of IKK- β and NF- κ B. *Nature medicine.*, 11(2): 183-90.
- Caselato, d. S. V. M., Freitas, d. S., Sgarbieri, V. C. (2011). The importance of prebiotics in functional foods and clinical practice. *Food and Nutrition Sciences*, 2 (2):133-144.
- CDC, (2013). State Indicator report on fruits and vegetables, <http://www.cdc.gov/nutrition/downloads/state-indicator-report-fruits-vegetables-2013.pdf>. Accessed on Jan 9, 2015.
- Delzenne, N. M., Neyrinck, A. M., Bäckhed, F., Cani, P. D. (2011). Targeting gut microbiota in obesity: effects of prebiotics and probiotics. *Nature Reviews Endocrinology*, 7(11): 639-46.
- Dumas, M., Barton, R. H., Toye, A., Cloarec, O., Blancher, C., Rothwell, A., et al. (2006). Metabolic profiling reveals a contribution of gut microbiota to fatty liver phenotype in insulin-resistant mice. *Proceedings of National Academy of Science*, 103(33):12511-12516.
- Fadigas, J. C., dos Santos, A. M., de Jesus, R. M., Lima, D. C., Fragoso, W. D., David, J. M., Ferreira, S. L. (2010). Use of multivariate analysis techniques for the characterization of

- analytical results for the determination of the mineral composition of kale. *Microchemical Journal*, 96(2):352-356.
- Feinberg, M., San-Redon, J., Assié, A. (2009). Determination of complex polysaccharides by HPAE-PAD in foods: validation using accuracy profile. *Journal of chromatography B*, 877(23): 2388-2395.
- Johnson, C. R., Thavarajah, D., Combs, Jr., G. F., Thavarajah P. (2013). Lentil (*Lens culinaris* L.): A prebiotic-rich whole food legume. *Food Research International*, 51(1):107–113.
- Johnson, C. R., Thavarajah, D., Thavarajah, P., Payne, S., Moore, J., Ohm, J. B. (2015). Processing, cooking, and cooling affect prebiotic concentrations in lentil (*Lens culinaris* Medikus). *Journal of Food Composition and Analysis*, 38:106-111.
- Johnston, K. L., Thomas, E. L., Bell, J. D., Frost, G. S., Robertson, M. D. (2010). Resistant starch improves insulin sensitivity in metabolic syndrome. *Diabetic Medicine*, 27 (4):391-397.
- Kawashima, L. M., Soares, L. M. V. (2003). Mineral profile of raw and cooked leafy vegetables consumed in Southern Brazil. *Journal of Food Composition and Analysis*, 16(5):605-611.
- Lee, Y. K., Mazmanian, S. K. (2010). Has the microbiota played a critical role in the evolution of the adaptive immune system? *Science*, 330 (6012):1768-1773.
- Lefsrud, M. G., Kopsell, D. A., Kopsell, D. E., Randle, W. M. (2006). Kale carotenoids are unaffected by, whereas biomass production, elemental concentrations, and selenium

- accumulation respond to, changes in selenium fertility. *Journal of Agricultural and Food chemistry*, 54(5):1764-1771.
- Ley, R. E., Bäckhed, F., Turnbaugh, P., Lozupone, C. A., Knight, R. D., Gordon, J. I. (2005). Obesity alters gut microbial ecology. *Proceedings of National Academy of Science*, 102:11070-11075.
- Muir, J. G., Rose, R., Rosella, O., Liels, K., Barrett, J. S., Shepherd, S. J., Gibson, P. R. (2009). Measurement of short-chain carbohydrates in common Australian vegetables and fruits by high-performance liquid chromatography (HPLC). *Journal of Agricultural and Food Chemistry*, 57(2): 554-565.
- National Academy of Sciences, Institute of Medicine, Food and Nutrition Board, (2004). Retrieved from: <http://www.iom.edu/Activities/Nutrition/SummaryDRIs>. Accessed on Jan 9 2016.
- Noia, J. D. (2014). Defining powerhouse fruits and vegetables: a nutrient density approach, *Preventing Chronic Disease*, 11:1-5.
- Roberfroid, M. (2007). Prebiotics: the concept revisited. *The Journal of nutrition*, 137(3): 830S-837S.
- SAS Institute. (2012). User's Guide: Statistics SAS Institute (Version 9.4). In. Cary, NC: SAS Institute.
- Schmidt-Rohr, K. (2015). Why Combustions Are Always Exothermic, Yielding About 418 kJ per Mole of O₂. *Journal of Chemical Education*, 92: 2094–2099.

Slavin, J. L., Lloyd, B. (2012). Health Benefits of Fruits and Vegetables, *Advances in Nutrition: An International Review Journal*, 3(4):506-516.

Thavarajah, D., Thavarajah, P., Sarker, A., Vandenberg, A. (2009). Lentils (*Lens culinaris* Medikus Subspecies *culinaris*): A Whole Food for Increased Iron and Zinc Intake. *Journal of Agricultural and Food Chemistry*, 57:5413-5419.

USDA, National Nutrient Database for Standard Reference Release 28. (2015).

<http://ndb.nal.usda.gov/ndb/foods?qlookup=kale>. Accessed on Jan 9, 2016.

USDA (2015). Dietary Guidelines for Americans, U.S. Department of Agriculture U.S.

Department of Health and Human Services, www.dietaryguidelines.gov. Accessed on Jan 9, 2016.

WHO, (2015). UNICEF-WHO-World Bank Group, Joint child malnutrition estimates.

<http://www.who.int/nutrition/topics/ida/en/>. Accessed on Jan 9, 2016.

Wu, G. D., Chen, J., Hoffmann, C., Bittinger, K., Chen, Y., Keilbaugh, S. A. et al. (2011).

Linking long-term dietary patterns with gut microbial enterotypes. *Science*, 334 (6052): 105-108.

Table 1: Kale genotypes grown in Pelion, SC, USA in the fall of 2014.

Market Class	Genotype
Curly	Darkibor, Dwarf Green Curled Afro, Pentlang Brig, Red Russian, Redbor, Reflex, Ripbor, Scarlet, Star & Stripes, Starbor, Vates, Winterbor, <i>withheld</i> (cultivar name withheld by grower), Blue Ridge, Blue Knight, Maribor
Portuguese	Beira, Dauro
Dinosaur	Black Magic, Bonanza, Italian Kale, Lacinato
Ornamental	Fizz
Mustard	Frizzy Joe, Frizzy Lizzy

Table 2: Nutritional profile (protein, energy, minerals, and prebiotic carbohydrates) of kale genotypes grown in SC, USA.

Nutrient	Concentration ^a		USDA ^b	Genotype effect ^c
	Range	Mean		
Protein (g/100 g)	1.6-5.9	4.2	4.28	*
Energy (Kcal/100 g)	36-98	66	49	*
Minerals (mg/100 g)				
K	188-873	488	491	*
Ca	35-300	106	150	*
Mg	20-100	44	47	*
Fe	0.5-2.3	1.1	1.47	*
Zn	0.2-1.6	0.7	0.56	*
Mn	0.2-2.3	0.8	0.65	*
Cu (µg/100 g)	2-116	55	149	*
Se (µg/100 g)	ND ^e -17	2.3	65	*
Prebiotic carbohydrates (mg/100 g)				
Sugar alcohol				
Sorbitol	ND-59.8	24.5	-	*
Mannitol	ND-45.4	17.9	-	*
Simple sugars				
Glucose	69-3348	993	-	*
Fructose	29-1933	545	-	*
Sucrose	0.4-212	39.3	-	*
Hemicellulose				
Arabinose	ND-245	73.5	-	*
Mannose	ND-703	241	-	*
Xylose	ND-320	59.9	-	*
Total identified prebiotic carbohydrates (g/100 g)	0.4-6.7	1.9	-	
Other prebiotic carbohydrates (g/100 g)	5.0-8.0	5.5	-	

^a Values based on 100 data points [fresh weight, 85% moisture]] from current study.

^b Values based on a single data point [fresh weight, 84% moisture] from USDA Standard Nutrient Data base

^c Genotype effect is significant at $P < 0.05$ (n=100).

^e ND, not detectable levels

Table 3: % RDA of different minerals from a 100 g serving of fresh kale grown in SC, USA.

Element	%RDA from 100g^a	Recommendation
K	5.4-14.3	Male/Female (4.7 g)
Ca	5.2-17.2	Male/Female (1 g)
Mg	6.3-16.5	Male (400-420 mg)
	8.1-21.3	Female (310-320 mg)
Fe	6.8-20.3	Male (8 mg)
	3.0-9.0	Female (18 mg)
Zn	3.4-10.5	Male (11 mg)
	4.6-14.4	Female (8 mg)
Mn	18.3-60	Male (2.3 mg)
	23.3-77	Female (1.8 mg)
Cu	2.8-11	Male/Female (900 µg)
Se	0.7-26.7	Male/Female (55 µg)

^a Percent recommended daily allowance (%RDA) was calculated based on National Academy of Sciences, 2004. Values are based on fresh weight (85% moisture), n=100 per nutrient

Figure 1 Variation in energy, protein, and total sugar concentration of 20 kale genotypes grown in SC, USA

Figure 2: Genetic variation of mineral concentrations in 25 kale genotypes: (a) K; (b) Ca and Mg; (c) Fe and Zn. Means with in the bars (element) followed by different letters are significant at $P<0.05$ (differences were shown only for highest and lowest values). Vertical bars indicate kale genotypes that provide 10% of the recommended daily intake (%RDA) of minerals.

Figure 3: Genetic variation of carbohydrate concentrations in 20 kale genotypes: (a) sugar alcohols; (b) simple sugars; and (c) hemicellulose.