**ACTIVE DEFORMATION IN THE NORTHERN SIERRA DE VALLE
FÉRTIL, SIERRAS PAMPEANAS, ARGENTINA**

Gustavo Ortiz^{1*}, Patricia Alvarado¹, Julie C. Fosdick², Laura Perucca¹, Mauro Saez¹,
Agostina Venerdini¹

*1 Centro de Investigaciones de la Geósfera y la Biósfera (CIGEOBIO). Facultad de
Ciencias Exactas, Físicas y Naturales, Universidad Nacional de San Juan - Consejo
Nacional de Investigaciones Científicas y Técnicas (CONICET). Meglioli 1160 S (5406)
Rivadavia, San Juan, Argentina. gfortiz@unsj-cuim.edu.ar, alvarado@unsj-cuim.edu.ar,
lperucca@unsj-cuim.edu.ar, msaez@unsj-cuim.edu.ar, agostinaliavenerdini@gmail.com*

*2 Department of Geological Sciences, Indiana University, 1001 East Tenth Street,
Bloomington, IN 47405, USA. jfosdick@indiana.edu*

**Corresponding author at: CIGEOBIO-Grupo de Sismotectónica. Facultad de Ciencias
Exactas, Físicas y Naturales, Universidad Nacional de San Juan, Meglioli 1160 Sur, Rivadavia
(5406), San Juan, Argentina. Tel.: +54 264 4234129 (203). E-mail address: gfortiz@unsj-
cuim.edu.ar*

Abstract

The Western Sierras Pampeanas region in the San Juan Province is characterized by thick-skinned deformation with approximately N-S trending ranges of average heights of 2500 m and a high frequency occurrence of seismic activity. Its location to the east of the mainly thin-skinned tectonics of the Argentine Precordillera fold-and-thrust belt suggests that at 30°S, deformation is concentrated in a narrow zone involving these two morphostructural units. In this paper, we present new apatite (U-Th)/He results (AHe) across the northern part of the Sierra de Valle Fértil (around 30°S) and analyze them in a framework of thermochronologic available datasets. We found Pliocene AHe results for Carboniferous and Triassic strata in the northern Sierra de Valle Fértil consistent with the hypothesis of recent cooling and inferred erosional denudation concentrated along the northern end of this mountain range. Our analysis shows that this northern region may have evolved under different conditions than the central part of the Sierra de Valle Fértil. Previous studies have observed AHe ages consistent with Permian through Cretaceous cooling, indicating the middle part of the Sierra de Valle Fértil remained near surface before the Pampean slab subduction flattening process. Those studies also obtained ~ 5 My cooling ages in the southern part of the Sierra de Valle Fértil, which are similar to our results in the northern end of the range. Taken together, these results suggest a pattern of young deformation in the northern and southern low elevation ends of the Sierra de Valle Fértil consistent with regions of high seismic activity, and Quaternary active faulting along the western-bounding thrust fault of the Sierra de Valle Fértil.

Keywords: Sierra de Valle Fértil, basement-cored uplift, thermochronology, flat-slab subduction

Introduction

Basement-involved deformation is an unusual feature in retroarc foreland regions of orogenic belts, and for this reason comparisons between ancient and modern basement-cored uplifts (like the Northamerican Laramide Rocky Mountains versus Argentinean Sierras Pampeanas) have been performed in order to better understand the mountain building processes responsible for this deformation (Jordan and Allmendinger, 1986; Ramos, 2000). Many years of investigations in worldwide tectonics has led scientists to propose flat-slab subduction as the mechanism that drives strain into the foreland, and hence deformation. This link between the timing of slab shallowing and the record of orogenic exhumation in the Argentine broken-foreland basin has been well documented in a continuous study of its structural, magmatic, tectonic and sedimentary history (Jordan et al., 1983; Strecker et al., 1989; Allmendinger et al., 1990; Kay et al., 1991; Ramos et al., 2002). In light of the fact that thermochronology may place quantitative constraints on the timing and magnitude of rock exhumation, relatively few low-temperature thermochronology studies have been conducted in the Argentinean Sierras Pampeanas. It is worth to note that the Sierra de Valle Fértil (SVF) lying above the flat-slab subduction of the Nazca Plate offers an ideal scenario to test signals of uplift/faulting and exhumation.

The aim of this paper is to identify and characterize the style of deformation in the northern SVF, which is part of the western Sierras Pampeanas (Fig. 1) combining new low-temperature thermochronology data, neotectonic and seismic activity evidences. Constraints on the timing and magnitude of exhumation compared to current deformation data in this region can provide insights about mechanisms of basement-cored uplifts into the Andean foreland, which are typically exposed in the flat slab subduction region of Argentina.

Recent investigations of exhumation patterns in the Sierras Pampeanas using low-temperature thermochronological techniques depict overall low (<2-3 km) magnitudes of exhumation, most of which has been attributed to pre-Cenozoic events and to a lesser degree Cenozoic exhumation (Coughlin et al., 1998; Löbens et al., 2011; Löbens et al., 2013; Bense et al., 2013). These studies used zircon and apatite (U-Th-Sm)/He (ZHe and AHe, respectively), zircon and apatite fission track (ZFT and AFT, respectively) thermochronology and K-Ar dating of fault gouge in order to better estimate the timing and rates of exhumation and inferred uplift of this wide morphotectonic area in central Argentina. Previous results from Coughlin et al. (1998) show two pre-Andean cooling events during late Paleozoic and late Mesozoic related to a Permian-Triassic flat-slab subduction and Mesozoic rifting events, respectively. This late Paleozoic flat-slab subduction is supported by evidence of an eastward propagation of calc-alkaline magmatism followed by extensional collapse of previous orogenic building (Ramos et al., 1994; Martinez et al., 2006). In addition, two phases of Andean events were recognized by Coughlin et al. (1998): an Early Miocene heating event corresponding to foreland basin sedimentation loading and a Miocene-Pliocene exhumation starting at ca. 10 Ma which involved some Precordillera front uplift and sediment recycling (Jordan et al., 1983; Coughlin et al., 1998; Fosdick and Carrapa, 2012). Dávila and Carter (2013) performed studies at the base of the ~10 km thick exposed Cenozoic deposits of the Vinchina Basin, showing unreset pre-Cenozoic thermochronology data, which remain inconsistently. According to their AFT data these pre-Cenozoic patterns of deformation were related to burial heating followed by rapid uplift and unroofing, which consequently may have lead to an uncompleted reset of the system. Although AFT data alone do not constrain the timing of exhumation for shallower surfaces, Bense et al. (2013) have used a different and more appropriate technique based on AHe dating obtaining non-reset AHe ages. Taken together,

these results suggest that some positive topography may have already existed in this region prior to Neogene flat-slab subduction. Superimposed, a causal relationship has been proposed between southeastward propagation of exhumation in the Sierras Pampeanas and the shallowing of the subducted slab probably enhanced by the buoyancy of the Juan Fernández Ridge (JFR) lying on top of the Nazca plate (Coughlin et al., 1998; Yañez et al., 2001; Kay and Mpodozis, 2002).

In this paper we present apatite (U-Th)/He data from the Carboniferous-Triassic basin fill that is exposed in the structural hangingwall block of the northern SVF (Fig. 2) to complement the existing datasets in the Sierra de Valle Fértil. Existing AHe data in the northern, central and southern sector of the Sierra de Valle Fértil (Bense et al., 2013; Fosdick et al., 2015) are also used in this work. In addition, seismicity distribution patterns and neotectonic features are described to put constraints on how the SVF evolved in the last 5 Myr and how it is actually deforming.

FIGURE 1

Geological Framework

The Sierras Pampeanas geological province is a wide morphotectonic region in the western-central part of Argentina, where continental deformation is observed more than 700 km east of the Chilean trench into the Andean foreland (González Bonorino, 1950; Jordan and Allmendinger 1986; Kay and Mpodozis 2002). In addition, there is a volcanic gap and a horizontal subduction of the slab between 27°S-33°S with the most recent volcanism recorded in Pocho and Morro at 4 Ma and 1.9 Ma, respectively (Gordillo and Lencinas, 1981; Llambías and Brogioni, 1981) (Fig 1). The Sierras Pampeanas region consists of a series of elongated N–S trending basement-cored uplifts that have structurally segmented

the previously continuous Andean foreland basin (González Bonorino, 1950; Cardozo and Jordan, 2001). The whole morphostructural unit has been divided in western (WSP) and eastern (ESP) Sierras Pampeanas domains according to their lithology and metamorphic grade of the exposed rocks (Gonzalez Bonorino, 1950; Caminos, 1979). The ESP are composed of metamorphic and igneous rocks developed by accretion of different allochthonous and parautochthonous terranes during the late Proterozoic and the early Paleozoic that have been amalgamated into a larger cratonic block defined as the Pampia terrane (Ramos et al., 2002, 2010; Steenken et al., 2004; Miller and Söllner, 2005). This terrane underwent a complex history of deformation, metamorphism, and magmatism after its collision with the Rio de La Plata craton at the end of the Proterozoic. West of the Pampia terrane, the subsequent collision of the Precordilleran (Cuyania) terrane of Laurentian affinities (Astini et al., 2003) generated the Early Ordovician Famatinian arc in the south-western boundary of Gondwana (Rapela et al., 2007; Otamendi et al., 2009). Vestiges of this ancient magmatic arc are grouped as part of the WSP.

The SVF is one of the westernmost expressions of the WSP in Argentina (Fig 2). The north and south margins of the crystalline basement-cored of the SVF are overlain by thick sequences of Triassic sedimentary rocks corresponding to the ~2500-m-thick Ischigualasto and ~1500-m-thick Marayes basins, respectively (Milana and Alcober, 1994; Malizia et al., 1995; Spalletti, 1999). The western flank of the range is bounded by the east-dipping Valle Fértil reverse fault system that places in contact basement rocks over the Bermejo valley strata (Snyder et al., 1990; Zapata and Allmendinger, 1996). The same region has been proposed to be the place of a suture between the Cuyania and Pampia terranes (Snyder et al., 1990; Ramos, 1994; Castro de Machuca et al., 2008).

Andean deformation associated with the Sierras Pampeanas region propagated from west to east during the past 10 Myr (Malizia et al., 1995; Ramos et al., 2002) and agrees well with the migration of magmatism away from the trench as the slab flattened (Kay and Abbruzzi 1996; Kay and Mpodozis, 2002). It is worth to note , that deformation and magmatism not only had migrated laterally from west to east but also from north to south in association with the shallowing of the subducted slab (Coughlin et al., 1998) and the path of the JFR beneath South America in the last 15 Myr (Yañez et al., 2001; Ramos et al., 2002). Because the SVF extends more than 180 km along a NW-SE trending orientation within the WSP, we suggest its deformational and exhumational history could help to test for any lateral migration of deformation along-strike.

The Sierras Pampeanas have been proposed as uplifted by deep high-angle reverse faults, which correspond to the inversion of former Triassic and Cretaceous extensional faults (Jordan and Allmendinger, 1986; Introcaso et al., 1992; Ramos et al., 2002). The extensional faults and associated rift basins are related with the opening of the South Atlantic Ocean (Uliana et al., 1989; Rossello and Mozetic, 1999). According to Ramos et al. (2002), the master extensional faults were generated by reactivation of the sutures between accreted terranes, resulting in formation of continental rift basins, usually with half-graben geometries and most of the deformation taking place in hangingwall blocks. Recent seismological results have imaged some decollement mid-crustal levels in the SVF region (Gallardo, 2011; Ammirati et al., 2015). Thus, the Sierra de Valle Fértil fault in the WSP and the related Ischigualasto-Villa Unión and Marayes eastern basins could be good candidates for accommodating actual deformation (Fig. 1) (Criado Roque et al., 1981; Aceñolaza and Toselli, 1988; Ramos et al., 2002).

FIGURE 2

Analytical Methodology

Apatite (U-Th-Sm)/He thermochronology

(U-Th-Sm)/He thermochronology is based on the accumulation of ^4He produced during the radioactive decay of ^{238}U , ^{235}U , and ^{232}Th in apatite crystals (Zeitler et al., 1987; Wolf et al., 1996; Farley, 2002). ^4He gas is highly mobile at temperatures above 80°C and, due to its mobility, is lost through diffusion out of the crystal lattice. However, below 55°C , apatite begins to retain daughter ^4He and consequently it is accumulated in the crystal lattice during rock exhumation to the surface. This range of temperature, where ^4He is partially retained, is known as the Partial Retention Zone (PRZ) (Wolf et al., 1996; Farley, 2002; Reiners et al., 2004). Conventionally, He diffusion in apatite is assumed to be sensitive to temperatures from $\sim 70\text{--}30^\circ\text{C}$ (Wolf et al., 1998). However, recent studies demonstrated that He retentivity increases with the accumulation of He and associated radiation damage in the apatite crystal, such that an apatite's effective closure temperature evolves through time (Farley, 2000). Forward models predict that for some thermal histories and apatite suites, the radiation-damage effect will be manifested as a span of (U-Th)/He dates positively correlated with both He concentration [He] and effective U concentration [eU]. The latter parameter [eU] weights the decay of U and Th for their He productivity, and is computed as $([\text{U}] + 0.235 \times [\text{Th}])$. Our data shows a positive correlation between date and [eU] (Fig. 3), which develops because an older apatite with higher [eU] and more radiation damage has a higher closure temperature than a lower [eU] (younger) apatite with less radiation damage, despite experiencing the same thermal history.

Due to the thermal sensitivity (55°C - 80°C) of the apatite (U-Th-Sm)/He system, subsequent heating due to burial or elevated geothermal gradient can cause He loss and resetting of the sample. This property is particularly useful for sedimentary samples in this study, because the thermochronometric results can be used to evaluate the post-depositional heating of basin infill and subsequent cooling and inferred exhumation during uplift. If a sedimentary sample yields a completely reset AHe date, it means that post-depositional burial heating was sufficient to degas the sample well above the closure temperature, which causes complete resetting of the original cooling signal due to the diffusive loss of He. On the other hand, a sedimentary sample that has undergone only moderate heating to within the temperatures of the PRZ is characterized by a range of apparent dates that are younger than the depositional age. Finally, an unreset sample represents a sedimentary sample that has remained below the PRZ temperature window and, therefore, will preserve information about the cooling of the source rock from which the sediment was eroded.

In all the cases described above, we obtained the thermal history using HeFTy code by Ketchum (2005), which generates both forward and inverse models for low-temperature thermochronometric systems. The forward model allows the user to calculate the expected data distribution for any given thermal history, whereas the inverse model finds the thermal history that best matches the input data. Only samples with acceptable results of helium and effective uranium concentration, as well as, a factor correction ($F_t \geq 4$) were modeled (Table 1). Initially, forward modeling of single grain age was performed. Constraints involved uncorrected age, alpha calculation, radius of the mineral, among other factors. Once thermal history is generated for each grain, they are run synchronously with the purpose to do an inversion for the time-temperature history, thus generating a set of thermal histories that are considered consistent with observations for each sample. This is evaluated using the goodness-of-fit (GOF) parameter between predicted and observed data

distributions (Ketchman, 2005) (Fig. 4). Because no thermal history ever uniquely defines a set of AHe data, the better fitting thermal histories must always be considered in the context of local geological constraints. For that reason user can set initials and final conditions, like constraints on the time and temperature of deposition or crystallization and ending temperature-time (e.g. samples extracted for boreholes have a different temperature constrain from samples at the surface). User can include geologic evidences at the field and other thermochronological data. Furthermore, user can also decide how many paths want to tried before start modeling. The more paths tried, the more accurate the T-t history obtained will be. Finally, the program allows to display the thermal history as paths, path envelopes or constraining points (Fig. 4).

Sample Suite

For this study, samples were collected in a west-east transect of the northern part of SVF in order to assess changes in timing and rate of cooling across the SVF hangingwall block. Although more than ten samples were collected in the field, only few of them have met the quality requirement criteria, i.e. containing large euhedral and inclusion-free apatite grains suitable for analysis. Obtained grain radius range from 32.8 μ m to 55.3 μ m (Table 1 and Fig. 3). Samples consist mainly of sedimentary rocks from the Paleo-Mesozoic Ischigualasto-Villa Unión basin in the north of Sierra de Valle Fértil (Figs. 2 and 5). Sample 12-SVF-02 was obtained in outcrops near the sierra de Valle Fértil thrust fault corresponding to fluvial-deltaic sandstone beds from the Carboniferous Guandacol Formation, while sample 12-SVF-11 was obtained from outcrops of fluvial sandstone strata from the Triassic Ischigualasto Formation, exposed near Ischigualasto basin. We also compiled a synthesis of

existing low-temperature thermochronology data from nearby sampled outcrops by Bense et al. (2013) and Fosdick et al. (2015).

Sample Preparation

Sample preparation and gas extraction were carried out in the Arizona Radiogenic Helium Dating Laboratory (ARHDL) at the University of Arizona (U.S.A.). Apatite grains were isolated using standard magnetic and density separation techniques and were carefully hand-selected under cross-polarized light to avoid U- and Th-rich mineral and fluid inclusions (e.g., zircon, monazite) that may produce errors in the apatite ages due to excess of helium. Images of selected grains were captured by a closed-circuit digital video camera mounted on the microscope and measured using analysis techniques for the purposes of an alpha ejection correction calculation (Farley et al., 1996). Selected crystals (n= 2-3 grains per sample) were degassed under vacuum using laser heating and analyzed for He using a quadrupole mass spectrometer. Following He measurements, the crystals were analyzed for determining U and Th abundances by using inductively coupled plasma-mass spectroscopy (ICP-MS) through isotope dilution following Reiners and Nicolescu (2007)'s methods.

Thermochronologic Data Modeling

Inverse modeling of AHe data from the SVF help us to better understand the Temperature-time (T-t) history of the region. As stated above, AHe ages were modeled using the HeFTy program (Ketcham, 2005), which can calculate solutions for up to seven different thermochronometers simultaneously. The kinetic model used for apatite (U-Th-Sm)/He is the radiation damage accumulation and annealing model (RDAMM) after Flowers et al. (2009). Modeled Temperature-time paths for two samples taken from the north of the SVF are shown in Figure 4. Input data for the modeling were the apatite (U-Th)/He dates, U and

Th concentrations, and effective grain size. We assumed T-t conditions for the modeling on a base of the geological history from field observations and previous ZFT, ZHe, AFT and AHe data. We performed four different models for two key samples from the SVF. The main difference between the models is the used time lag constrain. Thus, for sample 12-SVF-02 (Table 1 and Fig. 2) we have used in the modeling a time period of 350 Ma to present (Fig. 4A), and another younger time period corresponding to the last 20 Myr (Fig. 4C). The same tests were performed for sample 12-SVF-11 (Figs. 4B and 4C).

Two grains from sample 12-SVF-02 from Carboniferous Guandacol Fm. Sandstone were modeled simultaneously at different selected time intervals: an older T-t window (180 – 20°C for 350 – 250 Ma), which includes the sediment deposition; an intermediate T-t window (180 – 60°C for 250 – 100 Ma); and a shorter T-t window (160 – 20°C for 100 – 0 Ma). Modeling of two grains for sample 12-SVF-11 from Triassic Ischigualasto Fm. sandstone followed a similar analysis. In this case we tested three different T-t intervals: an older T-t window of 180 – 20°C for 260 – 200 Ma, which covers its depositional age; an intermediate T-t window of 180 – 40°C for 200 – 60 Ma; and a younger T-t window of 160 – 20°C for 60 – 0 Ma. A final temperature at the surface of 20°C was considered for both sample analyses.

Results

Our results for two samples from the northern region of the Sierra de Valle Fértil using apatite (U-Th-Sm)/He dates (AHe dates) help to complete a low-temperature thermochronologic dataset between 30°S and 31.5°S for this range (Figs. 1-5). Errors are quoted throughout as two standard deviations (2σ) as uncertainties of weighted mean ages. The main difference between AHe ages obtained in this study and those from Bense et al. (2013)'s dataset is that we calculate weighted mean dates from the single-grain dates,

whereas ages from Bense et al. (2013) are a simple average. Thus, our ‘mean dates’ are actually weighted mean dates, so grains with smaller uncertainties (i.e. better measurements) will have higher control over calculated mean. Measured single-grain AHe dates from Samples 12-SVF-02 and 12-SVF-11 (n=2 grains per sample) showed relatively young cooling ages (Table 1), indicating a complete post-depositional resetting prior to Pliocene-Pleistocene exhumation. Westernmost sample 12-SVF-02 yields a weighted mean AHe age of 4.7 ± 0.5 Ma. At the eastern border of the SVF close to Ischigualasto (see location in Figs. 2 and 5), sample 12-SVF-11 yields a weighted mean AHe age of 2.2 ± 0.4 Ma.

Modeled T-t paths for the two samples in this study at northern SVF clearly indicate both samples share a recent thermal history (Figs. 4A and 4B). For this reason, a second modeled T-t history was performed within the last 20 Myr to gain more insight on time of final cooling (Figs. 4C and 4D). Comparatively, modeling history from sample 12-SVF-02 shows more heating in early stages during late Paleozoic – Mesozoic than that one for sample 12-SVF-11; this agrees well with their different burial vertical stratigraphic position caused by deposition of Triassic sediments from the Ischigualasto basin. Thus, the westernmost sample 12-SVF-02 reached deeper depth levels than the easternmost sample 12-SVF-11. At Cenozoic, both samples followed similar periods of monotonic cooling in late Cretaceous – early Cenozoic, although at higher temperatures for sample 12-SVF-11. We note, however, that best fit paths exhibit an abrupt cooling period within the last 2 Myr.

Since AHe dates are substantially younger than depositional ages, we state all AHe dates to have been completely reset, yielding closely clustered dates as seen in Fig. 3 and Table 1.

FIGURE 3

TABLE 1

Previous Sierra de Valle Fértil Thermochronology Observations

Previous low-temperature thermochronology studies in the SVF document a complex thermal history with multiple phases of cooling and possible reheating episodes. Apatite (U-Th-Sm)/He, Zircon (U-Th-Sm)/He (Bense et al., 2013) and apatite fission track (AFT) (Coughlin et al., 1998) results from the central part of the Ordovician crystalline cored basement of the SVF suggest that this region of the Sierras Pampeanas underwent cooling through these closure systems ca. 293-63 Ma, 353-245 Ma and 305-193 Ma, respectively. These pre-Cenozoic thermochronology determinations provide a first-order constraint on the magnitude of cooling and exhumation related to the thermal condition of the samples prior to Andean deformation. These pre-Cenozoic AHe, ZHe and AFT ages may be showing an inherited age that has not been resetting or annealing and hence those rocks remained near surface prior to Andean exhumation. Furthermore, a sample from Carboniferous strata from the northern foothills of the SVF hanging wall showed a Permian ZHe date, suggesting that the basin infill has not been sufficiently buried to reset ZHe ages as occurred for AHe, ZHe and AFT in the central part of the SVF. However latest Miocene –Pliocene cooling and exhumation were showed for this Carboniferous sample through both the AFT and AHe partial retention zones occurring between 5.8 ± 0.8 Ma and 3.8 ± 0.5 Ma, respectively (Fosdick et al., 2015).

FIGURE 4

FIGURE 5

Discussion

The northern termination of the SVF and other minor ranges extending northward have been propose to disrupt the already formed Bermejo foreland basin during the Neogene

forming the current configuration for Bermejo and Ischigualasto-Villa Unión adjacent basins. Malizia et al. (1995), Jordan et al. (2001), among others have roughly constrained the ages of exhumation of these western Pampean Ranges by conducting chronostratigraphic, paleoenvironmental, and tectonics analyses on Neogene sedimentary deposits. The Pliocene AHe ages in this work allow us to better constrain the exhumation history of the SVF Pampean range, which is in good agreement with the hypothesis of the uplift of the Sierras Pampeanas region during slab flattening of the Nazca plate beneath South America. In this view, enhanced coupling between the plates may have (re)activated Pampean reverse faults, causing increasing topography that was subject to erosion and continue active to the present as shown by high level of crustal seismic activity (Alvarado et al., 2005) and neotectonic features (Bastias, 1985; Costa et al., 2000; Siame et al., 2005; Perucca et al., 2015) (Figs. 6 and 7).

In the last decades a wide variety of regional studies carried out in the Sierras Pampeanas region of west-central Argentina allowed to constrain a roughly eastward propagation of increased exhumation within the south-central Andes backarc. Stratigraphic data (Strecker, 1987) and AFT studies (Coughlin et al., 1998) have focused around 28°S and estimate the main uplift of the Sierra de Aconquija between 7.6–6.0 and 4.0–3.4 Ma (Ramos et al., 2002). Southward (29°S), magnetostratigraphic (Tripaldi et al., 2011) and isotopic records (Losada-Calderón et al., 1994; Tosselli, 1996) suggest the beginning of the uplift at Sierra de Famatina (Fig. 2) between 4.5 and 4.2 Ma (Ramos et al., 2002). In the north area of the Sierra de Valle Fértil, foreland basin studies in the adjacent Bermejo and Ischigualasto-Villa Unión basins (Johnson et al., 1986; Malizia et al., 1995) recorded an interruption in the sedimentation of the distal wedge of Bermejo basin at about 4.3 Ma likely associated to the uplift of the SVF and other minor ranges extending northward. Recent studies by Siame et

al. (2014) more to the south in the Western Sierras Pampeanas (31° S), used in-situ cosmogenic radionuclide and geomorphological data from the Sierra de Pie de Palo (see Figs. 1, 2 and 6) to determine the initiation of uplift between 4-6 Ma; the same studies suggest that it continues to present day with a shortening rate of about 4 mm/yr and an uplift rate of 1 mm/yr (Ramos and Vujovich, 2000). All these previous results are in agreement with the broken foreland of Argentina hypothesis proposed for the last 15 Myr (Zapata and Allmendinger, 1996; Ramos and Folguera, 2009), the migration of magmatism in the Andean backarc (Kay and Mpodozis, 2002) and subduction in a more west-east orientation of the JFR in the last 10 Ma (Yañez et al., 2001).

The Precordillera -Western Sierras Pampeanas interaction is well known by being an active seismogenic region in western Argentina. This zone has produced the most devastating earthquakes in Argentina history (Langer and Hartzell, 1996; Alvarado and Beck, 2006). Larger magnitude-sized earthquakes have produced relatively large fault rupture segments in the basement; in addition small to moderate seismicity continuously produces earthquake deformation consistent with an east-west compression stress regime in the Andean backarc (Siame et al., 2005; Monsalvo et al., 2013; Allmendinger and Judge, 2014). Although very sparse geological studies have focused in the Sierra de Valle Fértil, we provide evidence that north and south low elevation edges of the SVF shared not only young AHe ages but also higher levels of seismicity (Fig. 6) and Quaternary alluvial fans being affected by neotectonic features (Fig. 7); hence the importance of conducting thermochronology studies in the topographically subdued but active range terminations, both in the crystalline and sedimentary rocks.

Low-temperature Thermochronology and Tectonic Interpretation

377 Given a geothermal gradient of 20-30°C/km, the closure temperature of ~80°C for (U-Th-
378 Sm)/He system will be located around 2-3 km below surface. Due to the reset ages obtained
379 on AHe data for this and previous studies, we propose that at least 2-3 km of erosion of
380 Neogene sediments north and south of the SVF is clearly associated with Cenozoic Andean
381 exhumation, assuming monotonic cooling. In correspondence with our determinations,
382 Jordan et al. (2001), making use of compiled data from well sections and industry seismic
383 lines, inferred the northern termination of SVF being buried below >3 km of Neogene
384 sediments during the formation of the Bermejo foreland basin between 20-2.8 Ma, i.e.
385 before to be divided by the north SVF and other minor northern Pampean range uplifts (e.g.,
386 Johnson et al., 1986; Reynolds, 1987; Malizia et al., 1995).

387 For samples that yield older AHe ages at the central region of the SVF (Bense et al., 2013),
388 we hypothesize that they could represent (a) areas of only partial resetting during protracted
389 residence in AHe PRZ prior to the onset of cooling, and/or (b) these regions were remnant
390 positive relief prior to Cenozoic Andean tectonics.

391 (a) Structurally, the SVF and northern minor Pampean ranges shared the feature of the
392 doubly plunging asymmetrical folded terminations. In addition, besides sharing an analogue
393 basement-involved deformation, the SVF in the Sierras Pampeanas also shares a similar
394 pattern of deformation observed for some ranges in the Rocky Mountains of North America.
395 Crowley et al. (2002) carried out apatite (U-Th)/He thermochronology studies in the
396 Bighorn Mountains range, and found old pre-Laramide AHe ages in the center of the range
397 that decrease toward the edges. In their investigations a structural model for the range
398 involves its basement folded along-strike and parts of the range bearing the pre-Laramide
399 AHe remained near surface and below apatite (U-Th)/He closure temperature (~70 °C) prior
400 to Laramide exhumation; whereas the edges remained above apatite PRZ (~85 °C), resetting

the AHe system and showing Laramide ages. It is worth to note that this antiform-like style of deformation is shared among the Western Sierras Pampeanas of San Juan and La Rioja province. As the antiform grows and is uplifted, the exhumation begins at the center first and then progressively exhumes toward its edges. According to this pattern of exhumation the range will show a progressive age decreasing from the oldest ages in the center to the youngest ages toward the edges. The hypothesis of a long permanence in the PRZ is reliable but is hard to prove it with scarce AHe data.

(b) The other hypothesis that the Sierras Pampeanas and other regions of South America have developed a positive relief during most of the Phanerozoic time has attracted attention in the last decades. Evidence for Gondwana paleosurfaces has been proposed on a base of multiple studies as development of deep weathering soils profiles and formation of etchplains (Rabassa et al., 2010), recognition of uncovered relict landscapes and planation surfaces (Rabassa et al., 2005), low-temperature thermochronology (Dávila and Carter, 2013; Löbens et al., 2013; Bense et al., 2013), among others. The creation of this paleorelief has been estimated between the middle Jurassic and the Paleogene. Due to Neogene sediments are not found in the central body of the SVF range, it is possible that this central peak area may be acting as a sediment source for the adjacent basins by the time of Bermejo foreland basin was under formation. The fact that Pliocene AHe ages are found in the northern and southern terminations of the SVF (Bense et al., 2013 and this study), as well as in the southern edge of the Sierra de Pie de Palo (Löbens et al., 2013) provide evidence for the influence of Andean tectonics and exhumation at the edges of these western Pampean Ranges.

Seismicity and New Hints on the West Piedmont of the Sierra de Valle Fértil Deformation

The Western Sierras Pampeanas represent one of the most seismically active regions in Argentina, generating damaging shallow crustal earthquakes that have caused deaths and serious economic consequences in Argentinean history (INPRES, 2015). The latest large earthquake has occurred in the Sierra de Pie de Palo eastern flank near Caucete, San Juan on 23 November 1977, killing 63 people (INPRES, 2015; Figs. 2 and 6). Teleseismic waveform modeling from this (Mw 7.5) earthquake showed its seismic source was composed of two shocks rupturing multiple faults and basement blocks, and separated by a horizontal distance of about ~65 km and 20.8 seconds (Kadinsky-Cade, 1985; Langer and Hartzell, 1996). In addition, those studies illustrate how their aftershocks and associated coseismic deformation propagated across a broad area involving the Sierra de La Huerta, which is the southern prolongation of the SVF (Figs. 1 and 6). Estimation of displacements after the 1977 earthquake using precise leveling data indicated not only the Sierra de Pie de Palo but also the Sierra de La Huerta itself located more to the east is responsible for about 0.5 m of slip accommodation on a high-angle (~65°), shallow (~6 km) east dipping fault (Kadinsky-Cade et al., 1985; Langer and Bollinger, 1988).

Modern seismicity studied by local experiments using local temporary and permanent seismic stations in the region shows a higher concentration of relocated epicenters in the north and south SVF crust (Venerdini, 2014). Seismic moment tensor inversion solutions for two moderate crustal earthquakes in 2002 shown in Fig. 6 using complete broadband three-component regional waveform modeling recorded by the CHARGE experiment (Beck et al., 2001) indicate magnitudes and focal depths of Mw=4.2; h=25 km for Event 1-2002 in the north SVF, and Mw=5.1; h=21 km for Event 2-2002 in the SVF (31°S), respectively (Alvarado et al. 2005). Interestingly, their focal mechanism solutions and focal depths agree with seismically active mid-crustal levels within a ~50-km-thick crust constrained by

receiver functions, surface wave dispersion (Alvarado et al., 2009; Gans et al., 2011) and oil seismic reflection analyses (Zapata and Allmendinger, 1996). Total crustal thickness varies from about 50 km in the WSP to 35 km to the east of SVF (Ammirati et al., 2015) and several basement structures near the SVF fault like the buried north Pie de Palo anticline (Zapata, 1998) (Fig. 6) seem to be seismically active. In addition, two discontinuities in seismic velocities have been observed at ~12 km and ~28 km depths beneath the Sierra de Valle Fértil (Gallardo; 2011). Bouguer anomaly studies in the Bermejo basin and SVF by Lince et al. (2008) are consistent with previous structural models for the SVF proposed by Rosello et al. (1996) illustrating a flower structure with a detachment level located around 22-km depth. In their interpretation, this structure likely inherited from Triassic rifting in the region could have been reactivated later by the Andean compression (Ramos, 1994; Ramos and Folguera, 2009).

Comparison of GPS velocity vectors in the region clearly shows an important decay from west to east across the north SVF fault (Brooks et al., 2003) (Fig 6). Horizontal shortening of about 86 km in the past 13 m.y. as well as thickening seem to be occurring beneath the thin-skinned Precordillera at ~30°S. Average strain rate from Miocene to Holocene fault-slip data used in balanced cross-sections is indistinguishable from GPS strain rate (Allmendinger and Judge, 2014). Observation from those studies and Fosdick et al. (2015) using independent data indicate a progressive eastward migration of the thrust front through time as earlier proposed by Jordan et al. (1993) and Brooks et al. (2003) (Fig. 6).

FIGURE 6

Quaternary shortening in northern Sierra de Valle Fértil produced reverse-displacement reactivation of Mesozoic normal faults and structural inversion of Ischigualasto basin strata (Zapata and Allmendinger, 1996). Carboniferous and Triassic rocks are exposed in the

hangingwall of the Sierra de Valle Fértil Fault interpreted as reverse reactivated Triassic normal faults (Milana and Alcober 1994; Zapata and Allmendinger 1996). The ~N30°W trending Sierra de Valle Fértil Fault system seem to have controlled the continuous basement uplift of the Sierra de Valle Fértil-La Huerta and other ranges along 600 km in the Western Sierras Pampeanas in the modern Pampean flat slab segment of the Andes (Snyder et al., 1990).

Neotectonic structures and Quaternary active faulting are shown along the northern and southern western-bounding fault of the Sierra de Valle Fértil-La Huerta (Perucca and Vargas, 2014; Perucca et al., 2015). Specifically, at the northwestern SVF piedmont a variety of discontinuous parallel or suparallel to the mountain front fault traces, are observed disrupting Quaternary alluvial fans with scarps facing west (Figure 7). The fault scarp heights vary along the fault system between few centimeters to ~10 m, showing the highest elevations in its central section.

The best exposed geomorphic evidence and most recurrent expressions of Quaternary tectonic activity from northern SVF are anomalies in drainage patterns, scarps and alluvial terraces, which are entirely located in the hanging block (Fig. 7). The SVF fault system traces exhibit a trend perpendicular to the streams coming from the mountain, so incision in the hanging block has occurred in response to recent uplift (Fig. 7b). Besides, terraces successively increase in height above present channels (Fig. 7 g). Fig. 7e shows at least five subparallel scarps, characteristic of thrust where slip is distributed on multiple imbricate faults (McCalpin, 2009). In the southern prolongation of the SVF, the Holocene Las Chacras active fault strongly affects the western piedmont of this range (Rothlis et al., 2012) (Fig. 6). Its fault traces can be followed for a distance of about 40 km. Taken together, all the observations described above suggest that current deformation on the SVF and, at least during the last 5 Ma, is accommodated along its northern and southern terminations where

we see a conjunction of young AHe (this study; Bense et al 2013), bracketed seismicity (Venerdini, 2014) and neotectonic features affecting Quaternary piedmonts (Perucca et al. 2015) in both sectors.

FIGURE 7

Conclusions

We present in this work low thermochronological dating from northern SVF that allows us to complete AHe dataset for this western Pampean range. Comparison with existing data enabled us to constrain the tectonic evolution of this region within the Sierras Pampeanas of west-central Argentina. We obtained Pliocene AHe ages providing evidence for the influence of Neogene exhumation of the region at about 5 Ma. This might be related to migration of the deformation during the slab flattening of Nazca Plate beneath South America. We note that Permian through Cretaceous AHe ages in the central region of SVF (Coughlin et al., 1998; Bense et al., 2013) could represent either a protracted cooling and limited exhumation (less of 1-3km) in the last 20 Myr or that the central part of the range may have had existing topographic relief prior to reverse faulting and flat slab subduction of the Nazca plate during the Andean orogeny. The northern region of the SVF, in contrast, shows young AHe (this study and Fosdick et al., 2015), as well as young AFT ages. Assuming a geothermal gradient of 20°C/km, the Partial Annealing Zone for this last technique is located ~3-4 km below surface. Thus, this would suggest that Andean tectonic exhumed at least 3-4 km of rock at the northern SVF during the last 5-6 Ma.

In a regional context, our results suggest an eastward migration of exhumation and agree with the hypothesis of an original single foreland Bermejo basin that was later divided into two basins on this northern area by exhumation and uplift of the SVF around 3.8 - 2.6 Ma (Jordan et al., 1988; Cardozo and Jordan, 2001). Finally, a comparison of our results with

geodetic, geophysical and neotectonic observations indicates that both terminations, at the north and south of the Sierra de Valle Fértil, are seismically active in agreement with regions of Pliocene cooling ages. However, further determinations are needed to constrain a more complete thermochronological record of key regions all over the SVF and involving AFT, ZHe and AHe observations before we can assess a pattern of deformation within this Pampean range.

Acknowledgments

This research was supported by the Agencia Nacional de Promoción Científica y Tecnológica of Argentina (PICT2011-160), the Secretaría de Ciencia, Estado, Tecnología e Innovación (SECITI) de San Juan and the U. S. National Science Foundation (grant EAR-1049605). We gratefully acknowledge the analytical support from the laboratory personnel at the Arizona Radiogenic Helium Dating Laboratory at the University of Arizona. Critical reviews from two anonymous reviewers helped to improve this paper.

References

- Aceñolaza, G., Toselli, A.J., 1988. El Sistema de Famatina, Argentina: su interpretación como orógeno de margen continental activo. In: V Chilean Geological Congress, 1, 55-67.
- Alvarado, P., Beck, S., Zandt, G., Araujo, M., Triep, E., 2005, Crustal deformation in the south-central Andes backarc terranes as viewed from regional broad-band seismic waveform modeling, *Geophysical Journal International*, 163, 580–598.
- Alvarado, P., Pardo, M., Gilbert, H., Miranda, S., Anderson, M., Saez, M., Beck, S. 2009. Flat-slab subduction and crustal models for the seismically active Sierras Pampeanas region of Argentina. In *Backbone of the Americas: Shallow Subduction, Plateau*

uplift, and Ridge and Terrane Collision, Geological Society of America Memoir
204, 261-278.

Ammirati, J.B., Alvarado, P., Beck, S., 2015. A lithospheric velocity model for the flat slab
region of Argentina from joint inversion of Rayleigh wave phase velocity dispersion
and teleseismic receiver functions. *Geophysical Journal International*, 202, 224-241.

Anderson, M., Alvarado, P., Zandt, G., Beck, S., 2007. Geometry and brittle deformation of
the subducting Nazca Plate, Central Chile and Argentina, *Geophys. J. Int.* (2007)
171, 419–434.

Astini, R.A., 2003, The Ordovician proto-Andean basins, *in* Benedetto, J.L., ed., Ordovician
Fossils of Argentina: Córdoba, Secretaría de Ciencia y Tecnología, Universidad
Nacional de Córdoba, 1–74.

Beck, S., Zandt, G., Wallace, T., Anderson, M., Fromm, R., Shearer, T., Wagner, L., Koper,
K., Alvarado, P., Triep, E., Lince Klinger, F., Araujo, M., Bufaliza, M., Campos, J.,
Kausel, E., Ruiz Parades, J., 2001. CHARGE, The CHile ARgentinian Geophysical
Experiment: Imaging the south central Andean lithosphere using passive broadband
seismology. American Geophysical Union, Fall meeting 2001, abstract #T31A-
0828.

Bense, F. A., Löbens, S., Dunkl, I., Wemmer, K., Siegesmund, S. 2013. Is the exhumation
of the Sierras Pampeanas only related to Neogene flat-slab subduction? Implications
from a multi-thermochronological approach. *Journal of South American Earth
Sciences*, 48, 123-144

Braun, J., van der Beek, P., Valla, P., Robert, X., Herman, F., Glotzbach, C., Pedersen, V.,
Perry, C., Simon-Labric, T., Prigent, C., 2012. Quantifying rates of landscape

567 evolution and tectonic processes by thermochronology and numerical modeling of
568 crustal heat transport using PECUBE: *Tectonophysics*, 524-525, 1–28.

569 Brooks, B.A., Bevis, M., Smalley, R.J., Kendrick, E., Manceda, R., Lauría, E., Maturana,
570 R., Araujo, M., 2003, Crustal motion in the Southern Andes (26°–36°S): Do the
571 Andes behave like a microplate? *Geochemistry Geophysics Geosystems*, 4, 1–14.

572 Caminos, R., 1979, Cordillera Frontal: Córdoba, *in* Castellanos, T.G., Sersic, J.L.,
573 Amuchastegui, S., Caputto, R., Cocucci, A.E., Fuchs, G.L., Gordillo, C.E., and
574 Melo, C.R., eds., 2nd Simposio de Geología Regional Argentina: Córdoba,
575 Argentina, Academia Nacional de Ciencias, 1, 394–453.

576 Cardozo, N., Jordan, T., 2001. Causes of spatially variable tectonic subsidence in the
577 Miocene Bermejo foreland basin, Argentina. *Basin Research* 13, 335-357.

578 Castro de Machuca, B., Arancibia, G., Morata, D., Belmar, M., Previley, L., Pontoriero, S.,
579 2008. P-T-t evolution of an Early Silurian medium-grade shear zone on the west side
580 of the Famatinian magmatic arc, Argentina: implications for the assembly of the
581 Western Gondwana margin. *Gondwana Research* 13, 216–226.

582 Coughlin, T.J., P.B. O’Sullivan, B.P. Kohn, R.J. Holcombe. 1998. Apatite fission-track
583 thermochronology of the Sierras Pampeanas, central western Argentina:
584 Implications for the mechanisms of plateau uplift in the Andes: *Geology*, 26, 999-
585 1002.

586 Costa, C., Machette, M. N., Dart, R., Bastias, H., Paredes, J., Perucca, L., Tello, G. and
587 Haller, K., 2000. Map and Database of Quaternary Faults and Folds in Argentina.
588 USGS, Open-File Report 00-0108

589 Criado Roque, P., Mombrú, C., Ramos, V.A., Yrigoyen, M.R., 1981. Estructura e
 590 interpretación tectónica. In: VIII Argentinean Geological Congress, San Luis,
 591 Argentina, 155-192.

592 Crowley, P., Reiners, P., Reuter, J and Kaye, G., 2002. Laramide exhumation of the
 593 Bighorn Mountains, Wyoming: An apatite (U-Th)/He thermochronology study.
 594 Geology, 30, 27-30.

595 Dávila, F.M., Carter, A., 2013. Exhumation history of the Andean broken foreland revisited.
 596 Geology 41 (4), 443-446.

597 Farley KA. 2002. (U-Th)/He dating: techniques, calibrations, and applications. In Noble
 598 Gases in Geochemistry and Cosmochemistry, Reviews in Mineralogy and
 599 Geochemistry, 47, 819–44.

600 Farley, K.A., Wolf, R.A., 1996. The effects of long alpha-stopping distances on (Ue Th)/He
 601 ages. Geochimica et Cosmochimica Acta 60 (21), 4223-4229.

602 Flowers, R.M., Ketcham, R.A., Shuster, D.L., Farley, K.A., 2009, Apatite (U–Th)/He
 603 thermochronometry using a radiation damage accumulation and annealing model:
 604 Geochmica et Cosmochimica Acta, 73(8), 2347–2365, doi:
 605 10.1016/j.gca.2009.01.015.

606 Fosdick, J. C., Carrapa, B., Ortiz, G., 2015. Diachronous faulting and erosion in the
 607 Argentine Precordillera: reconciling bedrock cooling and basin provenance records
 608 of deformation during changes in subduction regime. Earth and Planetary Science
 609 Letters. (in revision).

610 Fosdick, J. C. and Carrapa, B., 2012. Paleogene initiation of Andean foreland sedimentation
 611 in the Bermejo basin and Precordillera thrust belt of NW Argentina from detrital
 612 geochronology and thermochronology: Geological Society of America. Abstracts
 613 with Programs, 44, 7, 72.

614 Gallardo, G., 2011. Determinación de la estructura de velocidades sísmicas a partir de la
 615 función del receptor en el flanco oriental de las Sierras de Valle Fértil–La Huerta,
 616 Final Thesis Undergraduate Work in Geophysics. Universidad Nacional de San
 617 Juan, Argentina. 86 p.

618 Gans, C.R., Beck, S.L., Zandt, G., Gilbert, H., Alvarado, P., Anderson, M., Linkimer, L.,
 619 2011. Continental and oceanic crustal structure of the Pampean flat slab region,
 620 western Argentina, using receiver function analysis: new high-resolution results,
 621 *Geophys. J. Int.*, 186

622 Gonzalez Bonorino, F., 1950. Algunos problemas geológicos de las Sierras Pampeanas,
 623 Revista de la Asociación Geológica Argentina, 5, 81-110.

624 Gordillo, C.E. Lencinas, A., 1981. Geocronología y petrografía de las vulcanitas terciarias
 625 del Departamento de Pocho. Provincia de Córdoba. Revista de la Asociación
 626 Geológica Argentina, 36, 380–388.

627 INPRES, Instituto Nacional de Prevención Sísmica, 2015. Argentina. Earthquake catalog
 628 (www.inpres.gov.ar).

629 Introcaso, A., Pacino, M.C., Fraga, H., 1992. Gravity, isostasy and Andean crustal
 630 shortening between latitudes 30 and 35°S, *Tectonophysics*, 205, 31–48

631 Johnson, N. M., T. E. Jordan, P. A. Johnson, Naeser, C.W., 1986, Magnetic polarity
 632 stratigraphy, age and tectonic setting of fluvial sediments in an eastern Andean
 633 foreland basin, San Juan Province, Argentina, in P. Allen and P. Homewood, eds.,
 634 Foreland basins: International Association of Sedimentologists, Special Publication
 635 8, 63–75.

636 Jordan, T. E., Isacks, B., Ramos, V. A. Allmendinger, R. W., 1983. Mountain building in
 637 the Central Andes. Episodes 1983(3): 20-26, Ottawa, USA.

638 Jordan, T.E., Isacks, B.L., Allmendinger, R.W., Brewer, J.A., Ramos, V.A., Ando, C.J.,
 639 1983, Andean tectonics related to geometry of subducted Nazca plate: Geological
 640 Society of America Bulletin, 94, 341–361.

641 Jordan, T., Allmendinger, R., 1986. The Sierras Pampeanas of Argentina: a modern
 642 analogue of Laramide deformation. American Journal of Science, 286, 737-764.

643 Jordan, T.E., Allmendinger, R.W., Damanti, J.F., and Drake, R., 1993, Chronology of
 644 motion in a complete thrust belt: The Precordillera, 30–31°S, Andes Mountains:
 645 Journal of Geology, v. 101, p. 135–156, doi: 10.1086/648213 .

646 Jordan, T.E., Schlunegger, F., Cardozo, N., 2001. Unsteady and spatially variable evolution
 647 of the Neogene Andean Bermejo foreland basin, Argentina: Journal of South
 648 American Earth Sciences, v. 14, p. 775– 798, doi: 10.1016/S0895-9811(01)00072
 649 -4.

650 Kadinsky-Cade, K., R. Reilinger, Isacks, B., 1985. Surface deformations associated with the
 651 November 23, 1977, Cauçete, Argentina, earthquake sequence, J. Geophys. Res., 90,
 652 12,691– 12,700.

653 Kay, S. M., Gordillo, E., 1994. Pocho volcanic rocks and the melting of depleted
 654 continental lithosphere above a shallowly dipping subduction zone in the Central
 655 Andes. *Contributions to Mineralogy and Petrology*, 117, 25-44.

656 Kay, S.M., Mpodozis, C. 2002. Magmatism as a probe to the Neogene shallowing of the
 657 Nazca plate beneath the modern Chilean flat-slab: *Journal of South American Earth*
 658 *Sciences*, 15, 39–57, doi: 10.1016/S0895-9811(02)00005-6.

659 Ketcham, R.A., 2005. Forward and inverse modeling of low-temperature
 660 thermochronometry data. *Reviews in Mineralogy and Geochemistry* 58, 275-314.

661 Langer, C.J., Bollinger, G.A., 1988. Aftershocks of the western Argentina (Caucete)
 662 earthquake of 23 November 1977: some tectonic implications, *Tectonophysics*, 148,
 663 131–146.

664 Langer, C.J., Hartzell, S., 1996. Rupture distribution of the 1977 western Argentina
 665 earthquake, *Physics of the Earth and Planetary Interiors*, 94, 121–132.

666 Lince, F., Gimenez, M., Martinez, P., Introcaso, A., 2008. Las estructuras de la cuenca de
 667 Bermejo y Sierra de Valle Fértil a partir de los métodos Deconvolución de Euler y
 668 Señal Analítica. *Revista de la Asociación Geológica Argentina*, 63, 281-287.

669 Llambias E. J., Brogioni N., 1981. Magmatismo Mesozoico y Cenozoico. In: *Geología de*
 670 *La Provincia de San Luis. VIII Argentinean Geological Congress. Relatorio, San*
 671 *Luis*, pp 101-115.

672 Löbens, S., Bense, F.A., Wemmer, K., Dunkl, I., Costa, C.H., Layer, P., Siegesmund, S.,
 673 2011. Exhumation and uplift of the Sierras Pampeanas: preliminary implications

674 from KeAr fault gouge dating and low-T thermochronology in the Sierra de
675 Comechingones (Argentina). *Int. J. Earth Sci.* 100 (2-3), 671-694.

676 Löbens, S., Sobel, E.R., Bense, F.A., Wemmer, K., Dunkl, I., Siegesmund, S., 2013.
677 Refined thermochronological aspects of the Northern Sierras Pampeanas. *Tectonics*
678 32 (3), 453-472.

679 Losada-Calderón, A., McBride, S., Bloom, M. 1994. The geology and $^{40}\text{Ar}/^{39}\text{Ar}$
680 geochronology of magmatic activity and related mineralization in the Nevados del
681 Famatina Mining District, La Rioja province, Argentina: *Journal of South American*
682 *Earth Sciences*, 7 (1), 9-24.

683 Malizia, D. C., Reynolds, J. H., Tabbutt, K., 1995. Cronología de la sedimentación
684 Neógena, tectonismo y edad de la estructuración en el Campo de Talampaya, Sierras
685 Pampeanas, Provincia de La Rioja, Argentina, YPF, BIP Junio, p. 78-105.

686 Mc Calpin, J. P. 1996. *Paleoseismology*, Academic Press Inc., 587 p., California. USA.

687 Milana, J. P., Alcober, O., 1994. Modelo tectosedimentario de la cuenca Triásica de
688 Ischigualasto (San Juan, Argentina). *Asociación Geológica Argentina* 49 (3-4), 217-
689 235.

690 Miller, H., Söllner, F., 2005. The Famatinian complex (NW Argentina): back-docking of an
691 island arc or terrane accretion? Early Palaeozoic geodynamics at the western
692 Gondwana margin. *Terrane Processes at the Margins of Gondwana: Geological*
693 *Society, London, Special Publications*, 246, 241–256.

694 Monsalvo, V.G., Alvarado, P., Saez, M., Linkimer, L., Bilbao, I., 2014. Deformación
695 sísmica reciente de la sierra de Pie de Palo, Provincia de San Juan, Revista de la
696 Asociación Geológica Argentina 71(2): 260-266.

697 Otamendi, J.E., Vujovich, G.I., de la Rosa, J.D., Tibaldi, A.M., Castro, A., Martino, R.D. y
698 Pinotti, L.P. 2009. Geology and petrology of a deep crustal zone from the
699 Famatinian paleoarc, Sierras de Valle Fértil and La Huerta, San Juan, Argentina.
700 Journal of South American Earth Sciences, 27, 258-279.

701 Perucca, L. P., N. Vargas, 2014. Neotectónica de la provincia de San Juan, centro-oeste de
702 Argentina, *Bol. de la Soc. Geol. Mexicana*, 66(2), 291–304.

703 Perucca, L. Espejo, K., Esper Angillieri, M., Vargas, M. 2015. Neotectonic controls on the
704 evolution of a river catchment: a case study in the Agua de la Peña river basin,
705 Western Pampean Ranges, Argentina. Geomorphology. In press

706 Ramos, V.A., 1994. Terranes of southern Gondwanaland and their control in the Andean
707 structure (30-33°S lat.). In: Reutter, K.J, Scheuber, E., Wigger, P.J. (Eds.). Tectonics
708 of the Southern Central Andes, Structure and Evolution of an Active Continental
709 Margin. Springer, Berlin, Germany p. 249-261.

710 Ramos, V. A., 2000. Evolución tectónica de la Argentina. In: Caminos, R. (Ed.). Geología
711 Argentina. Instituto de Geología y Recursos Minerales, Anales 29 (24), 715-784.

712 Ramos, V., and G. Vujovich, (2000), Hoja Geológica San Juan, escala 1:250.000, *Tech.*
713 *Rep.*, Servicio GeológicoMinero Argentino Boletín, BuenosAires.

714 Ramos, V.A., Cristallini, E.O., Pérez, D.J., 2002. The Pampean flat-slab of the central
715 Andes. Journal of South American Earth Sciences 15 (1), 59–78.

716 Ramos, V.A., Folguera, A., 2009. Andean flat-slab subduction through time. Geological
717 Society, London, Special Publications 327, 31–54.

718 Ramos, V.A., Vujovich, G., Martino, R., Otamendi, J., 2010. Pampia: a large cratonic block
719 missing in the Rodinia supercontinent. *Journal of Geodynamics* 50, 243–255.

720 Rapela, C.W., Pankhurst, R.J., Casquet, C., Fanning, C.M., Baldo, E.G., González-Casado,
721 J.M., Galindo, C., Dahlquist, J., 2007. The Río de la Plata Craton and the assembly
722 of SW Gondwana. *Earth-Science Review*, 83, 49-82.

723 Regnier, M., Chiu, J.M., Smalley, R., Jr., Isacks, B.L., Araujo, M., 1994, Crustal thickness
724 variation in the Andean foreland, Argentina, from converted waves: *Bulletin of the*
725 *Seismological Society of America*, 84, 1097–1111.

726 Reiners P.W., Spell T.L., Nicolescu S., Zanetti K.A., 2004. Zircon (U-Th)/He
727 thermochronometry: He diffusion and comparisons with $^{40}\text{Ar}/^{39}\text{Ar}$ dating.
728 *Geochim. Cosmochim. Acta* 68, 1857–87.

729 Reiners, P.W., Nicolescu, S., 2007, ARHDL Report 1: Measurement, no. December 2006.

730 Reynolds, J., 1987. Chronology of Neogene tectonics in the Central Andes (27°-33° S) of
731 western Argentina based on the magnetic polarity stratigraphy of foreland basin
732 sediments. Ph.D. Dissertation, Dartmouth College, 353 pp., (unpublished), New
733 Hampshire, USA.

734 Rosello, E., Mozetic, M., Cobbold, P., Marc de Urreiztieta, Gapais, D. and López-Gamundí,
735 O. 1996. The Valle Fértil flower structure and its relationship with the Precordillera
736 and Pampean Ranges (30-32°S, Argentina). *Third ISAG* 9, 481-484.

737 Rossello, E., Mozetic, M.E., 1999. Caracterización estructural y significado geotectónico de
 738 los depocentros cretácicos continentales del centro-oeste Argentino. 5° Simposio
 739 sobre o Cretáceo do Brasil, Serra Negra. Boletín 107–113.

740 Rothlis, M., Santi Malnis P., Pantano, A., Perucca, L., 2012. Evidencias de reactivación
 741 cuaternaria de la falla maestra de la cuenca triásica de Marayes, al norte del cerro
 742 Pan de Azúcar, Sierras Pampeanas occidentales, San Juan, Argentina. p-131. XV
 743 Reunión de Tectónica y IV Reunión de Campo. San Juan. Argentina.

744 Siame, L. L., O. Bellier, M. Sébrier, Araujo, M., 2005, Deformation partitioning in flat
 745 subduction setting: Case of the Andean foreland of western Argentina (28°S–33°S),
 746 Tectonics, 24(5), 1–24, doi:10.1029/2005TC001787.

747 Snyder, D.B., Ramos, V.A., Allmendinger, R.W., 1990. Thick-skinned deformation
 748 observed on deep seismic reflection profiles in Western Argentina. Tectonics 9, 773-
 749 788.

750 Spalletti, L. A., 1999. Cuencas triásicas del Oeste argentino: origen y evolución. Acta
 751 Geológica Hispánica, 32, 29-50.

752 Steenken, A., López de Luchi, M.G., Siegesmund, S., Wemmer, K., Pawlig, S., 2004.
 753 Crustal provenance and cooling of the basement complexes of the Sierra de San
 754 Luis: an insight into the tectonic history of the proto-Andean margin of Gondwana.
 755 Gondwana Research 7 (4), 1171–1195.

756 Strecker, M. R., 1987. Late Cenozoic landscape development, the Santa María Valley,
 757 Northwest Argentina, PhD. Thesis, pp1-262, Cornell University, Ithaca, N.Y. USA.

758 Tosselli, A. (1996), *Volcanismo Andino*, Aceñolaza, F. G., H. Miller, and A. J. Toselli (eds),
 759 A25 pp., Geología del Sistema de Famatina, Münchner Geologische Heste.
 760 Germany.

761 Tripaldi, A., L. Net, L. Limarino, S.Marensi, G. Re, and A. Caselli, 2011. Paleoambientes
 762 sedimentarios y procedencia de la Formación Vinchina, Mioceno, noroeste de la
 763 prov. de La Rioja, *Revista de la Asociación Geológica. Argentina*, 56(4), 443–465.

764 Uliana, M.A., Biddle, K.T., Cerdan, J., 1989. Mesozoic extension and the formation of
 765 Argentine sedimentary basins. AAPG Memoir 46, 599–614.

766 Venerdini, A., 2014. Modelo cortical de velocidades sísmicas para la región comprendida
 767 entre 30,0°S - 32,5°S y 67,0°O – 68,5°O. Final Thesis Undergraduate Work in
 768 Geophysics Universidad Nacional de San Juan, Argentina. 110 p.

769 Vita-Finzi, C., 2009. Pie de Palo, Argentina: A cataclastic diapir, *Geomorphology* 104,
 770 317–322.

771 Wolf, R.A., Farley, K.A., Silver, L.T. 1996. Helium diffusion and low-temperature
 772 thermochronometry of apatite: *Geochimica et Cosmochimica Acta*, 60(21), 4231-
 773 4240.

774 Yañez, G., Ranero, G.R., Huene, R., von, Díaz, J., 2001. Magnetic anomaly interpretation
 775 across a segment of the Southern Central Andes (32-34°S): implications on the role
 776 of the Juan Fernández Ridge in the tectonic evolution of the margin during upper
 777 Tertiary. *J. Geophys. Res.* 106, 6325-6345.

778 Zapata, 1998. Crustal structure of the Andean thrust front at 30 degrees S latitude from
779 shallow and deep seismic reflection profiles, Argentina. Journal of South American
780 Earth Sciences, 11, 131-151.

781 Zapata, T.R., Allmendinger, R.W., 1996, Thrust front zone of the Precordillera, Argentina:
782 A thick-skinned triangle zone: American Association of Petroleum Geologists
783 Bulletin, 80, 359–381.

784 Zeitler PK, Herczeg AL, McDougall I, Honda M. 1987. U-Th-He dating of apatite: a
785 potential thermochronometer. Geochimica et Cosmochimica Acta 51, 2865–68.

786 **Figures Captions**

787 **Figure 1.** Regional map of the Pampean flat-slab subduction zone showing terrane
788 boundaries (dashed lines) and main morphostructural units from Ramos et al. (2002) and
789 Rapela et al. (2007). Red triangles are expiring volcanic centers in the last 5 M.a. (Kay and
790 Gordillo, 1994). Slab contours to the top of the Wadati-Benioff zone from Anderson et al.
791 (2007) and the Juan Fernández Ridge (JFR) path according to Yañez et al. (2001) are also
792 indicated. The red dashed-line rectangle represents approximate location of the study area
793 shown in more detail in Figure 5.

794 **Figure 2. A)** Simplified schematic structural cross section along west-east line X-X' in the
795 northern Sierra de Valle Fértil based on Zapata and Allmendinger (1996); Rosello et al.
796 (1995) and Lince et al. (2008). **B).** Digital elevation map of the Andean retroarc region
797 around the Sierra de Valle Fértil (SVF) in west-central Argentina (Fig. 1) showing highest
798 peak (Mount Famatina) in the western Sierras Pampeanas region, main Andean faults and
799 terrane sutures after Vujovich et al. (2004). Solid black line indicates the location of the
800 transect X-X' along the northern Sierra de Valle Fértil shown in (A). Thick dashed line

corresponds to region of leveling measurements in the epicentral area of the 1977 earthquake (star) by Kadinsky-Cade et al. (1985). Key to symbols: EP: Eastern Precordillera, WSP: Western Sierras Pampeanas, M: Sierra de Maz, U: Sierra de Umango, F: Sierra de Famatina, I-VU: Ischigualasto-Villa Unión basin, B: Bermejo basin, SVF: Sierra de Valle Fértil, PP: Sierra de Pie de Palo (see geological references in the text).

Figure 3. Apatite (U-Th)/He results for Samples 12-SVF-02 (blue) and 12-SVF-11 (red) from this study at the northern Sierra de Valle Fértil (see Fig. 2A for location). (Left panel) Relationships between obtained dates and effective uranium concentration (eU). (Right panel) Same ages versus effective grain radius. Positive correlation between AHe date and effective uranium concentration (eU) probably indicating that burial heating was sufficient to partially reset apatites with low eU values.

Figure 4. Four inverse modeling results for two key samples (A) 12-SVF-02 and (B) 12-SVF-11 from the Sierra de Valle Fértil hanging wall (see location of the samples in Figs. 2 and 5). The results are displayed as Temperature versus time (T-t) envelope paths for good (magenta), acceptable (green) and best fit (solid black line) histories. Models (A) and (B) exhibit attempted 100,000 and 50,000 T-t paths, whereas (C) and (D) represent 50,000 and 10,000 T-t paths, respectively. Dark grey boxes indicate the imposed T-t constraints, which were chosen following geological evidence criteria. For (A) and (B), initial modeled boxes consider depositional ages and surface Temperatures; for (C) and (D), initial modeled boxes consider starting Temperatures ≥ 60 °C due the resetting of the system, and final boxes with present day surface temperatures. See text for more details.

Figure 5. Simplified geological map of the Sierra de Valle Fértil (modified from Otamendi et al., 2009) showing sample locations and AFT, ZHe and AHe ages discussed in the main

text. Errors quoted throughout as two standard deviations (2σ) of weighted mean AHe ages as well as number of analyzed grains (n) are also indicated for samples in the north of SVF.

Figure 6. Local crustal seismicity of magnitude $M \geq 3$ from INPRES (2015) during three years (red dots). Also shown, are focal mechanisms by Alvarado et al. (2005) obtained using broadband seismic waveform modeling; earthquakes are represented by year of occurrence, and magnitude Mw-focal depth in km in parenthesis. The 1977 event corresponds to the large (Mw 7.5) Cauçete crustal earthquake (Langer and Hartzell, 1996). The pentagon near Marayes shows location for accommodation of 0,4 m slip vertical motion along the SVF thrust fault after the 1977 earthquake according to leveling measurements (Kadinsky-Cade et al., 1985). GPS velocity vectors are from Brooks et al. (2003). Abbreviations, other symbols and references as in Figure 3. Location of Figure 7 (red dashed-line rectangle) showing neotectonic observations is also indicated.

Figure 7. Top right figure shows *Google Earth* image of the northern end of the Sierra de Valle Fértil Fault System in the western piedmont of the SVF mountain range as indicated in Fig. 6. A zoom for the area around each pin from northwest to southeast is shown in figures (A) to (G) using white arrows to denote recent scarps affecting alluvial deposits: **A)** Preferential slope in $\sim 330^\circ$ -trending scarps affecting Pleistocene-Holocene alluvial fans; **B)** N-S trending on west-facing scarp located to the south of the Agua de La Peña river. We note that temporary streams coming from the Sierra de Valle Fértil are definitely more incised in the hanging block; **C)** $N10^\circ$ -trending fault scarp affecting Holocene alluvial deposits; **D)** Parallel traces of 170° trending affecting several Quaternary alluvial levels. Unpaired terraces are associated to the hanging block; **E)** Discontinuous trace of a 330° trending preferential-slope scarp; **F)** Fault scarp of N-S trending, showing its free face to the

847 W with the maximum height of the fault system; **G)** Approximate N-S trending scarp
848 affecting at least two Quaternary terraces and alluvial levels.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

TABLES

Table 1. Apatite Thermochronology Data

Sample	Latitude (°S)	Longitude (°W)	Elevation (m)	Grain	Lithologic unit	effective radius (μm)	U (ppm)	Th (ppm)	Sm (ppm)	eU (ppm)	Th/U	He (nmols/g)	Ft	Single grain age (Ma ± 2σ)		Weighted mean age (Ma ± 2σ)	
12-SVF-02	-30,14255	-68,03188	898	Ap1	Carboniferous Paganzo Group Sandstone	35,3	13,79	7,70	52,2	15,6	0,558	0,15	0,61	2,9 ± 0,2		4,7 ± 0,5	
				Ap4		32,8	169,42	246,96	180,0	227,5	1,458	3,78	0,59	5,4 ± 0,1			
12-SVF-11	-30,16310	-67,84159	1374	Ap2	Triassic Ischigualasto Fm. Sandstone	55,3	1,32	6,30	4,3	2,8	4,778	0,02	0,74	2,2 ± 0,2		2,2 ± 0,4	
				Ap3		36,8	10,73	35,55	66,3	19,1	3,314	0,14	0,63	2,2 ± 0,1			