Abstract

The error-related negativity (ERN) is a neuroelectric signature of performance monitoring during speeded response time tasks. Previous studies indicate that individuals with anxiety disorders show ERN enhancements that correlate with the degree of clinical symptomology. Less is known about the error monitoring system in post-traumatic stress disorder (PTSD). PTSD is characterized by impairments in the regulation of fear and other emotional responses, as well as deficits in maintaining cognitive control. Here, combat Veterans with PTSD were compared to control Veterans in two different versions of the flanker task ($n=13$ or 14 per group). Replicating and extending previous findings, PTSD patients showed an intact ERN in both experiments. In addition, task performance and error compensation behavior were intact. Finally, ERN amplitude showed no relationship with self-reported PTSD, depression, or post-concussive symptoms. These results suggest that error monitoring represents a relative strength in PTSD that can dissociate from cognitive control functions that are impaired, such as response inhibition and sustained attention. A healthy awareness of errors in external actions could be leveraged to improve interoceptive awareness of emotional state. The results could have positive implications for PTSD treatments that rely on self-monitoring abilities, such as neurofeedback and mindfulness training.

Keywords: Error-related negativity; Performance monitoring; Self-monitoring; Mild traumatic brain injury; Anterior cingulate cortex; Cognitive control; Anxiety

1. Introduction

The capacity for self-monitoring is critical for effective performance in everyday tasks. If we make mistakes, the ability to detect and correct our errors helps us stay on track. Altered self-monitoring is a core feature of many psychiatric disorders, with deficient monitoring seen in schizophrenia (Frith and Done, 1989; Turken et al., 2003), and excessive monitoring in obsessive compulsive disorder (OCD) (Salkovskis, 1999; Gehring et al., 2000). Numerous studies have implicated the medial frontal cortex (MFC), especially the anterior cingulate cortex (ACC), in this crucial executive control function (Carter et al., 2001; Rushworth et al. 2004; Posner et al. 2007; Ridderinkhof et al. 2007; Ullsperger et al., 2014).

A key brain indicator of performance monitoring is the error-related negativity (ERN), a prominent electrophysiological response that occurs around the time an error is made (Falkenstein et al., 1991; Gehring et al., 1993). This event-related potential (ERP) component peaks about 50-100 ms after an erroneous response, showing maximal amplitude at fronto-central midline scalp electrodes (reviewed in Gehring et al., 2012). It is generally larger when accuracy is emphasized over speed, and when errors carry greater motivational significance. Lesion and dipole modeling studies suggest that the dorsal ACC is a major neural generator of the ERN (Dehaene et al., 1994; Holroyd et al., 1998; Swick and Turken, 2002; Stemmer et al., 2004), with contributions from lateral prefrontal cortex (Gehring and Knight, 2000; Ullsperger et al., 2002) and orbitofrontal cortex (Turken and Swick, 2008; Solbakk et al., 2014) as well.

Parallel to the alterations in self-monitoring behavior, clinical populations can exhibit changes in ERN amplitude. Individuals with schizophrenia show reductions in the ERN (Alain et al., 2002; Mathalon et al., 2002), whereas those with generalized anxiety disorder (GAD) and OCD show increases (Gehring et al., 2000; Endrass and Ullsperger, 2014; Xiao et al., 2011). The

findings in major depressive disorder are mixed, with some studies showing increases in ERN amplitude (Chiu and Deldin, 2007; Holmes and Pizzagalli, 2008; Aarts et al., 2013) and others decreases (Ruchow et al., 2006) or no difference (Compton et al., 2008; Schrijvers et al., 2009). Co-morbid conditions that occur with depression may account for some of the mixed results. For instance, Weinberg et al. (2012) replicated the typical finding of ERN enhancement in GAD, but found that patients with co-morbid anxiety and depression did not differ from controls.

Post-traumatic stress disorder (PTSD) has traditionally been classified as an anxiety disorder that can develop after a traumatic event (DSM-TR-IV: American Psychiatric Association, 2000). According to those criteria, PTSD is characterized by three symptom clusters: (1) re-experiencing of intrusive memories, such as flashbacks and nightmares; (2) avoidance of stimuli associated with the trauma and feelings of emotional numbing; and (3) hyperarousal symptoms such as hypervigilance to surroundings, increased startle, and insomnia.¹ Because of the serious hyperarousal symptoms in particular, one might predict that PTSD would be associated with larger ERNs and an exaggerated sensitivity to errors, accompanied by maladaptive signs of hypervigilance in everyday life. Thus, an important line of inquiry is whether individuals with PTSD show changes in ERN amplitude. This is particularly salient in light of the structural and functional alterations in MFC observed in this disorder (Hamner et al., 1999; Shin et al., 2001; Rauch et al., 2003; Kim et al., 2005).

Two previous studies have addressed whether the electrophysiological indices of error monitoring are altered in PTSD (Clemans et al., 2012; Rabinak et al., 2013). The first paper focused on time-frequency analysis but informally reported an intact ERN in civilians with PTSD (Clemans et al., 2012). The second study found comparable ERNs in healthy civilian controls and combat Veterans with PTSD, strongly suggesting that PTSD does not affect ERN

amplitude (Rabinak et al., 2013). In contrast, combat-exposed Veterans without PTSD showed a drastic reduction in ERN relative to both the PTSD and the civilian control groups (Rabinak et al., 2013). However, the paper did not report whether the Veterans had sustained traumatic brain injuries (TBIs) during their service in Iraq or Afghanistan. Approximately 20% of Operation Enduring Freedom/Operation Iraqi Freedom (OEF/OIF) Veterans have sustained TBIs, the majority of which are mild in severity (Tanielian and Jaycox, 2008; Fischer, 2014). TBI can be associated with reductions in ERN (Larson et al. 2007, 2009), so one cannot exclude the possibility that the combat-exposed control Veterans experienced more severe brain injuries than the Veterans with PTSD.

Three recent studies have looked at the effects of mTBI (or concussion) on the ERN in young athletes, with mixed results (Pontifex et al., 2009; Larson et al., 2012; De Beaumont et al., 2013). Pontifex and colleagues (2009) tested athletes with self-reported mTBI(s) and observed a reduction in ERN peak amplitude at the FCz electrode site. In contrast, Larson et al. (2012) tested a somewhat similar population of young adults with self-reported mTBI(s) due to sports injuries (69%) and other causes. No difference in ERN amplitude was found, even when measured in the same way as in the Pontifex et al. (2009) study. Finally, De Beaumont et al. (2013) tested athletes with two or more mTBIs, who showed ERN reductions in two separate experiments. It's unclear why the results were discrepant in these studies, but differences in the populations tested (including age at injury and time post-injury) and the EEG recording and analysis methods may have contributed.

To our knowledge, no studies have specifically examined the effects of mTBI on the ERN in OEF/OIF Veterans. Deployment-related changes in self-monitoring could hinder optimal transition to civilian life, particularly in military personnel exposed to blasts and other traumatic

events. The brain injuries caused by blast waves may alter brain structure and function differently than impact-related injuries (Fischer et al., 2014), so the effects on the ERN are unknown.

Our previous results indicated this group is impaired on a number of cognitive control tasks, showing more errors of commission in a Go/NoGo task (Swick et al., 2012), more variable RTs (Swick et al., 2013), greater interference in an emotional Stroop task using trauma-related words (Ashley et al., 2013), and decreases in working memory performance and ERPs in a dual-task condition (Honzel et al., 2014). More pronounced deficits in these tasks were associated with greater PTSD symptom severity. Thus, one line of reasoning suggests that PTSD would be associated with impaired error monitoring and decreased ERNs, another sign of difficulties with cognitive control.

Our original prediction for the present study was that Veterans with PTSD would show *increased* ERNs, due to elevated vigilance and hyperarousal. This would agree with findings in other anxiety disorders. However, the majority of patients seen clinically and recruited into the study had both PTSD and mTBI, which is consistent with the observations of other investigators (Carlson et al., 2011; Taylor et al., 2012). Thus, the question is whether combat Veterans with PTSD (most with mTBI as well, but some with PTSD-only) will show deficits in error monitoring as indexed by the ERN.

Another error-related component, the error positivity (Pe), occurs after the ERN (Overbeek et al., 2005). The Pe is observed between 200-400 ms following an error, with a central-parietal scalp topography. The Pe reflects post-error processing of some sort, although the exact nature of the psychological processes is unclear (Falkenstein et al., 2000). The existing literature suggests that neither PTSD nor mTBI affect the Pe component. Rabinak et al. (2013) did not find a

difference in Pe among the combat-exposed PTSD, combat-exposed control, and civilian control groups. Likewise, three separate studies failed to find a difference between mTBI patients and controls (Pontifex et al., 2009; Larson et al., 2012; De Beaumont et al., 2013).

The current study administered two different versions of the Eriksen flanker task (Eriksen and Eriksen, 1974) to Veterans with PTSD $\pm$ mTBI and Veterans without PTSD or mTBI. We initially predicted that PTSD would be associated with an exaggerated sensitivity to errors, as indicated by larger ERNs and maladaptive signs of hypervigilance in everyday life. The co-existence of mTBI could result in either a dampening of the hyperactive ERN (i.e., no net effect) or even a reduction in ERN relative to controls. The recent results of Rabinak et al. (2013) align more with the former prediction: that the ERN would be unchanged in the present population of combat Veterans with PTSD. Given the previous null findings for the Pe (Rabinak et al., 2013), we did not expect to see a difference between groups for this component.

2. Methods

2.1. Participants

Experiment 1: Participants were 18 Iraq and Afghanistan combat Veterans diagnosed with combat-related PTSD (17 male, 1 female) and 16 control Veterans matched in age and gender (15 male, 1 female). None of the enrolled participants reported significant substance abuse or a history of other psychological disorders, excluding depression. Six participants (four in the patient group and two in the control group) were excluded due to incorrect task performance that resulted in an average of 247 errors. These six participants mistakenly responded to the arrow in the center of the flanker array (the relative center) and not to the arrow in the center of the screen, and were therefore excluded. The final analysis yielded $n=14$ for each

group.

All of the trauma-exposed Veterans with PTSD had been deployed to Iraq or Afghanistan. Only five of the control Veterans had been deployed (and trauma-exposed), which is one limitation of the study. Ten out of 14 participants with PTSD reported a history consistent with mild TBI (mTBI); however, none reported a loss of consciousness greater than 1-2 min or any other pre-existing neurological condition. A semi-structured clinical interview was conducted by a neurologist, and mTBI was diagnosed based on patient self-report of the following criteria from the VA/DoD Clinical Practice Guidelines – loss of consciousness (LOC) 30 min or less or altered mental status (e.g., feeling dazed, disoriented, or confused), with post-traumatic amnesia less than 24 hrs (Management of Concussion/mTBI Working Group, 2009). PTSD diagnosis was based on semi-structured clinical interview using DSM-IV criteria. The diagnoses of mTBI and PTSD were corroborated with available VA medical records, to the fullest extent possible.

All participants completed the PTSD Checklist, Military Version (PCL-M) for DSM-IV (Weathers, et al., 1994) and the Beck Depression Inventory (BDI). The PCL-M is a 17-item self-report questionnaire that is used to establish the presence of PTSD symptoms in combat-exposed Veterans (Blanchard, et al., 1996). Symptoms are rated on a 1 to 5 scale. A score of 50 is used as a cutoff for PTSD diagnosis in Veteran populations (Forbes et al., 2001). The BDI is a 21-item test which measures the presence and degree of depression in adolescents and adults (Beck et al., 1988). As expected, the two groups showed highly divergent scores on these questionnaires (Table 1), indicating greater levels of PTSD and depression symptoms in the patients. The groups did not differ significantly in age [$t(26) = 1.44, p=0.16$] or education [$t(26) = 1.59, p=0.12$]. Some of the participants were receiving treatment (medication and/or psychotherapy), but most were not. See Table 1 for details on demographic data.

The Institutional Review Board of the VA Northern California Health Care System approved the experimental protocol, and all participants gave informed consent prior to beginning the experiment. They were paid for transportation plus \$20/h for their participation.

Experiment 2: The participants were 13 Veterans with PTSD and 13 control Veterans drawn from the same population as Experiment 1, using the same selection criteria. Ten patients reported a history of mTBI, diagnosed as reported above. The groups did not differ significantly in age [$t(24) = 0.21, p=0.84$] or education [$t(24) = 1.81, p=0.08$]. Seven patients and nine controls participated in both studies, an average of 4.8 months and 6.2 months apart, respectively. Both experiments were completed before the findings of Rabinak et al. (2013) were published. Full demographic information is available in Table 1.

2.2. Stimuli and tasks

Experiment 1: The study consisted of the single-task condition of an arrow flanker task, described in Honzel et al. (2014). Each flanker stimulus array was presented for 200 ms. Participants responded with a button press to indicate whether the target arrowhead (always in the center of the screen) pointed to the left or the right. Flanking arrowheads could point in either the same (congruent) direction (40% of trials) or the opposite (incongruent) direction (60% of trials). To increase error rates, the array could consist of five vertically arranged arrows (two above and two below) or three vertically arranged arrows (two above, or two below, the central target; illustrated in Pratt et al., 2011). The next trial began 300 to 500 ms after a response was made. If there was no response, the next trial began 900 ms after stimulus onset. Participants were instructed to respond as quickly and accurately as possible. A block of 20 practice trials preceded the main experiment, when subjects completed 10 blocks consisting of 60 trials each.

Experiment 2: The study used a modification of the design used by Kopp et al. (1996). Each trial began with a one second warning period, which consisted of a central fixation cross (a plus sign shown for 1000 ms). Fixation onset coincided with a 100 ms tone (frequency = 1000 Hz), accompanied by a temporary brightening of the stimulus. The fixation cross was removed from the screen 900 ms later. Then the flanker stimuli (two large arrows) appeared above and below the fixation position for 100 ms, in advance of when the smaller central target arrow appeared. The entire flanker display remained on the screen for 50 ms longer (total display time = 150 ms). Half of the flanker stimuli were congruent and half were incongruent with the target, presented in random order.

Each trial was terminated by a button press response indicating the direction of the target arrow. If no response was made, the trial advanced 1500 ms later. At trial termination, a small fixation square remained on the screen for 1888, 2021, 2144, or 2272 ms (randomly jittered), at which time the one second warning period began again. Participants were instructed to respond as quickly as possible and encouraged to correct their errors with a second response. A block of 24 practice trials with performance feedback preceded the main experiment, when subjects completed 10-12 blocks consisting of 53 trials each.

2.3 Procedure

Participants were tested in a darkened, sound-attenuating room while EEG was recorded. In both tasks, participants were instructed to fixate at the center of the screen and to blink as little as possible. In Exp. 1, each participant completed both single and dual task flanker conditions as well as a single task Sternberg condition in one session (Honzel et al., 2014). The task order was counterbalanced across participants. Here we present ERP data from only the single task flanker,

which is most comparable to Exp. 2. Following the Exp. 1 testing session, participants completed the Post-Concussion Symptom Checklist (Gouvier et al., 1992), which asked about the frequency, intensity and duration of specific symptoms for today only. They also completed the Neurobehavioral Symptom Inventory (Cicerone, 1995). This checklist evaluated a variety of items for symptom severity over the last month. The total time required for each experimental session was approximately 2 to 2.5 h. All data were collected between February 2010 and March 2012.

2.4. EEG recording

EEG was recorded from 64 scalp electrodes and from two external electrodes placed on the left and right mastoids using the ActiveTwo Biosemi electrode system (BioSemi, Amsterdam, the Netherlands). Two electrooculogram (EOG) electrodes were placed lateral to and below each eye to record blinks and eye movements. The EEG was sampled at 512 Hz and recorded onto a computer using ActiView software (BioSemi). Off-line analysis was completed using Brain Vision Analyzer software (Brain Products, Gilching, Germany). Data were referenced to the average of the two mastoid electrodes and bandpass filtered from 0.1 to 30 Hz (48 dB/octave). The EEG was segmented for each trial beginning 200 ms before the trial and extending to 900 ms post-response; the ERP averages were time locked to the button response. EEG was corrected for ocular artifacts using the algorithm of Gratton et al. (1983). Trials with other artifacts exceeding 150 microvolts were rejected. If a participant had fewer than 20 error trials in their averaged waveforms, they were excluded from the ERP analysis. All averages were baseline corrected using the -200 ms to 0 ms pre-response interval (our pre-planned parameter settings). We also calculated difference waves (error trials – correct trials), and made topographic maps

based on the difference waves for the ERN and Pe intervals. Finally, we include data averaged and analyzed using alternate pre-response baselines (-200 ms to -100 ms and -500 to -300 ms, the latter as in Rabinak et al., 2013) in the Supplementary Materials.

2.5. Statistical analysis

2.5.1. Behavioral performance: Initial behavioral analyses examined the effect of stimulus congruity on reaction time (RT) and accuracy using 2×2 repeated measures analyses of variance (ANOVAs). The within factor was Congruence (congruent, incongruent) and the between factor was Group (controls, patients). Only correct responses were used in this analysis. The effects of response accuracy on RTs were further examined in a 2 (congruent, incongruent) $\times 2$ (correct, error) $\times 2$ (controls, patients) repeated measures ANOVA. In addition, we were interested in the phenomenon of post-error slowing, where RTs after an incorrect response increase on the next trial (Rabbitt, 1966). The post-error slowing data were submitted to an additional ANOVA with Previous Response (correct, error) $\times$ Group (controls, patients) as factors. Only correct responses were used, i.e., we looked at RTs on correct trials that followed error trials (post-error) vs. correct trials that followed correct trials (post-correct). Follow-up t tests investigated any significant interactions, with Bonferroni correction as appropriate. Some of the behavioral results from Experiment 1 have been reported previously, namely the flanker interference effect (Honzel et al., 2014). However, the error RTs, post-error slowing, and ERP data have not been reported elsewhere. Error correction data were obtained in Exp. 2. The percentage of error trials that were corrected by controls and patients was compared using a t test.

2.5.2. ERPs: Initial repeated measures ANOVA examined the scalp distribution of ERN and Pe at six midline electrodes (Fz, FCz, Cz, CPz, Pz, POz), with additional factors of Response

(correct, error) and Group (controls, patients), using Greenhouse-Geisser correction when appropriate. To replicate Rabinak et al. (2013), the ERN was quantified as mean voltage amplitude between 0-100 ms post-response at electrode FCz. The Pe was measured as mean amplitude between 200-400 ms at Pz.. These values were entered into ANOVAs that examined Response (correct, error) and Group (controls, patients). Follow-up t tests investigated any significant interactions, using Bonferroni correction for multiple comparisons. Spearman correlation analyses were used to test the relationship between ERN amplitude and the severity of self-reported PTSD (PCL-M), depression (BDI), and post-concussion symptoms (PCSC), using appropriate corrections for multiple comparisons.

3. Results

3.1. Behavioral

3.1.1. Experiment 1: There were no significant group differences in RT, accuracy, or post-error slowing (see Table 2).

Reaction time: Participants showed the standard flanker interference effect, and the performance of Veterans with PTSD did not differ from that of controls (also see Honzel et al., 2014). A main effect of Congruence [$F(1,26) = 218.65, p < 0.0001$] indicated that participants were significantly faster in responding to target arrows when the flanking arrows were congruent, compared to incongruent. There were no main [$F(1,26) = 0.003, p = 0.96$] or interactive [$F(1,26) = 0.70, p = 0.41$] effects of Group. An additional ANOVA compared RTs for correct and error responses. Participants were faster on error trials, and this was true to a greater extent in the incongruent condition. This was indicated by a main effect of Response [$F(1,26) = 46.08, p < 0.0001$] and a Response $\times$ Congruence interaction [$F(1,26) = 50.73, p < 0.0001$]. These

effects did not interact further with Group [$F(1,26) = 0.22, p=0.64$; $F(1,26) = 1.50, p=0.23$, respectively].

Accuracy: The same pattern of results was observed for accuracy. A main effect of Congruence [$F(1,26) = 94.51, p<0.0001$] indicated that participants were more accurate when target arrows were congruent with the flankers, compared to incongruent. Again, there were no main [$F(1,26) = 1.41, p=0.25$] or interactive [$F(1,26) = 0.625, p=0.44$] effects of Group.

Post-error slowing: The standard post-error slowing effect was observed. Participants had slower RTs on correct trials immediately following an incorrect response, compared to a correct response, as shown by the main effect of Previous Response [$F(1,26) = 30.52, p<0.0001$]. There were no main [$F(1,26) = 0.025, p=0.88$] or interactive [$F(1,26) = 0.144, p=0.71$] effects of group.

3.1.2. Experiment 2. There were no significant behavioral performance differences between Veterans with PTSD and control Veterans, replicating Exp. 1 (see Table 2).

Reaction time: Participants again showed the standard flanker interference effect. The main effect of Congruence [$F(1,24) = 481.41, p<0.0001$] indicated that subjects were faster in responding to target arrows when the flanking arrows were congruent, compared to incongruent. There were no main [$F(1,24) = 1.48, p=0.24$] or interactive [$F(1,24) = 0.069, p=0.80$] effects of Group. An additional ANOVA compared RTs for correct and error responses. This analysis included only eight patients and seven controls, because the other subjects did not have any errors on congruent trials. For the remaining individuals, RTs were faster on errors, and this was true to a greater extent in the incongruent condition. This was indicated by a main effect of Response [$F(1,13) = 33.88, p<0.0001$] and a Response $\times$ Congruence interaction [$F(1,13) =$

13.91, $p=0.003$]. These effects did not interact further with Group, [$F(1,13) = 0.619, p=0.45$] and [$F(1,13) = 0.005, p=0.95$], respectively. Notably, RTs were over 100 ms faster in this study than in Exp. 1, and error rates on congruent trials were lower (see Table 2). This is because the location of the flankers was unpredictable in Exp. 1, but fixed in Exp. 2.

Accuracy: The same pattern of results was observed for accuracy. A main effect of Congruence [$F(1,24) = 77.93, p<0.0001$] indicated that participants were more accurate when target arrows were congruent with the flankers, relative to incongruent. There were no main [$F(1,24) = 0.001, p=0.98$] or interactive [$F(1,24) = 0.00003, p=0.995$] effects of Group.

Post-Error Slowing: In contrast to Exp. 1, post-error slowing was not observed in this experiment. RTs did not differ for correct trials that were immediately after incorrect, vs. correct, responses [$F(1,24) = 0.75, p=0.40$]. The main effect of group was not significant either [$F(1,24) = 0.88, p=0.36$], but the interaction between Previous Response and Group was significant [$F(1,24) = 5.48, p=0.03$]. This interaction indicated that the pattern of change in RTs after errors, relative to correct trials, differed in controls (6 ms slowing) and patients (13 ms speeding). However, neither of these values was a significant change in RT when examining the within-group comparisons: [$t(12) = -1.42, p=0.18$] for controls and [$t(12) = 1.88, p=0.09$] for patients. The lack of post-error slowing in Exp. 2 agrees with prior studies showing a smaller or non-existent effect when the interval from response to the next stimulus is long (Jentzsch and Dudschig, 2009; Danielmeier and Ullsperger, 2011).

Error correction: Participants in Exp. 2 had the opportunity to correct their errors with a second button press. The rate of error correction was low and variable ($46.3 \pm 38.0\%$ for controls, $28.4 \pm 31.5\%$ for patients), and did not differ between groups [$t(24) = 1.31, p=0.20$].

3.2. ERPs

3.2.1. *Experiment 1.* The amplitude of the ERN and Pe components did not differ between groups (Fig. 1, top). The mean number of error trials in the ERP averages was 69 ± 23 for the patients and 84 ± 37 for the controls, which did not differ significantly [$t(26)=1.36, p=0.19$]. For correct trials, the means were 519 ± 29 for patients and 516 ± 35 for controls [$t(26)=-0.22, p=0.83$].

ERN: The standard ERN effect was observed. Errors elicited a larger negative peak than correct responses, as indicated by the main effect of Response [$F(1,26)=62.53, p<0.0001, \eta_p^2=0.71$]. The Electrode $\times$ Response interaction [$F(5,130)=39.89, p<0.0001, \eta_p^2=0.61$] reflected the well-known finding that ERN is largest at fronto-central midline sites (Fig. 3, top left). The critical Group $\times$ Response interaction was not significant [$F(1,26)=0.01, p=0.94, \eta_p^2=0.0002$]. Similar results were seen for the mean amplitude measure of ERN at the FCz electrode, with a main effect of Response [$F(1,26)=56.46, p<0.0001, \eta_p^2=0.69$] but no Group $\times$ Response interaction [$F(1,26)=0.35, p=0.56, \eta_p^2=0.01$]. Fig. 2 (top left) shows the striking similarity of the ERN in controls vs. patients. These results also held for the ERN measured using alternate baseline procedures (see Figs. S1 and S2, and Supplementary Results).

Pe: There was a main effect of Response [$F(1,26)=25.76, p<0.0001, \eta_p^2=0.50$], with a more positive-going waveform on error trials compared to correct trials. The Electrode $\times$ Response interaction [$F(5,130)=4.46, p=0.01, \eta_p^2=0.15$] reflected a broad central-posterior scalp distribution (Fig. 3, top right). The Group $\times$ Response interaction was not significant [$F(1,26)=2.07, p=0.16, \eta_p^2=0.07$]. At Pz, there was again a main effect of Response [$F(1,26)=54.97, p<0.0001, \eta_p^2=0.68$]. The marginal Group $\times$ Response interaction [$F(1,26)=3.76, p=0.06, \eta_p^2=0.13$] reflected a numerically larger Pe effect in the patients. However, when the -500 to -300 baseline of Rabinak et al. (2013) was used, the Group $\times$

Response interaction was not significant (see Supplementary Materials).

Bayesian Analysis: Because there was no significant difference between the groups for either the ERN or the Pe, we wanted to determine how confident we could be in the null hypothesis (H0) in each case. Therefore, we performed Bayesian hypothesis tests and report Bayes Factors (BF). BF was calculated using the online Bayesian calculator provided by the Perception and Cognition Laboratory at the University of Missouri (<http://pcl.missouri.edu/bayesfactor>). We report the Scaled JZS Bayes Factors (Wagenmakers et al., 2011) in Table 3, along with the mean amplitudes of the ERPs on error trials, correct trials, and the error-correct difference (measured from 0-100 ms for ERN at FCz, and 200-400 ms for Pe at the Pz electrode). $BF > 1$ indicates that the result is more likely to occur under H0 than H1, and $BF < 1$ indicates the result is more likely under H1 than H0 (Wagenmakers et al., 2011). For example, $BF = 2.8$ for the ERN in Exp. 1 means that H0 is 2.8 times more likely than H1. In all cases (except for one), the null was more likely to occur than the alternative hypothesis, i.e. that PTSD patients differed from controls. The exception was for Pe amplitude, with $BF = 1.03$. This suggests it was not possible to determine whether H0 or H1 was more likely, and appears to reflect the numerically larger Pe in the patients (Fig. 1, top).

Correlations between ERN and questionnaires: The associations between the ERN and self-reported levels of PTSD, depression, and post-concussion symptoms were determined using Spearman Rank Correlations (corrected at $p < 0.01$). We did not find a correlation between ERN mean amplitude (error–correct difference at FCz) and symptom severity on any of the measures. Scores on the PCL-M, BDI, PCSC, and NSI did not show a relationship with error monitoring: ERN was not correlated with the PCL-M ($\rho = -0.13, p = 0.50$), BDI ($\rho = -0.19, p = 0.32$), PCSC ($\rho = -0.14, p = 0.47$) or NSI ($\rho = -0.23, p = 0.24$). This also held when the patients were

considered separately: PCL-M ($\rho = -0.07, p = 0.80$), BDI ($\rho = -0.06, p = 0.85$), PCSC ($\rho = -0.15, p = 0.58$) or NSI ($\rho = -0.32, p = 0.25$).

3.2.2. Experiment 2. PTSD patients and controls did not show any significant differences in ERN and Pe amplitudes, replicating Exp. 1 (Fig. 1, bottom). The mean number of error trials in the ERP averages was 62 ± 48 for the patients and 57 ± 34 for the controls [$t(24) = -0.31, p = 0.76$], and for correct trials the means were 524 ± 60 for patients and 505 ± 69 for controls [$t(24) = -0.73, p = 0.47$]. Two patients and two controls had less than 20 error trials each and were excluded from the grand averages. This brought the number of error trials to 72 ± 44 in the patients and 65 ± 29 in the controls [$t(20) = -0.44, p = 0.67$]. However, the results did not change if these participants were included.

ERN: The main effect of Response [$F(1,20) = 36.08, p < 0.0001, \eta_p^2 = .64$] indicated that errors elicited a larger negative component than correct responses. The Electrode $\times$ Response interaction [$F(5,100) = 20.77, p < 0.0001, \eta_p^2 = 0.51$] reflected the typical ERN scalp distribution: greatest at fronto-central midline sites (Fig. 3, bottom left). The Group $\times$ Response interaction was not significant [$F(1,20) = 0.32, p = 0.58, \eta_p^2 = 0.02$]. Similar results were seen for mean amplitude at FCz, with a main effect of Response [$F(1,20) = 32.66, p < 0.0001, \eta_p^2 = 0.62$] but no Group $\times$ Response interaction [$F(1,20) = 0.28, p = 0.60, \eta_p^2 = 0.01$]. As in Exp. 1, the ERP overlay in Fig. 2 (bottom left) shows the between-group similarity of the ERN. The same results were seen for the ERN measured using alternate baseline procedures (see Figs. S1 and S2, and Supplementary Results).

Pe: As in Exp. 1, there was a main effect of Response [$F(1,26) = 92.61, p < 0.0001, \eta_p^2 = 0.82$], with a more positive-going waveform on error trials compared to correct trials. The

Electrode $\times$ Response interaction [$F(5,130)=9.22, p=0.0004, \eta_p^2=0.32$] reflected the central-parietal distribution for the Pe (Fig. 3, bottom right). The Group $\times$ Response interaction was not significant [$F(1,26)=1.79, p=0.20, \eta_p^2=0.08$]. Similar results were seen at the Pz electrode, with a main effect of Response [$F(1,26)=96.48, p<0.0001, \eta_p^2=0.83$] but no Group $\times$ Response interaction [$F(1,26)=0.05, p=0.82, \eta_p^2=0.003$]. This also held for the Pe measured using the alternate baseline procedures (see Supplementary Materials).

Bayesian analysis: Similar to the results from Exp. 1, the Bayes Factors for the ERP measures were all > 1 , which indicates that H_0 was more likely to occur than H_1 (Table 3).

4. Discussion

Combat Veterans with PTSD showed intact error monitoring in two separate experiments, indicating a relative strength in this critical executive control function. The presence of PTSD altered neither the electrophysiological nor the behavioral indices of error monitoring, which contrasts with findings in participants with OCD and GAD (Gehring et al., 2000; Endrass and Ullsperger, 2014; Xiao et al., 2011). Specifically, PTSD patients and controls showed no differences in the ERN and Pe components recorded during a flanker interference task, replicating previous results (Rabinak et al., 2013). An extension of this observation was the failure to find any relationship between the severity of self-reported PTSD symptoms and the size of the ERN. Another novel finding in the present study was the lack of any difference in post-error slowing and error correction behaviors. Thus, PTSD differs from some anxiety disorders, but not others, in its lack of detectable impact on the error monitoring system.²

The ability to resolve response conflict was also intact in the current patients. Performance on two different versions of the arrow flanker task was unaffected by PTSD. This contrasts with the

deficits shown by these same patients in tasks tapping emotional interference control (Ashley et al., 2013), response inhibition (Swick et al., 2012), response consistency (Swick et al., 2013), and dual task performance (Honzel et al., 2014). Taken together, the behavioral results support the general notion of separable executive control functions (Miyake et al., 2000; Stuss, 2011).

4.1. Error-related negativity and PTSD

The present study provides tentative support for an intact ERN component in combat Veterans with PTSD, in agreement with Rabinak and colleagues (2013). The intact ERN generalized across two different task variants that yielded divergent RTs and error rates, suggesting that differences in speed and accuracy did not affect the results. Furthermore, scores on the PCL-M did not correlate with ERN mean amplitude. These results could have implications for a comparative view of PTSD in relation to other disorders (such as GAD) that are divergent for error monitoring measures, yet share other overlapping features. This fits with the notion of cognitive phenotypes (Bilder et al., 2009) or endophenotypes (Manoach and Agam, 2013), which are key to the Research Domain Criteria (RDoC) goal of developing a new, biologically based system for identifying and treating specific symptoms of psychiatric illnesses (Insel and Cuthbert, 2009). Our findings raise the possibility of using the ERN to differentiate anxiety disorders like GAD from fear-based or trauma-based disorders such as specific phobias and PTSD. A healthy self-awareness of committing errors in ongoing behavior could be exploited to help improve internal awareness of emotional state.

However, we acknowledge the limitations in interpreting a null effect. Therefore, we conducted Bayesian two-sample *t*-tests to quantify the confidence with which we could accept the null hypothesis (Boekel et al., 2015; Rotteveel et al., 2015). These tests indicated that the

present results were more likely to occur if the null hypothesis is true, but the evidence was only “anecdotal” (Wagenmakers et al., 2011). Additional studies with larger sample sizes will allow more firm conclusions in the future. A meta-analysis that included data from Exp. 1, Exp. 2, and Rabinak et al. would likely yield Bayes Factors over 3, which would provide substantial evidence for H0.³

An unexpected aspect of the earlier study was a diminished ERN in trauma-exposed control Veterans without PTSD (Rabinak et al., 2013). This ERN reduction was seen relative to Veterans with PTSD, as well as to civilian controls. The authors proposed that the reduced ERN was a potential marker of resilience to trauma, a loss of motivation to perform the task, or a lower significance of errors. In our view, the results suggest that for those control Veterans, either: (1) traumatic experiences during combat exposure impaired their error monitoring systems but did not result in PTSD, which is the opposite of what was seen in PTSD patients; (2) a pre-existing abnormality in the error monitoring system made them more resilient to trauma exposure; or (3) other unmeasured variables contributed to ERN reduction in this group.

4.2. Error-related negativity and mild TBI

One unreported variable in that study was exposure to blasts and other events that resulted in probable concussion or mTBI. As mentioned previously, one or more sports concussions may reduce the size of the ERN (Pontifex et al., 2009), although others have found no such effect (Larson et al., 2012; Bailey et al., 2015). In comparison to the nature of the impact-related mTBIs described in the sports injury literature, mTBI in the majority of OEF/OIF Veterans is due to blast exposure, which may produce different effects on the brain (Fischer et al., 2014). Thus, whether blast-related mTBI would have the same effect on the ERN as a sports concussion

is currently unknown. In our population, most of the PTSD patients also had mTBI, but a few did not. Informal comparison of these subgroups revealed no ERN differences between them, either in Exp. 1 [$t(12)=0.015, p=0.988$] or Exp. 2 [$t(9)=1.15, p=0.28$]. The present results strengthen the findings of Rabinak et al. (2013) by suggesting that OEF/OIF Veterans with PTSD have an intact ERN, regardless of whether they had an mTBI or not. This information was missing from the earlier paper, so the potential effects of mTBI in that study were unknown. However, this interpretation is quite tentative because of the low number of PTSD patients without mTBI in the present study.

We also note that there was no relationship between scores on the post-concussion checklists and ERN amplitude. This suggests that the severity of persistent symptoms following mTBI did not influence the size of the ERN. Furthermore, scores on the PCL-M correlated with post-concussion symptoms reported on the PCSC/day ($\rho=0.63, p=0.001$) and the NSI/month ($\rho=0.88, p<0.0001$) checklists. These results indicate that post-concussion symptoms were highly correlated with PTSD symptoms, and that the tight interrelationship between these self-report measures precluded separating the potential effects of each individually.

4.3. Error-related negativity and depression

Major depression is another condition that can affect the size of the ERN (Ruchow et al., 2006; Chiu and Deldin, 2007; Holmes and Pizzagalli, 2008), although not always (Schrijvers et al., 2009; Bailey et al., 2015). We saw no relationship between ERN amplitude and the severity of depression symptoms reported on the BDI. In fact, it was impossible to disentangle the effects of depression from those of PTSD, because of statistical collinearity. Accordingly, treating BDI scores as a covariate to parcel out the effects of depression is not recommended (Miller and

Chapman, 2001). As in our previous studies (Swick et al., 2012, 2013), we saw a very high correlation between the self-reported symptoms of PTSD and depression in this population ($\rho=0.90, p<0.0001$).

Although DSM-5 now classifies PTSD as a trauma-based disorder, the diagnosis still includes symptoms that overlap with major depression (avoidance, negative alterations in cognition) and anxiety (alterations in arousal and reactivity). The RDoC approach of studying specific symptoms and neural markers that cut across diagnostic lines may be informative here (Insel and Cuthbert, 2009). Vaidyanathan et al. (2011) reviewed evidence suggesting that the ERN could be useful in identifying the shared and distinctive features of different internalizing disorders, i.e. disorders involving anxious and/or depressive symptomatology. For example, an overactive error monitoring system is common to GAD and OCD, but this feature is not shared by phobic anxiety disorders that are strictly fear-based.

4.4. Anterior cingulate cortex and PTSD

The ERN results are interesting in light of the ACC alterations reported in previous structural and functional neuroimaging studies of PTSD (for reviews, see Hayes et al., 2012; Kuhn and Gallinat, 2013). Hamner and colleagues (1999) initially proposed that dysfunction within the ACC may result in exaggerated emotional and behavioral responses in individuals with PTSD. They suggested that ACC gates responses to external and internal stimuli, and an impairment in this top-down control function leads to problems with emotion regulation and attention (Hamner et al., 1999). A subsequent focus of research has been to compare the integrity of rostral/ventral (“affective”) vs. caudal/dorsal (“cognitive”) divisions of the ACC (Bush et al., 2000). This is especially relevant because the amygdala, often reported to be overactive in PTSD

(Shin et al., 2006), receives top-down projections from only the affective portion of the ACC (Devinsky et al., 1995).

Structural MRI studies have observed reductions in gray matter volume in participants with PTSD, in *both* rostral (Rauch et al. 2003; Woodward et al., 2006; Kasai et al., 2008; Herringa et al. 2012) *and* caudal (Woodward et al., 2006; Chen et al., 2012; Chao et al., 2013) subdivisions of the ACC. In many cases, these volume reductions correlated with PTSD symptom severity. Although we don't know if the current patients show ACC volume reductions, what might such changes mean for the generation of the ERN? Based on lesion evidence, damage to either dorsal or ventral regions of the ACC can result in reduced ERN amplitude (Swick and Turken, 2002, 2008). However, any structural alterations in PTSD are likely to be much more subtle than a focal lesion, with an uncertain impact upon the ERN. In an analogous example, individuals with schizophrenia showed no relationship between ACC volume and ERN amplitude, but control participants showed a *negative* correlation between the two (Araki et al., 2013). In another study, PTSD patients showed reductions in ACC volume and P300 amplitude (which tended to show a positive correlation), but controls showed no association between these two measures (Araki et al., 2005). These examples suggest a complicated relationship between ACC volume and the amplitude of specific ERP components. Thus, we cannot make inferences about the absence of structural alterations in the ACC (or any other ERN generator) based on the current results. Future studies should continue to investigate the relationship between changes in ACC volume and error monitoring as they relate to the ERN.

The majority of functional MRI studies in PTSD have presented emotional stimuli or used a symptom provocation design (e.g., Rauch et al., 1996; Bremner et al., 1999; Kim et al., 2008). More relevant to the present study are fMRI experiments that presented non-emotional stimuli in

cognitive control tasks. Here the results are mixed. Activation in dorsal ACC was decreased during the detection of neutral targets (Morey et al. 2008), normal in the conflict condition of the counting Stroop task (Shin et al., 2007), and increased in the multi-source interference task (Shin et al. 2011). On the basis of these conflicting fMRI results, it would be difficult to make a prediction about any larger implications for the error monitoring system. However, one recent fMRI paper reported that combat Veterans with a history of suicidal ideation (and diagnoses of PTSD, depression, and/or mTBI) showed hyperactivation of the ACC in response to errors, compared to similar Veterans without suicidal ideation (Matthews et al., 2012). This could point to an important avenue of ERN research in psychiatric populations at risk for suicide.

4.5. Limitations

Although we replicated the previous finding of an unaltered ERN in combat Veterans with PTSD (Rabinak et al., 2013), there are limitations in interpreting a null effect. Nonetheless, the evidence we obtained in two separate experiments indicated that the present results were more likely to occur if the null hypothesis is true. We acknowledge that Bayes Factors < 3 provide only “anecdotal” support for the null hypothesis (Wagenmakers et al., 2011), and that our sample sizes (and Rabinak et al.’s) are relatively small. But the current results add to the literature suggesting that PTSD does not affect error monitoring, which is an important outcome.

The current study had other limitations that can be addressed in future investigations. Most of the Veterans in our control group were not exposed to combat in Iraq or Afghanistan, which differs from the trauma-exposed Veteran controls of Rabinak et al. (2013). Having two well-matched control groups, one with no trauma exposure and the other with trauma-exposure but no PTSD, is the best way to disentangle the effects of combat experiences from the effects of PTSD

symptoms. In practice, however, it can be difficult to recruit enough individuals to constitute adequate groups for this comparison. This leads to another drawback, the relatively small size of the groups under study here, which limits the strength of evidence in favor of the null effect. Nonetheless, well-conducted studies with small sample sizes can still contribute to future meta-analyses of the ERN, and we propose that combining the current datasets with those of Rabinak et al. (2013) would yield convincing evidence in favor of the null hypothesis (see Footnote 3).

The most straightforward interpretation of the results is that neither PTSD nor mTBI affected the ERN in this study.⁴ However, we were unable to dissociate the potential effects of mTBI from those of PTSD. Thus, results from this military population may not generalize to civilian PTSD. Despite these limitations, we emphasize the importance of characterizing the cognitive strengths and weaknesses of OEF/OIF Veterans, regardless of the non-unitary nature of their diagnoses, to better re-integrate them into civilian life.

4.6. Conclusions

Post-traumatic stress disorder did not affect the ability to detect errors in performance and adjust behavior accordingly. In addition, PTSD symptom severity did not influence the size of the ERN component elicited by the commission of errors in an attentionally demanding task. ERN amplitude was also unaffected by the degree of self-reported depression and post-concussion symptoms. These findings were replicated in two separate experiments, and support previous observations of intact ERN and Pe components in combat Veterans with PTSD (Rabinak et al., 2013). The results could have positive implications for PTSD treatments that rely on self-monitoring abilities, such as neurofeedback and mindfulness training.

Acknowledgements

We are grateful to Dr. Andrew Kayser for patient referrals and to all participants for taking part in the study. We thank Victoria Ashley, Julien Cayton, Dr. Jary Larsen, and Dr. Adrian Willoughby for their assistance in various stages of the project. This work was supported by the U.S. Army Medical Research and Materiel Command under W81XWH-08-2-0086 (D.S.), a VA Merit Review grant (I01CX000566) from Clinical Science Research and Development (D.S.), and a VA Career Development Award (A.T.). The contents reported within do not represent the views of the Department of Veterans Affairs or the United States Government. The authors declare that they have no conflicts of interest.

Contributors

D.S. consulted on study design, analyzed data, and wrote the manuscript; N.H. and A.T. designed the experiments (Exp. 1 and Exp. 2, respectively), collected and analyzed data, and edited the manuscript.

Footnotes

¹ We note, however, that PTSD is no longer classified as an anxiety disorder in DSM-5 (American Psychiatric Association, 2013), which was released after we conducted these experiments.

² Among anxiety disorders, both OCD and GAD are associated with heightened ERN responses, but specific phobias are not (Hajcak et al., 2003).

³ For illustrative purposes, we ran exploratory Bayesian analyses on the combined dataset of Exp. 1 and Exp. 2 ($n=25$ for each group; see Table S1). The Bayes factors were all greater than 3.5, which provides “substantial” evidence in favor of the null hypothesis (Wagenmakers et al., 2011).

⁴ We also did ERN analyses that grouped participants by whether or not they had mTBI (regardless of PTSD). Similar to the results for PTSD, there were no differences in ERN amplitude based on mTBI status. ERN amplitude at FCz: [$t(26) = -0.01, p=0.99$] for Exp. 1; [$t(20) = 0.07, p=0.95$] for Exp. 2. ERN–CRN difference: [$t(26) = 0.45, p=0.66$] for Exp. 1; [$t(20) = 0.24, p=0.81$] for Exp. 2.

References

- Aarts, K., Vanderhasselt, M.A., Otte, G., Baeken, C., Pourtois, G., 2013. Electrical brain imaging reveals the expression and timing of altered error monitoring functions in major depression. *Journal of Abnormal Psychology* 122, 939-950.
- American Psychiatric Association, 2013. *Diagnostic and Statistical Manual of Mental Disorders* (5th ed.). Author, Washington, DC.
- Araki, T., Kasai, K., Yamasue, H., Kato, N., Kudo, N., Ohtani, T., Nakagome, K., Kirihara, K., Yamada, H., Abe, O., Iwanami, A., 2005. Association between lower P300 amplitude and smaller anterior cingulate cortex volume in patients with posttraumatic stress disorder: a study of victims of Tokyo subway sarin attack. *Neuroimage* 25, 43-50.
- Araki, T., Niznikiewicz, M., Kawashima, T., Nestor, P.G., Shenton, M.E., McCarley, R.W., 2013. Disruption of function-structure coupling in brain regions sub-serving self monitoring in schizophrenia. *Schizophrenia Research* 146, 336-343.
- Ashley, V., Honzel, N., Larsen, J., Justus, T., Swick, D., 2013. Attentional bias for trauma-related words: exaggerated emotional Stroop effect in Afghanistan and Iraq war veterans with PTSD. *BMC Psychiatry* 13, 86.
- Bailey, N.W., Hoy, K.E., Maller, J.J., Upton, D.J., Segrave, R.A., Fitzgibbon, B.M., Fitzgerald, P.B., 2015. Neural evidence that conscious awareness of errors is reduced in depression following a traumatic brain injury. *Biological Psychology* 106, 1-10.
- Beck, A.T., Steer, R., Gabin, M., 1988. Psychometric properties of the BDI: twenty-five years of evaluation. *Clinical Psychological Review* 8, 77-100.
- Bilder, R.M., Sabb, F.W., Parker, D.S., Kalar, D., Chu, W.W., Fox, J., Freimer, N.B., Poldrack, R.A., 2009. Cognitive ontologies for neuropsychiatric phenomics research.

Cognitive Neuropsychiatry 14, 419-450.

Blanchard, E.B., Jones-Alexander, J., Buckley, T.C., Forneris, C.A., 1996. Psychometric properties of the PTSD Checklist (PCL). Behaviour Research and Therapy 34, 669-673.

Boekel, W., Wagenmakers, E.J., Belay, L., Verhagen, J., Brown, S., Forstmann, B.U., 2015. A purely confirmatory replication study of structural brain-behavior correlations. Cortex 66, 115-133.

Bremner, J.D., Staib, L.H., Kaloupek, D., Southwick, S.M., Soufer, R., Charney, D.S., 1999. Neural correlates of exposure to traumatic pictures and sound in Vietnam combat veterans with and without posttraumatic stress disorder: A positron emission tomography study. Biological Psychiatry 45, 806-816.

Bush, G., Luu, P., Posner, M.I., 2000. Cognitive and emotional influences in anterior cingulate cortex. Trends in Cognitive Science 4, 215-222.

Carlson, K.F., Kehle, S.M., Meis, L.A., Greer, N., Macdonald, R., Rutks, I., Sayer, N.A., Dobscha, S.K., Wilt, T.J., 2011. Prevalence, assessment, and treatment of mild traumatic brain injury and posttraumatic stress disorder: a systematic review of the evidence. Journal of Head Trauma Rehabilitation 26, 103-115.

Carter, C.S., MacDonald, A.W., 3rd, Ross, L.L., Stenger, V.A., 2001. Anterior cingulate cortex activity and impaired self-monitoring of performance in patients with schizophrenia: an event-related fMRI study. American Journal of Psychiatry 158, 1423-1428.

Cicerone, K.D., Kalmar, K., 1995. Persistent postconcussion syndrome: The structure of subjective complaints after mTBI. Journal of Head Trauma Rehabilitation 10, 1-17.

Clemans, Z.A., El-Baz, A.S., Hollifield, M., Sokhadze, E.M., 2012. Single trial time-frequency domain analysis of error processing in post-traumatic stress disorder. Neuroscience Letters

525, 105-110.

- Compton, R.J., Lin, M., Vargas, G., Carp, J., Fineman, S.L., Quandt, L.C., 2008. Error detection and posterror behavior in depressed undergraduates. *Emotion* 8, 58-67.
- Danielmeier, C., Ullsperger, M., 2011. Post-error adjustments. *Frontiers in Psychology* 2, 233.
- De Beaumont, L., Beauchemin, M., Beaulieu, C., Jolicoeur, P., 2013. Long-term attenuated electrophysiological response to errors following multiple sports concussions. *Journal of Clinical Experimental Neuropsychology* 35, 596-607.
- Dehaene, S., Posner, M.I., Tucker, D.M., 1994. Localization of a neural system for error detection and compensation. *Psychological Science* 303-305.
- Devinsky, O., Morrell, M.J., Vogt, B.A., 1995. Contributions of anterior cingulate cortex to behaviour. *Brain* 118, 279-306.
- Endrass, T., Ullsperger, M., 2014. Specificity of performance monitoring changes in obsessive-compulsive disorder. *Neuroscience & Biobehavioral Reviews* 46, 124-138.
- Eriksen, B.A., Eriksen, C.W., 1974. Effects of noise letters upon identification of a target letter in a nonsearch task. *Perception & Psychophysics* 16, 143-149.
- Falkenstein, M., Hohnsbein, J., Hoormann, J., Blanke, L., 1991. Effects of crossmodal divided attention on late ERP components. II. Error processing in choice reaction tasks. *Electroencephalography and Clinical Neurophysiology* 78, 447-455.
- Falkenstein, M., Hoormann, J., Christ, S., Hohnsbein, J., 2000. ERP components on reaction errors and their functional significance: a tutorial. *Biological Psychology* 51, 87-107.
- Fischer, B.L., Parsons, M., Durgerian, S., Reece, C., Mourany, L., Lowe, M.J., Beall, E.B., Koenig, K.A., Jones, S.E., Newsome, M.R., Scheibel, R.S., Wilde, E.A., Troyanskaya, M., Merkley, T.L., Walker, M., Levin, H.S., Rao, S.M., 2014. Neural activation during response

- inhibition differentiates blast from mechanical causes of mild to moderate traumatic brain injury. *Journal of Neurotrauma* 31, 169-179.
- Fischer, H., 2014. A Guide to U.S. Military Casualty Statistics: Operation New Dawn, Operation Iraqi Freedom, and Operation Enduring Freedom. Congressional Research Service Report for Congress.
- Forbes, D., Creamer, M., Biddle, D., 2001. The validity of the PTSD checklist as a measure of symptomatic change in combat-related PTSD. *Behaviour Research and Therapy* 39, 977-986.
- Frith, C.D., Done, D.J., 1989. Experiences of alien control in schizophrenia reflect a disorder in the central monitoring of action. *Psychological Medicine* 19, 359-363.
- Gehring, W.J., Goss, B., Coles, M.G.H., Meyer, D.E., Donchin, E., 1993. A neural system for error detection and compensation. *Psychological Science* 4, 385-390.
- Gehring, W.J., Himle, J., Nisenson, L.G., 2000. Action-monitoring dysfunction in obsessive-compulsive disorder. *Psychological Science* 11, 1-6.
- Gehring, W.J., Liu, Y., Orr, J.M., Carp, J., 2012. The error-related negativity (ERN/Ne), in: Luck, S.J., Kappenman, E.S. (Eds.), *The Oxford Handbook of Event-Related Potential Components*. Oxford University Press, Oxford, pp. 231-291.
- Gouvier, W.D., Cubic, B., Jones, G., Brantley, P., Cutlip, Q., 1992. Postconcussion symptoms and daily stress in normal and head-injured college populations. *Archives of Clinical Neuropsychology* 7, 193-211.
- Gratton, G., Coles, M.G., Donchin, E., 1983. A new method for off-line removal of ocular artifact. *Electroencephalography and Clinical Neurophysiology* 55, 468-484.
- Hajcak, G., McDonald, N., Simons, R.F., 2003. Anxiety and error-related brain activity.

- Biological Psychology 64, 77–90.
- Hamner, M.B., Lorberbaum, J.P., George, M.S., 1999. Potential role of the anterior cingulate cortex in PTSD: review and hypothesis. *Depression & Anxiety* 9, 1-14.
- Hayes, J.P., Hayes, S.M., Mikedis, A.M., 2012. Quantitative meta-analysis of neural activity in posttraumatic stress disorder. *Biology of Mood & Anxiety Disorders* 2, 9.
- Holmes, A.J., Pizzagalli, D.A., 2008. Spatiotemporal dynamics of error processing dysfunctions in major depressive disorder. *Archives of General Psychiatry* 65, 179 –188.
- Holroyd, C.B., Dien, J., Coles, M.G., 1998. Error-related scalp potentials elicited by hand and foot movements: evidence for an output-independent error-processing system in humans. *Neuroscience Letters* 242, 65-68.
- Honzel, N., Justus, T., Swick D., 2014. Post-traumatic stress disorder is associated with limited executive resources in a working memory task. *Cognitive, Affective, & Behavioral Neuroscience* 14, 792-804.
- Insel, T.R., Cuthbert, B.N., 2009. Endophenotypes: Bridging genomic complexity and disorder heterogeneity. *Biological Psychiatry* 66, 988-989.
- Jentzsch, I., Dudschig, C., 2009. Why do we slow down after an error? Mechanisms underlying the effects of posterror slowing. *The Quarterly Journal of Experimental Psychology* 62, 209-218.
- Kim, M.J., Chey, J., Chung, A., Bae, S., Khang, H., Ham, B., Yoon, S.J., Jeong, D.U., Lyoo, I.K., 2008. Diminished rostral anterior cingulate activity in response to threat-related events in posttraumatic stress disorder. *Journal of Psychiatric Research* 42, 268-277.
- Kim, M.J., Lyoo, I.K., Kim, S.J., Sim, M., Kim, N., Choi, N., Jeong, D.U., Covell, J., Renshaw, P.F., 2005. Disrupted white matter tract integrity of anterior cingulate in trauma survivors.

- Neuroreport 16, 1049-1053.
- Kopp, B., Rist, F., Mattler, U., 1996. N200 in the flanker task as a neurobehavioral tool for investigating executive control. *Psychophysiology* 33, 282-294.
- Kühn, S., Gallinat, J., 2013. Gray matter correlates of posttraumatic stress disorder: a quantitative meta-analysis. *Biological Psychiatry* 73, 70-74.
- Larson, M.J., Clayson, P.E., Farrer, T.J., 2012. Performance monitoring and cognitive control in individuals with mild traumatic brain injury. *Journal of the International Neuropsychological Society* 18, 323-333.
- Larson, M.J., Kaufman, D.A., Kellison, I.L., Schmalfluss, I.M., Perlstein, W.M., 2009. Double jeopardy! The additive consequences of negative affect on performance-monitoring decrements following traumatic brain injury. *Neuropsychology* 23, 433-444.
- Larson, M.J., Kaufman, D.A., Schmalfluss, I.M., Perlstein, W.M., 2007. Performance monitoring, error processing, and evaluative control following severe TBI. *Journal of the International Neuropsychological Society* 13, 961-971.
- Manoach, D.S., Agam, Y., 2013. Neural markers of errors as endophenotypes in neuropsychiatric disorders. *Frontiers in Human Neuroscience* 18, 350.
- Mathalon, D.H., Fedor, M., Faustman, W.O., Gray, M., Askari, N., Ford, J.M., 2002. Response-monitoring dysfunction in schizophrenia: an event-related brain potential study. *Journal of Abnormal Psychology* 111, 22-41.
- Matthews, S., Spadoni, A., Knox, K., Strigo, I., Simmons, A., 2012. Combat-exposed war veterans at risk for suicide show hyperactivation of prefrontal cortex and anterior cingulate during error processing. *Psychosomatic Medicine* 74, 471-475.
- Miller, G.A., Chapman, J.P., 2001. Misunderstanding analysis of covariance. *Journal of*

Abnormal Psychology 110, 40-48.

Miyake, A., Friedman, N.P., Emerson, M.J., Witzki, A.H., Howerter, A., Wager, T.D., 2000.

The unity and diversity of executive functions and their contributions to complex "frontal lobe" tasks: a latent variable analysis. *Cognitive Psychology* 41, 49-100.

Morey, R.A., Petty, C.M., Cooper, D.A., Labar, K.S., McCarthy, G., 2008. Neural systems for executive and emotional processing are modulated by symptoms of posttraumatic stress disorder in Iraq War veterans. *Psychiatry Research* 162, 59-72.

Overbeek, T.J.M., Nieuwenhuis, S., Ridderinkhof, K.R., 2005. Dissociable components of error processing: on the functional significance of the Pe vis-a-vis the ERN/Ne. *Journal of Psychophysiology* 19, 319–329.

Pontifex, M.B., O'Connor, P.M., Broglio, S.P., Hillman, C.H., 2009. The association between mild traumatic brain injury history and cognitive control. *Neuropsychologia* 47, 3210-3216.

Posner, M.I., Rothbart, M.K., Sheese, B.E., Tang, Y., 2007. The anterior cingulate gyrus and the mechanism of self-regulation. *Cognitive, Affective, & Behavioral Neuroscience* 7, 391-395.

Rabbitt, P. M., 1966. Errors and error correction in choice-response tasks. *Journal of Experimental Psychology* 71, 264–272.

Rabinak, C.A., Holman, A., Angstadt, M., Kennedy, A.E., Hajcak, G., Phan, K.L., 2013. Neural response to errors in combat-exposed returning veterans with and without post-traumatic stress disorder: a preliminary event-related potential study. *Psychiatry Research* 213, 71-78.

Rauch, S.L., Shin, L.M., Segal, E., Pitman, R.K., Carson, M.A., McMullin, K., Whalen, P.J., Makris, N., 2003. Selectively reduced regional cortical volumes in post-traumatic stress disorder. *Neuroreport* 14, 913-916.

Rauch, S.L., van der Kolk, B.A., Fisler, R.E., Alpert, N.M., Orr, S.P., Savage, C.R., Fischman,

- A.J., Jenike, M.A., Pitman, R.K., 1996. A symptom provocation study of posttraumatic stress disorder using positron emission tomography and script-driven imagery. *Archives of General Psychiatry* 53, 380-387.
- Ridderinkhof, K.R., Nieuwenhuis, S., Braver, T.S., 2007. Medial frontal cortex function: an introduction and overview. *Cognitive, Affective, & Behavioral Neuroscience* 7, 261-265.
- Rotteveel, M., Gierholz, A., Koch, G., van Aalst, C., Pinto, Y., Matzke, D., Steingroever, H., Verhagen, J., Beek, T.F., Selker, R., Sasiadek, A., Wagenmakers, E.J., 2015. On the automatic link between affect and tendencies to approach and avoid: Chen and Bargh (1999) revisited. *Frontiers in Psychology* 6, 335.
- Ruchow, M., Herrnberger, B., Beschoner, P., Gron, G., Spitzer, M., Kiefer, M., 2006. Error processing in major depressive disorder: evidence from event-related potentials. *Journal of Psychiatry Research* 40, 37-46.
- Rushworth, M.F., Walton, M.E., Kennerley, S.W., Bannerman, D.M., 2004. Action sets and decisions in the medial frontal cortex. *Trends in Cognitive Science* 8, 410-417.
- Salkovskis, P.M., 1999. Understanding and treating obsessive-compulsive disorder. *Behaviour Research & Therapy* 37, S29-S52.
- Schrijvers, D., De Bruijn, E.R., Maas, Y.J., Vancoillie, P., Hulstijn, W., Sabbe, B.G., 2009. Action monitoring and depressive symptom reduction in major depressive disorder. *International Journal of Psychophysiology* 71, 218-224.
- Shin, L.M., Bush, G., Milad, M.R., Lasko, N.B., Brohawn, K.H., Hughes, K.C., Macklin, M.L., Gold, A.L., Karpf, R.D., Orr, S.P., Rauch, S.L., Pitman, R.K., 2011. Exaggerated activation of dorsal anterior cingulate cortex during cognitive interference: a monozygotic twin study of posttraumatic stress disorder. *American Journal of Psychiatry* 168, 979-985.

- Shin, L.M., Bush, G., Whalen, P.J., Handwerker, K., Cannistraro, P.A., Wright, C.I., Martis, B., Macklin, M.L., Lasko, N.B., Orr, S.P., Pitman, R.K., Rauch, S.L., 2007. Dorsal anterior cingulate function in posttraumatic stress disorder. *Journal of Traumatic Stress* 20, 701-712.
- Shin, L.M., Rauch, S.L., Pitman, R.K., 2006. Amygdala, medial prefrontal cortex, and hippocampal function in PTSD. *Annals of the New York Academy of Science* 1071, 67-79.
- Shin, L.M., Whalen, P.J., Pitman, R.K., Bush, G., Macklin, M.L., Lasko, N.B., Orr, S.P., McInerney, S.C., Rauch, S.L., 2001. An fMRI study of anterior cingulate function in posttraumatic stress disorder. *Biological Psychiatry* 50, 932-942.
- Solbakk, A.K., Funderud, I., Lovstad, M., Endestad, T., Meling, T., Lindgren, M., Knight, R.T., Kramer, U.M., 2014. Impact of orbitofrontal lesions on electrophysiological signals in a stop signal task. *Journal of Cognitive Neuroscience* 26, 1528-1545.
- Stemmer, B., Segalowitz, S.J., Witzke, W., Schonle, P.W., 2004. Error detection in patients with lesions to the medial prefrontal cortex: an ERP study. *Neuropsychologia* 42, 118-130.
- Stuss, D.T., 2011. Functions of the frontal lobes: relation to executive functions. *Journal of the International Neuropsychological Society* 17, 759-765.
- Swick, D., Honzel, N., Larsen, J., Ashley, V., 2013. Increased response variability as a marker of executive dysfunction in veterans with post-traumatic stress disorder. *Neuropsychologia* 51, 3033-3040.
- Swick, D., Honzel, N., Larsen, J., Ashley, V., Justus, T., 2012. Impaired response inhibition in veterans with post-traumatic stress disorder and mild traumatic brain injury. *Journal of the International Neuropsychological Society* 18, 917-926.
- Tanielian, T.L., Jaycox, L., Rand Corporation, 2008. Invisible wounds of war: psychological and cognitive injuries, their consequences, and services to assist recovery. RAND, Santa Monica,

CA.

- Taylor, B.C., Hagel, E.M., Carlson, K.F., Cifu, D.X., Cutting, A., Bidelsbach, D.E., Sayer, N.A., 2012. Prevalence and costs of co-occurring traumatic brain injury with and without psychiatric disturbance and pain among Afghanistan and Iraq War Veteran V.A. users. *Medical Care* 50, 342-346.
- Turken, A.U., Vuilleumier, P., Mathalon, D.H., Swick, D., Ford, J.M., 2003. Are impairments of action monitoring and executive control true dissociative dysfunctions in patients with schizophrenia? *American Journal of Psychiatry* 160, 1881-1883.
- Turken, A.U., Swick, D., 2008. The effect of orbitofrontal lesions on the error-related negativity. *Neuroscience Letters* 441, 7-10.
- Ullsperger, M., Fischer, A.G., Nigbur, R., Endrass, T., 2014. Neural mechanisms and temporal dynamics of performance monitoring. *Trends in Cognitive Science* 18, 259-267.
- Ullsperger, M., von Cramon, D.Y., Muller, N.G., 2002. Interactions of focal cortical lesions with error processing: Evidence from event-related brain potentials. *Neuropsychology* 16, 548-561.
- Vaidyanathan, U., Nelson, L.D., Patrick, C.J., 2012. Clarifying domains of internalizing psychopathology using neurophysiology. *Psychological Medicine* 42, 447-459.
- Wagenmakers, E.J., Wetzels, R., Borsboom, D., van der Maas, H.L., 2011. Why psychologists must change the way they analyze their data: the case of psi. Comment on Bem (2011). *Journal of Personality and Social Psychology* 100, 426-432.
- Weathers, F.W., Litz, B.T., Huska, J.A., Keane, T.M., 1994. PTSD Checklist—Military Version (PCL-M) for DSM-IV. National Center for PTSD—Behavioral Science Division, Boston, MA.
- Weinberg, A., Klein, D.N., Hajcak, G., 2012. Increased error-related brain activity distinguishes

generalized anxiety disorder with and without comorbid major depressive disorder. *Journal of Abnormal Psychology* 121, 885-896.

Woodward, SH, Kaloupek, DG, Streeter, CC, Martinez, C, Schaer, M, Eliez, S, 2006. Decreased anterior cingulate volume in combat-related PTSD. *Biological Psychiatry* 59, 582-587.

Xiao, Z., Wang, J., Zhang, M., Li, H., Tang, Y., Wang, Y., Fan, Q., Fromson, J.A., 2011. Error-related negativity abnormalities in generalized anxiety disorder and obsessive-compulsive disorder. *Progress in Neuro-Psychopharmacology & Biological Psychiatry* 35, 265-272.

Figure Legends

Fig. 1. ERPs are shown at six midline electrodes for Control participants (left) and PTSD participants (right). Response onset is at the vertical bar, and ERPs are shown from 200 ms pre-response to 500 ms post-response (x -axis is time in ms). The baseline interval is -200 to 0 ms pre-response. Scale for the y -axis is -10 μ V to +15 μ V in all cases, with negative voltage plotted upwards. The ERN (0-100 ms) is labeled at FCz, and the Pe component (200-400 ms) is labeled at Pz. *Top:* Grand average ERPs in Experiment 1. *Bottom:* Grand average ERPs in Experiment 2.

Fig. 2. ERP overlay of controls (blue lines) vs. PTSD participants (green lines) for Error trials (left) and Correct trials (right) at electrode FCz. Response onset is at the vertical bar, and ERPs extend from 200 ms pre-response to 900 ms post-response, with a -200 to 0 ms baseline. Scale for the y -axis is -10 μ V to +15 μ V, with negative plotted up. *Top:* Experiment 1. *Bottom:* Experiment 2.

Fig. 3. Voltage maps displaying the scalp topography for the Error-Correct difference waves for the ERN (measured as mean amplitude between 0 and 100 ms) and the Pe (mean amplitude between 200-400 ms) in Controls and PTSD patients. Left lateral electrodes are shown on the left side of each map, and right is on the right. *Top:* Topographic maps in Experiment 1. Scale bar extends from -5.5 μ V to +3.5 μ V. *Bottom:* Topographic maps in Experiment 2. Scale bar extends from -8 μ V to +8 μ V.

Table 1. Demographic information and self-rating scores for Controls and Patients with PTSD in Experiment 1 (left) and Experiment 2 (right)

	Experiment 1		Experiment 2	
	Controls (<i>n</i> =14)	Patients (<i>n</i> =14)	Controls (<i>n</i> =13)	Patients (<i>n</i> =13)
Age (years)	36.0 (9.1)	31.7 (6.4) n.s.	36.9 (9.5)	36.2 (9.4) n.s.
Education (years)	15.0 (2.1)	13.9 (1.3) n.s.	15.1 (1.9)	13.9 (1.4) n.s.
Handedness	13 R, 1 L	13 R, 1 L	12 R, 1 L	12 R, 1 L
Deployed	5	14	5	13
Years post	--	4.4 (1.7)	--	5.5 (1.9)
Medications (<i>n</i>)	1	2	0	5
Therapy (<i>n</i>) [#]	1	5	1	6
PCL-M	25.8 (7.2)	56.2 (12.9) ***	26.1 (9.2)	61.3 (9.5) ***
BDI	5.6 (5.5)	17.1 (6.6) ***	5.3 (5.4)	23.8 (9.6) ***
PCSC	36.8 (5.3)	54.9 (24.6) *	--	--
NSI	9.2 (6.2)	33.8 (14.3) ***	--	--

Note: The means (standard deviations) are given for age, education, estimated years post-event(s), PCL-M, BDI, PCSC, and NSI. n.s. = not significantly different from controls; * significantly different from controls at $p < .05$; *** significantly different from controls at $p \leq 0.001$. Therapy indicates number of participants undergoing individual psychotherapy before (completed) or during the date of testing (ongoing).[#] Data were not available for 9 controls, 1 patient in Exp. 1 and 5 controls, 1 patient in Exp. 2. R = right, L = left; PCL-M = PTSD checklist, military version; BDI = Beck Depression Inventory; PCSC = Postconcussion Syndrome Checklist; NSI = Neurobehavioral Symptom Inventory.

Table 2

Table 2. Behavioral performance in Experiment 1 (left) and Experiment 2 (right) for controls and PTSD patients

		Exp. 1		Exp. 2	
	Condition	Controls Mean (SD)	Patients Mean (SD)	Controls Mean (SD)	Patients Mean (SD)
Conflict effect	RT Congr	452.8 (51.7)	454.6 (66.4)	303.2 (38.1)	329.1 (68.4)
	RT Incon	508.6 (51.6) ***	504.4 (66.1) ***	387.6 (30.2) ***	415.4 (77.6) ***
PE slowing	RT after correct	481.2 (51.9)	479.9 (66.8)	340.2 (34.6)	369.3 (73.0)
	RT after error	517.7 (56.3) ***	511.7 (71.5) ***	346.2 (34.3) n.s.	356.1 (62.9) n.s.
Conflict effect	Error rate Congr	4.5 (2.8)	3.5 (2.1)	1.2 (2.0)	1.1 (1.7)
	Error rate Incon	16.4 (7.7) ***	13.6 (6.3) ***	18.7 (9.7) ***	18.6 (12.6) ***

Note: The means (standard deviations) are in msec for RT (reaction time) and in percentages for error rate. Congr = congruent trials; Incon = incongruent trials; Conflict Effect = Incon > Congr; PE slowing (post-error slowing) = RT after error > RT after correct; n.s. = not significant for within-group comparison with row above; *** $p < .00001$ for within-group comparison with row above. No between-group differences were observed.

Table 3. ERP mean amplitude measures in Experiment 1 (top) and Experiment 2 (bottom) for controls and PTSD patients

Exp. 1	Controls Mean (SD)	Patients Mean (SD)	Bayes factor
ERN	-2.80 (2.50)	-2.60 (3.41)	2.798
CRN	1.97 (2.73)	2.98 (2.03)	1.785
Δ ERN	-4.77 (3.15)	-5.58 (4.08)	2.483
Pe	1.79 (1.74)	2.87 (1.68)	1.030
Pc	0.16 (1.59)	0.09 (1.76)	2.815
Δ Pe	1.63 (1.26)	2.78 (1.83)	1.374
Exp. 2			
ERN	-2.96 (3.33)	-3.40 (4.33)	2.536
CRN	3.77 (4.65)	2.20 (3.81)	1.988
Δ ERN	-6.73 (4.63)	-5.59 (5.45)	2.351
Pe	7.10 (2.96)	5.58 (2.55)	1.452
Pc	1.57 (1.70)	0.31 (2.83)	1.468
Δ Pe	5.53 (2.60)	5.28 (2.57)	2.552

Note: The Bayes Factor indicates the decisiveness of evidence for or against the null hypothesis (Boekel et al., 2015). Means (standard deviations) are in microvolts for these ERP components: the Error-Related Negativity (ERN), the Correct-Related Negativity (CRN), Error – Correct difference (Δ ERN), the Error Positivity (Pe), Correct Positivity (Pc), and Error – Correct difference (Δ Pe). ERN was measured in the 0-100 ms post-response window at FCz, and Pe

measured from 200-400 ms at Pz. Bayes Factor > 1 indicates that the result is more likely to occur under the null hypothesis (H_0) than the alternate hypothesis (H_1). Bayes Factor < 1 indicates the result is more likely under H_1 than H_0 (Wagenmakers et al., 2011).

Figure 1

Exp. 1

Exp. 2

— Correct
— Error

Figure 2

Figure 3