Genomic diversity and phylogenetic relationships in the genus *Parthenium*

Daniel C. Ilut^a, Paul L. Sanchez^b, Denise E. Costich^c, Bernd Friebe^d, Terry A. Coffelt^b, John M. Dyer^b, Matthew A. Jenks^e, Michael A. Gore^{a,*}

^aPlant Breeding and Genetics Section, School of Integrative Plant Science, Cornell University, Ithaca, NY, 14853, USA

^bUSDA-ARS, U.S. Arid-Land Agricultural Research Center, 21881 North Cardon Lane, Maricopa, AZ, 85138, USA

^cInternational Maize and Wheat Improvement Center (CIMMYT), Texcoco, Mexico

^dWheat Genetics Resource Center, Department of Plant Pathology, Kansas State University, Manhattan, KS, 66506, USA

^eDivision of Plant and Soil Sciences, West Virginia University, Morgantown, WV 26506, USA

Abstract

Guayule (*Parthenium argentatum* A. Gray) is a perennial woody shrub native to the North American Chihuahuan Desert that holds promise as a sustainable source of natural rubber and hypoallergenic latex. The improvement of guayule for commercial-scale production could be accelerated through genomics-assisted breeding, but such a strategy is severely limited by the paucity of available genomic tools and well-characterized genetic resources. To that end, we used genotyping-by-sequencing (GBS) to simultaneously identify and genotype tens of thousands of single-nucleotide polymorphism (SNP) markers across 62 plant samples from seven wild and cultivated guayule, three *Parthenium* interspecific hybrid, four mariola (*Parthenium incanum* Kunth), and one wild quinine (*Parthenium integrifolium* L.) accession(s) that have been characterized for ploidy level and nuclear genome size in this and a prior study. Phylogenetic analysis using the SNP data identified at least two distinct sources of guayule breeding material in the cultivated accessions, previously unknown multi-species

*Corresponding author

Email addresses: dci11@cornell.edu (Daniel C. Ilut), paul.sanchez@ars.usda.gov (Paul L. Sanchez), d.costich@cgiar.org (Denise E. Costich), friebe@ksu.edu (Bernd Friebe), terry.coffelt.ars@gmail.com (Terry A. Coffelt), john.dyer@ars.usda.gov (John M. Dyer), majenks@mail.wvu.edu (Matthew A. Jenks), mag87@cornell.edu (Michael A. Gore)

Preprint submitted to Industrial Crops and Products

July 9, 2015

hybrids within two *Parthenium* hybrid cultivars, and guayule/mariola hybrids within one guayule and one mariola wild collected accession. Similar to previously reported results for guayule and mariola, we observed aneuploidy and multiple ploidy levels among individual plants (mixed ploidy) within three *Parthenium* interspecific hybrid accessions newly characterized in this study. Nuclear genome size characterization of wild quinine, a first for this species, found an estimated haploid nuclear genome size (5757 Mb) for the tetraploid ($2n = 4x = 72$) accession that was more than 1.5-fold larger than that of tetraploid ($2n = 4x = 72$) guayule or mariola. Together, these results further underscore the need for a comprehensive characterization of available guayule germplasm and sister taxa with both SNP markers and flow cytometry, illustrate the novel utility of GBS for the genus *Parthenium*, and lay the foundation for genomics-assisted breeding in guayule.

Keywords: Guayule, Mariola, *Parthenium*, Genotyping-by-sequencing, Polyploidy, Apomixis, Nuclear genome size

1. Introduction

Natural rubber, one of the essential raw materials of modern society, is a major international trade commodity with worldwide imports of 8.5 million metric tons valued at over 24 billion USD in 2013 (DESA/UNSD, United Nations Comtrade database; <http://comtrade.un.org>). Nearly all of it is derived from the rubber tree [*Hevea brasiliensis* (Willd. ex A. Juss.) Müll. Arg.] with more than two thirds being produced in Thailand and Indonesia (FAOSTAT database; <http://faostat3.fao.org>). The geographical concentration of natural rubber production and the vulnerability of monoculture rubber tree plantations to South American leaf blight caused by the fungal pathogen *Microcyclus ulei*

Abbreviations

GBS: genotyping-by-sequencing, GRIN: Germplasm Resources Information Network, SNP: single-nucleotide polymorphism, NCBI: National Center for Biotechnology Information, NPGS: USDA-ARS National Plant Germplasm System, NALPGRU: USDA-ARS National Arid Land Plant Genetics Resources Unit

11 (Rivano, 1997) make the nearly exclusive reliance on *H. brasiliensis* plantations
12 for the world’s natural rubber needs a precarious situation. In combination
13 with the projected shortage of rubber by the next decade (Mann, 2009), this
14 has led to a renewed interest in the development of alternative sources of natural
15 rubber.

16 The woody perennial shrub guayule (*Parthenium argentatum* A. Gray), a
17 native plant of the desert regions of northern Mexico and southwestern United
18 States, has been intermittently improved and cultivated as a source of natu-
19 ral rubber over the past century (reviewed in Hammond and Polhamus, 1965;
20 Thompson and Ray, 1989). However, development of high rubber yielding cul-
21 tivars has met with only partial success, in large part due to a combination
22 of limited genetic diversity in the source of germplasm (Thompson and Ray,
23 1989), extensive polyploidy variation within and among the available guayule
24 accessions (Gore et al., 2011), and a complex mode of reproduction that is
25 characterized by sporophytic self-incompatibility in diploid plants and faculta-
26 tive apomixis (diplospory type) in polyploid plants (reviewed in Ray et al., 2005;
27 Thompson and Ray, 1989). Recently, Gore et al. (2011) successfully adapted au-
28 tomated flow cytometry methods to efficiently characterize ploidy level variation
29 in guayule breeding programs. However, in order to enable modern genomics-
30 assisted breeding approaches for the accelerated improvement of this nascent
31 crop, a thorough characterization of both polyploidy variation and genetic di-
32 versity within the guayule germplasm pool is essential.

33 In addition to rubber content, improvements in other agronomically im-
34 portant traits such as disease resistance, cold tolerance, biomass, and vigor
35 could significantly contribute to the suitability of guayule as a commercial crop.
36 Although guayule is the only species in the genus *Parthenium* that produces
37 significant quantities of natural rubber, interspecific hybrids with other *Parthe-*
38 *nium* species are a potential source of such agronomic improvements (reviewed
39 in Ray et al., 2010; Thompson and Ray, 1989), and several *Parthenium* species
40 such as mariola (*Parthenium incanum* Kunth) and wild quinine (*Parthenium*
41 *integrifolium* L.) have been shown to form natural and/or artificial interspecific

hybrids with guayule (Powers and Rollins, 1945; Rollins, 1946; Hashemi et al., 1989). Mariola, a perennial shrub native to Mexico and southwestern United States (Rollins, 1950), is a facultative apomict with a natural ploidy series and considered the closest related taxon to guayule (Powers and Rollins, 1945). It has been suggested as a source of vigor and disease resistance (Thompson and Ray, 1989), and its habitat range (Figure B.5) suggests a broader climatic tolerance than guayule. Wild quinine is a perennial herbaceous plant native to central and eastern United States (Mears, 1975; Rollins, 1950) and a potential source of cold resistance and non-arid adaptation (Thompson and Ray, 1989). However, with the exception of a recent ploidy and nuclear genome size evaluation of mariola accessions (Sanchez et al., 2014), minimal genomic information is available for these two *Parthenium* species, and the availability of this information is expected to greatly facilitate interspecific breeding programs.

Building upon previously published data for 10 guayule, mariola, and *Parthenium* hybrid accessions (Gore et al., 2011; Sanchez et al., 2014), we used flow cytometry to expand ploidy and nuclear genome size characterization to four interspecific hybrids between guayule and other *Parthenium* species. Additionally, we estimated the nuclear genome size and ploidy level of a wild quinine accession, a first for this *Parthenium* species. We also used genotyping-by-sequencing (GBS; Elshire et al., 2011) for simultaneously identifying and scoring tens of thousands of single-nucleotide polymorphism (SNP) markers on 62 plant samples of the 15 *Parthenium* accessions that captured a range of ploidy levels to i) characterize genomic diversity among and within these *Parthenium* species and interspecific hybrids, ii) identify cryptic hybrids, and iii) estimate the heterozygosity level of *Parthenium* accessions.

2. Materials and methods

2.1. Plant material and growth conditions

Achenes (hereafter seeds) of one wild collected accession (R1110) and three interspecific hybrids of guayule and other *Parthenium* species (CAL-1, CAL-2,

71 CAL-3) were obtained from the USDA-ARS National Plant Germplasm System
 72 (NPGS; www.ars-grin.gov/npgs) through the National Arid Land Plant Genet-
 73 ics Resources Unit (NALPGRU) in Parlier, CA. In addition, commercial seed
 74 of a wild quinine accession was obtained from Prairie Nursery, Inc., Westfield,
 75 WI, because this taxon was not available from NPGS. The pedigrees for all
 76 accessions in this study are summarized in Figure 1, and more extensive infor-
 77 mation is available in Table A.2 and references therein. Plants were grown as
 78 described in Sanchez et al. (2014). All experimental procedures that do not
 79 differ significantly from Sanchez et al. (2014) are described in AppendixC for
 80 reference.

81 *2.2. Collection locales and habitat range*

82 Collection locales for the Powers 1942 historic accessions 4254 and 4265
 83 were identified from literature: “area southwest of Mapimí, Durango, Mexico”
 84 (Powers, 1942; Bergner, 1946) and “from 5 plants growing in an isolated area
 85 along the road from La Mancha to Pasaje” (Powers, 1942; Johnson, 1950) re-
 86 spectively. Locales relevant to collection sites or accession development were
 87 geolocated using the Google Maps Javascript API v1.3 ([https://developers.](https://developers.google.com/maps/documentation/javascript)
 88 [google.com/maps/documentation/javascript](https://developers.google.com/maps/documentation/javascript)), and areas of germplasm collec-
 89 tion were geocoded using locale descriptions or recorded GPS coordinates. Maps
 90 of herbarium specimen collections for mariola, guayule, and wild quinine were
 91 obtained from a genus treatment by Rollins (1950), and approximate habitat
 92 ranges were derived by expanding outward from each collection point. A map
 93 of the northernmost guayule habitat line by Bullard (1946) was obtained from
 94 its reproduction in Hammond and Polhamus (1965). The collection maps and
 95 guayule habitat line were overlaid on a geographic map of North America using
 96 the Google Maps Javascript API v1.3.

97 *2.3. Sample collection and analysis of ploidy level with flow cytometry*

98 Leaf tissue samples were collected and prepared for ploidy analysis for each
 99 of the five accessions (CAL-1, CAL-2, CAL-5, R1110, wild quinine) as part of

an experiment that also included 17 additional guayule, mariola, and interspecific hybrid accession(s) (Sanchez et al., 2014) following procedures described therein. A total of 12 or 15 plants were evaluated for each of the five *Parthenium* accessions, and samples were run in random order by assigning a plant from each accession with equal probability to one of 15 blocks (runs) on the flow cytometer. Only 12 plants were evaluated for the wild quinine accession due to its relatively lower germination and seedling survival rate. The entire experiment was replicated twice, for a total of two measurements per plant, and the ploidy level was calculated as described in Sanchez et al. (2014).

2.4. Cytological preparation and image analysis

Following Gore et al. (2011), we selected representative plants within a ploidy level of the five accessions and conducted mitotic chromosome counts in root tip cells as described in Sanchez et al. (2014). Counts of metaphase chromosomes from images were independently confirmed by two people (authors Paul L. Sanchez and Bernd Friebe).

2.5. Sample preparation and analysis of nuclear genome size with flow cytometry

Leaf tissue samples were collected and prepared for genome size analysis from 3, 4, or 6 plants each of the five *Parthenium* accessions, for a total of 22 plants. This analysis was performed as part of an experiment that had also included 12 additional guayule, mariola, interspecific hybrid, sunflower, and lettuce accessions (Sanchez et al., 2014) following procedures described therein.

2.6. Sample collection, preparation, and DNA sequencing

Leaf tissue samples consisting of multiple young leaves for each plant were collected from 62 single plants and the tissue was lyophilized for long-term storage. For each plant sample, 17 mg to 22 mg of lyophilized leaf tissue was ground separately with a Talboys High Throughput Homogenizer (Troemner LLC, Thorofare, NJ, USA) and total genomic DNA was isolated from the

127 powdered lyophilized leaf tissue using the DNeasy Plant 96 Kit (Qiagen In-
 128 corporated, Valencia, CA, USA). The isolated genomic DNA was prepared ac-
 129 cording to sequencing center instructions ([http://www.biotech.cornell.edu/brc/](http://www.biotech.cornell.edu/brc/genomic-diversity-facility/gbs-sample-submission)
 130 [genomic-diversity-facility/gbs-sample-submission](http://www.biotech.cornell.edu/brc/genomic-diversity-facility/gbs-sample-submission)). Briefly, to ensure that the
 131 isolated total genomic DNA samples were neither contaminated nor degraded,
 132 100 ng of each sample was run on a 1 % agarose gel with 500 ng of two λ HindIII
 133 size/mass standards. Trial digestions with the EcoRI restriction enzyme were
 134 performed on 10 % of the samples, and sample concentration was quantified
 135 using the Quant-iT PicoGreen dsDNA Assay Kit (Life Technologies, Carlsbad,
 136 CA, USA). The DNA samples for the 62 plants were arrayed on a 96 well plate
 137 into 83 sequencing samples as follows: i) 42 plant samples were placed in one
 138 well each using 750 ng of DNA per well, producing 42 sequencing samples; ii) 17
 139 plant samples were placed in two distinct wells each as technical replicates using
 140 750 ng of DNA per well, producing 34 sequencing samples; iii) 2 plant samples
 141 were placed in two distinct wells each as technical replicates using 750 ng of DNA
 142 for the first well and 625 ng of DNA for the second well, producing 4 sequencing
 143 samples; iv) 1 plant sample was placed in three distinct wells as technical repli-
 144 cates using 750 ng of DNA per well, producing 3 sequencing samples. One well
 145 was left blank as a negative control, and the remaining 12 wells were populated
 146 with *Parthenium* samples not included in this study.

147 The DNA samples were sequenced at the Cornell Genomic Diversity Facil-
 148 ity (Cornell University, Ithaca, NY, USA) using the GBS procedure of Elshire
 149 et al. (2011) with minor modifications. Briefly, for each sequencing sample total
 150 genomic DNA was digested with the EcoT22I restriction enzyme and Illumina
 151 Read 1 sequencing primers plus a barcode adapter were then ligated to the
 152 fragment ends. After adapter ligation, GBS libraries were generated by PCR
 153 amplification, and the resulting libraries were sequenced on a single Illumina
 154 HiSeq 2000 flow cell (Illumina Incorporated, San Diego, CA, USA).

155 2.7. GBS data processing

156 All raw data processing and downstream analysis of GBS sequences, includ-
157 ing de-novo reference assembly, SNP calling, genotyping, heterozygosity level
158 estimation, and phylogenetic inference, was performed in a UNIX environment
159 (Linux Mint 17; www.linuxmint.com). Custom scripts for Bash (version 4.3.11)
160 and Perl (version 5.18.2) were used unless otherwise specified. FASTQ format-
161 ted (Cock et al., 2009) raw sequence data consisting of 101 bp long nucleotide
162 sequences (reads) and corresponding quality values was processed by assign-
163 ing each read to a sequencing sample and evaluating reads based on nucleotide
164 sequence quality, content, and frequency. Only reads that were assigned to a
165 specific sample and passed all of the evaluation steps (hereafter ‘filtered reads’)
166 were used in subsequent analysis. Sequence evaluation criteria were optimized
167 for de-novo reference assembly and SNP genotyping.

168 Raw sequence data was first demultiplexed following standard procedures
169 (AppendixC.8). Due to variation in the length of the sample-specific sequence
170 tags, the resulting reads ranged in length from 86 to 92 bp and were further
171 trimmed to a maximum length of 86 bp by removing nucleotides from the end
172 of the read as necessary. For each trimmed read, the corresponding quality
173 values were trimmed as well. The sequencing quality of each read assigned to a
174 sequencing sample was evaluated, and only reads that met all of the following
175 criteria were kept for further analysis: i) the quality value for the last base of
176 the read must not be the special Illumina code ‘2’, corresponding to unreliable
177 nucleotide assignments at the end of the read; ii) the read must not contain any
178 nucleotides encoded as ‘N’, corresponding to unknown nucleotides; iii) the read
179 must not contain homopolymers (i.e., sequence of identical bases) longer than 9
180 bp; iv) the average quality value for the read must be ≥ 30 , corresponding to an
181 average probability of erroneous base call per nucleotide of ≤ 0.001 ; and v) the
182 read must not contain low complexity regions longer than 25 % of the read length
183 based on the empirical distribution. Low complexity regions were identified
184 with the DustMasker (Morgulis et al., 2006) software component of the National
185 Center for Biotechnology Information (NCBI) C++ Toolkit (version 12.0.0; ftp:

186 //ftp.ncbi.nih.gov/toolbox/ncbi_tools++/CURRENT/) using a window length
187 of 86 bp, a threshold level of 20, and a linker value of 1.

188 The reads that met all of the sequencing quality criteria above were further
189 evaluated for sequence content and frequency of sampling in order to identify
190 and eliminate potential contamination, sequencing error, and reads derived from
191 multiple genomic loci (e.g. transposable elements). A FASTA format (Lipman
192 and Pearson, 1985) file containing the subset of distinct nucleotide sequences
193 was generated, and for each distinct sequence the number of times it was ob-
194 served (read depth) was calculated by counting the number of reads having
195 that particular sequence. In order to reduce the possibility of sequencing error
196 (AppendixC.7) a minimum read depth of two, corresponding to two indepen-
197 dent reads confirming the sequence, was required for each distinct sequence,
198 and any distinct sequences that did not satisfy this criterion were eliminated
199 from further analysis. Distinct sequences with a read depth of two or higher
200 were screened against the nucleotide sequences in the NCBI UniVec dataset
201 (build 8; <http://www.ncbi.nlm.nih.gov/tools/vecscreen/univec>), which among
202 other vector sequences contains entries for the specific PCR vectors used in
203 the GBS library preparation protocol (UniVec identifiers ‘gnl|uv|NGB00362.1:1-
204 61’ and ‘gnl|uv|NGB00360.1:1-58’). They were further screened against the
205 NCBI artificial vector sequence dataset (taxon id 29278; [http://www.ncbi.nlm.](http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=29278)
206 [nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=29278](http://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?mode=Info&id=29278)), the guayule
207 chloroplast sequence (NCBI gi 281190713; Kumar et al., 2009), and the “Rep-
208 resentative Genomes” subset of the NCBI Microbial Genomes database (build
209 of May 28, 2014; [ftp://ftp.ncbi.nlm.nih.gov/blast/db/Representative_Genomes.](ftp://ftp.ncbi.nlm.nih.gov/blast/db/Representative_Genomes.*tar.gz)
210 [*tar.gz](ftp://ftp.ncbi.nlm.nih.gov/blast/db/Representative_Genomes.*tar.gz)). The screening step was done using the blastn component of BLAST+
211 (version 2.2.28; Camacho et al., 2009) with the ‘blastn-short’ task preset, and a
212 successful match required 90 % identity over the matched region and an E-value
213 of 0.01 or less. All distinct sequences matching any of these target data sets were
214 removed from further analysis. Finally, we identified in the remaining data set
215 the top 0.01 % most frequently observed distinct sequences and removed them
216 as potential multi-locus genomic sequences.

217 2.8. *De-novo genomic reference assembly*

218 In order to facilitate SNP identification, a de-novo guayule pseudo-reference
 219 was created using the two plant samples from the diploid guayule accession
 220 W6 429. All the filtered reads from these two samples were combined and,
 221 following Ilut et al. (2014), distinct sequences were aligned to each other using
 222 the global alignment algorithm implemented by SlideSort (version 2; Shimizu
 223 and Tsuda, 2011). Allowing for a maximum pairwise edit distance of 20 and
 224 considering both forward and reverse matches, alignments were generated using
 225 the default scoring matrix and edit distances calculated by SlideSort were further
 226 adjusted as follows: i) potential insertion-deletion polymorphisms (indels) were
 227 encoded as single events by assigning a value of 1 to the cost of an indel and
 228 removing from the calculated edit distance any contribution from the length
 229 of the indel sequence; ii) indel penalties due to global alignment were removed
 230 by modifying the edit distance such that indels at the edges of the sequences
 231 were ignored. Next, for a range of maximum pairwise edit distances (d) from
 232 1 to 20, distinct sequences with adjusted pairwise edit distances less than d
 233 were grouped together into putative locus clusters using transitive clustering.
 234 For each value of d , 90 % or more of the putative locus clusters had four or
 235 fewer distinct sequences. Given that at most four alleles can be expected at a
 236 locus when aligning reads from two diploid samples, putative locus clusters were
 237 further limited to those containing 4 or fewer distinct sequences, and for each
 238 value of d from 1 to 20 the distribution of pairwise distance within each of these
 239 putative locus clusters was evaluated. Regardless of the maximum threshold d ,
 240 80 % or more of the reads within a putative locus cluster differed by 3 or fewer
 241 mismatches, and therefore the set of putative locus clusters corresponding to
 242 $d = 3$ was selected for the de-novo pseudo-reference assembly. For each putative
 243 locus cluster, the distinct sequence with the highest read depth was selected as
 244 a representative nucleotide sequence.

245 2.9. SNP identification and genotype assignment

246 We created a reference index from the de-novo guayule pseudo-reference
247 sequences using the ‘panati-build’ component of the PANATI package (<http://panati.sourceforge.net>; Ilut et al., 2012; Spindel et al., 2013) with a word size
248 parameter of 12. Next, for each plant sample, we aligned all reads from that
249 sample that passed quality control against the reference index and generated
250 genotype calls using the ‘panati’ and ‘combine-samples’ functions of the PA-
251 NATI package. The software parameters used applied the following filters to
252 the input sequences: i) trim edge bases with quality of 30 or less, ii) remove
253 sequences with a probability of error $\geq 1\%$ after trimming, iii) and remove se-
254 quences of < 75 bp length after trimming. The alignment parameters required
255 that at least 80 % of the query sequence was aligned and there were at most 5 %
256 mismatched bases between query and target sequence. The SNP identification
257 parameters required a maximum posterior error probability for a SNP call of
258 10^{-12} and a maximum posterior error probability for a genotype call of 1 %.
259 We retained only biallelic SNPs, excluding from the genotype matrix all indels
260 and SNPs with more than 2 alleles. The top 5 % most polymorphic reference
261 loci, representing loci with more than 10 SNPs per locus (Figure B.4), were also
262 removed from the genotyping process.

264 2.10. Estimation of heterozygosity levels

265 In order to improve confidence in homozygous genotype calls, the genotype
266 matrix was modified such that only genotype calls that had a read depth of at
267 least 10 were considered to be genotyped in a particular sample. A SNP was
268 considered heterozygous in a sample if two alleles were detected, and the level of
269 heterozygosity for a given sample was calculated as the number of heterozygous
270 genotype calls divided by the total number of aligned nucleotides. For polyploid
271 samples, reads from multiple copies of the genome were considered together and
272 the calculated heterozygosity was not modified to account for the ploidy level
273 or relative frequency of allelic loci (gene dosage).

274 2.11. Phylogenetic network analysis

275 For the phylogenetic analysis, the genotype matrix was modified such that
276 only SNPs which had at least 4 reads from a given plant sample aligned to them
277 were considered to be genotyped in that sample, and only SNPs that were geno-
278 typed in at least 70 % of the plant samples were retained. The genotype calls
279 were re-coded as pairs of binomial characters, using 00 for genotypes homozy-
280 gous for the reference allele, 11 for genotypes homozygous for the alternative
281 allele, and 01 for heterozygous genotypes. The re-coded genotype matrix was
282 then converted into a concatenated character matrix, saved as a NEXUS format
283 (Maddison et al., 1997) file, and imported into SplitsTree (version 4.1.1; Huson
284 and Bryant, 2006). A phylogenetic network was computed by the Neighbour-
285 Net (Bryant and Moulton, 2004) distance method. For each pair of samples, the
286 distance metric was computed as the number of characters in the re-coded geno-
287 type matrix at which the two samples differed divided by the total number of
288 characters genotyped in both samples. Variance was estimated via the ordinary
289 least squares method, bootstrap support was calculated using 1000 replicates,
290 and network splits with bootstrap support below 90 % were removed.

291 3. Results and discussion

292 3.1. Sampling rationale, species selection, and genotyping

293 The guayule germplasm accessions available in the NPGS collection span
294 over 70 years of cultivar development and collection from wild populations. We
295 report here for the first time a genetic characterization of plant samples from
296 this collection with SNP markers. Multiple guayule and interspecific guayule
297 hybrid accessions were selected from a temporal transect of cultivar develop-
298 ment and wild collection programs between 1942 and 2008 as a representative
299 sample of the genetic diversity available in the germplasm collection (Figure 1).
300 Additionally, we selected for characterization recent wild collections of mariola,
301 the *Parthenium* species most closely related to guayule (Rollins, 1950; Kumar
302 et al., 2009) and a common guayule interspecific hybridization partner in the

303 wild (Rollins, 1950). Contemporary nursery material of wild quinine, a *Parthe-*
 304 *nium* species distantly related to guayule (Rollins, 1950; Kumar et al., 2009),
 305 was selected as the outgroup for phylogenetic analysis. A subset of these acces-
 306 sions has been previously analyzed for ploidy, chromosome number, and nuclear
 307 genome size (Gore et al., 2011; Sanchez et al., 2014) and this study expands
 308 those analyses to a further five accessions, providing ploidy, chromosome num-
 309 ber, and nuclear genome size information for the accessions that were genetically
 310 characterized with SNP markers (Table 1). From the 62 plant samples analyzed
 311 via GBS, we obtained 48 million reads that passed quality control. Of these
 312 reads, 2.4 million were derived from the two samples of diploid guayule (ac-
 313 cession W6 429), and they were assembled into a de-novo pseudo-reference of
 314 76 991 putative loci, each 86 bp in length. This pseudo-reference was used to
 315 align all 48 million reads and identify 30 051 variable loci with 10 or fewer SNPs
 316 per locus, for a total of 93 558 SNP markers.

317 From this total set of SNP markers and associated genotype calls, we gener-
 318 ated three distinct subsets suited for different measurements and analysis meth-
 319 ods. The first subset, consisting of all genotype calls that had a read depth of
 320 10 or higher, was used for heterozygosity level estimates since the read depth
 321 at each genotype call is critical for the reliability of homozygous genotype infer-
 322 ence. The number of SNP markers with the required depth was variable among
 323 samples (Table A.3), but all 93 558 SNP markers had the required depth in at
 324 least one sample. The second subset, consisting of 19 043 SNP markers that
 325 were genotyped in at least 70 % of the samples with a read depth per sample of
 326 4 or higher, was used for phylogenetic analysis because genotyping each locus
 327 across many samples is crucial to the reliability of inferred relationships among
 328 samples. Finally, the third subset of 3330 high confidence SNP markers, defined
 329 as markers genotyped in at least 70 % of the samples with a read depth of 10 or
 330 higher was not used for analyses in this study but rather generated as a poten-
 331 tial resource for guayule breeders to construct a custom SNP genotyping array
 332 and provided as an online supplemental file.

333 *3.2. Genomic characterization of guayule germplasm*

334 A network of phylogenetic relationships, when combined with site collection
335 and breeding records, serves as a tool to provide evolutionary insights into ge-
336 netic similarities as well as reveal inconsistencies and potential misclassifications
337 in the identity of germplasm bank accessions. Phylogenetic network analysis of
338 the 62 plant samples in this study identified four major groups, with the ma-
339 jority of samples representing guayule cultivars and wild collections clustered
340 in group I (Figure 2). This group can be generally divided into four distinct
341 clades, marked in Figure 2 with the letters A, B, C, and D at the root node of
342 each respective clade.

343 Clade A contains the two plant samples of diploid guayule (W6 429), all the
344 samples from accession 11591, the tetraploid samples of N566, and all the sam-
345 ples of the wild collected accession R1110. It should be noted that accessions
346 11591 and N566 were both derived from the same historic accession (4265-I;
347 Thompson and Ray, 1989), which in turn represented one of three predominant
348 types identified by Johnson (1950) based on morphological and biochemical
349 characteristics of historic accession 4265 (Figure 1). Accession 4265 was derived
350 from a bulk seed collection of five individual plants from eastern Durango, Mex-
351 ico in 1942 (Figure 3). Therefore 4265-I, the accession that gave rise to both
352 11591 and N566 as well as another 13 of the 26 core USDA guayule accessions
353 (Thompson and Ray, 1989), likely captured a very limited number of alleles.
354 The W6 429 accession is likewise derived from a selection obtained during the
355 same collection expedition (1942) and from roughly the same geographical area
356 (Figure 3; Bergner, 1946; Thompson and Ray, 1989). Given the historical origins
357 of these three accessions their genetic similarity is not surprising, and it is prob-
358 ably representative of the original germplasm collected in 1942 from Durango,
359 Mexico.

360 The fourth accession in clade A, R1110, was collected in the wild in 1979
361 from Mesa, AZ, an area that is not part of the native range for guayule (Figure
362 B.5; Rollins, 1950). However, Arizona has been an area of active guayule cul-
363 tivation and germplasm development by both private and public entities since

the early 20th century (Hammond and Polhamus, 1965). Beginning in 1914, the University of Arizona operated several experimental agricultural stations in Mesa and surrounding areas (University of Arizona Agricultural Experiment Station records. Special Collections, University of Arizona Libraries). Although all activities from these stations were transferred in 1983 to the Maricopa Agricultural Center in Maricopa, AZ, the experimental stations were still active when accession R1110 was collected in 1979. Together with the genetic similarity between this accession and historical guayule breeding lines, this suggests that plants in accession R1110 are derived from historic breeding lines rather than native stands of wild guayule. The plants in this accession were originally identified as a hybrid of guayule and *P. tomentosum* var. *stramonium* (Green) Rollins based on morphological characteristics, but the phylogenetic placement of R1110 in clade A rather than the clade containing artificial hybrids between guayule and *P. tomentosum* var. *stramonium* (clade D, below) does not support this particular hybrid classification.

As previously reported (Sanchez et al., 2014), ploidy level variation exists within the 11591 and N566 accessions, with 11591 containing triploid and pentaploid samples and N566 containing tetraploid and pentaploid samples. Accession 4265, which served as the source of these two accessions, was originally characterized as a triploid (Johnson, 1950), although chromosome counts higher than 54 (the triploid chromosome number for guayule) have been reported for this accession (Gerstel, 1948; Johnson, 1950). In facultatively apomictic plants such as guayule, generation of progeny with increased chromosome numbers can occur when the unreduced megaspore mother cell is fertilized (reviewed in Thompson and Ray, 1989, Table 4.1). Within accession 11591, the low frequency occurring pentaploid sample is genetically very similar to the triploid samples, suggesting a change in ploidy from triploid to pentaploid as a result of fertilization between plants from this or a closely related accession rather than hybridization with a different accession or species. Somewhat surprisingly, the tetraploid samples of accession N566 are genetically distinct from other members of clade A, including 11591, yet the accession was derived from the same

395 historic accession as 11591. Whether the tetraploid samples of N566 are the
396 result of selection of uncharacterized tetraploids from the source germplasm,
397 or the change from triploid to tetraploid occurred later within the accession,
398 the phylogenetic placement suggests that the pollen donor responsible for the
399 ploidy increase was genetically distinct from the plants from which 11591 was
400 derived. Whereas 14 of 15 samples evaluated for ploidy from accession 11591
401 were triploid, the predominant ploidy level for N566 (12 of 15 samples) appears
402 to be pentaploid (Table A.4; Sanchez et al., 2014) and all the pentaploid sam-
403 ples of N566 in this study form a single clade distinct from clade A (Figure
404 2, Clade B), sharing at least 1578 SNP alleles that do not exist in any of the
405 tetraploid samples from this accession. This suggests that pentaploid samples
406 of N566 were the result of hybridization between tetraploid N566 plants (which
407 belong to the A clade) and plants from an accession outside the A clade, a
408 hybridization that also resulted in a change of ploidy. This is consistent with
409 heterozygosity level estimates (Table 1), which show that pentaploid plants of
410 N566 have slightly higher allelic diversity than the pentaploid plant of 11591.

411 Clade C consists of samples from the breeding line AZ-5 developed by the
412 University of Arizona and the USDA (Ray et al., 1999), the wild collected acces-
413 sion R1109 from Coahuila, Mexico, and one individual from accession CFS-21
414 collected in the wild near Fort Stockton, Texas. Within this group, individual
415 plant samples from accession AZ-5 are genetically more similar to plant samples
416 from R1109 than to other AZ-5 plants. Likewise, plants from R1109 are genet-
417 ically more similar to plants from AZ-5. The clade is homogeneous in terms of
418 cytogenetic and genomic features, with all samples being tetraploid and hav-
419 ing very similar nuclear genome sizes and heterozygosity level estimates (Table
420 1). As a whole, clade C is genetically distinct from historic guayule accessions
421 11591, N566, and W6 429 (clade A), containing 759 shared SNP alleles that are
422 not found in any of the samples from clade A. Combined, these features suggest
423 that both accessions were derived from a recent common ancestor, and that the
424 two accessions are genetically indistinguishable.

425 In order to explain how accessions from such disparate sources could be so

426 closely related genetically, it is important to understand the historical context
427 of their collection and maintenance. Accession R1109 was collected in the wild
428 in September 1976 by R. Rollins near the locality of Parras in the southern
429 part of Coahuila, Mexico (Figure 3; Table A.2) and deposited into the National
430 Plant Germplasm System in 1989 by Dr. D. D. Rubis along with 134 other lines
431 from his collection. For AZ-5, the oldest traceable source of germplasm consists
432 of selections made by Rubis from a mix of USDA lines and unknown sources,
433 with open-pollinated seed of these plants subsequently planted at Marana, AZ,
434 in 1982 and 1983 (Figure 3; Dierig, 1987). After Dr. Rubis' retirement, open-
435 pollinated seeds from 6 selected plants in this field were collected in 1986 and
436 planted at Maricopa, AZ, in 1988. Further rounds of single-plant selections of
437 open-pollinated seed and progeny evaluation based on rubber and resin content
438 resulted in the release of six guayule breeding lines (including AZ-5) in 1997
439 (Ray et al., 1999). Given that R1109 was collected in 1976, it is likely that this
440 or another related accession was already maintained by Rubis prior to the 1982
441 and 1983 transplants that eventually resulted in the AZ-5 accession. It should
442 also be noted that accession R1109 had been classified as a hybrid of guayule
443 and mariola based on morphological characteristics. However, the phylogenetic
444 analysis in this study, which included samples of mariola, does not support this
445 hybrid classification.

446 Clade C also contains one of the four plants from accession CFS-21 that were
447 collected in the wild in Pecos County, TX, on October 29, 2008 (Figure 3; Table
448 A.2). Although this location is more than 350 miles (563 km) away from the
449 collection location for R1109 (the other wild collected accession in this clade),
450 previous sampling of guayule specimens (Figure B.5; Rollins, 1950) indicated
451 the presence of guayule plants throughout the intervening region. Therefore,
452 we cannot exclude the possibility that this sample and those from R1109 are
453 part of a large interbreeding population. However, another possible source of
454 this CFS-21 germplasm consists of a former guayule research station originally
455 operated by Texas A&M University in partnership with the Firestone Tire and
456 Rubber Company (and subsequently Bridgestone Corporation after its aquisi-

tion of Firestone Tire and Rubber Company in 1983). The site is located near Fort Stockton, TX, less than 100 miles (161 km) away from the collection locale of CFS-21. Although guayule research at the site was discontinued in the late 1990s, Bridgestone Corporation continues to operate the site and records for samples collected at that location in 2005 describe the field as follows: “*The Firestone Natural Rubber and Latex Company (a division of the Firestone Tire and Rubber Company) established 180 acres of guayule in 1978-1981 by transplants originating from several seed sources. Approximately 15 acres contained 23 USDA and Anderson [accession 593] lines, while 163 acres were devoted to stands derived from bulk seed collected in native populations of north-central Mexico, and from bulk collections at the Los Angeles State and County Arboretum, Arcadia, CA. About 2 acres were devoted to research plots. Firestone closed the site in 1981, and the Texas Agricultural Experiment Station leased the area in 1983 and conducted research there until 1997.*” (NPGS collection records for CFS18-2005 [PARL 805]). The vast majority of plantings appear to have been derived from north-central Mexico collections, likely from the germplasm collected by R.C. Rollins in 1976 (which included accession R1109) and Naqvi & Hanson in 1977 (reviewed in Thompson and Ray, 1989). Thus, it is plausible that plants with an allelic composition similar to those of R1109 were introduced in Texas at this location and subsequently spread into the wild, either displacing or admixing with native guayule plants. Finally, given that we only found one sample from this accession with a genotype similar to AZ-5 and R1109, it is also possible that a sample from one of those two accession was erroneously labeled as CFS-21. Following up on these results, we plan to genotype further samples from accession CFS-21 and other wild collected guayule from the area in a future experiment. However, regardless of the status of this particular sample, the historical presence of north-central Mexican guayule germplasm at the Firestone plantation indicates that such guayule genotypes are expected to be present in the area.

The last major clade in Group I, clade D, consists of triploid and tetraploid samples from accessions CAL-1, CAL-2, and CAL-5. Ploidy level variation was

488 detected among the CAL-5 samples of this clade, and aneuploidy was identified
 489 in all three accessions (Table 1). These accession were generated via interspecific
 490 crosses between diploid guayule germplasm and *P. tomentosum* var *tomentosum*
 491 DC, *P. fruticosum* Less., and *P. tomentosum* var *stramonium* respectively, and
 492 progeny were selected for resistance to *Verticillium* wilt (Estilai, 1985; Tysdal
 493 et al., 1983). The crosses were produced at the Los Angeles State and County
 494 Arboretum using a common diploid guayule maternal parent (Waln et al., 1983)
 495 and therefore many alleles are expected to be shared among them, resulting in
 496 the complex phylogenetic relationships identified in this clade. Compared to
 497 the triploid samples, the three tetraploid samples of these interspecific hybrids
 498 are more closely related to the guayule clades, suggesting an increase in ploidy
 499 resulting from a maternal interspecific hybrid being fertilized by haploid pollen
 500 from the diploid guayule parent. It should be noted that accession CAL-5,
 501 which is the source of all three tetraploid samples, was developed from open-
 502 pollinated seed of early generation, segregating progeny plants that were grown
 503 alongside the diploid parents for 3 to 4 years prior to seed selection, providing
 504 ample opportunity for further crossing with the diploid parent.

505 3.3. Genomic diversity among wild mariola samples

506 The mariola germplasm sampled for this study (Figure 2, Group III) was
 507 all collected at the same time and from the same geographical region (Figure
 508 3). With the exception of one plant (CFS01-2005 pentaploid in Group II),
 509 the phylogenetic analysis places all of these samples into a single clade, with
 510 samples from the same location (same germplasm identifier) and of the same
 511 ploidy level forming monophyletic groups. All the samples analyzed are likely
 512 pure mariola accessions, and the phylogenetic and collection data suggest that
 513 they are part of a large interbreeding population of wild mariola growing in
 514 southwest Texas. Although aneuploidy and variation in ploidy levels were both
 515 detected among these samples (Table 1), close phylogenetic relationships of
 516 samples with different ploidy levels (Figure 2, Group III) and high similarity in
 517 heterozygosity levels across ploidy levels (Table 1) suggest that ploidy changes

518 resulted from the fertilization of unreduced megaspore mother cells by pollen of
519 mariola plants from the same population. The nuclear genome size of mariola
520 appears to be slightly larger than that of guayule at equivalent ploidy levels
521 (Table 1). Furthermore, we detected a small but significant (t-test, $P < 10^{-6}$)
522 difference between the average chromosome size of mariola (mean of 97.98 Mb /
523 chromosome) and guayule (mean of 88.19 Mb / chromosome) when considering
524 samples with exact chromosome counts (Table A.3; Figure B.6).

525 3.4. *Cryptic Parthenium interspecific hybrids*

526 Unlike traditional bifurcating phylogenetic trees, a phylogenetic network
527 presents all conflicting phylogenetic relationships concurrently, highlighting pu-
528 tative hybrid samples by their well-supported placement as sister to two or
529 more other taxa. Several samples (Figure 2, Group II) were thus identified as
530 containing such hybrid genotypes. Four samples, previously classified as wild-
531 collected guayule (CFS-21) or wild-collected mariola (CFS01-2005), appear to
532 be hybrids between guayule and mariola. Given the sympatric distribution of
533 guayule and mariola in the wild (Figure B.5; Rollins, 1950) it is not surpris-
534 ing that wild-collected samples of guayule and mariola contain morphologically
535 cryptic hybrids.

536 In addition, four samples from interspecific guayule hybrid lines (CAL-1,
537 CAL-2) appear to be the result of further hybridization with an unknown lin-
538 eage, resulting in an increased ploidy level from triploid to pentaploid (Table 1,
539 Figure 2). This suggests that hybridization and corresponding increase in ploidy
540 level occurred via fertilization of an unreduced megaspore mother cell of the
541 interspecific CAL-1 and CAL-2 hybrids by pollen from a triploid or tetraploid
542 plant of unknown species. Estimates of heterozygosity levels (Table 1) show that
543 these four pentaploid samples have the highest levels of any samples evaluated
544 in this study, further supporting the hypothesis of a subsequent hybridization
545 event.

546 3.5. Nuclear genome size and allelic divergence of wild quinine

547 The wild quinine accession was obtained from a commercial nursery and
548 represents a member of the *Parthenium* genus that is not limited to arid lands,
549 but is found throughout the Eastern half of the United States (Figure B.5;
550 Rollins, 1950). The long private branch of this clade in the phylogenetic analysis
551 (Figure 2) indicates that it is genetically quite divergent from both guayule and
552 mariola. The haploid nuclear genome size for tetraploid *P. integrifolium* is 5757
553 Mb, much larger than that of guayule (3071 Mb) or mariola (3433 Mb) plants
554 with the same number of chromosomes (Table A.3, Figure B.6 A). Although its
555 adaptation to colder climates presents it as a tempting candidate for improving
556 the cold resistance of guayule germplasm, the large divergence and difference in
557 nuclear genome size suggest successful hybridization between this species and
558 guayule would be difficult.

559 3.6. Genomic diversity within the germplasm collection

560 Our results show that ploidy variation in several accessions is closely related
561 to genetic differentiation between ploidy levels, suggesting further separation
562 of germplasm from these accessions in order to maintain genetically consistent
563 cultivars. In contrast with the genetic heterogeneity observed in some of the
564 accessions, other distinct accessions show high genetic similarity, indicating that
565 further genomic characterization of the extant wild and cultivated *Parthenium*
566 germplasm pool is necessary to construct a genetically representative diversity
567 panel for genome-wide association studies in guayule. Such a panel would sig-
568 nificantly reduce redundancy in effort and maximize potential payoffs for future
569 genetic studies and crop improvement efforts in this species. Specifically, SNP
570 genotyping of other accessions derived from 4265-I should help elucidate the
571 historic gene flow among these closely related accessions and identify the most
572 genetically diverse subset. Similarly, the SNP genotyping of other accessions
573 from Rubis' collection would further refine the pedigree of AZ-5 and other cul-
574 tivars derived from the 1982 single-plant selections and help identify sources of
575 novel alleles from wild collections subsequent to the Emergency Rubber Project

576 of the 1940s. Genotyping of other *Parthenium* species with SNP markers, in-
 577 cluding the non-guayule parents of the CAL-1, CAL-2, and CAL-5, would help
 578 identify the unknown species involved in generating the pentaploid samples from
 579 these accessions as well as quantify the genetic contribution of guayule and other
 580 species in these interspecific hybrids. Finally, in order to assess the potential for
 581 novel germplasm present in wild populations of guayule from the Big Bend area
 582 of Texas, SNP genotyping of wild collections of guayule from this area as well
 583 as accessions derived from the Rollins and Naqvi & Hanson collections should
 584 help elucidate these relationships and assess the degree of admixture between
 585 introduced guayule plants from north-central Mexico and native Texas plants.

586 **4. Conclusions**

587 This study represents the first use of GBS for the simultaneous discovery
 588 and scoring of SNP markers in *Parthenium* germplasm, and the first nuclear
 589 genome size estimation for wild quinine, a *Parthenium* species with a haploid
 590 nuclear genome size almost twice as large as guayule. We have characterized the
 591 genomic diversity of several accessions of guayule germplasm and other *Parthe-*
 592 *nium* species, identified the 1976 guayule collections by R. C. Rollins from the
 593 southern part of Coahuila, Mexico as the likely source of the AZ-5 cultivar,
 594 and have shown that they provided novel alleles compared to McCallum's 1942
 595 collections from central Durango, Mexico from which most classic cultivars of
 596 guayule are derived. We identified several plants of unknown hybrid parent-
 597 age that were sampled from CAL-1 and CAL-2 cultivars, as well as sampled
 598 mariola/guayule hybrid plants from among the wild collections of these species.
 599 Additionally, we demonstrated the utility of GBS for the phylogenetic classifi-
 600 cation of *Parthenium* accessions even in the absence of a reference genome se-
 601 quence. Finally, we have shown the need for further SNP genotyping of guayule
 602 germplasm in order to optimize genetic studies and breeding efforts in guayule.

603 **5. Data availability**

604 All sequence data has been deposited under BioProject PRJNA289254 in
605 the NCBI Sequence Read Archive (SRA; Wheeler et al., 2008). For each plant
606 sample, the corresponding SRA accession number is indicated in Table A.3.
607 Nucleotide sequences for all the de-novo assembled reference loci, unfiltered
608 genotype calls, and locus sequences and SNPs for the high confidence SNP
609 dataset are provided as an online supplemental data set.

610 **Acknowledgements**

611 We thank the National Arid Land Plant Genetics Resources Unit at Parlier,
612 CA, USA for providing seeds. Also, we wish to thank Greg Leake, Stephanie
613 Johnson, and Kristen Cox for assistance in maintaining plants in the greenhouse
614 and collecting leaf tissue for flow cytometry experiments. Preparation of leaf
615 tissue samples for nuclear genome size analysis was carried out by Patrick Red-
616 mond and Michael Boerman at the Institute for Genomic Diversity at Cornell
617 University, and instrument coordination by Lavanya Sayam at the Biomedical
618 Sciences Flow Cytometry Core Laboratory at Cornell University. Preparation of
619 total genomic DNA samples for GBS was carried out by Catherine B. Kandianis
620 and Elodie E. Gazave in the Gore laboratory at Cornell University. This work
621 was funded and supported by the USDA-ARS and USDA-NIFA/DOE Biomass
622 Research and Development Initiative (BRDI) Grant No. 2012-10006. Mention
623 of trade names or commercial products in this publication is solely for the pur-
624 pose of providing specific information and does not imply recommendation or
625 endorsement by the United States Department of Agriculture. The USDA is an
626 equal opportunity provider and employer.

627 **References**

628 Arumuganathan, K., Earle, E. D., 1991. Estimation of nuclear DNA content of
629 plants by flow cytometry. *Plant Molecular Biology Reporter* 9 (3), 229–241.

631 Bergner, D. A., 1946. Polyploidy and aneuploidy in guayule. United States
632 Department of Agriculture Technical Bulletin (918). 5, 14

633 Bryant, D., Moulton, V., 2004. Neighbor-Net: an agglomerative method for
634 the construction of phylogenetic networks. *Molecular Biology and Evolution*
635 21 (2), 255–265. 12

636 Bullard, W. E., 1946. Climate and guayule culture. United States Department
637 of Agriculture, unpublished report. 5, 33

638 Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer,
639 K., Madden, T. L., 2009. BLAST+: architecture and applications. *BMC*
640 *Bioinformatics* 10, 421. 9

641 Cock, P. J. A., Fields, C. J., Goto, N., Heuer, M. L., Rice, P. M., 2009.
642 The Sanger FASTQ file format for sequences with quality scores, and the
643 Solexa/Illumina FASTQ variants. *Nucleic Acids Research* 38 (6), 1767–1771.
644 8

645 Costich, D. E., Friebe, B., Sheehan, M. J., Casler, M. D., Buckler, E. S., 2010.
646 Genome-size variation in switchgrass (*Panicum virgatum*): flow cytometry
647 and cytology reveal rampant aneuploidy. *The Plant Genome Journal* 3 (3),
648 130–141. 35

649 Dierig, D. A., 1987. Contribution of yield components to rubber production in
650 guayule. Doctoral dissertation, University of Arizona. 17

651 Doležel, J., Greilhuber, J., Suda, J., 2007. Estimation of nuclear DNA content
652 in plants using flow cytometry. *Nature Protocols* 2 (9), 2233–2244. 35

653 Elshire, R. J., Glaubitz, J. C., Sun, Q., Poland, J. A., Kawamoto, K., Buckler,
654 E. S., Mitchell, S. E., 2011. A robust, simple genotyping-by-sequencing (GBS)
655 approach for high diversity species. *PLOS ONE* 6 (5), e19379. 4, 7

656 Estilai, A., 1985. Registration of Cal-5 guayule germplasm. *Crop Science* 25 (2),
657 369. 19

- 658 Gerstel, D. U., 1948. Mid-year report of the genetics section, rubber plant in-
659 vestigations. United States Department of Agriculture, unpublished report.
660 15
- 661 Gore, M. A., Coyle, G., Friebe, B., Coffelt, T. A., Salvucci, M. E., 2011. Complex
662 ploidy level variation in guayule breeding programs. *Crop Science* 51 (1), 210–
663 216. 3, 4, 6, 13, 34, 36
- 664 Hammond, B. L., Polhamus, L. G., 1965. Research on guayule (*Parthenium*
665 *argentatum*), 1942-1959. United States Department of Agriculture. 3, 5, 15,
666 33
- 667 Hashemi, A., Estilai, A., Naqvi, H. H., 1989. Intersectional hybrids between
668 induced tetraploid guayule (*Parthenium argentatum*) and natural tetraploid
669 *P. intergrifolium* (Asteraceae). *Botanical Gazette* 150 (1), pp. 78–83. 4
- 670 Huson, D. H., Bryant, D., 2006. Application of phylogenetic networks in evolu-
671 tionary studies. *Molecular Biology and Evolution* 23 (2), 254–267. 12
- 672 Ilut, D. C., Coate, J. E., Luciano, A. K., Owens, T. G., May, G. D., Farmer, A.,
673 Doyle, J. J., 2012. A comparative transcriptomic study of an allotetraploid
674 and its diploid progenitors illustrates the unique advantages and challenges
675 of RNA-seq in plant species. *American Journal of Botany* 99 (2), 383–396. 11
- 676 Ilut, D. C., Nydam, M. L., Hare, M. P., 2014. Defining loci in restriction-based
677 reduced representation genomic data from nonmodel species: sources of bias
678 and diagnostics for optimal clustering. *BioMed Research International* 2014,
679 e675158. 10
- 680 Johnson, B. L., 1950. High rubber yielding selections from a natural population
681 of guayule. *Agronomy Journal* 42 (7), 345–350. 5, 14, 15
- 682 Kumar, S., Hahn, F. M., McMahan, C. M., Cornish, K., Whalen, M. C.,
683 2009. Comparative analysis of the complete sequence of the plastid genome of
684 *Parthenium argentatum* and identification of DNA barcodes to differentiate
685 *Parthenium* species and lines. *BMC Plant Biology* 9 (1), 131. 9, 12, 13

686 Lipman, D. J., Pearson, W. R., 1985. Rapid and sensitive protein similarity
687 searches. *Science* 227 (4693), 1435–1441. 9

688 Loureiro, J., Rodriguez, E., Doležel, J., Santos, C., 2007. Two new nuclear
689 isolation buffers for plant DNA flow cytometry: a test with 37 species. *Annals*
690 *of Botany* 100 (4), 875–888. 34, 36

691 Maddison, D. R., Swofford, D. L., Maddison, W. P., 1997. Nexus: an extensible
692 file format for systematic information. *Systematic Biology* 46 (4), 590–621.
693 12

694 Mann, C. C., 2009. Addicted to rubber. *Science* 325 (5940), 564–566. 3

695 Mears, J. A., 1975. The taxonomy of *Parthenium* section *Partheniastrum* DC.
696 (*Asteraceae-Ambrosiinae*). *Phytologia* 31, 463–482. 4, 30

697 Morgulis, A., Gertz, E. M., Schäffer, A. A., Agarwala, R., 2006. A fast and
698 symmetric DUST implementation to mask low-complexity DNA sequences.
699 *Journal of Computational Biology* 13 (5), 1028–1040. 8

700 Powers, L., 1942. Seed collection of guayule from Mexico and the Trans-Pecos
701 area (Big Bend Country) of Texas. United States Department of Agriculture,
702 unpublished report. 5

703 Powers, L., Rollins, R. C., 1945. Reproduction and pollination studies on
704 guayule, *Parthenium argentatum* Gray and *P. incanum* H. B. K. *Agronomy*
705 *Journal* 37 (2), 96–112. 4

706 Ray, D. T., Coffelt, T. A., Dierig, D. A., 2005. Breeding guayule for commercial
707 production. *Industrial Crops and Products* 22 (1), 15–25. 3

708 Ray, D. T., Dierig, D. A., Thompson, A. E., Coffelt, T. A., 1999. Registration of
709 six guayule germplasms with high yielding ability. *Crop Science* 39 (1), 300.
710 16, 17

- 711 Ray, D. T., Foster, M. A., Coffelt, T. A., McMahan, C., 2010. Guayule: culture,
712 breeding and rubber production. In: Singh, B. P. (Ed.), Industrial Crops and
713 Uses. CABI, pp. 384–410. 3
- 714 Rivano, F., 1997. South american leaf blight of Hevea I. Variability of *Mi-*
715 *crocyclus ulei* pathogenicity. Plantations, Recherche, Développement 4 (2),
716 104–114. 3
- 717 Rollins, R. C., 1946. Interspecific hybridization in Parthenium II. Crosses in-
718 volving *P. argentatum*, *P. incanum*, *P. stramonium*, *P. tomentosum* and *P.*
719 *hysterophorus*. American Journal of Botany 33 (1), 21–30. 4
- 720 Rollins, R. C., 1950. The guayule rubber plant and its relatives. Contributions
721 from the Gray Herbarium of Harvard University 172, 1–72. 4, 5, 12, 13, 14,
722 17, 20, 21, 33
- 723 Sanchez, P. L., Costich, D. E., Friebe, B., Coffelt, T. A., Jenks, M. A., Gore,
724 M. A., 2014. Genome size variation in guayule and mariola: fundamental
725 descriptors for polyploid plant taxa. Industrial Crops and Products 54, 1–5.
726 4, 5, 6, 13, 15, 16, 30, 34, 36, 37
- 727 Shimizu, K., Tsuda, K., 2011. SlideSort: all pairs similarity search for short
728 reads. Bioinformatics 27 (4), 464–470. 10
- 729 Spindel, J., Wright, M., Chen, C., Cobb, J., Gage, J., Harrington, S., Lorieux,
730 M., Ahmadi, N., McCouch, S., 2013. Bridging the genotyping gap: using geno-
731 typing by sequencing (GBS) to add high-density SNP markers and new value
732 to traditional bi-parental mapping and breeding populations. Theoretical and
733 Applied Genetics 126 (11), 2699–2716. 11
- 734 Thompson, A. E., Ray, D. T., 1989. Breeding guayule. In: Janick, J. (Ed.),
735 Plant Breeding Reviews. John Wiley & Sons, Inc., pp. 93–165. 3, 4, 14, 15,
736 18

737 Tysdal, H. M., Estilai, A., Siddiqui, I. A., Knowles, P. F., 1983. Registration of
 738 four guayule germplasms (Reg. No. GP1 to GP4). *Crop Science* 23 (1), 189.
 739 19

740 Waln, K., Toukdarian, K., Elder, N., Hanson, G., 1983. Interspecific hybridiza-
 741 tion. *Proceedings of the 3rd International Guayule Conference*, 317–326. 19

742 Wheeler, D. L., Barrett, T., Benson, D. A., Bryant, S. H., Canese, K.,
 743 Chetvernin, V., Church, D. M., Dicuccio, M., Edgar, R., Federhen, S., Feolo,
 744 M., Geer, L. Y., Helmberg, W., Kapustin, Y., Khovayko, O., Landsman, D.,
 745 Lipman, D. J., Madden, T. L., Maglott, D. R., Miller, V., Ostell, J., Pruitt,
 746 K. D., Schuler, G. D., Shumway, M., Sequeira, E., Sherry, S. T., Sirotkin, K.,
 747 Souvorov, A., Starchenko, G., Tatusov, R. L., Tatusova, T. A., Wagner, L.,
 748 Yaschenko, E., 2008. Database resources of the National Center for Biotech-
 749 nology Information. *Nucleic Acids Research* 36 (Database issue), D13–21. 23

Table 1: Phylogenetic group, ploidy level, chromosome number, nuclear genome size, and heterozygosity estimates for 15 *Parthenium* accessions. Standard deviation in measurements is denoted by $\pm$. For accession W6 429, the heterozygosity was estimated using two plants, whereas nuclear genome size and ploidy values in this table were estimated using one plant each. The assigned ploidy number is based on flow cytometry estimates from a larger sample for each accession (Table A.4). Ploidy level for the *P. integrifolium* sample was assigned based on chromosome counts and literature (Mears, 1975).

CLADE	SAMPLE INFORMATION			CYTOGENETIC ANALYSIS		GENOMIC ANALYSIS				
	SPECIES RECORD	OF	ACCESSION NAME	ASSIGNED PLOIDY	NO. OF PLANTS	CHROMOSOME COUNTS	NO. OF PLANTS	HAPLOID NUCLEAR GENOME SIZE (Mb)	ESTIMATED PLOIDY	HETEROZYGOSITY (%)
I-A	<i>P. argentatum</i>		W6 429 ^e	2 <i>x</i>	1	2 <i>n</i> = 36	1,2	1624	2.00	0.53 ± 0.01
	Hybrid ^a		R1110	4 <i>x</i>	1	2 <i>n</i> = 72	3	3047 ± 73	3.85±0.08	0.99 ± 0.12
	<i>P. argentatum</i>		11591 ^e	3 <i>x</i>	1	2 <i>n</i> = 54	3	2435 ± 20	3.04±0.01	0.98 ± 0.10
	<i>P. argentatum</i>		11591 ^e	5 <i>x</i>			1	3858	4.92	1.33
	<i>P. argentatum</i>		N566 ^e	4 <i>x</i>	1	2 <i>n</i> = 72	3	3071 ± 38	3.92±0.03	1.11 ± 0.10
I-B	<i>P. argentatum</i>		N566 ^e	5 <i>x</i>	1	2 <i>n</i> = 90	3	3939 ± 26	4.99±0.04	1.45 ± 0.05
I-C	<i>P. argentatum</i>		AZ-5 ^e	4 <i>x</i>	1	2 <i>n</i> = 72	3	3205 ± 29	4.01±0.02	1.58 ± 0.03
	<i>P. argentatum</i>		CFS-21 ^e	4 <i>x</i>			1	3237	3.98	1.56
	Hybrid ^b		R1109 ^e	4 <i>x</i>	2	2 <i>n</i> = 70, 73 ^g	3	3179 ± 23	4.01±0.01	1.54 ± 0.06
I-D	Hybrid ^c		CAL-1	3 <i>x</i>	2	2 <i>n</i> = 55 ^g	3	2533 ± 39	3.05±0.01	1.48 ± 0.05
	Hybrid ^d		CAL-2	3 <i>x</i>	2	2 <i>n</i> = 55, 56 ^g	3	2580 ± 17	3.02±0.03	1.36 ± 0.02
	Hybrid ^a		CAL-5	3 <i>x</i>	2	2 <i>n</i> = 55 ^g	3	2521 ± 22	3.04±0.01	1.48 ± 0.15
	Hybrid ^a		CAL-5	4 <i>x</i>	1	2 <i>n</i> = 74 ^g	3	3169 ± 71	4.01±0.03	1.60 ± 0.02
II	Hybrid ^c		CAL-1	5 <i>x</i>	1	2 <i>n</i> = 90	3	4223 ± 155	4.95±0.02	2.24 ± 0.02
	Hybrid ^d		CAL-2	5 <i>x</i>	1	2 <i>n</i> = 90	1	4201	4.95	2.17
	<i>P. argentatum</i>		CFS-21 ^e	4 <i>x</i>			2	3190 ± 33	3.98±0.03	1.48 ± 0.01
	<i>P. argentatum</i>		CFS-21 ^e	6 <i>x</i>	1	2 <i>n</i> = 108	1	4812	5.88	1.37
	<i>P. incanum</i>		CFS01-2005 ^e	5 <i>x</i>	1	2 <i>n</i> = 90	1	4181	4.92	1.79
III	<i>P. incanum</i>		CFS11-2005 ^e	3 <i>x</i>	1	2 <i>n</i> = 55 ^g	3	2616 ± 77	3.12±0.05	0.98 ± 0.04
	<i>P. incanum</i>		CFS11-2005 ^e	5 <i>x</i>	1	2 <i>n</i> = 84 ^g	1	3990	4.70	0.99
	<i>P. incanum</i>		CFS01-2005 ^e	4 <i>x</i>			3	3472 ± 62	4.05±0.04	1.08 ± 0.22
	<i>P. incanum</i>		CFS04-2005 ^e	3 <i>x</i>	1	2 <i>n</i> = 54	3	2704 ± 15	3.11±0.02	1.03 ± 0.13
	<i>P. incanum</i>		CFS04-2005 ^e	4 <i>x</i>	1	2 <i>n</i> = 76 ^g	1	3751	4.40	0.94
	<i>P. incanum</i>		CFS06-2005 ^e	3 <i>x</i>	1	2 <i>n</i> = 55 ^g	3	2680 ± 68	3.14±0.01	1.01 ± 0.14
	<i>P. incanum</i>		CFS06-2005 ^e	4 <i>x</i>	1	2 <i>n</i> = 72	3	3433 ± 26	4.13±0.03	1.09 ± 0.08
IV	<i>P. integrifolium</i>		<i>P. integrifolium</i> ^f	4 <i>x</i>	1	2 <i>n</i> = 72	3	5757 ± 96	6.56±0.04	1.28 ± 0.23

^a Interspecific hybrid of *P. argentatum* A. Gray and *P. tomentosum* var. *stramonium* (Green) Rollins

^b Interspecific hybrid of *P. argentatum* A. Gray and *P. incanum* Kunth

^c Interspecific hybrid of *P. argentatum* A. Gray and *P. tomentosum* var. *tomentosum* DC

^d Interspecific hybrid of *P. argentatum* A. Gray and *P. fruticosum* Less.

^e Nuclear genome size and ploidy from Sanchez et al. (2014)

^f Unofficial accession designation used in this study

^g Aneuploid plant(s)

751 **Figures**

Figure 1: Sources for *Parthenium* accessions used in this study. Programs responsible for maintaining the germplasm are indicated by the rectangular blue boxes, and labeled at the top of the figure. The blue horizontal ribbon represents the time axis and indicates the year of accession collection, selection, or generation. Clouds indicate unspecified sources of accessions. The asterisk indicates cryptic hybrids within accessions that were identified by the current study.

Figure 2: Phylogenetic network of 62 plant samples from 15 accessions. Each sample is labeled with the accession name and the assigned ploidy level (number in square brackets). Group **I** indicates samples of guayule and known interspecific *Parthenium* hybrids. Group **II** indicates interspecific hybrid samples newly identified in this study. Group **III** indicates mariola samples. Group **IV** indicates wild quinine samples. Capital letters indicate the clades of Group **I** detailed in section 3.2

Figure 3: Geographical location of origin (white circles) and inferred movement of plant material (blue lines) for the guayule (labels with orange border) and mariola (labels with red border) accessions used in this study. The dashed label border indicates plant material of unknown origin. Asterisks indicate historic accessions from which current cultivars are derived as indicated in Figure 1. Cultivars are indicated by angle brackets around the accession name.

Figure 1

Figure 2

Figure 3