Serology in the 21st Century: The Molecular-Level Analysis of the Serum Antibody Repertoire

Yariv Wine¹, Andrew P. Horton^{3,6}, Gregory C. Ippolito⁴ and George Georgiou²⁻⁵

¹ Department of Molecular Microbiology and Biotechnology, Faculty of Life Sciences, Tel Aviv University, Tel Aviv 69978, Israel

² Department of Chemical Engineering, University of Texas at Austin, Austin, Texas, USA.

³ Department of Biomedical Engineering, University of Texas at Austin, Austin, Texas, USA.

⁴ Department of Molecular Biosciences, University of Texas at Austin, Austin, Texas, USA.

⁵ Institute for Cell and Molecular Biology, University of Texas at Austin, Austin, Texas, USA

⁶ Center for Systems and Synthetic Biology, University of Texas at Austin, Austin, Texas, USA

Abstract

The ensemble of antibodies found in serum and secretions represents the key adaptive component of B-cell mediated humoral immunity. The antibody repertoire is shaped by the historical record of exposure to exogenous factors such as pathogens and vaccines, as well as by endogenous host-intrinsic factors such as genetics, self-antigens, and age. Thanks to very recent technology advancements it is now becoming possible to identify and quantify the individual antibodies comprising the serological repertoire. In parallel, the advent of high throughput methods for antigen and immunosignature discovery opens up unprecedented opportunities to transform our understanding of numerous key questions in adaptive humoral immunity, including the nature and dynamics of serological memory, the role of polyspecific antibodies in health and disease and how protective responses to infections or vaccine challenge arise. Additionally, these technologies also hold great promise for therapeutic antibody and biomarker discovery in a variety of settings

Antibody Serology: Past Achievements

Serology is classically defined as the study of proteins, predominantly antibodies, found in blood and secretions such as saliva. The genesis of serology dates to the end of the 19th century and the pioneering “serum therapy” of Emil von Behring and Paul Ehrlich, followed for decades by elegant studies on the specificity of serological reactions by Karl Landsteiner. But it would take Landsteiner until the twilight of his career to formally demonstrate that an anti-serum does not comprise merely a single antibody but rather a mixture of different antibody populations of unknown complexity[1]. It was another decade before the plasma cell, which is responsible for the secretion of antibodies, was discovered, and then it was only 50 years ago, in 1965 that it was convincingly shown that antibodies are produced by B lymphocytes [2]. This latest discovery coincided with the development of new technologies in protein chemistry and the advent of molecular biology that, together, catalyzed a remarkable pace of progress in the understanding of B cell development and antibody formation. We now know that long-lived plasma cells constitute the (most likely) irreversible end-point of B cell development, show little or no

evidence of proliferation and produce copious amounts of antibodies for years, and quite possibly for decades, in humans [3]. Long-lived plasma cells reside predominantly but not exclusively in the bone marrow, surviving within specialized anatomical niches with the help of anti-apoptotic signals provided by stromal cells [4]. Of note, a fraction of bone marrow plasma cells have been recently reported to lack CD19 expression and to be protected from mobilization and replacement by newly formed antibody producing cells following infection[5], underscoring the heterogeneity of the long-lived compartment of plasma cells and, by extension, the pool of serum immunoglobulins.

The compendium of antibodies produced first by long-lived plasma cells and second by transient waves of short-lived plasma cells or plasmablasts (elicited in response to pathogen, vaccine or autoantigen stimulation) constitutes the two main components of antibody serological immunity. A third component is contributed by natural antibodies which recognize common pathogen antigens such as galactose- α -1,3-galactose (Anti-Gal) and have an innate-like protective function [6,7]. The relative contribution of antibodies from long-lived plasma cells, transient plasmablasts and by “natural antibody”-producing cells (rodent B1 and marginal zone B cells, and perhaps the human analogues of B1 cells, although these have yet to be definitively identified) to the serological immunity is poorly understood. The antibody concentration in blood is homeostatically controlled. For example, the Immunoglobulin G (IgG) concentration in healthy adults is maintained at roughly between 7-17 mg/ml. The level of IgG produced by long-lived plasma cells changes very slowly over time [8]. Plasmablast bursts following vaccine or infection are accompanied by a dramatic increase in the antigen-specific antibody titer but generally seem to have little impact on the total concentration of immunoglobulin in blood. It is noteworthy that the fate of the large number of plasmablasts present in circulation at the peak of the plasmablast wave has never been quantitatively followed although it is well established that the majority apoptose and only a relatively small fraction mature to long lived plasma cells and are able to take residence in niches so that they can contribute to serological memory[9]. Finally, the fraction of natural antibodies in blood or secretions is not known, nor is it known how the total level of natural antibody (comprising mostly IgM but also some IgG and IgA) vary as a function of time in health or in inflammation.

Antibody Serology: Future Directions

The essence of serological immunity is predicated on the existence of a diverse repertoire of antibodies, elicited over the life of the host and representing the integrated response to numerous antigenic stimuli. Due to the complexity and temporally dynamic nature of the antibody repertoire, the identification of its component immunoglobulins represents a formidable challenge. All serological studies to date have relied on the detection of an *ensemble* of antibodies that either could be resolved by a certain analytical technique or bound to a specified antigen (**Fig. 1**). Among the most useful metrics for assessing humoral immunity, the presence of neutralizing antibodies in the serum following vaccination or infection represents the best correlate for vaccine efficacy and for protection during invasive infections [10,11,12]. The limitations imposed by the inability to resolve complex serum antibody mixtures into their constituent clonal representatives and the need to have pre-established the identity of antigens of potential interest have obscured central questions of profound basic and clinical significance, some of which are outlined below:

- First and foremost, there is nearly no information on the number of sequences (clonal diversity), functions and relative concentrations of the individual antibodies in serum. Upon reflection it is surely striking that after over a century of immunology research that the clonality of serum antibodies (number of distinct antibody clonotypes having same IGHV and IGHJ segments and highly homologous CDR-H3 sequences) in blood and in secretions has not been established experimentally even to a first approximation, although a remarkably prescient estimate of antibody diversity was first proposed by Talmage based on theoretical arguments, over half a century ago [13]. Nor do we know anything about how the clonality of serum antibodies changes as a function at extreme ages and disease status, or about the concentration distribution of individual antibodies in biological fluids. Without a detailed description of the identities and relative amounts of the antibodies that comprise the serological repertoire it is not possible to establish precisely how the B cell developmental program shapes protective immunity.

Information on the serological antibody repertoire is particularly pertinent to vaccine design/evaluation and to antibody discovery from patients. In an effort to better delineate the serological response to influenza by taking into account antigenic variation among strains circulating in humans, Fonville et al. sought to establish correlations between serum mediated neutralization against panels of different hemagglutinins and their phylogenetic distance [14]. The authors further developed a visualization tool for describing complex serological data by plotting antibody-mediated immunity as a function of the antigenic relationships among viruses (landscape plot).

For rapidly evolving viral pathogens such as HIV-1 or influenza, immortalization techniques and single cell cloning coupled with high throughput micro-neutralization assays of peripheral plasmablasts or memory B cells have led to the discovery of numerous broadly neutralizing antibodies (bNAbs) displaying neutralization breadth towards many or even nearly all known clinical isolates of a particular virus [15•,16•,17,18•,19]. To provide a first-order approximation of component antibody neutralization specificities in sera, Georgiev et al.[20•] proposed matching bNAb and serological neutralization potency fingerprints towards HIV-1 isolates.

Identification of bNAbs coupled with the reconstruction of the evolutionary route that led to the development of phylogenetically- related antibodies from an un-mutated common ancestor (UCA) is essential for the design of vaccine immunogens with the potential of eliciting broad protection. However, the fact that a DNA encoding a bNAb has been isolated from peripheral B cells does not mean that the respective antibody is present in serum or that it is produced at physiologically relevant amounts (i.e. at a concentration near the equilibrium dissociation constant such that the antibody can bind antigen). This divergence was described initially by Burnet stating that there is no requirement in the clonal selection theory that every cellular receptor has a corresponding secreted globulin [21]. As a matter of fact, considering that the BCR repertoire diversity in the memory and plasmablast compartments is orders of magnitude greater than that of the serological repertoire [22•,23] it follows that the overwhelming majority of peripheral B cell-encoded antibodies are unlikely to be present in detectable amounts as soluble proteins in blood or secretions and thus could not have contributed to humoral immunity.

- Second, while it is well established that a significant fraction of antibodies display polyreactivity and that these antibodies have important physiological functions in processes such as the clearance of cell debris and in pathogen recognition [24,25], there is a paucity of methods for quantifying and characterizing the polyreactive fraction of the serological response. It is noteworthy that, in addition to natural antibodies, which are typically polyreactive, IgG antibodies encoded by human memory B cells [26] and even bone marrow plasma cells and also bNAbs elicited by anti-viral immune responses in response to HIV-1 infection [27] often recognize multiple antigens. There is a clear need to understand the mechanisms that drive polyreactivity and its implications in health and disease. One possible explanation is that polyreactivity originates from B cells that were not removed from the repertoire during B-cell development. For some pathogens, notably HIV, polyreactivity may confer a selective advantage to pathogen-specific antibodies [26,28].
- Third, in many instances the antigens that are recognized by serum antibodies are not known *a priori*. The significance of identifying antigens, antigen surrogates (i.e. antigen-mimics that can be chemically distinct from the antigens that elicited an antibody response) and immunosignatures [29] for disease diagnosis is being increasingly recognized. Additionally, mapping the serum antigen reactivity profile in a comprehensive manner is key to understanding which environmental exposures play a more dominant role in shaping humoral immunity.

Antigen and Immunosignatures Discovery: Earlier methods for antigen discovery include the use of peptide libraries either immobilized in array format or displayed on M13 phage and serological analysis of recombinant cDNA expression libraries (SEREX), which employs cDNA libraries from tumor tissues for cancer antigen discovery and finally, antigen arrays [29,30,31,32]. More recently advances in library screening methodologies and separately, in high throughput gene and peptide synthesis technologies have led to an explosion of high-resolution methods for the discovery of autoantigens and antigen surrogates. Daugherty and coworkers developed a technology that capitalized on *E.coli* display [33] of very large random peptide libraries for the identification of peptide motifs that bound to antibodies from patient sera. This technique

led to the discovery of a diagnostically -relevant epitope for celiac disease that exhibited a high degree of amino acid identity to deamidated gliadin, a key antigen in celiac disease. In a separate study the same group reported a 7 peptide panel of highly statistical diagnostic significance for preeclampsia [34,35]. Using massively parallel DNA synthesis Larman et al. [36] constructed libraries of >410,000 overlapping 26-mer peptides spanning all ORFs in the human genome on T7 phage. Phage was immunoprecipitated with patient sera and diagnostic immunosignatures were determined by NextGen sequencing. In an alternative approach very high density printed arrays of synthetic random peptide arrays were used for immunosignature discovery and were reported to exhibit higher accuracy than tiled peptide epitope arrays [37,38]. Since peptides may not be able to capture antibodies that bind to post-translational modifications and conformational epitopes Kodadek and coworkers pioneered the use of libraries of unnatural molecules and specifically peptoids (N-substituted oligoglycine polymers) for the discovery of surrogate antigen biomarkers from patient sera [29,39]. Other powerful methods that are currently used for antigen discovery include antigen microarrays [40,41], nucleic acid programmable protein arrays (NAPPAs) produced using *in-vitro* transcription/translation[42], peptide arrays [43] or finally arrays based on a different targets such as DNA, peptides and recombinant/native proteins [44].

Finally, for very complex conformational antigens on proteins that cannot be immobilized on arrays (e.g. integral membrane proteins) or contain complex post-translational modifications or for which a peptide or synthetic surrogate is not available, novel solution phase assays are being developed for diagnostic purposes. For example radio-binding assays (RBA) [45] using *in vitro* transcription/translation RBAs have been used for detection of autoantibodies. In one notable example Joseph and co-workers, studied the association of scleroderma with an humoral immune response to cancer. They found an association of genetic alterations the RNA polymerase III subunit (RPC1) encoding gene (POLR3A) in patients with autoantibodies to RPC1 suggesting that POLR3A mutations triggered cellular immunity and cross-reactive humoral immune responses [46]. Alternatively, light-emitting recombinant antigens (LIPS) [47] have been developed and used for the profiling of 16 autoantigens in SLE patients [48]. Solution phase assays are useful for diagnostic purposes but are difficult to implement for antigen

discovery since thousands of soluble proteins having a native conformation would need to first be produced and validated

The Serological Antibody Repertoire: Until recently determining the sequence and relative concentration of the antibodies in the serum repertoire was considered a nearly impossible task: biological fluids contain many thousands of different antibodies all of which are chemically very similar, having an overall high degree of sequence identity, and whose concentrations can vary by several orders of magnitude in a dynamic fashion. Starting in 2008 LC-MS/MS instrumentation began to be used to detect immunoglobulin tryptic fragments in serum samples [49,50]. However, the Ig-derived peptides detected in these earlier studies, were overwhelmingly derived from the framework regions and did not provide sufficient information to piece together complete antibody sequences. By restricting the diversity of the antigen-specific antibody pool from the serum of immunized rabbits using antigen-affinity chromatography under stringent elution conditions, Polakiewicz and coworkers succeeded in using MS to identify overlapping antibody peptides that could help assemble a complete V gene. Combinatorial pairing of separate VH and VL sequences deduced from LC-MS/MS was then used then used to produce several antibodies displaying high affinity for antigen, first from rabbits and subsequently from humans [51,52]. In an alternate approach, dating to 2011, our lab invented a technology for determining the serological repertoire to a specified antigen [53] by combining: (i) NextGen V gene sequencing from peripheral memory B cells and plasmablasts to first create an archive of the antibodies encoded by an individual; (ii) enrichment of the pool of antigen-specific antibodies by affinity chromatography; (iii) determining the CDR-H3 peptides and other informative peptides as needed (e.g. CDR-H2) by LC-MS/MS bottom up proteomics; and (iv) the use of stringent informatics filters to assign the informative mass spectra to the entire VH gene with the help of the sequenced antibody gene archive from (i) above [54,55]. In parallel we developed methods for the massive sequencing of the natively paired VH:VL repertoire, first in 10s of thousands and later on, in millions of peripheral B cells[56,57••] In this manner, the

mass spectrometric identification of a unique peptide in a serum antibody could be mapped into the paired VH:VL repertoire archive to reveal the complete sequence of that antibody which then could be produced and studied *in vitro*. Thanks to the exquisite sensitivity of modern MS instrumentation, individual serum antibodies can be detected semi-quantitatively at levels as low as 0.4 ng/ml[22••]. Given that for an antibody to bind to antigen it has to be present at a concentration at or above its equilibrium dissociation constant, which is estimated to have a ceiling of around 0.1 nM or [58] the approach outlined above has more than adequate resolution for the detection of the repertoire of physiologically relevant antibodies in a sample.

As with any analytical technique, the experimental pipeline described above has certain limitations. First, a fraction of antibodies in the repertoire may evade discovery due to constraints inherent in LC-MS/MS proteomics. Such fault may arise when proteolysis fails to produce unique peptides for some sequences, physicochemical properties such as exceptional length limit peptide observation, or MS/MS spectral identification is ambiguous or incorrect. Additionally, unanticipated post-translational modifications or an incomplete sequence database may prevent assignment of some spectra. We estimate that the fraction of CDR-H3 peptide spectra that cannot be assigned is <20%, based on semiautomated inspection of the MS/MS data. Antibodies that evade detection using the proteomic approach outlined above may potentially be identified through use of alternative proteases and instrument fragmentation methods[59]Second, bottom up proteomic approaches reveal the antibody clonotypes (i.e. the family of antibodies having near identical CDR-H3s and V and J segments and likely derived from the same precursor B cell) in the repertoire. Somatic variants of antibodies having the same CDR-H3 are determined based on the NextGen sequencing data but cannot be unambiguously quantified proteomically. Third, the analysis of the antibody repertoire is both technically demanding and expensive, requiring access to VH:VL paired repertoire sequencing, high-end (Thermo-Electron Orbitrap or equivalent mass accuracy) LC-MS/MS instrumentation and finally the requisite informatic and data storage infrastructure. Nonetheless, the above technical limitations notwithstanding, we now have the capability to delineate the serum antibody repertoire with medium-high resolution and with excellent sensitivity.

What's Next? Addressing Fundamental Questions in Humoral Immunity: What kind of information can we glean by utilizing these new tools? We are at the dawn of a new era in the molecular-level understanding of the serological response and numerous questions wait to be answered. Examples of some major outstanding issues we expect to be addressed in the near future are listed below:

- The size of the serological antibody repertoire: We have argued based on the amount of total IgG in healthy humans and the fact that physiologically relevant antibodies must exist at concentrations at or above K_D , that a possible upper bound for the number of distinct clonotypes in human blood can be estimated to be $\geq 10^4$ [23]. Preliminary MS analyses by our lab place the size of the IgG repertoire for a middle age adult at <20,000 distinct clonotypes consistent with previous numbers presented by Talmage during the late 50's [13]. To put this number in perspective, the serological memory repertoire to vaccines we have analyzed so far such as tetanus[22••] TIV seasonal influenza, inactivated polio and pneumococcal polysaccharide typically comprises between 50-400 antibody clonotypes each. In turn these results suggest that as much as 10-15% of the blood IgG pool corresponds or is "taken up" by vaccine-induced antibodies. How clonotypic diversity of the IgG pool changes as a function of age especially in the elderly and during early development and the relationship between overall antibody diversity and susceptibility to infection will need to be determined. Also it will be interesting to determine whether vaccination with a new vaccine impacts the established serological memory antibody repertoire to an earlier vaccine.

- Concentration distribution of circulating antibodies: In biological fluids individual antibodies are present at widely different concentrations. For example we have found that out of the 80-120 antibody clonotypes that typically constitute the serum titer to the tetanus toxoid vaccine a single clonotype can account for as much as 15-30% of antigen-specific response [22••]. We have observed a similar polarization of the serological repertoire, manifest by the presence of one or a few highly abundant antibodies with a number of vaccine antigens. Obviously, such antibodies must be the product of large plasma cell expansions. The mechanisms that dictate the extent of expansion of certain B cell clones following antigen stimulation are not clear. Since highly abundant

antibodies display comparable binding affinity as much less prevalent clones, differences in the degree of plasma cell expansions observed in peripheral blood at day 7 (the typical peak of plasmablast response in serum) clearly cannot be solely due to antigen affinity. Of note, extremely highly abundant antibody clones (clinically defined as globulin level >30 mg/ml) are observed in about 3% of asymptomatic adults over 50 yr old and in multiple myeloma patients[60]. There is a documented reciprocal change with age in the antibody titer to extrinsic and intrinsic antigens [61], but whether this is related to the presence of age-associated highly expanded serum antibodies remains to be determined.

- The relationship and contribution of the different isotype repertoires to humoral immunity: In addition to IgG, human blood contains an appreciable level of IgM and IgA. IgD is present at a level 1/100 of that of IgA and IgE is found at an even lower concentration [62]. The relationships among the isotypic repertoires in serum need to be elucidated. For example to what degree is the antigen-specific IgG repertoire clonally related to the respective IgA repertoire? Additionally, there might be isotypic “layers” to human B cell memory, as recently described in the mouse, wherein particular memory B-cell subsets are predisposed to differentiate into plasma cells while others re-enter germinal center reactions [63]. Are features such as degree of somatic hypermutation, V gene usage, CDR-H3 length etc comparable among the class switched repertoires? While some of these issues are beginning to be explored for peripheral B cells through the use of NextGen sequencing approaches [64,65] there is presently no information for the repertoire at the protein level. A related immunological puzzle which could be clarified using Ig-seq proteomics is the extent to which peripheral blood IgM⁺IgD⁺CD27⁺ circulating B cells are germinal center-derived memory B cells or are instead related to marginal zone B cells found in human spleen, and whether, analogous to the mouse, they constitute a distinct B-cell lineage harboring a pre-diversified repertoire that is responsible for predominating anti-polysaccharide responses against encapsulated bacteria such as *S. pneumonia*. Finally an equally intriguing question is the degree to which the antigen-specific repertoires in blood and in secretions are clonally related and whether they are directed to the same epitopes.

- Significance of antibody post-translational modifications: Post-translational modifications impact all aspects of antibody function including antigen-binding, stability

to aggregation and degradation, interaction with Fc binding proteins and effector functions and half-life[66•]. Antibody post-translational modifications have been studied in the context of quality control of therapeutic antibodies, a key issue for the biopharmaceutical industry. How post-translational modifications influence the selection and function of circulating antibodies is not known. Certainly, oxidation must be quite prevalent in circulating antibodies since they originate from oxidatively stressed plasma cells and then are exposed to the high redox potential of the serum for many days. It is quite plausible that surface Ig on B cells is also subject to oxidation and perhaps oxidized residues may contribute to antigen affinity and B cell selection during the germinal center reaction. Fc glycosylation is well established to impact the effect of antibodies on the innate immune system via the modulation of Fc receptor interactions[67]. While glycosylation heterogeneity in monoclonal antibodies has been studied extensively [66•] recent advances now enable deep analyses of human glycosylation in complex mixtures. For example, Pučić et al. used hydrophilic interaction chromatography and mass spectrometry to investigate correlation between gender, age and heredity on the relative abundance of serum IgG N-glycan structures for a cohort of 2298 individuals[68]. Though this work focused solely on the characterization of the pooled N-glycans from each sample, MS based methods may enable the future study of serum Ig glycosylation at the individual antibody level[69].

- The polyspecific antibody repertoire: As mentioned above, the immunological significance of antibody polyreactivity or polyspecificity is being increasingly appreciated [24,70]. However, polyspecificity as a feature of the serum antibody response has not been investigated in a systematic way. Part of the problem is inherent to the diversity of the antibody repertoire of normal serum. Perhaps this is an area where lessons and tools from the biotherapeutics industry which has developed a number of widely used metrics and probes to evaluate polyspecificity in the context of therapeutic antibody development can be applied to molecular immunology[71]. Cell-based studies have shown that up to 50% of B cells found in cord blood of human newborns, and approximately 20% of peripheral blood B cells circulating in human adults encode polyreactive antibodies[72,73]. Furthermore, although significantly reduced by putative checkpoint mechanisms, the terminal repertoire of bone marrow plasma cells still

encompasses $\geq 10\%$ polyreactive antibodies [74] which suggests that a similar observation awaits discovery at the level of the serum repertoire (**Fig. 1**).

Conclusions

We are on the verge of a new and exciting research era in serology. For the first time, newly developed technologies provide us with the means to interrogate the adaptive immune response at unprecedented resolution to determine both the spectrum of antigens towards which the serological response is directed and the antibodies that comprise the respective polyclonal antibody pools towards these antigens. Together with new tools for the analysis of B cell subsets and BCR repertoires we can now begin to envision the possibility of a fairly complete description of the nature of antibody responses in health and disease. Needless to say numerous challenges remain: First off, the molecular level dissection of humoral immunity is technically demanding and requires a combination of tools that are currently available only in a very small set of laboratories. It will be interesting to see how and whether these techniques may be “industrialized” to become available for clinical use and for use by the wider research community. Second, these studies result in very large and complex data files and a major effort will be required for data storage, annotation, storage and meta-analysis. Third, numerous experimental challenges still need to be addressed: how to develop appropriate probes for the quantification and for characterizing the composition of the polyspecific antibody repertoire, how to best characterize the antigenic determinants in complex pathogens (e.g. *S. aureus* and other bacterial pathogens, parasites etc) how to efficiently identify the neutralizing antibodies present in serum (as contrasted to relying on B cell cloning which as was argued above may not necessarily yield the biologically relevant antibodies in a patient) and many others. Finally and perhaps most importantly the deluge of data that can be generated by the tools described above and by other systems immunology approaches should by no means overshadow the need for first formulating testable hypotheses and well posed questions that in turn can provide definitive insights into the working of the immune system. Critical thinking, appreciation of the biological context and, for disease biology a clear understanding of the clinical need must be the driving force behind any large scale science including in this case modern serology.

Acknowledgments

Work on immune profiling in our lab was supported by grants HDTRA1-12-C-0105 from DTRA, RO1CA106006 and 5U19AI057234-09 from NIH and by the Clayton Foundation.

Figure 1 caption: Approaches for the analysis of human antibody repertoires.

Isolated B cells are sorted into several subsets based on expressed cell markers that correspond to the developmental stage of the B cell. These populations can be further processed for high-throughput sequencing to generate the antibody repertoire encoded by B cells (cellular repertoire, left side of the figure). The corresponding serum immunoglobulins are isolated from the samples and can be analyzed by various methods including well established technologies such as 2D gels or by recently established methodologies such as high resolution shotgun proteomics (serological repertoire, left side of the figure). The methodologies for serological immunoglobulin analysis can be broadly based upon the phenotype of an antibody subpopulation (e.g., ELISA titer of antigen-specific fraction) or upon decipherment of the molecular identity and sequence determination of an antibody subpopulation (e.g., LC-MS/MS immunoglobulin sequencing, Ig-seq).

References

1. K. Landsteiner, J. van der Scheer
On cross reactions of egg albumin sera
J Exp Med, 71 (1940), pp. 445-454
2. M.D. Cooper
The early history of B cells
Nat Rev Immunol, 15 (2015), pp. 191-197
3. T. Yoshida, H.E. Mei, T. Dörner, F. Hiepe, A. Radbruch, S. Fillatreau, B.F. Hoyer
Memory B and memory plasma cells
Immunol Rev, 237 (2010), pp. 117-139

4. K. Kometani, T. Kurosaki

Differentiation and maintenance of long-lived plasma cells

Curr Opin Immunol, 33c (2015), pp. 64-69

5. * H.E. Mei, I. Wirries, D. Frolich, M. Brisslert, C. Giesecke, J.R. Grun, T. Alexander, S. Schmidt, K. Luda, A.A. Kuhl, R. Engelmann, M. Durr, T. Scheel, M. Bokarewa, C. Perka, A. Radbruch, T. Dörner

A unique population of IgG-expressing plasma cells lacking CD19 is enriched in human bone marrow

Blood, 125 (2015), pp. 1739-1748

On mature B cells, CD19 is a co-receptor of the B cell receptor and regulates its signaling. This study characterizes distinct subsets of human plasma cells (PC) differing by CD19 expression. Using a combination of FACS-sorting, single cell sequencing, EliSpot and statistical analysis, the researchers demonstrated that the CD19- B cell population in the BM are independent from being replaced by new PC subset and comprise a lower level of somatic hypermutation. The authors concluded that this PC subset are differentiated in early life.

6. • S. Panda, J.L. Ding

Natural antibodies bridge innate and adaptive immunity

J Immunol, 194 (2015), pp. 13-20

Natural Abs, mostly isotypes IgM, IgG3, and IgA, were discovered nearly half a century ago. Despite knowledge about the role of the polyreactive natural IgM there is a lack of clarity about the physiological role of natural IgG and natural IgA because they appear incapable of recognizing Ags on their own and are perceived as nonreactive. In this review, the authors describe the findings on natural Abs, focusing on the role of natural IgM, recent findings on natural IgG, which collaborates with serum lectins, and how this immune complex links innate immunity to adaptive immunity.

7. U. Galili

Anti-Gal: an abundant human natural antibody of multiple pathogeneses and clinical benefits

Immunology, 140 (2013), pp. 1-11

8. I.J. Amanna, N.E. Carlson, M.K. Slifka

Duration of humoral immunity to common viral and vaccine antigens

N Engl J Med, 357 (2007), pp. 1903-1915

9. A. Radbruch, G. Muehlinghaus, E.O. Luger, A. Inamine, K.G. Smith, T. Dorner, F. Hiepe

Competence and competition: the challenge of becoming a long-lived plasma cell
Nat Rev Immunol, 6 (2006), pp. 741-750

10. S.A. Plotkin

Vaccines: correlates of vaccine-induced immunity
Clin Infect Dis, 47 (2008), pp. 401-409

11. S.A. Plotkin

Correlates of Protection Induced by Vaccination
Clin Vaccine Immunol, (2010), pp. 1055-1065

12. R. Rappuoli

Bridging the knowledge gaps in vaccine design
Nat Biotechnol, 25 (2007), pp. 1361-1366

13. D.W. Talmage

Immunological Specificity: Unique combinations of selected natural globulins provide an alternative to the classical concept
Science, 129 (1959), pp. 1643-1648

14. J.M. Fonville, S.H. Wilks, S.L. James, A. Fox, M. Ventresca, M. Aban, L. Xue, T.C. Jones, N.M. Le, Q.T. Pham, N.D. Tran, Y. Wong, A. Mosterin, L.C. Katzelnick, D. Labonte, T.T. Le, G. van der Net, E. Skepner, C.A. Russell, T.D. Kaplan, G.F. Rimmelzwaan, N. Masurel, J.C. de Jong, A. Palache, W.E. Beyer, Q.M. Le, T.H. Nguyen, H.F. Wertheim, A.C. Hurt, A.D. Osterhaus, I.G. Barr, R.A. Fouchier, P.W. Horby, D.J. Smith

Antibody landscapes after influenza virus infection or vaccination
Science, 346 (2014), pp. 996-1000

15. • D. Corti, A. Lanzavecchia

Broadly neutralizing antiviral antibodies
Annu Rev Immunol, 31 (2013), pp. 705-742

The high variability of viral glycoproteins and sophisticated viral escape mechanisms presents a great challenge to the development of antibody-based therapies or vaccines that could confer broad and long-lasting protection. In this review the authors describe viral escape mechanisms and neutralization. The review focuses on new approaches that

were developed during the past five years that enabled the high efficiency investigation of human memory B cells and plasma cells and to identify and isolate several broadly neutralizing antiviral antibodies against highly variable pathogens such as HIV-1 and influenza virus.

16. • H.X. Liao, R. Lynch, T. Zhou, F. Gao, S.M. Alam, S.D. Boyd, A.Z. Fire, K.M. Roskin, C.A. Schramm, Z. Zhang, J. Zhu, L. Shapiro, J.C. Mullikin, S. Gnanakaran, P. Hraber, K. Wiehe, G. Kelsoe, G. Yang, S.M. Xia, D.C. Montefiori, R. Parks, K.E. Lloyd, R.M. Searce, K.A. Soderberg, M. Cohen, G. Kamanga, M.K. Louder, L.M. Tran, Y. Chen, F. Cai, S. Chen, S. Moquin, X. Du, M.G. Joyce, S. Srivatsan, B. Zhang, A. Zheng, G.M. Shaw, B.H. Hahn, T.B. Kepler, B.T. Korber, P.D. Kwong, J.R. Mascola, B.F. Haynes

Co-evolution of a broadly neutralizing HIV-1 antibody and founder virus
Nature, 496 (2013), pp. 469-476

In this study the authors provided data regarding the understanding of the interaction between HIV-1 virus evolution and maturing broad neutralizing antibody lineages in early infection. By providing data related to the viral and antibody evolution leading to induction of a lineage of HIV-1 broadly neutralizing antibodies, they provided insights into strategies to elicit similar antibodies by vaccination. The authors reported the isolation of the CH103 CD4-binding site broad neutralizing antibody clonal lineage from an African patient and show that the CH103 broad neutralizing antibody lineage is less mutated than most other CD4- binding site broad neutralizing antibody.

17. N.A. Doria-Rose, C.A. Schramm, J. Gorman, P.L. Moore, J.N. Bhiman, B.J. DeKosky, M.J. Ernandes, I.S. Georgiev, H.J. Kim, M. Pancera, R.P. Staube, H.R. Altae-Tran, R.T. Bailer, E.T. Crooks, A. Cupo, A. Druz, N.J. Garrett, K.H. Hoi, R. Kong, M.K. Louder, N.S. Longo, K. McKee, M. Nonyane, S. O'Dell, R.S. Roark, R.S. Rudicell, S.D. Schmidt, D.J. Sheward, C. Soto, C.K. Wibmer, Y. Yang, Z. Zhang, J.C. Mullikin, J.M. Binley, R.W. Sanders, I.A. Wilson, J.P. Moore, A.B. Ward, G. Georgiou, C. Williamson, S.S. Abdool Karim, L. Morris, P.D. Kwong, L. Shapiro, J.R. Mascola

Developmental pathway for potent V1V2-directed HIV-neutralizing antibodies
Nature, 509 (2014), pp. 55-62

18. • D.J. DiLillo, G.S. Tan, P. Palese, J.V. Ravetch

Broadly neutralizing hemagglutinin stalk-specific antibodies require FcγR interactions for protection against influenza virus in vivo
Nat Med, 20 (2014), pp. 143-151

In this study, the researches utilize anti-HA stalk broad neutralizing antibodies that neutralize a panel of H1 or of both H1 and H5 influenza viruses, to show that the anti-stalk bNAbs required Fc-FcγR interactions for maximum bNAb-mediated neutralization of influenza virus in vivo, whereas the strain-specific anti-HA head mAbs did not. The

anti-stalk bNAbs functioned after viral entry in vivo and were superior inducers of antibody-dependent cellular cytotoxicity (ADCC), suggesting a mechanism for their FcγR-dependent function in vivo. The authors underlined the potential of engineering the human IgG1 Fc to enhance binding to its activation receptors and augment antibody effector function that may results in the development of future therapies, including vaccination protocols to elicit bNAbs and passive antibody therapies using bNAbs.

19. J.R. Mascola, B.F. Haynes

HIV-1 neutralizing antibodies: understanding nature's pathways

Immunol Rev, 254 (2013), pp. 225-244

20. • I.S. Georgiev, N.A. Doria-Rose, T. Zhou, Y.D. Kwon, R.P. Staupe, S. Moquin, G.Y. Chuang, M.K. Louder, S.D. Schmidt, H.R. Altae-Tran, R.T. Bailer, K. McKee, M. Nason, S. O'Dell, G. Ofek, M. Pancera, S. Srivatsan, L. Shapiro, M. Connors, S.A. Migueles, L. Morris, Y. Nishimura, M.A. Martin, J.R. Mascola, P.D. Kwong
Delineating antibody recognition in polyclonal sera from patterns of HIV-1 isolate neutralization

Science, 340 (2013), pp. 751-756

In this paper the authors hypothesized that when the immune system is presented with a diverse set of HIV-1 isolates, it generates monoclonal antibodies with specific neutralization fingerprints and neutralization patterns of a polyclonal serum could be viewed as the combined effect of the neutralization fingerprints of the constituent monoclonal antibodies. They demonstrated the neutralization fingerprints for 30 neutralizing antibodies on a panel of 34 diverse HIV-1 strains and showed that similarity in neutralization fingerprint correlated with similarity in epitope. Furthermore, they used these fingerprints to delineate specificities of polyclonal sera from 24 HIV-1-infected donors. Collectively, they suggested that epitope delineation based on neutralization fingerprints may provide a transformative strategy for screening sera or characterizing antibody specificities induced upon infection or vaccination against HIV-1 as well as other viruses.

21. F.M. Burnet: *The Clonal Selection Theory of Acquired Immunity*: Vanderbilt University Press; (1959). pp 95

22. •• J.J. Lavinder, Y. Wine, C. Giesecke, G.C. Ippolito, A.P. Horton, O.I. Lungu, K.H. Hoi, B.J. DeKosky, E.M. Murrin, M.M. Wirth, A.D. Ellington, T. Doerner, E.M. Marcotte, D.R. Boutz, G. Georgiou

Identification and characterization of the constituent human serum antibodies elicited by vaccination

Proc Natl Acad Sci USA, 111 (2014), pp. 2259-2264

Most vaccines confer immunity by eliciting long-term production of antibodies that bind to and neutralize the vaccine antigen. Remarkably, very little is known regarding the

identities, sequence diversity, relative concentrations, or binding functionalities of the mAbs that comprise the serum repertoire elicited by vaccination. In this study the authors delineated the constituent antibodies of the human serum IgG repertoire after vaccination and examined their relationship to the antibody V gene repertoire encoded by circulating B cells. The results detail the molecular composition and characteristics of the vaccine-specific serum antibody repertoire and demonstrate differences between the end-point response (the serum antibodies) and the peripheral B cells responding to the vaccine.

23. J.J. Lavinder, A.P. Horton, G. Georgiou, G.C. Ippolito
Next-generation sequencing and protein mass spectrometry for the comprehensive analysis of human cellular and serum antibody repertoires
Curr Opin Chem Biol, 24c (2015), pp. 112-120

24. J.D. Dimitrov, C. Planchais, L.T. Roumenina, T.L. Vassilev, S.V. Kaveri, S. Lacroix-Desmazes
Antibody polyreactivity in health and disease: statu variabilis
J Immunol, 191 (2013), pp. 993-999

25. Z.H. Zhou, Y. Zhang, Y.F. Hu, L.M. Wahl, J.O. Cisar, A.L. Notkins
The broad antibacterial activity of the natural antibody repertoire is due to polyreactive antibodies
Cell Host Microbe, 1 (2007), pp. 51-61

26. T. Tiller, M. Tsuiji, S. Yurasov, K. Velinzon, M.C. Nussenzweig, H. Wardemann
Autoreactivity in human IgG+ memory B cells
Immunity, 26 (2007), pp. 205-213

27. M. Liu, G. Yang, K. Wiehe, N.I. Nicely, N.A. Vandergrift, W. Rountree, M. Bonsignori, S.M. Alam, J. Gao, B.F. Haynes, G. Kelsoe
Polyreactivity and autoreactivity among HIV-1 antibodies
J Virol, 89 (2015), pp. 784-798

28. H. Mouquet, M.C. Nussenzweig
Polyreactive antibodies in adaptive immune responses to viruses
Cell Mol Life Sci, 69 (2012), pp. 1435-1445

29. T. Kodadek
Chemical tools to monitor and manipulate the adaptive immune system
Chem Biol, 21 (2014), pp. 1066-1074

30. • Y. Weiss-Ottolenghi, J.M. Gershoni
Profiling the IgOme: meeting the challenge
FEBS Lett, 588 (2014), pp. 318-325

This paper reviews various methodologies that enable researchers to better understand the entire repertoire of antibodies in our serum (IgOme). They discuss various aspect of the development of the adaptive immune system and how “serum memory” is affected by immunization, boosts, encounters with pathogens, physiology and old age.

31. J. Sokolove, T.M. Lindstrom, W.H. Robinson
Development and deployment of antigen arrays for investigation of B-cell fine specificity in autoimmune disease
Front Biosci (Elite Ed), 4 (2012), pp. 320-330

32. J. Kügler, J. Zantow, T. Meyer, M. Hust
Oligopeptide M13 Phage Display in Pathogen Research
Viruses, 5 (2013), pp. 2531-2545

33. P.S. Daugherty, G. Chen, M.J. Olsen, B.L. Iverson, G. Georgiou
Antibody affinity maturation using bacterial surface display
Protein Eng, 11 (1998), pp. 825-832

34. J.T. Ballew, J.A. Murray, P. Collin, M. Mäki, M.F. Kagnoff, K. Kaukinen, P.S. Daugherty
Antibody biomarker discovery through in vitro directed evolution of consensus recognition epitopes
Proc Natl Acad Sci USA, 110 (2013), pp. 19330-19335

35. S.E. Elliott, N.F. Parchim, C. Liu, Y. Xia, R.E. Kellems, A.R. Soffici, P.S. Daugherty
Characterization of antibody specificities associated with preeclampsia
Hypertension, 63 (2014), pp. 1086-1093

36. H.B. Larman, Z. Zhao, U. Laserson, M.Z. Li, A. Ciccia, M.A.M. Gakidis, G.M. Church, S. Kesari, E.M. Leproust, N.L. Solimini, S.J. Elledge
Autoantigen discovery with a synthetic human peptidome
Nat Biotechnol, 29 (2011), pp. 535-541

37. P. Stafford, Z. Cichacz, N.W. Woodbury, S.A. Johnston
Immunosignature system for diagnosis of cancer.
Proc Natl Acad Sci USA, 111 (2014), pp. 3072-3080
38. K.A. Navalkar, S.A. Johnston, P. Stafford
Peptide based diagnostics: Are random-sequence peptides more useful than tiling proteome sequences?
J Immunol Methods, 417 (2015), pp. 10-21
39. T.M. Doran, S. Simanski, T. Kodadek
Discovery of Native Autoantigens via Antigen Surrogate Technology: Application to Type 1 Diabetes
ACS Chem Biol, 10 (2015), pp. 401-412
40. A. Yeste, F.J. Quintana
Antigen microarrays for the study of autoimmune diseases
Clin Chem, 59 (2013), pp. 1036-1044
41. P. Kloppot, M. Selle, C. Kohler, S. Stentzel, S. Fuchs, V. Liebscher, E. Muller, D. Kale, K. Ohlsen, B.M. Broker, P.F. Zipfel, B.C. Kahl, R. Ehricht, M. Hecker, S. Engelmann
Microarray based identification of human antibodies against Staphylococcus aureus antigens
Proteomics Clin Appl, (2015), pp.
42. S. Miersch, X. Bian, G. Wallstrom, S. Sibani, T. Logvinenko, C.H. Wasserfall, D. Schatz, M. Atkinson, J. Qiu, J. LaBaer
Serological autoantibody profiling of type 1 diabetes by protein arrays
J Proteomics, 94 (2013), pp. 486-496
43. B. Ayoglu, A. Haggmark, M. Khademi, T. Olsson, M. Uhlen, J.M. Schwenk, P. Nilsson
Autoantibody profiling in multiple sclerosis using arrays of human protein fragments
Mol Cell Proteomics, 12 (2013), pp. 2657-2672
44. J.V. Price, D.J. Haddon, D. Kemmer, G. Delepine, G. Mandelbaum, J.A. Jarrell, R. Gupta, I. Balboni, E.F. Chakravarty, J. Sokolove, A.K. Shum, M.S. Anderson, M.H. Cheng, W.H. Robinson, S.K. Browne, S.M. Holland, E.C. Baechler, P.J. Utz

Protein microarray analysis reveals BAFF-binding autoantibodies in systemic lupus erythematosus

J Clin Invest, 123 (2013), pp. 5135-5145

45. A.K. Shum, M. Alimohammadi, C.L. Tan, M.H. Cheng, T.C. Metzger, C.S. Law, W. Lwin, J. Perheentupa, H. Bour-Jordan, J.C. Carel, E.S. Husebye, F. De Luca, C. Janson, R. Sargur, N. Dubois, M. Kajosaari, P.J. Wolters, H.A. Chapman, O. Kampe, M.S. Anderson

BPIFB1 is a lung-specific autoantigen associated with interstitial lung disease

Sci Transl Med, 5 (2013), pp. 206ra139

46. C.G. Joseph, E. Darrah, A.A. Shah, A.D. Skora, L.A. Casciola-Rosen, F.M. Wigley, F. Boin, A. Fava, C. Thoburn, I. Kinde, Y. Jiao, N. Papadopoulos, K.W. Kinzler, B. Vogelstein, A. Rosen

Association of the autoimmune disease scleroderma with an immunologic response to cancer

Science, 343 (2014), pp. 152-157

47. J. Ustinova, E. Zusinaite, M. Utt, K. Metskula, K. Reimand, V. Huchaiah, A. Merits, R. Uibo

Development of a luciferase-based system for the detection of ZnT8 autoantibodies

J Immunol Methods, 405 (2014), pp. 67-73

48. K.H. Ching, P.D. Burbelo, C. Tipton, C. Wei, M. Petri, I. Sanz, M.J. Iadarola

Two Major Autoantibody Clusters in Systemic Lupus Erythematosus

PLoS ONE, 7 (2012), pp. e32001

49. B. Obermeier, R. Mentele, J. Malotka, J. Kellermann, T. Kumpfel, H. Wekerle, F. Lottspeich, R. Hohlfeld, K. Dornmair

Matching of oligoclonal immunoglobulin transcriptomes and proteomes of cerebrospinal fluid in multiple sclerosis

Nat Med, 14 (2008), pp. 688-693

50. M.M. VanDuijn, L.J. Dekker, L. Zeneyedpour, P.A. Smitt, T.M. Luider

Immune responses are characterized by specific shared immunoglobulin peptides that can be detected by proteomic techniques

J Biol Chem, 285 (2010), pp. 29247-29253

51. W.C. Cheung, S.A. Beausoleil, X. Zhang, S. Sato, S.M. Schieferl, J.S. Wieler, J.G. Beaudet, R.K. Ramenani, L. Popova, M.J. Comb, J. Rush, R.D. Polakiewicz
A proteomics approach for the identification and cloning of monoclonal antibodies from serum
Nat Biotechnol, 30 (2012), pp. 447-452

52. S. Sato, S.A. Beausoleil, L. Popova, J.G. Beaudet, R.K. Ramenani, X. Zhang, J.S. Wieler, S.M. Schieferl, W.C. Cheung, R.D. Polakiewicz
Proteomics-directed cloning of circulating antiviral human monoclonal antibodies
Nat Biotechnol, 30 (2012), pp. 1039-1043

53. S. Reddy, X. Ge, J. Lavinder, D. Boutz, A.D. Ellington, E.M. Marcotte, G. Georgiou:
Rapid Isolation of Monoclonal Antibodies from Animals. US Patent 2011.

54. • Y. Wine, J.J. Lavinder, D.R. Boutz, A.E. Miklos, R.A. Hughes, K.H. Hoi, S.T. Jung, A.P. Horton, E.M. Murrin, A.D. Ellington, E.M. Marcotte, G. Georgiou
Molecular deconvolution of the monoclonal antibodies that comprise the polyclonal serum response
Proc Natl Acad Sci USA, 110 (2013), pp. 2993-2998

This paper described how the monoclonal antibody composition was delineated using a methodology that integrates proteomic analysis of the serum- derived antigen-specific polyclonal antibody pool, NextGen sequencing of the immunoglobulin heavy chain variable region (VH gene) repertoire with advanced bioinformatics tools. After booster immunization in a rabbit, it was demonstrated that the antigen- specific serum immune response is oligoclonal, comprising antibodies encoding 34 different CDRH3s that group into 30 distinct antibody VH clonotypes. Of these 34 CDRH3s, 12 account for $\square 60\%$ of the antigen-specific CDRH3 peptide mass spectral counts. Proteomically identified antibodies were synthesized and shown to display subnanomolar affinities. The ability to deconvolute the polyclonal serum response is likely to be of key importance for analyzing antibody responses after vaccination and help answer fundamental questions related to the adaptive immune responses in health and disease.

55. D.R. Boutz, A.P. Horton, Y. Wine, J.J. Lavinder, G. Georgiou, E.M. Marcotte
Proteomic Identification of Monoclonal Antibodies from Serum
Anal Chem, 86 (2014), pp. 4758-4766

56. B.J. DeKosky, G.C. Ippolito, R.P. Deschner, J.J. Lavinder, Y. Wine, B.M. Rawlings, N. Varadarajan, C. Giesecke, T. Dörner, S.F. Andrews, P.C. Wilson, S.P. Hunicke-Smith, C.G. Willson, A.D. Ellington, G. Georgiou
High-throughput sequencing of the paired human immunoglobulin heavy and light chain repertoire.

Nat. Biotechnol., 31 (2013), pp. 166-169

57. •• B.J. DeKosky, T. Kojima, A. Rodin, W. Charab, G.C. Ippolito, A.D. Ellington, G. Georgiou

In-depth determination and analysis of the human paired heavy- and light-chain antibody repertoire

Nat Med, 21 (2015), pp. 86-91

Here the authors describe a technology that enables sequencing of the native paired VH-VL repertoire from millions of B cells within a few hours of experimental effort and using equipment that can be built inexpensively by most laboratories. Because the variable domains of antibody heavy and light chains (VH and VL, respectively) are encoded by different mRNA transcripts, until recently it was only possible to determine the paired VH-VL sequences for moderate numbers of cells, far fewer than the diversity contained reasonable samples size. Massive VH-VL repertoire analyses of three human donors provided new immunological insights including (i) the identity, frequency and pairing propensity of shared, or 'public', VL genes, (ii) the detection of allelic inclusion (an implicated autoimmune mechanism) in healthy individuals and (iii) the occurrence of antibodies with features, in terms of gene usage and CDR3 length, associated with broadly neutralizing antibodies to rapidly evolving viruses such as HIV-1 and influenza.

58. J. Foote, H.N. Eisen

Breaking the affinity ceiling for antibodies and T cell receptors

Proc Natl Acad Sci U S A, 97 (2000), pp. 10679-10681

59. D.L. Swaney, C.D. Wenger, J.J. Coon

Value of using multiple proteases for large-scale mass spectrometry-based proteomics

J Proteome Res, 9 (2010), pp. 1323-1329

60. E. Sigurdardottir, I. Turesson, S. Lund, et al.

The role of diagnosis and clinical follow-up of monoclonal gammopathy of undetermined significance on survival in multiple myeloma

JAMA Oncology, Published advanced online (2015), pp.

61. M. Rowley, H. Buchanan, I. Mackay

Reciprocal change with age in antibody to extrinsic and intrinsic antigens

The Lancet, 292 pp. 24-26

62. K. Murphy, P. Travers, M. Walport, C. Janeway: *Janeway's immunobiology*. New York: Garland Science; (2012). pp

63. T. Kurosaki, K. Kometani, W. Ise
Memory B cells
Nat Rev Immunol, 15 (2015), pp. 149-159
64. B.S. Briney, J.R. Willis, J.A. Finn, B.A. McKinney, J.E. Crowe, Jr.
Tissue-specific expressed antibody variable gene repertoires
PLoS One, 9 (2014), pp. e100839
65. G. Georgiou, G.C. Ippolito, J. Beausang, C.E. Busse, H. Wardemann, S.R. Quake
The promise and challenge of high-throughput sequencing of the antibody repertoire
Nat Biotech, 32 (2014), pp. 158-168
66. • L.K. Hmiel, K.A. Brorson, M.T. Boyne, 2nd
Post-translational structural modifications of immunoglobulin G and their effect on biological activity
Anal Bioanal Chem, 407 (2015), pp. 79-94
67. A. Pincetic, S. Bournazos, D.J. DiLillo, J. Maamary, T.T. Wang, R. Dahan, B.M. Fiebiger, J.V. Ravetch
Type I and type II Fc receptors regulate innate and adaptive immunity
Nat Immunol, 15 (2014), pp. 707-716
68. M. Pucic, A. Knezevic, J. Vidic, B. Adamczyk, M. Novokmet, O. Polasek, O. Gornik, S. Supraha-Goreta, M.R. Wormald, I. Redzic, H. Campbell, A. Wright, N.D. Hastie, J.F. Wilson, I. Rudan, M. Wuhrer, P.M. Rudd, D. Josic, G. Lauc
High throughput isolation and glycosylation analysis of IgG-variability and heritability of the IgG glycome in three isolated human populations
Mol Cell Proteomics, 10 (2011), pp. M111.010090
69. M. Thaysen-Andersen, N.H. Packer
Advances in LC-MS/MS-based glycoproteomics: getting closer to system-wide site-specific mapping of the N- and O-glycoproteome
Biochim Biophys Acta, 1844 (2014), pp. 1437-1452
70. F. Klein, H. Mouquet, P. Dosenovic, J.F. Scheid, L. Scharf, M.C. Nussenzweig
Antibodies in HIV-1 Vaccine Development and Therapy
Science, 341 (2013), pp. 1199-1204

71. Y. Xu, W. Roach, T. Sun, T. Jain, B. Prinz, T.-Y. Yu, J. Torrey, J. Thomas, P. Bobrowicz, M. Vásquez, K.D. Wittrup, E. Krauland
Addressing polyspecificity of antibodies selected from an in vitro yeast presentation system: a FACS-based, high-throughput selection and analytical tool
Protein Eng Des Sel, 26 (2013), pp. 663-670
72. H. Wardemann, S. Yurasov, A. Schaefer, J.W. Young, E. Meffre, M.C. Nussenzweig
Predominant autoantibody production by early human B cell precursors
Science, 301 (2003), pp. 1374-1377
73. Z.J. Chen, C.J. Wheeler, W. Shi, A.J. Wu, C.H. Yarboro, M. Gallagher, A.L. Notkins
Polyreactive antigen-binding B cells are the predominant cell type in the newborn B cell repertoire
Eur J Immunol, 28 (1998), pp. 989-994
74. J.F. Scheid, H. Mouquet, J. Kofer, S. Yurasov, M.C. Nussenzweig, H. Wardemann
Differential regulation of self-reactivity discriminates between IgG⁺ human circulating memory B cells and bone marrow plasma cells
P Natl Acad Sci USA, 108 (2011), pp. 18044-18048

Figure 1: Approaches for the analysis of human antibody repertoires.

Isolated B cells are sorted into several subsets based on expressed cell markers that correspond to the developmental stage of the B cell. These populations can be further processed for high-throughput sequencing to generate the antibody repertoire encoded by B cells (cellular repertoire, left side of the figure). The corresponding serum immunoglobulins are isolated from the samples and can be analyzed by various methods including well established technologies such as 2D gels or by recently established methodologies such as high resolution shotgun proteomics (serological repertoire, left side of the figure). The methodologies for serological immunoglobulin analysis can be broadly based upon the phenotype of an antibody subpopulation (e.g., ELISA titer of antigen-specific fraction) or upon decipherment of the molecular identity and sequence determination of an antibody subpopulation (e.g., LC-MS/MS immunoglobulin sequencing, Ig-seq).

Cellular

Serological

Phenotyping

Deciphering

B cell
Sequences
(BCR-seq)

VH
VL
VH:VL

Serum
Electrophoresis

SPEP
IEF
2D

ELISA

Bulk
Titers

Affinity
Purification

Antigen
Specificity

Functional
Assay

Neutralization
Agglutination
Precipitation

Antibody
Landscapes

Specificity
Titer

Antibody
Proteomics
(Ig-seq)

Specificity
Abundance
Molecular Identity