Lessons from Tissue Compartment-Specific Analysis of Androgen Receptor Alterations in Prostate Cancer

Mark Daniel^{a, b}, Scott M. Dehm^{b, c, d}

^aGraduate Program in Microbiology, Immunology, and Cancer Biology, University of Minnesota, Minneapolis, MN, USA

^bMasonic Cancer Center, University of Minnesota, Minneapolis, MN, USA

^cDepartment of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis, MN, USA

^dDepartment of Urology, University of Minnesota, Minneapolis, MN, USA

Correspondence to: Scott M. Dehm, Masonic Cancer Center, University of Minnesota, Mayo Mail Code 806, 420 Delaware Street SE, Minneapolis, Minnesota, USA. E-mail address:

dehm@umn.edu

Abstract

Androgen receptor (AR) splice variants (AR-Vs) are constitutively active transcription factors that function in the absence of ligand. AR-Vs represent one of several AR re-activation mechanisms utilized by prostate cancer to circumvent first-line androgen deprivation therapy. Second line therapies such as enzalutamide and abiraterone are treatments that re-target components of the androgen/AR axis. However, these second line therapies do not benefit all patients, and patients that do receive initial benefit can develop resistance rapidly. Alterations in components of the androgen/AR axis, including expression of AR-Vs, appear to be linked to primary as well as secondary resistance to second line therapies. However, some key conclusions appear to differ depending on the tissue compartment and measurement platform utilized for analysis. In this review, alterations in AR and the broader AR pathway will be examined in the context of primary prostate cancer tissue, metastatic castration-resistant prostate cancer tissue, circulating tumor cells, and circulating cell-free tumor DNA. Questions regarding the utility of AR-V measurements to provide prognostic information or predict patient responses to AR-targeted therapies will be addressed.

Keywords: androgen receptor splice variant; prostate cancer; castration-resistant prostate cancer; circulating tumor cell

Abbreviations: LHRH, luteinizing hormone releasing hormone; ADT, androgen depletion therapy; CRPC, castration-resistant prostate cancer; CYP17, cytochrome P450 c17; MR, mineralocorticoid receptor; ER, estrogen receptor; PR, progestins receptor; GR, glucocorticoid receptor; NTD, NH₂-terminal domain; AF-1, activation function 1; LBD, ligand bind domain; ARE, androgen response element; RIP, RNA co-immunoprecipitation; NCOA, nuclear receptor coactivator; NCOR, nuclear receptor corepressor; CTC, circulating tumor cell; ctDNA, circulating tumor DNA; FISH, fluorescence *in situ* hybridization; PSA, prostate-specific antigen; PMSA, prostate-specific membrane antigen.

Contents

1. Introduction

1.1 Brief Introduction to AR Structure/Function

1.2 Brief Introduction to AR Splice Variants (AR-Vs)

2. Alterations in the AR and AR-V pathways in primary prostate cancer tissue

2.1 Alterations in AR in primary prostate cancer tissue

2.2 AR-Vs in normal and primary prostate cancer tissue

2.3 Alterations in the broader AR pathway in primary prostate cancer tissue

3. Alterations in the AR and AR-V pathways in CRPC tissue

3.1 Alterations in AR in CRPC tissue

3.2 AR-Vs in CRPC tissue

4. Detection of Alterations in the AR and AR-V Pathways in Blood

4.1 Alterations in the AR in CTCs

4.2 Alterations in the AR in ctDNA

4.3 AR-Vs in CTCs

5. Functional Changes in AR and AR-V signaling during PCa progression

5.1 AR cistrome and transcriptome in normal tissue, primary prostate cancer, and CRPC

5.2 AR versus AR-V cistrome and transcriptome

6. Conclusion

1. Introduction

Prostate cancer accounts for a quarter of all new cancer diagnoses, and one in seven men in the United States will develop the disease. Of those men, roughly 27,000 will die each year, making prostate cancer the second leading cause of cancer related death in the United States. [1]. Prostate cancer is referred to as an androgen-dependent disease, which is based on the requirement of androgen-mediated activation of the androgen receptor (AR) for growth and survival of prostate cancer cells. As a result of androgen-dependence, the androgen/AR axis has been a target of treatment since the early 1940s when Huggins *et al.* established the beneficial effects of castration in men with metastatic prostate cancer [2]. Contemporary therapies for prostate cancer continue to rely on reducing the levels of circulating androgens and inhibiting AR function via orchiectomy or luteinizing hormone releasing hormone (LHRH) agonists. Antiandrogens such as bicalutamide are also used to competitively block the ability of androgens to bind and activate the AR. Patients that fail this first-generation of androgen depletion therapies (ADT) and progress to a disease stage referred to as castration-resistant prostate cancer (CRPC) are treated with second-generation AR-targeting therapies such as abiraterone acetate and enzalutamide (**Fig. 1A**). Abiraterone acetate is an inhibitor of the cytochrome P450 c17 (CYP17) testosterone synthesis enzyme [3], which can achieve a more robust inhibition of androgen production than castration. Enzalutamide is an antagonist that binds the AR with a higher affinity than bicalutamide, and can partially inhibit AR translocation to the nucleus, reduce binding of AR to DNA, and prevent recruitment of coactivators required for active transcription [4]. Despite these advances in treatment, CRPC remains a uniformly fatal disease. Importantly, persistent AR signaling can be maintained during and after treatment with these

1
2
3
4 androgen/AR-targeted therapies. The focus of this review will be observations made in
5
6 clinical tissues collected at various disease stages (**Fig. 1B**) that have illuminated key
7
8 mechanisms of AR re-activation that can occur in CRPC (**Fig. 1C**), with an emphasis on AR
9
10 splice variants (AR-Vs).
11
12
13
14

15 16 17 **1.1 Brief Introduction to AR Structure/Function**

18
19 The AR is a member of the class I nuclear steroid receptor family. Within this family
20
21 of steroid receptors are transcription factors regulated by ligands including
22
23 mineralocorticoids (MR), estrogens (ER), progestins (PR) and glucocorticoids (GR). All of
24
25 these steroid receptors are activated upon binding ligand, after which they engage their
26
27 cognate hormone response element DNA sequences located in promoter and enhancer
28
29 regions throughout the genome. The AR gene is located on the X chromosome at
30
31
32 cytogenetic position Xq11-12. This gene contains eight exons and encodes a modular
33
34 110kDa protein [5]. Exon 1 of AR encodes the NH₂-terminal domain (NTD), which is
35
36 intrinsically disordered and harbors the transcriptional activation function-1 (AF-1)
37
38 domain essential for the bulk of AR transcriptional activity. Exons 2 and 3 encode the DNA-
39
40 binding domain. The 5' section of exon 4 encodes the hinge region, which is a flexible
41
42 domain containing a nuclear localization signal (NLS) that becomes accessible upon ligand
43
44 binding. The 3' section of exon 4 along with exons 5-8 encode the AR ligand-binding
45
46 domain (LBD) and an AF-2 domain that is transcriptionally active but with weaker activity
47
48 than AF-1 [5]. Upon binding to androgen, AR undergoes a conformational change, which
49
50 exposes the NLS, and results in translocation to the nucleus. Once in the nucleus, AR binds
51
52 to androgen response elements (AREs) and recruits a wide variety of coregulatory proteins
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 with various scaffolding, enzymatic, and chromatin-modifying functions, ultimately leading
5
6 to a finely-tuned level of transcriptional output for target genes. These features of canonical
7
8 AR signaling are illustrated in **Figure 1C**.
9

10 11 12 13 14 **1.2 Brief Introduction to AR Splice Variants (AR-Vs)**

15
16 Over the past 8 years, AR-Vs have emerged as an important component of resistance
17
18 to therapies targeting the androgen/AR axis. Approximately 20 discrete AR-Vs have been
19
20 identified in cell- and animal-based models of CRPC progression, as well as tissues from
21
22 patients with prostate cancer [6, 7, 8, 9]. AR-Vs share the same NTD/DBD modular domain
23
24 structure as the full-length AR, but commonly lack the COOH-terminal LBD/AF-2 module.
25
26
27

28
29 **Figure 1C** illustrates the structure of select AR-Vs that will be detailed in this review.
30

31 Truncation of the AR LBD results in constitutive transcriptional activity of the AR
32
33 NTD/DBD core in the absence of ligand. Thus, most AR-Vs have been reported to function
34
35 as ligand-independent transcription factors, although discrepancies have been noted [10].
36
37 Originally, AR-Vs were reported to arise from proteolytic cleavage of full-length AR by
38
39 calpains, which are calcium-dependent proteinases [11]. These early studies found that the
40
41 AR hinge region harbored a calpain cleavage site, which could be targeted to eliminate the
42
43 AR LBD. However, subsequent studies in the 22Rv1 cell line, which was the same model
44
45 used to develop the proteolysis mechanism, revealed that truncated AR protein species
46
47 were differentially sensitive to small interference RNAs (siRNAs) compared with full-length
48
49 AR. This finding of differential siRNA sensitivity formed the basis for the concept that AR-
50
51 Vs were encoded by mRNAs that were separate from the mRNA encoding full-length AR
52
53
54
55
56
57
58
59 [12]. Subsequent studies demonstrated that different AR-V proteins were encoded by
60
61
62
63
64
65

1
2
3
4 mRNAs containing exons 1-3 or exons 1-4 of the AR gene, but variable 3' terminal exons
5
6
7 harboring in-frame translation termination signals [13]. Subsequent studies defined an
8
9 additional class of AR-V proteins that were encoded by skipping of exons encoding the AR
10
11 LBD, but terminating with an out-of-frame 3' terminal AR exon 8 [13].
12
13

14 One mechanism thought to underlie AR-V expression in prostate cancer is
15
16 alternative splicing. RNA co-immunoprecipitation (RIP) assays using enzalutamide-treated
17
18 VCaP cells revealed that splicing regulatory factors ASF/SF2, and U2AF65 were recruited to
19
20 pre-mRNA sequences aligning with a splice acceptor site in AR exon CE3, which is the 3'
21
22 terminal exon in the mRNA encoding AR-V7 [14]. In this scenario, it was concluded that
23
24 increased recruitment, but not expression, of splicing factors contributed to AR-V7 in these
25
26 cells. In line with this, the AR-V7-negative LNCaP cell line did not display the same pattern
27
28 of engagement with splicing factors seen in VCaP cells. Further, siRNA-mediated
29
30 knockdown of splicing factors in this model system reduced the expression of AR-V7 [14].
31
32
33 However, one caveat with comparing splicing factor recruitment to AR pre-mRNAs in VCaP
34
35 vs. LNCaP cells is that VCaP cells harbor approximately 20 more copies of the AR gene than
36
37 the hypotetraploid LNCaP cell line due to massive amplification of the AR gene [15]. In this
38
39 case, VCaP cells express much higher levels of AR mRNA and protein, and it would be
40
41 assumed that there would be higher sensitivity for detection of splicing regulatory factors
42
43 bound to AR pre-mRNA in VCaP cells.
44
45
46
47
48
49
50

51 An additional mechanism that has been shown to underlie AR-V expression in
52
53 prostate cancer is genomic rearrangements within the AR gene. For example, a 35kb
54
55 tandem duplication containing AR exon 3 as well as several cryptic exons including CE3 is
56
57 linked to the creation of truncated AR variants in the androgen-independent 22Rv1 cell line.
58
59
60
61
62
63
64
65

This tandem duplication extends the genomic span between exons 3 and 4, which may increase the probability of exon 3 splicing to exon CE3 or other cryptic exons, thereby increasing expression of AR-V7 as well as other AR-Vs relative to full-length AR [16, 17]. Moreover, patient derived xenograft models LuCaP 86.2 and LuCaP 136 express high levels of the ARv567es exon skipping AR splice variant resulting from splicing of AR exon 4 directly to AR exon 8. These patient derived xenografts harbor a deletion or inversion, respectively, of a genomic segment containing AR exons 5-7. These specific rearrangements were demonstrated to cause ARv567es expression following targeted manipulation of the AR gene in prostate cancer cell lines using genome engineering technology [17].

2. Alterations in the AR and AR-V pathways in primary prostate cancer tissue

2.1 Alterations in AR in primary prostate cancer tissue

Alterations in the AR gene are very rare in primary prostate cancer tissues, which is consistent with the lack of use of ADT for patients with localized disease. In a targeted sequencing study of archival, formalin-fixed specimens, only one AR point mutation was detected in primary prostate cancer tissue [18]. Other studies have reported a slightly higher mutational burden in primary prostate cancer specimens, but rates still remain relatively low [19]. The most comprehensive study of primary prostate cancer tissues to date, the cancer genome atlas (TCGA) project, reported zero somatic AR point mutations across a series of 333 primary prostate cancer tissues [19]. On the other hand, this TCGA study did report rare instances of AR gene amplification, but at less than 1% frequency.

Overall, these studies provide evidence of minimal direct AR alterations in primary prostate cancer.

2.2 AR-Vs in normal and primary prostate cancer tissue

Although mutation and amplification of the AR gene are rarely observed in primary prostate cancer, expression of AR-Vs has been observed in non-malignant prostate tissue as well as primary prostate cancer. Targeted qRT-PCR analysis of AR-V7 mRNA revealed that AR-V7 was expressed in 5 of 17 benign prostate tissues [8]. Analysis of RNA-seq data from peritumoral prostate tissues in the TCGA project also revealed frequent expression of AR-V7 as well as other AR-Vs in normal tissue [19]. The significance of AR-V mRNA expression in benign tissue is not clear, but is consistent with another study where mRNA expression of several AR-Vs was detected in normal prostate tissue from men who had been castrated with an LHRH receptor agonist or treated with DHT gel in a study testing male contraception [9]. Analysis of RNA-seq data from cancer tissues in the TCGA cohort indicated that AR-V7 was the most frequently expressed AR-V mRNA in primary prostate cancer tissue [19]. Additionally, a range of other AR-V mRNAs arising from cryptic exon inclusion as well as exon skipping events were also detected frequently in primary prostate cancer tissues from the TCGA study [19]. An unexpected finding from analysis of TCGA data was an overall lack of expression of the ARv567es exon skipping AR splice variant [19]. This finding was unexpected because a previous targeted RT-PCR study found that ARv567es expression mirrored that of AR-V7 and AR-V1 in normal prostate tissue as well as primary prostate cancer samples, with levels detectable but low relative to full length AR [9, 20]. In contrast, targeted qRT-PCR analysis of AR-V7 mRNA was largely in-line with

findings from the TCGA study, revealing that 34 of 82 hormone naïve prostatectomy specimens were positive for AR-V7 mRNA expression [8]. Interestingly, AR-V7 mRNA expression greater than the median across this cohort was associated with lower probability of PSA-progression free survival following surgical treatment [8]. Conversely, expression of AR-V1, composed of contiguously-spliced AR exons 1, 2, 3, and CE1, was not found to be associated with PSA-progression free survival. However, a subsequent study that evaluated AR-V1 and AR-V7 mRNA expression using a branched chain assay failed to identify an association between either of these AR-V species and prostate cancer progression [21]. As a result of these discordant findings, it remains unclear whether AR-V7 mRNA levels in primary prostate cancer tissue provide prognostic information.

The unique COOH-terminal extensions encoded by the unique 3' terminal exons in AR-Vs such as AR-V7 have provided opportunities for development of AR-V-specific antibodies [7, 8]. For example, staining of tissue microarrays with an antibody specific for AR-V7 revealed that AR-V7 protein was expressed in both the nucleus and cytoplasm of hormone naïve primary prostate cancer tissues [7]. Tissue microarrays also exhibited cytoplasmic AR-V7 staining in benign tissue but no nuclear signal [7]. AR-V7 cytoplasmic staining in human prostate tissue microarrays was found to be associated with shorter time to PSA recurrence [7]. The finding of cytoplasmic AR-V7 expression is unexpected, as biochemical studies have demonstrated this species to be a constitutively nuclear protein [22]. This raises concern about the performance of AR-V7 antibody reagents in tissue-based studies, particularly in benign prostate and hormone naïve prostate cancer, where AR-V7 expression is extremely low.

2.3 Alterations in the broader AR pathway in primary prostate cancer tissue

Primary prostate cancer has recently been classified based on seven distinct molecular subtypes. Four of these subtypes are defined by genomic fusions of ETS family genes (ERG, ETV1, ETV4, or FLI1), or mutations in SPOP, FOXA1, or IDH1. In the analysis of 333 primary tumors from the TCGA study, 74% of samples fell into one of these subclasses [19]. Interestingly, the majority of these molecular subclasses are mechanistically linked to the broader androgen/AR pathway in prostate cancer. For example, the most frequent ERG fusion is TMPRSS2:ERG, which results in androgen/AR-mediated overexpression of ERG mRNA and protein via the androgen/AR-regulated TMPRSS2 enhancer and promoter [19, 23]. TMPRSS2:ETV1 is the most frequent ETV fusion, which has a similar regulatory outcome of androgen/AR-mediated ETV1 mRNA and protein overexpression via the TMPRSS2 enhancer and promoter. Additional fusions to diverse promoters and enhancers have been characterized for ERG, ETV1, ETV4, and FLI1, the majority of which are highly-active in prostate cells by virtue of androgen/AR regulation. Impaired expression of these genetic drivers following inhibition of the androgen/AR axis provides a molecular basis for the robust response of prostate cancer cells to ADT. However, evaluation of whether TMPRSS2:ERG or other ETS fusions relay any prognostic information has not yielded clear results [24, 25].

In addition to gene fusion subclasses, the FOXA1 and SPOP mutant subclasses can also be considered as molecular alterations affecting the broader AR pathway. For instance, the FOXA1 gene encodes a pioneer transcription factor that facilitates the relaxation of compacted chromatin structures for AR and other transcription factors [19, 26, 27, 28]. Mutations in FOXA1 include missense, inframe deletions, and truncating mutations, which

are concentrated in the exons encoding the FOXA1 forkhead DNA binding domain [19]. Similarly, the SPOP gene, which encodes the substrate-binding subunit of a Cullin-based E3 ubiquitin ligase [29, 30], is also frequently mutated in primary prostate cancer [19, 28]. SPOP mutations are concentrated in exons encoding the MATH domain, which is responsible for substrate recruitment. Two substrates identified for SPOP are AR and the AR co-activator SRC-3, with mutant versions of SPOP displaying impaired binding to these substrates [31, 32, 33]. Therefore, AR and SRC-3 proteins are stabilized in cells harboring mutant SPOP. Interestingly, the SPOP MATH domain interacts with the AR LBD, and therefore AR-Vs appear to be resistant to SPOP-mediated degradation [31]. However, a separate study indicated that AR-Vs may remain susceptible to SPOP-dependent ubiquitination and degradation by virtue of heterodimerization with full-length AR [34]. Assignment of an AR activity score to primary prostate cancer tissues, which was based on expression levels of a set of 20 AR regulated genes, revealed that FOXA1 and SPOP subclasses had the highest levels of AR transcriptional output. This reinforces how intimately FOXA1 and SPOP are tied to the normal function of AR and also the disease-associated function of AR in primary prostate cancer.

3. Alterations in the AR and AR-V pathways in CRPC tissue

Most prostate tumors have an initial response to ADT because of the androgen-dependent nature of the disease. However, virtually all patients will eventually develop resistance. In this CRPC stage of the disease, most tumors remain dependent on activity of the androgen/AR axis [35]. Mechanisms underlying persistent activity of the androgen/AR

axis in CRPC have been reviewed in detail [36], and include AR mutations, AR amplification, expression of AR splice variants, and alterations in AR regulators (**Fig. 1C**). As discussed below, a critical role in CRPC is supported by the finding that these alterations occur with higher frequency in CRPC than in primary prostate cancer, although there are some notable discrepancies. Additionally, the proportion of the genome that is altered through copy number alteration and/or mutation is greater in patients with CRPC, indicating that androgen/AR-independent mechanisms also contribute to resistance [19]. Overall, CRPC is a complex disease state with a myriad of genomic alterations that enable continued growth and disease spread in the face of therapeutic pressure.

3.1 Alterations in AR in CRPC tissue

Point mutations in AR are more frequent in CRPC than primary prostate cancer. In a whole exome sequencing study of 25 tissues from patients with CRPC, AR point mutations were found to occur in 20% of samples [18]. Other studies with larger sample sizes detected AR point mutations at a frequency of roughly 10% [37, 38]. Among the mutations found in these studies were T878A, W742C, and L702H, which are located in the AR LBD and have been shown to convert a variety of AR antagonists into agonists [18, 38].

Although single point mutations are the most frequent occurrence, there have been reports where multiple point mutations have been detected in a single tissue sample, such as T878A and Q903H in a recent whole exome sequencing study of CRPC biopsies [38]. This study did not establish if these mutations were concurrent on the same AR allele, or whether this reflected intra-tumor heterogeneity. However, the latter scenario is likely given that the Q903H mutation was present at a lower allelic fraction than T878A.

1
2
3
4 The most common AR alteration in CRPC is AR gene amplification. In a biopsy-based
5
6 whole exome sequencing study of 150 metastatic CRPC bone and soft tissue tumor samples,
7
8 AR amplification occurred in more than 50% of samples [38]. In another study of CRPC
9
10 samples, AR copy number gain was detected in 25 of 50 cases [37]. Mechanistically, AR
11
12 amplification leads to overexpression of AR mRNA and protein, which is thought to
13
14 facilitate AR activation by a mass action effect under low androgen conditions [39, 40].
15
16 Given that AR amplification in primary prostate cancer is exceedingly rare, this highlights
17
18 the clinical importance of increased AR protein expression in CRPC [19, 41, 42, 43]. AR
19
20 gene amplification appears to occur in a mutually exclusive pattern with AR point
21
22 mutations, although rare exceptions have been noted that may reflect intra-tumor
23
24 heterogeneity [38].
25
26
27
28
29
30
31
32
33

34 **3.2 AR-Vs in CRPC tissue**

35
36 Because AR-Vs were discovered in cell line model systems of CRPC and were shown
37
38 in these models to function as ligand-independent transcription factors, most studies
39
40 interrogating AR-Vs in clinical material have focused on CRPC tissues. Although expression
41
42 of numerous AR-Vs has been detected in CRPC tissues [7, 8, 9, 20, 38], quantitative
43
44 assessments have often revealed that AR-Vs are expressed at a much lower level in clinical
45
46 material than in cell lines where they were initially discovered and characterized
47
48 functionally [7, 8, 9, 12]. This has complicated efforts to clearly define the importance of
49
50 AR-Vs in mediating resistance to ADT and second-generation AR-targeted therapies such as
51
52 abiraterone and enzalutamide. Despite these challenges, there is compelling evidence in
53
54 support of a strong link between AR-V expression and CRPC progression. For example, Hu
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 and colleagues evaluated 124 clinical samples representing normal prostate, hormone-
5
6 naïve primary prostate cancer, and CRPC for mRNA expression of AR-V7 (1/2/3/CE3
7
8 isoform) using an RT-PCR assay. They observed that 84% of CRPC specimens displayed AR-
9
10 V7 mRNA expression vs. 41% of hormone-naïve primary prostate cancer specimens [8].
11
12 This mRNA-based finding was supported by a subsequent immunohistochemistry analysis
13
14 of a human prostate tissue microarray consisting of hormone-naïve primary prostate
15
16 cancer and CRPC specimens [7]. Staining with an antibody specific for AR-V7 demonstrated
17
18 a significant increase in AR-V7 nuclear expression in CRPC tumors relative to hormone-
19
20 naïve primary prostate tumors. Surprisingly, 86% of hormone-naïve primary prostate
21
22 cancer tissues had AR-V7 protein expression in the cytoplasm compared to 100% in CRPC
23
24 samples [7]. However, as discussed earlier, cytoplasmic AR-V7 staining may reflect non-
25
26 specific reactivity in tissues expressing AR-V7 protein at a level below the threshold of
27
28 accurate detection. Overall, these data demonstrate that AR-V7 mRNA and protein is
29
30 expressed in treatment-naïve primary prostate cancer samples, but the transcriptional
31
32 effects of nuclear AR-V7 may not materialize until the disease has progressed to the CRPC
33
34 stage.
35
36
37
38
39
40
41
42

43
44 More recent RT-PCR-based studies have interrogated expression of multiple AR-Vs
45
46 simultaneously in CRPC tissues. For example, Sun and colleagues reported that 31 of 69
47
48 metastatic CRPC specimens were positive for either AR-V7 or ARv567es mRNA expression
49
50 [9]. Of note, ARv567es was detected at twice the frequency as AR-V7 in this study, which is
51
52 in opposition to more recent RNA-seq based studies that have concluded AR-V7 is the most
53
54 frequently-expressed AR splice variant in CRPC tissue [19, 38]. As expanded below, these
55
56 discrepancies may reflect the use of targeted RT-PCR vs. RNA-seq based assessments. In
57
58
59
60
61
62
63
64
65

1
2
3
4 this same study by Sun and colleagues, AR-V7 and ARv567es were co-expressed in 6 of 69
5
6 metastatic samples [9]. Hörnberg and colleagues performed a similar RT-PCR-based
7
8 analysis of AR-V1 and AR-V7 transcript expression in bone metastases from hormone-
9
10 naïve and CRPC patients [20]. In this study, AR-V1 expression was consistently detected in
11
12 CRPC bone metastases. AR-V1 positivity was also consistently observed in hormone-naïve
13
14 bone metastases and primary prostate tumors, although, AR-V1 expression in CRPC bone
15
16 metastases samples was higher relative to the other samples. Moreover, AR-V7 mRNA
17
18 displayed a similar expression pattern in this study, but was detected less frequently than
19
20 AR-V1 in both primary prostate tumor and hormone-naïve bone metastases [20].
21
22 Importantly, patients with bone metastases displaying the highest quartile of AR-V7 mRNA
23
24 expression had reduced cancer-specific survival compared with patients having lower AR-
25
26 V7 expression in bone metastases [20]. Collectively, these studies employing RT-PCR
27
28 detection have provided strong support for an important disease role of AR-Vs in CRPC.
29
30
31
32
33
34
35

36
37 Interestingly, RNA-seq assessments of AR-Vs in clinical CRPC have not been
38
39 confirmatory of the RT-PCR-based conclusions that AR-Vs play an important role in CRPC
40
41 progression. The common strategy for assessment of AR-V expression in RNA-seq datasets
42
43 has been to quantify the number of individual RNA-seq reads spanning exon/exon
44
45 boundaries that are common to all AR mRNA species as well as the number of individual
46
47 RNA-seq reads spanning exon/exon boundaries that are unique to AR and AR-Vs, and
48
49 developing ratios to reflect the fractional amount of overall AR expression that is
50
51 attributable to any particular AR species. This quantification strategy was used to analyze
52
53 RNA-seq data from 125 pre-abiraterone and -enzalutamide CRPC biopsy samples as well as
54
55 333 primary prostate cancer specimens [19, 38]. These data demonstrated that AR-V7
56
57
58
59
60
61
62
63
64
65

(defined by the exon 3/CE3 splice junction) and AR-V3 (defined by the exon 2/CE4 splice junction) were the most frequently expressed AR-V mRNAs in CRPC tissues, and displayed the highest fractional expression of all AR-V mRNAs as a function of overall AR expression. Remarkably, AR-V7 and AR-V3 were also the most frequently detected AR-Vs in primary prostate cancer specimens, and their fractional expression levels as a function of overall AR expression appeared to be largely unchanged compared to the measurements in CRPC tissues [38]. Additionally, ARv567es, defined by the exon 4/8 splice junction, was only detected in 4/125 CRPC specimens, despite having been found in previous RT-PCR studies of CRPC metastases to display a trend towards association with reduced cancer-specific survival [20]. Understanding the basis for this discordance in conclusions following RT-PCR measurements vs. RNA-seq measurements of AR-Vs in prostate cancer tissues will be important for developing a clearer picture of their role in CRPC progression. One possibility is that RT-PCR measurements typically measure the expression of a specific AR-V relative to an internal housekeeping gene whereas RNA-seq data has been evaluated based on fractional expression of an AR-V relative to overall AR expression. It is possible this RNA-seq data analysis strategy is not highlighting those tissues where overall AR expression is increased, leading to higher expression of AR-Vs as well as full-length AR. Additionally, data comparing the change in expression of AR-Vs after introduction of second line AR inhibitors would provide great insight into how AR-Vs respond, and thereby contribute to CRPC progression in different treatment contexts.

3.2 Alterations in the broader AR pathway in CRPC tissue

As is the case in primary prostate cancer tissue, alterations in AR-related pathway components are also observed in CRPC tissue. For example, FOXA1 has been found to be mutated in CRPC samples, at a slightly higher frequency than in primary prostate cancer [28, 37, 38]. Interestingly, there also appears to be a shift in the distribution of FOXA1 mutations in CRPC compared with primary prostate cancer. For example, in primary prostate cancer, FOXA1 mutations were concentrated in the Forkhead domain [19] whereas a study of CRPC tissue found that four of seven FOXA1 mutations were in the tail-anchored (TA) domain and only one (G87R) mutation was detected in the Forkhead domain [37]. Similar non-Forkhead domain mutations, most of which localized to a region between the Forkhead domain and tail-anchored domain, were also frequently observed in a biopsy-based study of 150 CRPC specimens [38]. SPOP mutations, like FOXA1 aberrations, are also present at similar frequency in CRPC as primary prostate cancer, but with no apparent differences in distribution [28, 38]. Beyond FOXA1 and SPOP, several other AR activators and repressors within the AR pathway are altered in CRPC. For instance, nuclear receptor coactivators 1 and 2 (NCOA1/2), which are known to facilitate AR transcriptional activity, are frequently overexpressed at the mRNA and protein levels in metastatic CRPC [41], while nuclear receptor corepressors 1 and 2 (NCOR1/2), which are AR repressors, are commonly inactivated by mutation (nonsense, frameshift, missense), gene fusion or decreased mRNA expression [38, 41]. The importance of these specific AR pathway alterations is underscored by the finding that they are more frequent in metastatic CRPC samples compared to primary prostate cancer [19, 28]. Taking into account the cumulative total of all the alterations identified in the AR and the broader AR pathway has indicated

1
2
3
4 that AR signaling is altered in 100% of metastatic CRPC cases [41]. These data underscore
5
6 the critical role of AR in prostate cancer progression and therapeutic resistance.
7
8
9

10 11 **4. Detection of Alterations in the AR and AR-V Pathways in Blood** 12 13

14
15
16
17 Circulating tumor cells (CTCs) and cell-free circulating tumor DNA (ctDNA) provide
18
19 great potential for determining cancer progression in real time with methods that are non-
20
21 invasive compared to biopsies, where sample yield is typically low. Observing proteomic,
22
23 transcriptomic, miRNA expression, and genotypic changes in CTCs or genomic changes in
24
25 ctDNA in response to various therapeutics enables analysis and understanding of tumor
26
27 evolution. As this technology matures, it may also lead to more nuanced care in the
28
29 developing age of personalized medicine. In the context of CRPC, CTC and ctDNA are
30
31 thought to represent cells or cellular material shed from metastases. This section of the
32
33 review will discuss how alterations in the AR and broader AR pathway have been evaluated
34
35 in CTCs and ctDNA, with emphasis on how major findings have been in concordance or
36
37 discordance with the tissue-based findings discussed in previous sections.
38
39
40
41
42
43
44
45

46 47 **4.1 Alterations in the AR in CTCs** 48

49
50 An early examination by Miyamoto *et al* into AR alterations in CTCs focused on
51
52 transcriptional function of androgen-activated AR [44]. Given that PSA is a gene target
53
54 transcriptionally activated by the AR, and PSMA is a gene target transcriptionally repressed
55
56 by AR, immunophenotyping for expression of these AR gene targets was used to classify
57
58 individual CTCs as “AR active” (PSA+/PSMA-), “AR inactive” (PSA-/PSMA+), or “mixed AR
59
60
61
62
63
64
65

1
2
3
4 signaling” (PSA+/PSMA+). Within this study, initial ADT resulted in a robust switch from
5
6 activated AR signaling to inactive AR signaling, whereas second line treatment with
7
8 abiraterone induced a more mixed signal [44]. These trends in AR activity within CTCs
9
10 suggest an evolution toward a more complex and heterogeneous disease state.
11
12 Understanding this evolution with respect to alterations in the AR gene and expression
13
14 profiles may reveal a mechanistic basis for this heterogeneity.
15
16
17
18

19 Shaffer *et al.* performed one of the earliest analyses of AR alterations in CTCs from
20
21 patients with CRPC. Within this study, Shaffer and colleagues used fluorescence *in situ*
22
23 hybridization (FISH) to examine chromosomal aberrations in nine patients. Among the
24
25 nine patients with CRPC, five of them exhibited amplification of AR [45]. Even within this
26
27 small set, AR amplification frequency in CTCs mirrored the frequencies seen in larger scale
28
29 tissue studies [38]. Related to this finding was a study that examined the AR gene in CTCs
30
31 from patients treated with abiraterone. In this study, AR amplification was heterogeneous
32
33 after second line therapy [46]. More recently, other AR alterations such as missense
34
35 mutations, deletions, and insertions were examined in CTCs. One study found an unusually
36
37 high percentage of missense mutations (15 of 35 CRPC patients) [47]. This frequency of AR
38
39 mutation was much higher than observed in tissue-based analyses of CRPC tissue. One
40
41 possible explanation is that tissue-based studies have routinely sampled a single metastatic
42
43 site, whereas the cells that are enriched during CTC collection procedures are thought to
44
45 originate from multiple metastatic sites. Given that sampling of several CRPC tissue sites
46
47 from individual patients has revealed multiple independent AR amplification and mutation
48
49 events [48], it is possible that CTC-based analysis reflects the totality of AR mutational
50
51 burden within a patient, leading to a higher estimate of AR mutation frequency.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 In a separate study that performed single-cell RNA-seq of CTCs, the T877A point
5
6 mutation associated with ligand promiscuity and therapy resistance was only detected in
7
8 CTCs from 1 of 13 patients with CRPC. An E709E mutation was also detected in 1/13 CRPC
9
10 patients in this study [49]. This AR mutational frequency (2/13 patients) is concordant
11
12 with the AR mutational frequency observed in larger-scale studies with CRPC tissues.
13
14 However, there was insufficient depth of AR mRNA sequencing coverage for many of the
15
16 individual CTCs in this study to rule out the possibility of a higher AR mutation frequency.
17
18 Overall, these studies have demonstrated proof-of-principle for detection of specific AR
19
20 gene alterations using various analysis platforms in CTCs, but the results have been highly
21
22 variable. Improving the sensitivity and specificity of these platforms for AR analyses may
23
24 provide a more accurate picture going forward.
25
26
27
28
29
30
31
32
33

34 **4.2 Alterations in the AR in ctDNA**

35
36 In line with the non-invasive principles of CTC analysis, interrogation of ctDNA is
37
38 another strategy for evaluating tumor evolution and tracking responses to therapy in
39
40 patients with CRPC. Romanel *et al.* reported targeted next-generation sequencing of ctDNA
41
42 from plasma samples collected longitudinally from 97 CRPC patients before and during
43
44 therapy with abiraterone [50]. Of 274 plasma samples analyzed, 41 displayed AR point
45
46 mutations known to associate with ADT resistance [50], which is in concordance with the
47
48 frequency observed in studies of CRPC tissue [37, 38]. Only 217 plasma samples from this
49
50 cohort had a sufficient cancer cell fraction to enable analysis of AR gene copy number. Of
51
52 these 217 samples, 80 demonstrated an AR copy number gain [50]. Importantly, detection
53
54 of AR point mutations or copy number gain in plasma ctDNA from patients pre-therapy was
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 associated with less favorable clinical outcomes including reduced PSA response rates,
5
6 shorter progression-free survival, and shorter overall survival compared to patients where
7
8 no AR alterations were detected [50].
9
10

11 12 13 14 **4.3 AR-Vs in CTCs** 15

16
17 Single-cell RNA-seq of CTCs from patients with CRPC has also enabled examination
18
19 of AR splicing patterns and detection of AR-V expression in this context. For instance, in a
20
21 study of individual CTCs isolated from 13 patients, 33/73 CTCs analyzed displayed
22
23 expression of at least one type of AR-V [49]. Consistent with RNA-seq studies of CRPC
24
25 tissue [38], AR-V7 was the most frequently detected AR-V (26 of 73 CTCs; 8 of 11 patients)
26
27 [49]. Expression of ARv567es was also detected with high frequency (18 of 73 CTCs; 8 of 11
28
29 patients) [49], which is in disagreement with the overall lack of ARv567es mRNA
30
31 expression observed in RNA-seq studies of CRPC tissue. Expression of AR-V1, AR-V3, and
32
33 AR-V4 (contiguously-splice AR exons 1/2/3/CE4) has also been detected by single-cell
34
35 RNA-seq of CTCs [49]. Interestingly, 13 of 73 individual CTCs expressed two or more
36
37 different AR-Vs, which demonstrates that multiple AR splicing patterns can exist in a single
38
39 cell. However, development of approaches for detection of AR-V protein expression in
40
41 individual CTCs will be necessary to understand whether these mRNA species are
42
43 translated to generate functional AR-V protein. Finally, partitioning the patients in this
44
45 study into those that had been treated with enzalutamide vs. those that had not been
46
47 treated with enzalutamide did not reveal enrichment for AR-V expression in the
48
49 enzalutamide-treated cohort [49].
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 While there is evidence of heterogeneous AR-V expression in CTCs before and after
5
6 treatment with enzalutamide, there is strong evidence that simple detection by RT-PCR of
7
8 AR-V7 mRNA in CTCs may serve as a predictive biomarker for patients treated with
9
10 second-generation AR-targeted therapies. Specifically, in a study of 62 CRPC patients
11
12 initiating therapy with abiraterone (31 patients) or enzalutamide (31 patients),
13
14 Antonarakis and colleagues reported that the presence of CTCs positive for AR-V7 mRNA
15
16 expression was associated with lower PSA response rates, diminished progression-free
17
18 survival, and reduced overall survival in both treatment groups [51]. These data imply a
19
20 clear association of AR-V7 mRNA with resistance to second line AR-targeted therapies.
21
22 Results in this study are in agreement with a study examining AR-V7 protein expression in
23
24 bone marrow aspirates from CRPC patients [52] as well as an independent study evaluating
25
26 CTCs from patients treated with enzalutamide [53]. Interestingly, these studies have also
27
28 reported that increased AR-V7 mRNA expression in bone marrow aspirates and CTCs
29
30 directly correlated with the number of prior therapies patients had received [51, 52, 53].
31
32 Indeed, patients with an AR-V7 mRNA negative CTC profile pre-therapy have been shown
33
34 to convert to an AR-V7 mRNA positive CTC profile during treatment with enzalutamide or
35
36 abiraterone [51, 53, 54, 55].
37
38
39
40
41
42
43
44
45

46 While AR-V7 mRNA positivity in CTCs may be predictive of lack of response to
47
48 abiraterone or enzalutamide therapy, AR-V7 positivity in CTCs does not appear to be
49
50 associated with lack of response to taxane chemotherapy. Specifically, patients with an AR-
51
52 V7 mRNA positive CTC profile pre-therapy displayed superior clinical outcomes when
53
54 treated with taxane chemotherapy compared with abiraterone or enzalutamide.
55
56 Conversely, clinical outcomes were indistinguishable for all three treatments in patients
57
58
59
60
61
62
63
64
65

1
2
3
4 with an AR-V7 mRNA negative CTC profile per-therapy [55]. This result may be unexpected
5
6 in light of mechanistic cell-based studies reporting that microtubule-dependent
7
8 translocation of AR to the nucleus can be impaired by taxane chemotherapy, whereas AR-
9
10 V7 translocation to the nucleus is taxane insensitive [56, 57]. However, subsequent studies
11
12 demonstrated this AR translocation effect may be an artifact of treating prostate cancer
13
14 cells with concentrations of taxanes higher than can be achieved clinically [58].
15
16
17 Interestingly, it has also been reported that patients with an AR-V7 mRNA positive CTC
18
19 profile pre-therapy can undergo reversion to AR-V7 mRNA negative CTC profile during
20
21 taxane chemotherapy [54,55]. This plasticity in AR-V7 expression further highlights the
22
23 complexity of AR-V expression in prostate cancer, and shows how various selection
24
25 pressures can modulate resistance mechanisms.
26
27
28
29
30
31
32
33

34 **5. Functional Changes in AR and AR-V signaling during PCa progression**

35
36
37
38

39 Since AR is a transcription factor, understanding functional changes in regard to
40
41 gene expression repertoire (ie the set of transcriptional targets regulated by AR, or the AR
42
43 “transcriptome”) and chromatin binding patterns (ie the collection of cis-elements bound
44
45 by AR, or the AR “cistrome”) during cancer progression is important. Understanding how
46
47 AR gene and splicing alterations, disease context, co-regulator expression, and treatment
48
49 history can influence AR chromatin binding sites and transcriptional targets is expected to
50
51 provide further insight to how aberrations in AR and the broader AR signaling pathway
52
53 drive cancer progression. Fine-tuned and expanded usage of techniques such as ChIP-seq,
54
55 and related ChIP-seq variations, has allowed for more advanced and context specific
56
57
58
59
60
61
62
63
64
65

mapping of AR binding. Furthermore, the complex nature of AR binding is inherent in binding sites where AREs can be found at sites distant from target genes. In this section, AR and AR-V cistromics and transcriptional programs will be discussed in further detail.

5.1 AR cistrome and transcriptome in normal tissue, primary prostate cancer, and CRPC

ChIP-seq and transcriptional profiling in prostate cancer cell lines has been a popular avenue for researchers to understand functional changes in AR action [59]. Moreover, these studies have highlighted mechanistic aspects of the AR cistrome and transcriptional regulation by FOXA1 and ERG [59, 60, 61]. In order to fully understand functional changes in AR signaling, it is important to analyze the transcriptome and cistrome of AR in prostate cancer tumor tissue. This allows for the verification of gene expression and binding data themes found in model systems, which, as a result, relay a more robust understanding of AR functional behavior in a variety of disease settings. For instance, comparison of AR ChIP-seq datasets derived from prostate cancer cell lines and clinical tissues demonstrated that AR chromatin binding events in prostate cancer cell lines more closely reflect the AR chromatin binding events that occur in CRPC tissues than the AR chromatin binding events that occur in primary prostate cancer. This is expected, given that the prostate cancer cell lines used in this study were derived from patients with CRPC. However, over 50% of AR chromatin binding sites observed in CRPC tissues were not detected in prostate cancer cell lines, highlighting the importance of studying tissues directly [62]. Recently, Pomerantz and colleagues used ChIP-seq to compare AR chromatin binding sites in primary prostate cancer and adjacent normal tissue [63]. Hierarchical

1
2
3
4 clustering of AR chromatin binding events revealed distinct groups of AR cistromes
5
6 between primary tumor samples and matched normal tissues. Inclusion of the AR positive
7
8 prostate cancer cell lines LNCaP and VCaP in hierarchical clustering resulted in
9
10 classification of a third distinct group, which most closely resembled primary tumors [63].
11
12 Overall, this study identified 9,179 AR chromatin binding sites that were enriched in tumor
13
14 tissue and 2,690 AR chromatin binding sites that were enriched in normal tissue.
15
16 Importantly, this study identified FOXA1 and HOXB13 as key AR-associated transcription
17
18 factors that ultimately determined tumor-enriched AR chromatin binding events.
19
20 Consistent with this finding, transfection of a normal prostate epithelial cell line with
21
22 lentivirus encoding FOXA1 and HOXB13 resulted in a pattern of AR chromatin binding that
23
24 resembled primary prostate cancer tissue [63]. Building on these same themes, comparison
25
26 of AR chromatin binding events in primary prostate cancer vs. CRPC specimens revealed
27
28 differences in AR chromatin binding associated with disease progression [64]. These
29
30 findings have indicated that AR chromatin binding undergoes extensive reprogramming at
31
32 various stages of disease progression. Understanding the mechanisms by which alterations
33
34 in the AR and broader AR pathway intersect with these chromatin binding alterations will
35
36 provide important insights to the genome-wide impacts of AR alterations in prostate
37
38 cancer.
39
40
41
42
43
44
45
46
47
48
49
50

51 **5.2 AR versus AR-V cistrome and transcriptome**

52
53
54 There has also been great interest in understanding the similarities and differences
55
56 in cistromes and transcriptomes regulated by full-length AR and AR-Vs. One early
57
58 observation was that AR-Vs uniquely regulated a set of G2/M-phase regulatory genes
59
60
61
62
63
64
65

1
2
3
4 including UBE2C [65]. However, more recent analysis of this set of G2/M-phase cell cycle
5
6
7 regulator genes revealed they were biphasic targets that could be induced by both
8
9
10 androgen/AR as well as AR-Vs at low levels of AR transcriptional output, but were
11
12 repressed when AR transcriptional output was increased [66]. A study by Lu and
13
14 colleagues compared the cistromes and transcriptomes regulated by AR and AR-Vs in the
15
16 22Rv1 prostate cancer cell line manipulated with siRNA to achieve cell preparations that
17
18 were either full length AR-/AR-V+ or full length AR-/AR-V- [67]. In this study, AR-Vs were
19
20 observed to regulate a unique gene set separate from full length AR in the absence of
21
22 androgen. Additionally, binding of AR-Vs didn't appear to be affected by loss of full length
23
24 AR or androgens [67]. However, it should be noted that full-length AR in 22Rv1 cells
25
26 harbors a tandem duplication of a zinc finger module in the AR DNA binding domain [16].
27
28 Therefore it is not clear whether the differences in AR vs. AR-V chromatin binding observed
29
30 in this study was due to differential DNA recognition by a larger 3-zinc finger AR DBD
31
32 module vs. a normal 2-zinc finger AR DBD module. Nevertheless, this study also found that
33
34 AR-V chromatin binding sites had 75% overlap with full-length AR chromatin binding sites
35
36 [67]. In a separate ChIP-seq study with cell lines engineered to express either full-length AR
37
38 or ARv567es from the same endogenous AR promoter, ARv567es was found to engage with
39
40 the same genomic sites engaged by full-length AR, albeit with lower affinity [66]. This study
41
42 further revealed that AR-Vs required dimerization to engage with androgen response
43
44 elements [66], which has since been confirmed by a separate study [68]. A more complete
45
46 picture of how AR-Vs engage chromatin will require analysis of additional cell line models
47
48 as well as clinical tissues expressing various levels of AR-Vs relative to full-length AR.
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

6. Conclusion

Interrogating multiple tissue compartments in patients at various stages of prostate cancer progression has provided key insights to the alterations that occur in the AR and broader AR signaling pathway. While mutation and amplification of the AR gene do not appear to play a major role in prostate tumorigenesis, there is ample evidence that other common genomic alterations including ETS gene fusions, SPOP mutations, and FOXA1 mutations can alter transcriptional regulation of the AR signaling axis. These alterations may be intimately linked to differences in transcriptional output and/or chromatin binding that have been noted in the AR in studies with primary prostate cancer and adjacent normal tissue. Alternatively, mutation and amplification of the AR gene are more frequent in CRPC tissue, and appear to be responses to therapeutic stress as opposed to alterations that underlie metastasis. Altered splicing of the AR gene, manifesting in expression of AR-Vs appears to occur at all stages of disease progression as well as normal prostate tissue, raising questions about the specific context in which these species may play a role in disease progression. Nevertheless, targeted interrogation approaches have revealed a trend towards increased expression of specific AR-Vs, such as AR-V7, in CRPC vs. primary prostate cancer tissue. This is underscored by the potential predictive utility of detecting AR-V7 mRNA in CTCs from patients with CRPC. Reconciling the discrepancies noted in AR-V7 expression patterns in tissue profiling (ie RNA-seq) vs. targeted (ie RT-PCR) studies with tissues or CTCs will be critical for future studies. Additionally, understanding the mechanistic role of AR-Vs in disease progression will require their evaluation in the context of other AR and AR pathway alterations known to occur during disease progression, as well as alterations in AR independent pathways. Moreover, development of inhibitors that can

selectively target the transcriptional activity of AR-Vs will provide important clinical information on their contributions at various disease stages.

Acknowledgements

SMD is currently funded by a Movember/Prostate Cancer Foundation Challenge Award,

American Cancer Society Research Scholar Grant RSG-12-031-01-TBE, NIH grant

R01CA174777, US Department of Defense Prostate Cancer Research Program grants

W81XWH-12-2-0093, W81XWH-13-1-0518, W81XWH-15-1-0633, and W81XWH-15-1-0501,

and a grant from the Minnesota Partnership for Biotechnology and Medical Genomics.

References:

- [1] R.L. Siegel, K.D. Miller, A. Jemal, Cancer statistics, 2015., CA. Cancer J. Clin. 65 (2015) 5–29.
doi:10.3322/caac.21254.
- [2] C. Huggins, C. V Hodges, Studies on prostatic cancer. I. The effect of castration, of estrogen and androgen injection on serum phosphatases in metastatic carcinoma of the prostate., CA. Cancer J. Clin. 22 232–40.
<http://www.ncbi.nlm.nih.gov/pubmed/4625049>.
- [3] C.J. Ryan, M.R. Smith, J.S. de Bono, A. Molina, C.J. Logothetis, P. de Souza, et al., Abiraterone in metastatic prostate cancer without previous chemotherapy., N. Engl. J. Med. 368 (2013) 138–48.
doi:10.1056/NEJMoa1209096.
- [4] H.I. Scher, K. Fizazi, F. Saad, M.-E. Taplin, C.N. Sternberg, K. Miller, et al., Increased survival with enzalutamide in prostate cancer after chemotherapy., N. Engl. J. Med. 367 (2012) 1187–97.
doi:10.1056/NEJMoa1207506.
- [5] E.P. Gelmann, Molecular biology of the androgen receptor., J. Clin. Oncol. 20 (2002) 3001–15.
<http://www.ncbi.nlm.nih.gov/pubmed/12089231>.
- [6] P.A. Watson, Y.F. Chen, M.D. Balbas, J. Wongvipat, N.D. Socci, A. Viale, et al., Constitutively active androgen receptor splice variants expressed in castration-resistant prostate cancer require full-length androgen receptor., Proc. Natl. Acad. Sci. U. S. A. 107 (2010) 16759–65. doi:10.1073/pnas.1012443107.
- [7] Z. Guo, X. Yang, F. Sun, R. Jiang, D.E. Linn, H. Chen, et al., A novel androgen receptor splice variant is up-regulated during prostate cancer progression and promotes androgen depletion-resistant growth., Cancer Res. 69 (2009) 2305–13. doi:10.1158/0008-5472.CAN-08-3795.

- [8] R. Hu, T.A. Dunn, S. Wei, S. Isharwal, R.W. Veltri, E. Humphreys, et al., Ligand-independent androgen receptor variants derived from splicing of cryptic exons signify hormone-refractory prostate cancer., *Cancer Res.* 69 (2009) 16–22. doi:10.1158/0008-5472.CAN-08-2764.
- [9] S. Sun, C.C.T. Sprenger, R.L. Vessella, K. Haugk, K. Soriano, E.A. Mostaghel, et al., Castration resistance in human prostate cancer is conferred by a frequently occurring androgen receptor splice variant., *J. Clin. Invest.* 120 (2010) 2715–30. doi:10.1172/JCI41824.
- [10] R. Hu, W.B. Isaacs, J. Luo, A snapshot of the expression signature of androgen receptor splicing variants and their distinctive transcriptional activities., *Prostate.* 71 (2011) 1656–67. doi:10.1002/pros.21382.
- [11] S.J. Libertini, C.G. Tepper, V. Rodriguez, D.M. Asmuth, H.-J. Kung, M. Mudryj, Evidence for calpain-mediated androgen receptor cleavage as a mechanism for androgen independence., *Cancer Res.* 67 (2007) 9001–5. doi:10.1158/0008-5472.CAN-07-1072.
- [12] S.M. Dehm, L.J. Schmidt, H. V Heemers, R.L. Vessella, D.J. Tindall, Splicing of a novel androgen receptor exon generates a constitutively active androgen receptor that mediates prostate cancer therapy resistance., *Cancer Res.* 68 (2008) 5469–77. doi:10.1158/0008-5472.CAN-08-0594.
- [13] S.M. Dehm, D.J. Tindall, Alternatively spliced androgen receptor variants., *Endocr. Relat. Cancer.* 18 (2011) R183–96. doi:10.1530/ERC-11-0141.
- [14] L.L. Liu, N. Xie, S. Sun, S. Plymate, E. Mostaghel, X. Dong, Mechanisms of the androgen receptor splicing in prostate cancer cells., *Oncogene.* 33 (2014) 3140–50. doi:10.1038/onc.2013.284.
- [15] Y. Li, T.H. Hwang, L.A. Oseth, A. Hauge, R.L. Vessella, S.C. Schmechel, et al., AR intragenic deletions linked to androgen receptor splice variant expression and activity in models of prostate cancer progression., *Oncogene.* 31 (2012) 4759–67. doi:10.1038/onc.2011.637.

- 1
2
3
4
5
6 [16] Y. Li, M. Alsagabi, D. Fan, G.S. Bova, A.H. Tewfik, S.M. Dehm, Intragenic rearrangement and altered
7 RNA splicing of the androgen receptor in a cell-based model of prostate cancer progression., *Cancer Res.* 71
8 (2011) 2108–17. doi:10.1158/0008-5472.CAN-10-1998.
9
10
11
12
13
14 [17] M.D. Nyquist, Y. Li, T.H. Hwang, L.S. Manlove, R.L. Vessella, K.A.T. Silverstein, et al., TALEN-
15 engineered AR gene rearrangements reveal endocrine uncoupling of androgen receptor in prostate cancer.,
16 *Proc. Natl. Acad. Sci. U. S. A.* 110 (2013) 17492–7. doi:10.1073/pnas.1308587110.
17
18
19
20
21
22 [18] H. Beltran, R. Yelensky, G.M. Frampton, K. Park, S.R. Downing, T.Y. MacDonald, et al., Targeted next-
23 generation sequencing of advanced prostate cancer identifies potential therapeutic targets and disease
24 heterogeneity., *Eur. Urol.* 63 (2013) 920–6. doi:10.1016/j.eururo.2012.08.053.
25
26
27
28
29
30 [19] A. Abeshouse, J. Ahn, R. Akbani, A. Ally, S. Amin, C.D. Andry, et al., The Molecular Taxonomy of
31 Primary Prostate Cancer, *Cell.* 163 (2015) 1011–1025. doi:10.1016/j.cell.2015.10.025.
32
33
34
35
36 [20] E. Hörnberg, E.B. Ylitalo, S. Crnalic, H. Antti, P. Stattin, A. Widmark, et al., Expression of androgen
37 receptor splice variants in prostate cancer bone metastases is associated with castration-resistance and short
38 survival., *PLoS One.* 6 (2011) e19059. doi:10.1371/journal.pone.0019059.
39
40
41
42
43
44 [21] H. Zhao, M.A. Coram, R. Nolley, S.W. Reese, S.R. Young, D.M. Peehl, Transcript levels of androgen
45 receptor variant AR-V1 or AR-V7 do not predict recurrence in patients with prostate cancer at indeterminate
46 risk for progression., *J. Urol.* 188 (2012) 2158–64. doi:10.1016/j.juro.2012.08.014.
47
48
49
50
51
52 [22] S.C. Chan, L.A. Selth, Y. Li, M.D. Nyquist, L. Miao, J.E. Bradner, et al., Targeting chromatin binding
53 regulation of constitutively active AR variants to overcome prostate cancer resistance to endocrine-based
54 therapies., *Nucleic Acids Res.* 43 (2015) 5880–97. doi:10.1093/nar/gkv262.
55
56
57
58
59
60
61
62
63
64
65

- [23] S.A. Tomlins, D.R. Rhodes, S. Perner, S.M. Dhanasekaran, R. Mehra, X.-W. Sun, et al., Recurrent fusion of TMPRSS2 and ETS transcription factor genes in prostate cancer., *Science*. 310 (2005) 644–8. doi:10.1126/science.1117679.
- [24] B.G. Barwick, M. Abramovitz, M. Kodani, C.S. Moreno, R. Nam, W. Tang, et al., Prostate cancer genes associated with TMPRSS2-ERG gene fusion and prognostic of biochemical recurrence in multiple cohorts., *Br. J. Cancer*. 102 (2010) 570–6. doi:10.1038/sj.bjc.6605519.
- [25] S. Minner, M. Enodien, H. Sirma, A.M. Luebke, A. Krohn, P.S. Mayer, et al., ERG status is unrelated to PSA recurrence in radically operated prostate cancer in the absence of antihormonal therapy., *Clin. Cancer Res.* 17 (2011) 5878–88. doi:10.1158/1078-0432.CCR-11-1251.
- [26] J.S. Carroll, X.S. Liu, A.S. Brodsky, W. Li, C.A. Meyer, A.J. Szary, et al., Chromosome-wide mapping of estrogen receptor binding reveals long-range regulation requiring the forkhead protein FoxA1., *Cell*. 122 (2005) 33–43. doi:10.1016/j.cell.2005.05.008.
- [27] C.S. Grasso, A.K. Cani, D.H. Hovelson, M.J. Quist, N.J. Douville, V. Yadati, et al., Integrative molecular profiling of routine clinical prostate cancer specimens., *Ann. Oncol.* 26 (2015) 1110–8. doi:10.1093/annonc/mdv134.
- [28] C.E. Barbieri, S.C. Baca, M.S. Lawrence, F. Demichelis, M. Blattner, J.-P. Theurillat, et al., Exome sequencing identifies recurrent SPOP, FOXA1 and MED12 mutations in prostate cancer., *Nat. Genet.* 44 (2012) 685–9. doi:10.1038/ng.2279.
- [29] Y. Nagai, T. Kojima, Y. Muro, T. Hachiya, Y. Nishizawa, T. Wakabayashi, et al., Identification of a novel nuclear speckle-type protein, SPOP., *FEBS Lett.* 418 (1997) 23–6. <http://www.ncbi.nlm.nih.gov/pubmed/9414087>.

- 1
2
3
4 [30] M. Zhuang, M.F. Calabrese, J. Liu, M.B. Waddell, A. Nourse, M. Hammel, et al., Structures of SPOP-
5 substrate complexes: insights into molecular architectures of BTB-Cul3 ubiquitin ligases., *Mol. Cell.* 36
6 (2009) 39–50. doi:10.1016/j.molcel.2009.09.022.
7
8
9
10
11
12 [31] J. An, C. Wang, Y. Deng, L. Yu, H. Huang, Destruction of full-length androgen receptor by wild-type
13 SPOP, but not prostate-cancer-associated mutants., *Cell Rep.* 6 (2014) 657–69.
14
15
16
17
18
19
20 [32] C. Geng, K. Rajapakshe, S.S. Shah, J. Shou, V.K. Eedunuri, C. Foley, et al., Androgen receptor is the key
21 transcriptional mediator of the tumor suppressor SPOP in prostate cancer., *Cancer Res.* 74 (2014) 5631–43.
22
23
24
25
26
27
28
29 [33] C. Geng, B. He, L. Xu, C.E. Barbieri, V.K. Eedunuri, S.A. Chew, et al., Prostate cancer-associated
30 mutations in speckle-type POZ protein (SPOP) regulate steroid receptor coactivator 3 protein turnover.,
31
32
33
34
35
36
37 [34] N. Mitsiades, C.C. Sung, N. Schultz, D.C. Danila, B. He, V.K. Eedunuri, et al., Distinct patterns of
38 dysregulated expression of enzymes involved in androgen synthesis and metabolism in metastatic prostate
39 cancer tumors., *Cancer Res.* 72 (2012) 6142–52. doi:10.1158/0008-5472.CAN-12-1335.
40
41
42
43
44
45 [35] H. Bonkhoff, R. Berges, From pathogenesis to prevention of castration resistant prostate cancer, *Prostate.* 70
46 (2010) 100–112. doi:10.1002/pros.21042.
47
48
49
50
51 [36] P.A. Watson, V.K. Arora, C.L. Sawyers, Emerging mechanisms of resistance to androgen receptor inhibitors
52 in prostate cancer., *Nat. Rev. Cancer.* 15 (2015) 701–11. doi:10.1038/nrc4016.
53
54
55
56
57
58
59
60
61
62
63
64
65

- [37] C.S. Grasso, Y.-M. Wu, D.R. Robinson, X. Cao, S.M. Dhanasekaran, A.P. Khan, et al., The mutational landscape of lethal castration-resistant prostate cancer., *Nature*. 487 (2012) 239–43. doi:10.1038/nature11125.
- [38] D. Robinson, E.M. Van Allen, Y.-M. Wu, N. Schultz, R.J. Lonigro, J.-M. Mosquera, et al., Integrative Clinical Genomics of Advanced Prostate Cancer, *Cell*. 161 (2015) 1215–1228. doi:10.1016/j.cell.2015.05.001.
- [39] M.J. Linja, K.J. Savinainen, O.R. Saramäki, T.L. Tammela, R.L. Vessella, T. Visakorpi, Amplification and overexpression of androgen receptor gene in hormone-refractory prostate cancer., *Cancer Res*. 61 (2001) 3550–5. <http://www.ncbi.nlm.nih.gov/pubmed/11325816>.
- [40] O.H. Ford, C.W. Gregory, D. Kim, A.B. Smitherman, J.L. Mohler, Androgen receptor gene amplification and protein expression in recurrent prostate cancer., *J. Urol*. 170 (2003) 1817–21. doi:10.1097/01.ju.0000091873.09677.f4.
- [41] B.S. Taylor, N. Schultz, H. Hieronymus, A. Gopalan, Y. Xiao, B.S. Carver, et al., Integrative genomic profiling of human prostate cancer., *Cancer Cell*. 18 (2010) 11–22. doi:10.1016/j.ccr.2010.05.026.
- [42] L. Bubendorf, J. Kononen, P. Koivisto, P. Schraml, H. Moch, T.C. Gasser, et al., Survey of gene amplifications during prostate cancer progression by high-throughout fluorescence in situ hybridization on tissue microarrays., *Cancer Res*. 59 (1999) 803–6. <http://www.ncbi.nlm.nih.gov/pubmed/10029066>.
- [43] P. Koivisto, J. Kononen, C. Palmberg, T. Tammela, E. Hyytinen, J. Isola, et al., Androgen receptor gene amplification: a possible molecular mechanism for androgen deprivation therapy failure in prostate cancer., *Cancer Res*. 57 (1997) 314–9. <http://www.ncbi.nlm.nih.gov/pubmed/9000575>.

- [44] D.T. Miyamoto, R.J. Lee, S.L. Stott, D.T. Ting, B.S. Wittner, M. Ulman, et al., Androgen receptor signaling in circulating tumor cells as a marker of hormonally responsive prostate cancer., *Cancer Discov.* 2 (2012) 995–1003. doi:10.1158/2159-8290.CD-12-0222.
- [45] D.R. Shaffer, M.A. Leversha, D.C. Danila, O. Lin, R. Gonzalez-Espinoza, B. Gu, et al., Circulating tumor cell analysis in patients with progressive castration-resistant prostate cancer., *Clin. Cancer Res.* 13 (2007) 2023–9. doi:10.1158/1078-0432.CCR-06-2701.
- [46] G. Attard, J.F. Swennenhuis, D. Olmos, A.H.M. Reid, E. Vickers, R. A’Hern, et al., Characterization of ERG, AR and PTEN gene status in circulating tumor cells from patients with castration-resistant prostate cancer., *Cancer Res.* 69 (2009) 2912–8. doi:10.1158/0008-5472.CAN-08-3667.
- [47] Y. Jiang, J.F. Palma, D.B. Agus, Y. Wang, M.E. Gross, Detection of androgen receptor mutations in circulating tumor cells in castration-resistant prostate cancer., *Clin. Chem.* 56 (2010) 1492–5. doi:10.1373/clinchem.2010.143297.
- [48] G. Gudem, P. Van Loo, B. Kremeyer, L.B. Alexandrov, J.M.C. Tubio, E. Papaemmanuil, et al., The evolutionary history of lethal metastatic prostate cancer, *Nature.* 520 (2015) 353–357. doi:10.1038/nature14347.
- [49] D.T. Miyamoto, Y. Zheng, B.S. Wittner, R.J. Lee, H. Zhu, K.T. Broderick, et al., RNA-Seq of single prostate CTCs implicates noncanonical Wnt signaling in antiandrogen resistance, *Science.* 349 (2015) 1351–1356. doi:10.1126/science.aab0917.
- [50] A. Romanel, D.G. Tandefelt, V. Conteduca, A. Jayaram, N. Casiraghi, D. Wetterskog, et al., Plasma AR and abiraterone-resistant prostate cancer, *Sci. Transl. Med.* 7 (2015) 312re10–312re10. doi:10.1126/scitranslmed.aac9511.

- [51] E.S. Antonarakis, C. Lu, H. Wang, B. Lubber, M. Nakazawa, J.C. Roeser, et al., AR-V7 and Resistance to Enzalutamide and Abiraterone in Prostate Cancer., *N. Engl. J. Med.* 371 (2014) 1028–38. doi:10.1056/NEJMoa1315815.
- [52] E. Efstathiou, M. Titus, S. Wen, A. Hoang, M. Karlou, R. Ashe, et al., Molecular characterization of enzalutamide-treated bone metastatic castration-resistant prostate cancer., *Eur. Urol.* 67 (2015) 53–60. doi:10.1016/j.eururo.2014.05.005.
- [53] J. Steinestel, M. Luedeke, A. Arndt, T.J. Schnoeller, J.K. Lennerz, C. Wurm, et al., Detecting predictive androgen receptor modifications in circulating prostate cancer cells., *Oncotarget.* (2015). <http://www.ncbi.nlm.nih.gov/pubmed/25970787>.
- [54] M. Nakazawa, C. Lu, Y. Chen, C.J. Paller, M.A. Carducci, M.A. Eisenberger, et al., Serial blood-based analysis of AR-V7 in men with advanced prostate cancer., *Ann. Oncol.* 26 (2015) 1859–65. doi:10.1093/annonc/mdv282.
- [55] E.S. Antonarakis, C. Lu, B. Lubber, H. Wang, Y. Chen, M. Nakazawa, et al., Androgen Receptor Splice Variant 7 and Efficacy of Taxane Chemotherapy in Patients With Metastatic Castration-Resistant Prostate Cancer., *JAMA Oncol.* 1 (2015) 582–91. doi:10.1001/jamaoncol.2015.1341.
- [56] M.S. Darshan, M.S. Loftus, M. Thadani-Mulero, B.P. Levy, D. Escuin, X.K. Zhou, et al., Taxane-induced blockade to nuclear accumulation of the androgen receptor predicts clinical responses in metastatic prostate cancer., *Cancer Res.* 71 (2011) 6019–29. doi:10.1158/0008-5472.CAN-11-1417.
- [57] M. Thadani-Mulero, L. Portella, S. Sun, M. Sung, A. Matov, R.L. Vessella, et al., Androgen receptor splice variants determine taxane sensitivity in prostate cancer., *Cancer Res.* 74 (2014) 2270–82. doi:10.1158/0008-5472.CAN-13-2876.

- [58] R. de Leeuw, L.D. Berman-Booty, M.J. Schiewer, S.J. Ciment, R.B. Den, A.P. Dicker, et al., Novel actions of next-generation taxanes benefit advanced stages of prostate cancer., *Clin. Cancer Res.* 21 (2015) 795–807. doi:10.1158/1078-0432.CCR-14-1358.
- [59] Q. Wang, W. Li, Y. Zhang, X. Yuan, K. Xu, J. Yu, et al., Androgen receptor regulates a distinct transcription program in androgen-independent prostate cancer., *Cell.* 138 (2009) 245–56. doi:10.1016/j.cell.2009.04.056.
- [60] Y. Chen, P. Chi, S. Rockowitz, P.J. Iaquina, T. Shamu, S. Shukla, et al., ETS factors reprogram the androgen receptor cistrome and prime prostate tumorigenesis in response to PTEN loss., *Nat. Med.* 19 (2013) 1023–9. doi:10.1038/nm.3216.
- [61] J. Yu, J. Yu, R.-S. Mani, Q. Cao, C.J. Brenner, X. Cao, et al., An integrated network of androgen receptor, polycomb, and TMPRSS2-ERG gene fusions in prostate cancer progression., *Cancer Cell.* 17 (2010) 443–54. doi:10.1016/j.ccr.2010.03.018.
- [62] N.L. Sharma, C.E. Massie, A. Ramos-Montoya, V. Zecchini, H.E. Scott, A.D. Lamb, et al., The androgen receptor induces a distinct transcriptional program in castration-resistant prostate cancer in man., *Cancer Cell.* 23 (2013) 35–47. doi:10.1016/j.ccr.2012.11.010.
- [63] M.M. Pomerantz, F. Li, D.Y. Takeda, R. Lenci, A. Chonkar, M. Chabot, et al., The androgen receptor cistrome is extensively reprogrammed in human prostate tumorigenesis, *Nat. Genet.* 47 (2015) 1346–1351. doi:10.1038/ng.3419.
- [64] S. Stelloo, E. Nevedomskaya, H.G. van der Poel, J. de Jong, G.J. van Leenders, G. Jenster, et al., Androgen receptor profiling predicts prostate cancer outcome., *EMBO Mol. Med.* 7 (2015) 1450–64. doi:10.15252/emmm.201505424.

- [65] R. Hu, C. Lu, E.A. Mostaghel, S. Yegnasubramanian, M. Gurel, C. Tannahill, et al., Distinct transcriptional programs mediated by the ligand-dependent full-length androgen receptor and its splice variants in castration-resistant prostate cancer., *Cancer Res.* 72 (2012) 3457–62. doi:10.1158/0008-5472.CAN-11-3892.
- [66] S.C. Chan, Y. Li, S.M. Dehm, Androgen receptor splice variants activate androgen receptor target genes and support aberrant prostate cancer cell growth independent of canonical androgen receptor nuclear localization signal., *J. Biol. Chem.* 287 (2012) 19736–49. doi:10.1074/jbc.M112.352930.
- [67] J. Lu, P.E. Lonergan, L.P. Nacusi, L. Wang, L.J. Schmidt, Z. Sun, et al., The cistrome and gene signature of androgen receptor splice variants in castration resistant prostate cancer cells., *J. Urol.* 193 (2015) 690–8. doi:10.1016/j.juro.2014.08.043.
- [68] D. Xu, Y. Zhan, Y. Qi, B. Cao, S. Bai, W. Xu, et al., Androgen Receptor Splice Variants Dimerize to Transactivate Target Genes., *Cancer Res.* 75 (2015) 3663–71. doi:10.1158/0008-5472.CAN-15-0381.

Fig. 1 Canonical/resistance AR signaling and AR-V structure. A. Hypothetical timeline representing progression of primary prostate cancer to castration resistant prostate cancer. B. Tissues and other clinical material (CTCs-circulating tumor cells; ctDNA-cell free, circulating tumor DNA) available for interrogation of alterations in the androgen/AR

1
2
3
4 pathway. C. Illustration of normal and aberrant modes of androgen/AR signaling. The
5
6 upper left quadrant summarizes canonical AR signaling that occurs in normal development
7
8 and prostate cancer. In this setting, DHT binds AR, which promotes AR translocation to the
9
10 nucleus and transcriptional activation. Components of this axis targeted by abiraterone and
11
12 enzalutamide are indicated. The top right quadrant illustrates how amplification of the AR
13
14 locus results in AR protein overexpression, which is highly sensitive to castrate levels of
15
16 androgen. The bottom right quadrant illustrates the domain structure of AR-Vs. The
17
18 expanded view displays exon composition of 5 separate mRNAs that encode discrete AR-Vs
19
20 that are detailed within the text. The bottom left quadrant shows how mutations in the AR
21
22 ligand binding domain can convert non AR ligands and AR antagonists to agonists, leading
23
24 to transcription of AR target genes.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 1