**Syntheses of chondroitin sulfate tetrasaccharide structures containing
4,6-disulfate patterns and analysis of their interaction with
glycosaminoglycan-binding protein**

Kento Miyachi,¹ Masahiro Wakao,^{1*} Yasuo Suda^{1,2*}

¹Department of Chemistry, Biotechnology, and Chemical Engineering, Graduate School of
Science and Engineering, Kagoshima University, 1-21-40 Kohrimoto, Kagoshima 890-0065,
Japan

²SUDx-Biotec Corporation, 1-41-1 Shiroyama, Kagoshima 890-0013, Japan

AUTHOR EMAIL ADDRESS

wakao@eng.kagoshima-u.ac.jp, ysuda@eng.kagoshima-u.ac.jp

AUTHOR PHONE&FAX

TEL: (+81)99-285-7843, FAX: (+81)99-285-7856 for M. Wakao

TEL & FAX: (+81)99-285-8369 for Y. Suda

Abstract

Chondroitin sulfate tetrasaccharide ligand conjugates, namely GlcA-GalNAc6S-GlcA-GalNAc4S6S (CS-C+E) **1**, GlcA2S-GalNAc6S-GlcA2S-GalNAc4S6S (CS-D+T) **2**, GlcA-GalNAc4S6S-GlcA-GalNAc4S (CS-E+A) **3**, GlcA-GalNAc4S6S-GlcA-GalNAc6S (CS-E+C) **4**, and GlcA-GalNAc4S6S-GlcA-GalNAc4S6S (CS-E+E) **5**, were systematically synthesized using a disaccharide building block **6**. Synthesized CS tetrasaccharide structures were immobilized onto gold-coated chips to prepare array-type sugar chips, and the binding properties of protein were evaluated by surface plasmon resonance imaging biosensor. CS-D+T, CS-E+A, CS-E+C, and CS-E+E showed greater affinity for basic fibroblast growth factor than did other tetrasaccharides (CS-C+D, C+E, D+D).

Keyword

basic fibroblast growth factor; chondroitin sulfate; glycosaminoglycan; sugar chips; surface plasmon resonance; synthesis

Text

Chondroitin sulfate (CS) is a glycosaminoglycan (GAG) that is widely distributed throughout various tissues as components of the cell membrane or extracellular matrix. CS possesses a repeating disaccharide structure that is composed of glucuronic acid (GlcA) and *N*-acetylgalactosamine (GalNAc), and it is often sulfated heterogeneously by multiple and random enzymatic modifications during its biosynthesis.¹ The structure of CS is classified on the basis of the disaccharide unit; type A (CS-A), type C (CS-C), type D (CS-D), and type E (CS-E) units are the major disaccharide structures known from natural sources to date. CS-E, containing a 4, 6-disulfated GalNAc moiety, interacts with various bioactive proteins, such as growth factors, cytokines, and extracellular matrix proteins, and regulates their biofunctions.²⁻⁸ Therefore, the structure–activity relationship of the CS-E unit at the molecular level has been intensively studied to gain understanding of its biofunction.

Although much effort⁹⁻²² has been invested to obtain a structurally defined CS-E oligosaccharide, the synthesis of oligosaccharides containing a CS-E unit is still challenging. In particular, the synthesis of a hetero-dimeric CS tetrasaccharide structure has not been reported. In this study, we attempted the systematic synthesis of CS tetrasaccharide ligand conjugates containing homo- or hetero-dimeric CS-E structures and their analogs. Additionally, we used surface plasmon resonance (SPR) imaging to investigate the binding properties of CS-binding proteins using the tetrasaccharide structures we had synthesized.

For efficient syntheses of the CS tetrasaccharide structures **1-5** (Figure 1), trifluoroimidates **9** and **13** were prepared as glycosyl donors; these compounds were utilized for the generation of CS-C and -D and CS-E units, respectively (Scheme 1). Trifluoroimide **9** was induced by selective ring-opening of the benzylidene group, whereas trifluoroimide **13** was induced by acidic hydrolysis of the benzylidene group. For both compounds, the

induction was followed by levunylation from the disaccharide building block **6**.

Synthesis of the acceptor moiety was carried out as shown in Scheme 2. Three kinds of acceptor moieties were derived from a common trisaccharide intermediate **14**. Desilylation of the 4" position yielded acceptor **15**, which was used for the construction of the CS-E and -T units. After selective ring-opening of the benzylidene group and levulinoylation, removal of the *tert*-butyldimethylsilyl (TBDMS) group yielded acceptors **18** and **20**, which were used for construction of the CS-A and CS-C units, respectively.

With these glycosyl donors and acceptors, we then attempted the synthesis of pentasaccharides and their ligand conjugates (Scheme 3). Glycosylation with an appropriate combination of glycosyl donor and acceptor was carried out in the presence of trimethylsilyl trifluoromethanesulfonate (TMSOTf) to produce the corresponding pentasaccharides. The pentasaccharides so obtained were then converted into CS tetrasaccharide structures and their ligand conjugates via a suitable deprotection and sulfation protocol. Pentasaccharides **21**, **22**, **23**, and **24** were treated with a Zn–Cu couple in AcOH, and the amino groups were acetylated. Selective deprotection of the levulinoyl (Lev) group was performed by using hydrazine hydrate in the presence of AcOH. The benzoyl (Bz) groups and methyl esters were hydrolyzed under basic conditions (with NaOH). The resulting free hydroxyl groups were sulfated with a sulfur trioxide-pyridine complex at 50 °C. After deprotection of the TBDMS group with HF•pyridine, all of the benzyl protective groups were removed by hydrogenolysis with a catalytic amount of Pd/C. Pentasaccharides **25**, **26**, **27**, **28**, and **29** were purified by ODS column chromatography, followed by gel filtration chromatography with Sephadex G-10 or G-15 fine to produce the desired pentasaccharides containing any of the following two disaccharide units: CS-C+E, CS-D+T, CS-E+A, CS-E+C, or CS-E+E. Ligand conjugates were synthesized with these pentasaccharides, according to previously described methods,^{23,24}

yielding CS-tetrasaccharide ligand conjugates **1**, **2**, **3**, **4**, and **5**. These compounds were purified by ODS chromatography and gel filtration chromatography, using Sephadex G-10 or G-15 fine, and their structures were confirmed by ^1H NMR and ESI-TOF/MS analyses.

Binding interactions were investigated using an SPR imaging biosensor, which allows real-time analysis of interactions without requiring labeling of analytes. Basic fibroblast growth factor (FGF-2) was chosen as the GAG-protein, because FGF-2 is known to interact strongly with the CS-E structure; we therefore investigated the structure–activity relationship of the CS-E unit-containing microdomain structure in detail. Specific binding interactions were clearly observed with ligand-conjugates containing the CS-T disaccharide²⁵, CS-E+A, CS-E+C, CS-E+D²⁶, and CS-E+E tetrasaccharides (Figure 2a). The kinetic parameters of these interactions are summarized in Table 1. It has been reported that FGF-2 is strongly recognized by tetrasaccharides containing CS-E unit structures.²⁷⁻³² To characterize the properties of CS tetrasaccharide structures in greater detail, sugar chips, immobilized with a mixture of CS ligand-conjugates prepared in this study and with a Glc ligand-conjugate **39**^{23,33} which does not bind FGF-2 (Figure 2b: 1:2 molar ratio, Figure 2c: 1:8 molar ratio), were fabricated and tested. FGF-2 bound strongly to the CS-T disaccharide and CS-D+T, -E+A, -E+C, -E+D, and -E+E tetrasaccharides (Figure 2b, 2c). The binding affinity of the CS tetrasaccharide structures differed markedly, depending on whether they had a CS-E unit (CS-E+A, -E+C, -E+D, and -E+E) or a CS-D unit (CS-D+D²⁶, CS-D+T) at the non-reducing end. This indicated that the CS disaccharide unit at the reducing end was also recognized by the protein. Furthermore, our sugar chip technique showed that the inverse sequence had different binding properties; the CS-E+C tetrasaccharide interacted strongly with FGF-2, whereas the CS-C+E tetrasaccharide did not.

In conclusion, we synthesized five types of homo- and hetero-dimeric CS

tetrasaccharide structures with a 4,6-disulfated pattern, viz., CS-C+E **1**, CS-D+T **2**, CS-E+A **3**, CS-E+C **4**, and CS-E+E **5**. In addition, we fabricated sugar chips with these synthesized CS ligand-conjugates and analyzed their binding interaction with FGF-2. The results showed that our sugar chip technology can be utilized for real-time analysis of the structure–function relationship of GAG-binding proteins at the molecular level in real-time, without requiring labeling of analytes.. Although construction of a CS tetrasaccharide library is underway, our CS sugar chips may be used for high-throughput analysis of the binding properties of protein samples in order to decipher the CS sugar code.

Acknowledgements

The present study was financially supported in part by grants from the Ministry of Education, Culture, Sports, Science, and Technology (MEXT; Grant-in-Aid for Young Scientists B, Kakenhi 21710233, to MW and Grant-in-Aid for Scientific Research B, Kakenhi 23300356, to YS), Japan Science and Technology Agency (JST) (A-Step, Post-venture Support, AS2416907P to YS), and Japan Ministry for Health, Labor, and Welfare (Nano-Medicine, H21-nano-ippan-002 to YS).

References

1. Silbert, J. E.; Sugumaran, G. *IUBMB Life* **2002**, *54*, 177.
2. Ueoka, C.; Kaneda, N.; Okazaki, I.; Nadanaka, S.; Muramatsu, T.; Sugahara, K. *J. Biol. Chem.* **2000**, *277*, 37407.
3. Takagaki, K.; Munakata, H.; Kakizaki, I.; Iwafune, M.; Itabashi, T.; Endo, M. *J. Biol. Chem.* **2002**, *277*, 8882.
4. Kawashima, H.; Atarashi, K.; Hirose, M.; Hirose, J.; Yamada, S.; Sugahara, K.; Miyasaka, M. *J. Biol. Chem.* **2002**, *277*, 12921.
5. Deepa, S.S.; Umehara, Y.; Higashiyama, S.; Itoh, N.; Sugahara, K. *J. Biol. Chem.* **2002**, *277*, 43707.
6. Sugahara, K.; Mikami, T.; Uyama, T.; Mizuguchi, S.; Nomura, K.; Kitagawa, H. *Curr. Opin. Chem. Biol.* **2003**, *13*, 612.
7. Hashiguchi, T.; Mizumoto, S.; Yamada, S.; Sugahara, K. *Glycoconj. J.* **2010**, *27*, 49.
8. Mizumoto, S.; Fongmoon, D.; Sugahara, K. *Glycoconj. J.* **2013**, *30*, 619.
9. Jacquinet, J. C. *Carbohydr. Res.* **1990**, *199*, 153.
10. Lubineau, A.; Bonnaffé, D. *Eur. J. Org. Chem.* **1999**, 2523.
11. Fujikawa, S.; Ohmae, M.; Kobayashi S. *Biomacromol.* **2005**, *6*, 2935.
12. Jacquinet, J.C.; Lopin-Bon, C.; Vibert, A. *Chem. Eur. J.* **2009**, *15*, 9579.
13. Tamura, J.; Neumann, K.W.; Ogawa, T. *Bioorg. Med. Chem. Lett.* **1995**, *5*, 1351.
14. Tamura, J.; Neumann, K. W.; Kurono, S.; Ogawa, T. *Carbohydr. Res.* **1998**, *305*, 43.
15. Karst, N.; Jacquinet, J.C. *Eur. J. Org. Chem.* **2002**, 815.
16. Tully, S. E.; Mabon, R.; Gama, C.I.; Tsai, S.M.; Liu, X.; Hsieh-Wilson, L. *J. Am. Chem.*

Soc. **2004**, 126, 7736.

17. Gama, C.I.; Tully, S.E.; Sotogaku, N.; Clark, P.M.; Rawat, M.; Vaidehi, N.; Goddard, W. A., III.; Nishi, A.; Hsieh-Wilson, L.C. *Nat. Chem. Biol.* **2006**, 2, 467.
18. Tamura, J.; Tsutsumisita-Nakai, N.; Nakao, Y.; Kawano, M.; Kato, S.; Takeda, N.; Nadanaka, S.; Kitagawa, H. *Bioorg. Med. Chem. Lett.* **2012**, 22, 1371.
19. Bedini, E.; Castro, C.D.; De Rosa, M.; Di Nola, A.; Restaino, O.F.; Schiraldi, C.; Parrilli, M. *Chem. Eur. J.* **2012**, 18, 2123.
20. Maza, S.; Mar Kayser, M.; Macchione, G.; López-Prados, J.; Angulo, J.; de Paz, J.L.; Nieto, P.M. *Org. Biomol. Chem.* **2013**, 42, 3510.
21. Macchione, G.; Maza, S.; Mar Kayser, M.; de Paz, J. L.; Nieto, P. M. *Eur. J. Org. Chem.* **2014**, 3868.
22. Jacquinet, J. C.; Lopin-Bon, C. *Carbohydr. Res.* **2015**, 402, 35.
23. Suda, Y.; Arano, A.; Fukui, Y.; Koshida, S.; Wakao, M.; Nishimura, T.; Kusumoto, S.; Sobel, M. *Bioconjugate Chem.* **2006**, 17, 1125.
24. Wakao, M.; Saito, A.; Ohishi, K.; Kishimoto, Y.; Nishimura, T.; Sobel, M.; Suda, Y. *Bioorg. Med. Chem. Lett.* **2008**, 18, 2499.
25. Wakao, M.; Obata, R.; Miyachi, K.; Kakitsubata, Y.; Kondo, T.; Sakami, C.; Suda, Y. *submitted*.
26. Miyachi, K.; Wakao, M.; Ichiki, Y.; Sakami, C.; Suda, Y. *submitted*.
27. Turnbell, J.E.; Fernig, D.G.; Ke, Y.; Wilkinson, M.C.; Gallagher, J.T. *J. Biol. Chem.* **1992**, 267, 10337.
28. Habuchi, H.; Suzuki, S.; Saito, T.; Tamura, T.; Harada, T.; Yoshida, K.; Kimata, K. *Biochem. J.* **1992**, 285, 805.
29. Nandini, C.D.; Mikami, T.; Ohta, M.; Itoh, N.; Akiyama-Nambu, F.; Sugahara, K. *J. Biol.*

Chem. **2004**, 279, 50799.

30. Bao, X.F.; Nishimura, S.; Mikami, T.; Yamada, S.; Itoh, N.; Sugahara, K. *J. Biol. Chem.* **2004**, 279, 9765.
31. Shipp, E.L.; Hsieh-Wilson, L.C. *Chem. Biol.* **2007**, 14, 195.
32. Rogers, C.J.; Clark, P. M.; Tully, S.E.; Abrol, R.; Garcia, K.C; Goddard III, W.A.; Hsieh-Wilson, L.C. *Proc. Nat. Acad. Sci.* **2011**, 108, 9747.
33. Nakamura-Tsuruta, S.; Kishimoto, Y.; Nishimura, T.; Suda, Y. *J. Biochem.* **2008**, 143, 833.

Legends to Schemes and Figures

Figure 1. Tetrasaccharide structures **1-5**.

Figure 2. Binding study of FGF-2 and chondroitin sulfate (CS) subunits using SPR imaging. Measurements were carried out with analytes in the concentration range between 8 and 500 nM. Bar graph profiles of different concentrations of FGF-2 at the maximum binding intensity of SPR imaging. The CS sugar chain and Glc ratio were (a) 1:0, (b) 1:2, and (c) 1:8. Abbreviations: CS-A = GlcA-GalNAc4S-Glc ligand conjugate **31** was immobilized; CS-C = GlcA-GalNAc6S-Glc ligand conjugate **32** was immobilized; CS-D = GlcA2S-GalNAc6S-Glc

ligand conjugate **33** was immobilized; CS-E = GlcA-GalNAc4S6S-Glc ligand conjugate **34** was immobilized; CS-T = GlcA2S-GalNAc4S6S-Glc ligand conjugate **35** was immobilized; CS-C+D = GlcA-GalNAc6S-GlcA2S-GalNAc6S-Glc ligand conjugate **36** was immobilized; CS-C+E = GlcA-GalNAc6S-GlcA-GalNAc4S6S-Glc ligand conjugate **1** was immobilized; CS-D+D = GlcA2S-GalNAc6S-GlcA2S-GalNAc6S-Glc ligand conjugate **37** was immobilized; CS-D+T = GlcA2S-GalNAc6S-GlcA2S-GalNAc4S6S-Glc ligand conjugate **2** was immobilized; CS-E+A = GlcA-GalNAc4S6S-GlcA-GalNAc4S-Glc ligand conjugate **3** was immobilized; CS-E+C = GlcA-GalNAc4S6S-GlcA-GalNAc6S-Glc ligand conjugate **4** was immobilized; CS-E+D = GlcA-GalNAc4S6S-GlcA2S-GalNAc6S-Glc ligand conjugate **38** was immobilized; CS-E+E= GlcA-GalNAc4S6S-GlcA-GalNAc4S6S-Glc ligand conjugate **5** was immobilized; Glc = Glc-Glc (maltose) ligand conjugate **39** was immobilized.

Scheme 1. Synthesis of glycosyl donor **9** and **13**. (a) Et_3SiH , PhBCl_2 in CH_2Cl_2 , -78°C , 10 min; LevOH, DMAP, DCC in CH_2Cl_2 , r.t., 2 h, 90% (2 steps); (b) $\text{Ir}[(\text{cod})(\text{PMePh}_2)_2]\text{PF}_6$, H_2 in THF, r.t., 30 min; I_2 , H_2O in THF, r.t., 1 h, 93% (2 steps); (c) $\text{CF}_3\text{C}(=\text{NPh})\text{Cl}$, K_2CO_3 in CH_2Cl_2 , 18 h; (d) TFA, H_2O in CH_2Cl_2 , 0°C , 2 h, 89%; (e) LevOH, DMAP, DCC in CH_2Cl_2 , r.t., 2 h, 97%; (f) $\text{Ir}[(\text{cod})(\text{PMePh}_2)_2]\text{PF}_6$, H_2 in THF, r.t., 30 min; I_2 , H_2O in THF, r.t., 1 h, 98% (2 steps); (g) $\text{CF}_3\text{C}(=\text{NPh})\text{Cl}$, K_2CO_3 in CH_2Cl_2 , 12 h.

Scheme 2. Syntheses of glycosyl acceptor **15**, **18**, and **20**. (a) HF•pyridine in THF, Pyr., 0 °C → r.t., 24 h, 97 %; (b) Et₃SiH, TfOH in CH₂Cl₂, -78 °C, 10 min, 82%; (c) LevOH, DMAP, DCC in CH₂Cl₂, r.t., 2 h, 84%; (d) HF•pyridine in THF, Pyr., 0 °C → r.t., 19 h, 94%; (e) Et₃SiH, PhBCl₂ in CH₂Cl₂, -78 °C, 10 min; LevOH, DMAP, DCC in CH₂Cl₂, r.t., 2 h, 90% (2 steps); (f) HF•pyridine in THF, Pyr., 0 °C → r.t., 24 h, 97%.

Scheme 3. (a) TMSOTf in CH₂Cl₂, -20 °C, 1 h, 50% (2 steps) for **9** + **15** to **21**, 45% (2 steps) for **13** + **18** to **22**, 53% (2 steps) for **13** + **20** to **23**, 40% (2 steps) for **13** + **15** to **24**; (b) Zn–Cu couple in AcOH, r.t., 1-3 h; Ac₂O in Pyr., 0 °C, 2-4 h, 47% (2 steps) for **21** to **25** and **26**, Zn–Cu couple in AcOH; Ac₂O in Pyr., 36% (2 steps) for **22** to **27**, 38% (2 steps) for **23** to **28**, 31% (2 steps) for **24** to **29**; (c) hydrazine hydrate, AcOH in Pyr., r.t., 0.5-1 h. 99% for **21** to **25** and **26**, hydrazine hydrate, AcOH in Pyr., 63% for **22** to **27**, 81% for **23** to **28**, 85% for **24** to **29**; (d) TFA, H₂O in CH₂Cl₂, 0 °C, 2-4 h, 62% (recovery 30%) for **21** to **25** and **26**, 70% (recovery 25%) for **24** to **29**; (e) NaOH aq., H₂O in THF, 0 °C, 4 h; NaOH aq. in MeOH/THF, 0 °C, 1 d, 92% (2 steps); (f) SO₃•Pyr. in Pyr., 55 °C, 10 h; HF•pyridine in Pyr., 0 °C, 3-4 d, 35% (2 steps); (g) SO₃•Pyr. in Pyr., 50-60 °C, 4-16 h; HF•pyridine in Pyr., 0 °C, 3 d; NaOH aq., H₂O in THF, 0 °C, 2-4 h; NaOH aq. in MeOH/THF, 0 °C, 1 d, 38% (4 steps) for **21** to **25**, 35% (4 steps) for **22** to **27**, 41% (4 steps) for **23** to **28**, 36% (4 steps) for **24** to **29**; (h) Pd/C, H₂ (7.5 atm) in MeOH/H₂O/AcOH, r.t., 3-4 d, 99% for **21** to **25**, 93% for **21** to **26**, 97% for **22**

to **27**, 86% for **23** to **28**, 97% for **24** to **29**; (i) **30** then NaBH₃CN in DMAc/H₂O/AcOH, 40 °C, 3-4 d, 45% (recovery: 50%) for **25** to **1**, 40% (recovery: 52%) for **26** to **2**, 32% (recovery: 65%) for **27** to **3**, 73% (recovery: 20%) for **28** to **4**, 25% (recovery: 60%) for **29** to **5**.

Table 1. Kinetic parameters of the interaction between FGF-2 and chondroitin sulfate (CS) oligosaccharides.

Unit structure	k_a ($M^{-1}s^{-1}$)	k_d (s^{-1})	K_D (nM)
CS-A	N.D.*	N.D.*	—
CS-C	1.2×10^5	4.2×10^{-3}	36
CS-D	9.9×10^4	2.3×10^{-3}	23
CS-E	9.1×10^4	1.5×10^{-3}	16
CS-T	3.8×10^4	8.2×10^{-4}	21
CS-C+D	N.D.*	N.D.*	—
CS-C+E	N.D.*	N.D.*	—
CS-D+D	7.9×10^4	3.8×10^{-3}	48
CS-D+T	4.7×10^4	7.7×10^{-4}	16
CS-E+A	1.0×10^5	5.6×10^{-4}	5.6
CS-E+C	5.6×10^4	4.6×10^{-4}	8.2
CS-E+D	6.9×10^4	3.2×10^{-4}	4.7
CS-E+E	8.5×10^4	4.4×10^{-4}	5.2

*N.D., not determined due to weak interaction

Graphical Abstract

Syntheses of chondroitin sulfate tetrasaccharide structures containing 4,6-disulfate patterns and analysis of their interaction with glycosaminoglycan-binding protein

Kento Miyachi,¹ Masahiro Wakao,^{1*} Yasuo Suda^{1,2*}

CS tetrasaccharide structures

CS-C+E (GlcA-GalNAc6S-GlcA-GalNAc4S6S)

CS-D+T (GlcA2S-GalNAc6S-GlcA2S-GalNAc4S6S)

CS-E+A (GlcA-GalNAc4S6S-GlcA-GalNAc4S)

CS-E+C (GlcA-GalNAc4S6S-GlcA-GalNAc6S)

CS-E+E (GlcA-GalNAc4S6S-GlcA-GalNAc4S6S)