Synthesis and P1' SAR Exploration of Potent Macrocyclic Tissue Factor-Factor VIIa Inhibitors

Vladimir (Uladzimir) Ladziata,^{*} Peter W. Glunz, Yan Zou,[†] Xiaojun Zhang, Wen Jiang, Swanee Jacutin-Porte, Daniel L. Cheney, Anzhi Wei, Joseph M. Luetzgen, Timothy M. Harper, Pancras C. Wong, Dietmar Seiffert, Ruth R. Wexler, E. Scott Priestley

Bristol-Myers Squibb Research & Development, Princeton, New Jersey 08543, United States

ARTICLE INFO

Article history:

Received

Revised

Accepted

Available online

Keywords:

Thrombosis

Anticoagulant

Factor VIIa-TF inhibitor

Macrocycle

ABSTRACT

Selective tissue factor-factor VIIa complex (TF-FVIIa) inhibitors are viewed as promising compounds for treating thrombotic disease. In this contribution, we describe multifaceted exploratory SAR studies of S1'-binding moieties within a macrocyclic chemotype aimed at replacing cyclopropyl sulfone P1' group. Over the course of the optimization efforts, the 1-(1*H*-tetrazol-5-yl)cyclopropane P1' substituent emerged as an improved alternative, offering increased metabolic stability and lower clearance, while maintaining excellent potency and selectivity.

2016 Elsevier Ltd. All rights reserved.

Thrombosis-related disorders, such as myocardial infarction (MI), thromboembolic stroke, deep vein thrombosis (DVT) and pulmonary embolism (PE), are the leading cause of death in the industrialized world.¹ Traditional therapies, which include low molecular weight heparin (LMWH) and warfarin, suffer from a number of drawbacks which limit their utility for long-term use. Namely, LMWH is administered only via parenteral route, while use of warfarin is complicated by a slow onset of action, food effects, and requires close monitoring due to a substantial bleeding risk.² Therefore, in the past several decades much research has focused on discovery and development of thrombin and factor Xa small molecule inhibitors with improved pharmacokinetic and pharmacodynamic parameters.³ Among approved novel oral anticoagulants (NOACs), which target these two specific blood coagulation cascade enzymes, are the thrombin inhibitor dabigatran etexilate (Pradaxa[®]),^{4a-d} and the factor Xa inhibitors rivaroxaban (Xarelto[®]),^{4e-h} apixaban (Eliquis[®]),^{4i-k} and edoxaban (Savaysa[®]).^{4l,m} The introduction of these novel antithrombotics into the clinical practice enables replacement of the traditional therapies in many patients.⁵

Given the fact that the tissue factor-factor VIIa complex (TF-FVIIa) is situated at a critical initiation point of the coagulation

cascade, factor VIIa inhibitors are regarded as a pharmacologically attractive class of anticoagulants, and much pharmaceutical research efforts over the past two decades have been devoted to finding orally bioavailable TF-FVIIa small molecule inhibitors.⁶ The TF-VIIa complex plays a key role in initiation of the extrinsic coagulation pathway: circulating zymogen FVII interacts with membrane-anchored TF, which is exposed upon pathophysiological vascular injury or atherosclerotic plaque rupture, thereby forming the proteolytically active TF-FVIIa complex.^{7a} The initial step is then amplified/propagated via activation of factors IX and X, leading to direct generation of additional FVIIa in the former case, and thrombin formation in the latter; thrombin in turn cleaves fibrinogen to fibrin and mobilizes platelets through protease-activated receptors, which ultimately results in stable clot formation.^{7b} Moreover, FVIIa target validation was accomplished in numerous *in vivo* models utilizing biological agents,^{8a-c} as well as multiple small molecule chemotypes,^{8d-1} where effective anticoagulation coupled with low risk of bleeding was consistently observed. Furthermore, apart from their apparent utility as antithrombotics, TF-FVIIa inhibitors may also possess clinical potential in tumor metastasis via inhibition of protease-activated receptor 2 (PAR2) signaling,⁹ and in inflammation.¹⁰

^{*} Corresponding author. Tel.: +1-609-466-5055; fax: +1-609-818-3550; e-mail: vladimir.ladziata@bms.com.

[†] Co-author deceased. Date of death: 03/21/2014.

Figure 1. Selected structures of disclosed TF-FVIIa inhibitors **1-8**.

A number of diverse series of small molecule TF-FVIIa inhibitors have been reported to date.¹¹ Based on the interaction with the polar primary specificity pocket, the reported FVIIa inhibitors could be nominally divided into two categories. The first group is comprised of compounds bearing strongly basic benzamidine P1 moieties,^{12a} which are characterized by high TF-FVIIa inhibitory potency, but exhibit low permeability and poor oral absorbance. Conversely, the second group includes compounds with less basic non-benzamidine P1 groups, which ordinarily possess improved permeability, albeit at the expense of TF-FVIIa binding affinity. Some representative examples of benzamidine (**1-4**)^{12b-f} and non-benzamidine (**5-8**)^{12g-j} small molecule TF-FVIIa inhibitors are shown in **Figure 1**. Very recently our research team reported novel 16-membered macrocyclic FVIIa inhibitors, bearing an aminoisoquinoline P1 group.^{12j} Macrocycle **8** is a potent FVIIa inhibitor with excellent selectivity (>500 fold) against primary serine protease anti-targets FIXa, FXa, thrombin and trypsin, and excellent FVII deficient prothrombin time¹³ EC_{2x} of 1.2 μ M. Furthermore, the macrocycle provided a rigid scaffold allowing for direct optimization of P1, P2 and P1' groups.

Considering TF-FVIIa inhibition is likely to provide an apparent therapeutic benefit for various thrombosis-related diseases, our research laboratories aimed at developing potent and selective TF-FVIIa inhibitors with improved ADME properties. Building on the initial success of the macrocyclization strategy applied to FVIIa inhibitors, Glunz *et al.* further enhanced this approach by utilizing rational structure-guided atropisomer control via incorporation of a P2 benzylic methyl group, and thereby locking macrocyclic scaffold in the bioactive conformation.¹⁴ From the resulting conformationally stable macrocyclic template, Zhang *et al.* subsequently undertook detailed structure-activity relationship (SAR) studies in the P1 and P1' regions of the macrocyclic core.¹⁵ After surveying a number of P1' moieties, cyclopropyl sulfone was identified as an optimal group. Namely, compound **9** demonstrated outstanding TF-FVIIa binding affinity and anti-clotting activity (K_i = 0.16 nM, EC_{2x} = 1.1 μ M) coupled with an excellent selectivity profile (**Figure 2**).

Despite having good metabolic stability (dog liver microsomes¹⁶ $t_{1/2}$ = 106 min), inhibitor **9** showed elevated dog clearance at 29 mL/min/kg, indicating rapid elimination of the compound. In addition, FVIIa inhibitor **9** demonstrated very low dog oral exposure, presumably due to limited permeability.¹⁵ Indeed, passive Caco-2 permeability¹⁷ for macrocycle **9** measured lower at 47 nm/s.

Figure 2. Comparison of sulfone P1' **9** and des-sulfone P1' **10**.

We envisioned that polar sulfone functionality in inhibitor **9** could be partially responsible for the poor dog pharmacokinetic (PK) profile. To that end, we synthesized analog **10**, where the $-SO_2-$ fragment is excised, and the cyclopropyl group is directly attached to the macrocyclic scaffold (**Figure 2**). We were delighted to learn that upon p.o. and i.v. administration of compound **10** in dogs oral bioavailability and clearance were much improved. It should be noted that whereas analog **10**, in comparison to sulfone counterpart **9**, possessed improved passive Caco-2 permeability and maintained good metabolic stability, TF-FVIIa K_i was reduced (~12 fold), along with selectivity against a number of serine proteases (data not shown). Therefore, we decided to modify macrocycle **10** at the cyclopropyl P1' region in order to improve TF-FVIIa inhibitory activity and selectivity, while at the same time maintaining the ADME profile. Equipped with structural information for the S1' binding pocket we obtained previously,^{12e,12j,18} we aimed at expanding the cyclopropyl ring in order to enhance hydrophobic interaction with the Cys42-Cys58 disulfide bridge, as well as engaging

Lys60A side chain via incorporating acidic groups as shown in **Figure 3**. Since we previously demonstrated that the 4-fluoroaminoisoquinoline P1 group imparts better permeability due to reduced basicity,¹⁵ we decided to use 4-fluoroaminoisoquinoline bearing macrocyclic scaffolds for the P1' SAR examination described below.

Figure 3. Design strategy for novel macrocyclic FVIIa inhibitors.

Synthesis of cyclopropyl P1' precursor **18** is depicted in **Scheme 1**. Commercially available 2-bromobenzaldehyde **11** was nitrated, and the obtained nitrobenzaldehyde **12** was treated sequentially with MeNH₂ and NaBH₄ in MeOH to afford N-methyl benzylamine **13** in good yield. Compound **13** was subsequently converted to the corresponding Boc- and Cbz-protected derivatives **14** and **15** under standard conditions. Synthesis of 4-cyclopropylaniline **18** was achieved in three steps starting with Boc-protected bromobenzylamine **14** via sequential Pd-catalyzed cyclopropanation,¹⁹ followed by Zn-mediated reduction of the nitro group, and Boc removal. It should be noted that reduction of **16** to **17** under hydrogenative conditions (H₂ (1 atm), Pd-C) resulted in a partial cyclopropyl ring opening giving rise to the respective *n*-propyl contaminant.

Scheme 1. Synthesis of 1-cyclopropyl P1' building block **18**. Reagents and conditions: (a) HNO₃, H₂SO₄, 0 °C, 2 h; (b) i. MeNH₂, MeOH, rt, 1 h; ii. NaBH₄, MeOH, 0 °C to rt, 5 h; (c) Boc₂O, TEA, THF, 40 °C, 2 h; (d) Cbz-Cl, TEA, THF, 0 °C to rt, 3 h; (e) *c*PrB(OH)₂, Pd(OAc)₂, PCy₃, K₃PO₄, PhMe-H₂O, 100 °C, 12 h; (f) Zn, NH₄Cl, MeOH-THF, rt then 40 °C, 3 h; (g) HCl, dioxane, 30 min.

In order to obtain advanced intermediates to explore the Lys60A binding vector as shown in **Figure 3**, we carried out a six step sequence starting with Cbz-protected bromoarene **15**. As shown in **Scheme 2**, CuI/(*S*)-Pro-catalyzed C-arylation of malonates, disclosed by Ma and co-workers,²⁰ afforded corresponding malonate **20** and cyanoacetate **21** in excellent yields. Intermediates **20** and **21** were decarboxylated in a facile manner under Krapcho conditions,²¹ and the subsequent succession of reduction and protecting group swap resulted in di-Boc protected intermediates **24** and **25**. Ensuing condensation with paraformaldehyde under basic conditions provided styrenes

26 and **27**. Gratifyingly, the double bonds of **26** and **27** were efficiently cyclopropanated under Corey-Chaykovsky conditions,²² and subsequent Boc removal provided building blocks **30** and **31** (**Scheme 2**).

Scheme 2. Synthesis of substituted cyclopropyl P1' building blocks **30** and **31**. Reagents and conditions: (a) CuI (0.2 equiv.), (*S*)-Proline (0.4 equiv.), Cs₂CO₃, DMSO, 50 °C, 3 d; (b) LiCl, DMSO-H₂O, 130 °C, 2 h; (c) i. Zn, NH₄Cl, MeOH-THF, rt, 2 h; (ii) Boc₂O, 85 °C; (iii) H₂, Pd-C, Boc₂O, MeOH, rt; (d) (CH₂O)_n, TDA-1, PhMe, 85 °C; (e) [Me₃SO]I (3.0 equiv.), NaH, DMSO, rt to 60 °C, 3 h; (f) HCl, dioxane, 2 h.

In agreement with the design strategy, we set out to synthesize an array of cycloalkyl P1' groups bearing carboxylate at the 1-position. However, initial attempts to C-alkylate arylacetate **22** were unsuccessful, presumably because of the poor nucleophilicity of the corresponding anion due to extensive delocalization on the nitroarene system. Thus, the synthetic sequence was modified to sidestep the aryl nitro precursor, which resulted in effectual delivery of substituted cycloalkyl carboxylates **42-45** (**Scheme 3**). Namely, nitroarene **22** was reduced to the respective aniline **32**, which was then reacted with nitrosonium tetrafluoroborate, and the intermediate diazonium species were treated with I₂/NaI to give rise to the required iodoarene precursor **33**. In this instance, treatment with NaH and the respective dibromo electrophiles resulted in formation of substituted aryl cycloalkanes **34-37** in moderate to good yields. Compounds **34-37** were then subjected to Cu-catalyzed amination conditions²³ to afford anilines **38-41**, which were conveniently deprotected under H₂ (1 atm) in the presence of Pd-C to furnish intended intermediates **42-45**.

Scheme 3. Preparation of carboxyethyl cycloalkyl P1' intermediates **42-45**. Reagents and conditions: (a) Zn, NH₄Cl, MeOH-THF, rt, 2 h; (b) [NO][BF₄], NaI, I₂, MeCN, -40 °C to 0 °C, 4 h; (c) Br-(L)-Br, NaH, DMF, 0 °C to rt; (d) acetimidamide, CuI (0.15 equiv.), (*S*)-Proline (0.3 equiv.), Cs₂CO₃, DMF, 100 °C, 2 d; (e) H₂ (1 atm), Pd-C, EtOH, rt, 1.5 h.

Having synthesized the necessary P1' intermediates **18**, **30-31**, and **42-45**, we were then equipped to assemble all the target macrocyclic TF-FVIIa inhibitors. Macrocycles **10** (Figure 2) and **46** (Table 1) were accessed starting from the precursor **18** utilizing a general synthetic sequence we have disclosed earlier.^{14,15} Additional macrocyclic structures **52-55** were constructed as outlined in Scheme 4 from previously described phenylglycine **47**.¹⁵ BOP coupling of the respective precursors afforded the required aminophenyl alcohols **48-52** in good yields. Our previously reported two-step carbamoylation protocol¹⁴ involving phosgene functionalization and subsequent slow addition to a dilute MeCN solution of an amine base afforded the corresponding macrocyclic carbamates, which after saponification and Boc removal yielded the desired macrocycles **53-57**.²⁴

Scheme 4. Synthesis of the target macrocycles **52-55**. Reagents and conditions: (a) BOP, TEA, DMF-MeCN, rt, 30 min. (b) i. COCl₂, MeCN-DMPU, 0 °C; ii. Et₃N, MeCN (~1 mM), 70 °C, 3 h; (c) chiral HPLC; (d) KO-TMS, THF, 60 °C, 18 h; (e) TFA, rt, 15 min.

Table 1. Structure-activity relationship of substituted cycloalkyl P1' macrocycles.

Compd	46	53	54	55	56	57	59
TF-FVIIa K _i 37 °C (nM) ^a	18	0.55	0.14	0.22	0.06	4.4	0.72
FXa K _i 25 °C (nM) ^b	3000	6650	850	1800	900	1400	1050
Thrombin K _i 25 °C (nM) ^b	18000	>13300	>13300	>13300	>13300	>13300	>13300
Trypsin K _i 25 °C (nM) ^b	>6200	2750	3400	2200	505	>6200	4900
HK-1 ^c K _i 37 °C (nM) ^b	15	1100	1150	1600	1200	5300	960
aPC ^d K _i 37 °C (nM) ^b	220	70	725	340	155	1600	175
FVII def. PT EC _{2X} (μM) ^e	32	4.7	7.9	6.6	18	11	10
Human PPB (% free)	-	4.4	3.3	2.8	1.7	5.8	1.8
PAMPA pH 5.5, 7.4 (nm/s)	-, -	337, 117	209, 34	211, 40	805, 80	26, 21	119, 48
HLM t _{1/2} (min)	23	100% rem. @ 10 min	>120	-	94	>120	114
Clearance (mL/min/kg)	-	-	56 (d)	54 (d)	-	-	7 (d)

^aTF-FVIIa assays were performed with recombinant human enzyme with FX as a substrate (n ≥ 2). ^bK_i's for the indicated enzymes were determined by chromogenic substrate assays (n = 2). ^cHuman tissue kallikrein-1. ^dActivated protein C. ^eSee reference 13 for a detailed description.

To further probe charged interactions with Lys60A, an acid surrogate was introduced at the P1' moiety. Cyclopropyl-bearing macrocycle **53** was selected as a parent structure, and the tetrazole moiety was chosen as an acid bioisostere (Scheme 5). Nitrile-containing macrocycle **58** was prepared using a synthetic route analogous to that outlined in Scheme 4. Subsequent treatment with tributyltin azide in PhMe at 100 °C affected conversion of the nitrile moiety to the required tetrazole, and the sequence was concluded with Boc removal to yield macrocyclic tetrazole **59**.²⁴

Scheme 5. Preparation of macrocycle **59**. Reagents and conditions: (a) Bu₃SnN₃, PhMe, 105 °C, 2 d; (b) TFA, 15 min.

The series of prepared macrocycles was evaluated for TF-FVIIa inhibition and compared to the corresponding parent macrocycle **46** bearing an unsubstituted cyclopropyl group at P1' region (Table 1). Gratifyingly, introduction of the carboxylic acid moiety in compound **53** resulted in an ~30 fold boost in inhibitory potency, and at the same time dramatically improved overall selectivity for the compound. For example, human tissue

kallikrein changed from being essentially equipotent for cyclopropane **46** (HK-1 K_i /TF-FVIIa K_i = 0.8) to being fully selective for acid analog **53** (HK-1 K_i /TF-FVIIa K_i = 2000). Progressive expansion of the ring size in homologs **53-56** produced a gradual increase in FVIIa inhibitory activity, culminating in picomolar inhibitor **56** (TF-FVIIa K_i = 60 pM). It should be noted that selectivity against protease anti-targets was generally preserved in the homologous series **53-56**. Of note, cyclobutyl analog **54** exhibited an optimal selectivity against aPC in the homologous **53-56** series, presumably due to differential steric requirements to bind to this anti-target. 4-THP derivative **57** was prepared to introduce a mild degree of polarity to the P1' ring system. As a result, **57** suffered ~70-fold loss in potency compared to the respective cyclohexyl analog **56**, indicating that polar substituents may not be tolerated. Lastly, cyclopropyl tetrazole **59** showed an outstanding level of TF-FVIIa inhibitory potency and selectivity, illustrating that this carboxylic acid surrogate represents a competent pharmacophore for the series.

Intrigued by the substantial increase in potency associated with the introduction of cycloalkyl acid P1' groups we decided to generate X-ray crystal structures of macrocycles **54** and **56** bound to FVIIa.^{25,26} The structures of acids **54** and **56** are presented in **Figure 4**. The 4-F aminoisoquinoline P1 moiety as well as the methylphenylene P2 moiety bound in a manner consistent with X-ray structures described previously.^{12i,14} In contrast, the S1' binding pattern was quite different from the alkyl sulfone moieties.^{18a} As anticipated, **54** carboxylate appeared to interact with the Lys60A side chain although via an intermediary water molecule. Unexpectedly, the carboxylate of **56** forms a water mediated interaction with Lys192 side chain. Both cyclobutyl (**54**) and cyclohexyl (**56**) groups are oriented nicely into the S1' Cys42-Cys58 pocket (3.8-5.1 Å), forming a hydrophobic edge to edge interaction with the disulfide bridge. Overlay of cyclobutyl and cyclohexyl P1' groups is depicted in **Figure 4c**. As can be seen from the superimposition, the six-membered ring occupies the binding pocket more fully, which may explain the exceptional FVIIa binding affinity observed for **56**. The abovementioned structural data could also explain for 4-THP analog **57** potency loss, where oxygen atom is presumably placed inside a highly hydrophobic pocket, and is thus disfavored. The interesting interactions of the carboxylic acids described above prompted us to obtain an additional crystal structure of a tetrazole acid surrogate, the des-F aminoisoquinoline analog of **59** (**Figure 5**). In this case, the tetrazole moiety is positioned in much closer proximity to the Lys60A side chain making a direct contact (2.9 Å), presumably due to the more spatially delocalized nature of the tetrazolide anion vs the carboxylate.²⁸ As observed for analogs **54** and **56**, the cyclopropyl group efficiently interacts with the Cys42-Cys58 disulfide bridge. Overall, P1' residues in **54**, **56** and **59** bind to FVIIa in a highly productive manner, providing the optimal level of ligand efficiency with almost every atom engaged in a beneficial interaction with the enzyme.

Further profiling of macrocycles **53-57** and **59** revealed that the compounds maintained good PAMPA permeability across the series, with the exception of analog **57** (**Table 1**). Lower PAMPA values in this case could be explained by increased polarity of the 4-THP moiety. In addition, all macrocyclic inhibitors exhibited excellent metabolic stability, and represented a substantial improvement over parent analog **46**. Measured FVII deficient prothrombin time EC_{2x} values¹³ showed a stronger correlation

Figure 4. (a) Crystal structure of **54** bound to FVIIa (PDB ID: 5L2Y), solved at 1.82 Å. The omit Fo-Fc electron density is displayed at 5 rmsd. (b) Crystal structure of **56** bound to FVIIa (PDB ID: 5L2Z), solved at 1.79 Å. The omit Fo-Fc electron density is displayed at 3.5 rmsd. (c) Overlay of **54** with **56** at the FVIIa binding site with surface. Graphics were generated with the program PyMOL.²⁷

Figure 5. X-ray crystal structure of the des-F aminoisoquinoline analog of **59** complexed with FVIIa (PDB ID: 5L30), solved at 1.73 Å. The omit Fo-Fc electron density is displayed at 3.5 rmsd. Graphics were generated with the program PyMOL.²⁷

with human plasma free fraction values rather than with the *in vitro* TF-FVIIa inhibition constants. For instance, the most potent analog *in vitro* (**56**; TF-FVIIa K_i = 60 pM) appeared to have the weakest anti-clotting activity due to a low free fraction. On the other hand, ~70-fold weaker inhibitor **57** showed a more robust FVII deficient prothrombin time increase due to reduced protein binding.

To further understand the effect of the novel P1' groups on compounds' overall properties, two acid-containing macrocycles **54** and **55** were advanced into i.v. PK studies (Table 1). Disappointingly, both compounds showed high clearance in dog, likely due to glucuronidation of the carboxylic acid moiety. We then decided to dose tetrazolyl analog **59** to investigate if an acid surrogate can exert a positive effect on the overall pharmacokinetic profile. We were pleased to discover that tetrazole **59** displayed a much lower clearance of 7 mL/min/kg after i.v. administration in dogs, however low oral bioavailability (F% = 2) was observed. The aforementioned data suggests that tetrazole derivative **59** may have less propensity for glucuronidation, and could be considered an excellent starting point for further potency and ADME optimization to reach oral bioavailability.

In conclusion, we have successfully designed and synthesized macrocyclic TF-FVIIa inhibitors bearing novel 1-(1*H*-tetrazol-5-yl)cyclopropyl and 1-carboxycycloalkyl P1' moieties as cyclopropyl sulfone replacements. The discovered scaffolds allowed for remarkable improvements both in FVIIa inhibitory potency (25-300 fold increase) as well as in overall protease selectivity (>1000 fold enhancement). Important highlights from the X-ray crystallography and SAR studies include: (1) carboxylic acid substituent interacts with both Lys60A and Lys192, (2) lipophilic cycloalkyl moieties are preferred, and bulkier alkyls interface more efficiently with Cys42-Cys58 motif resulting in potency boost, (3) tetrazolyl moiety performed as an acid bioisotere, allowing for similar interactions with FVIIa enzyme and yielding outstanding inhibitory potency. Ultimately, compound **59** had the optimal balance of TF-FVIIa potency, selectivity, physicochemical properties and clearance, representing an optimized and structurally diverse alternative to cyclopropyl sulfone P1'.

Acknowledgements

The authors would like to thank Dr. Rick Ewing and Dr. Donald Pinto for their valuable advice during preparation of this manuscript. Jeffrey Bozarth, Randi Brown, Tara Peterson, Mojgan Abousleiman, Frank Barbera and Yiming Wu are acknowledged for obtaining enzyme inhibition and clotting assay data. We also would like to thank Dr. Steven Sheriff for the help with preparation and PDB deposition of coordinates and data for FVIIa complexes with compounds **54**, **56** and the des-F aminoisoquinoline analog of **59**. X-ray data were collected by Rick Walter and Gina Ranieri of Shamrock Structures LLC using the IMCA beamlines, 17-BM and 17-ID, at the Advanced Photon Source. Use of the Advanced Photon Source was supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under Contract No. DE-AC02-06CH11357. Use of the IMCA-CAT beamline 17-ID at the Advanced Photon Source was supported by the companies of the Industrial Macromolecular Crystallography Association through a contract with Hauptman-Woodward Medical Research Institute.

References and notes

- (a) Braunwald, E.; Califf, R. M.; Cannon, C. P.; Fox, K. A.; Fuster, V.; Gibler, W. B.; Harrington, R. A.; King, S. B., 3rd; Kleiman, N. S.; Theroux, P.; Topol, E. J.; Van de Werf, F.; White, H. D.; Willerson, J. T. *Am. J. Med.* **2000**, *108*, 41; (b) Cushman, M.; Tsai, A. W.; White, R. H.; Heckbert, S. R.; Rosamond, W. D.; Enright, P.; Folsom, A. R. *Am. J. Med.* **2004**, *117*, 19.
- Ansell, J.; Hirsh, J.; Poller, L.; Bussey, H.; Jacobson, A.; Hylek, E. *Chest* **2004**, *126*, 204S.
- De Caterina, R.; Husted, S.; Wallentin, L.; Andreotti, F.; Arnesen, H.; Bachmann, F.; Baigent, C.; Huber, K.; Jespersen, J.; Kristensen, S. D.; Lip, G. Y.; Morais, J.; Rasmussen, L. H.; Siegbahn, A.; Verheugt, F.W.; Weitz, J. I. *J. Am. Coll. Cardiol.* **2012**, *59*, 1413.
- (a) Huel, N. H.; Nar, H.; Priepeke, H.; Ries, U.; Stassen, J. M.; Wienen, J. *Med. Chem.* **2002**, *45*, 1757; (b) Weitz, J. I.; Linkins, L. A. *Expert Opin. Invest. Drugs* **2007**, *16*, 271; (c) Haas, S.; Schellong, S. *Hamostaseologie* **2007**, *27*, 41; (d) Eriksson, B. I.; Smith, H.; Yasothan, U.; Kirkpatrick, P. *Nat. Rev. Drug Discov.* **2008**, *7*, 557; (e) Roehrig, S.; Straub, A.; Pohlmann, J.; Lampe, T.; Pernerstorfer, J. *J. Med. Chem.* **2005**, *48*, 5900; (f) Lassen, M. R.; Ageno, W.; Borris, L. C.; Lieberman, J. R.; Rosencher, N.; Bandel, T. J.; Misselwitz, F.; Turpie, A. G. N. *Eng. J. Med.* **2008**, *358*, 2776; (g) Straub, A.; Roehrig, S.; Hillisch, A. *Curr. Top. Med. Chem.* **2010**, *10*, 257; (h) Perzborn, E.; Roehring, S.; Straub, A.; Kubitz, D.; Misselwitz, F. *Nat. Rev. Drug Discov.* **2011**, *10*, 61; (i) Pinto, D.; Orwat, M.; Koch, S.; Rossi, K.; Alexander, R.; Smallwood, A.; Wong, P.; Rendina, A.; Luetgen, J. M.; Knabb, R. M.; Xin, B.; Wexler, R. R.; Lam, P. *J. Med. Chem.* **2007**, *50*, 5339; (j) He, K.; Luetgen, J. M.; Zhang, D.; He, B.; Grace, J. E. Jr.; Xin, B.; Pinto, D. J. P.; Wong, P. C.; Knabb, R. M.; Lam, P. Y. S.; Wexler, R. R.; Grossman, S. J. *Eur. J. Drug Metab. Pharm.* **2011**, *36*, 129; (k) Pinto, D. J. P.; Smallheer, J. M.; Cheney, D. L.; Knabb, R. M.; Wexler, R. R. *J. Med. Chem.* **2010**, *53*, 6243; (l) Furugohri, T.; Isobe, K.; Honda, Y.; Kamisato-Matsumoto, C.; Sugiyama, N.; Nagahara, T.; Morishima, Y.; Shibano, T. *J. Thromb. Haemost.* **2008**, *6*, 1542. (m) Bounameaux, H.; Camm A. J.; *Drugs* **2014**, *74*, 1209.
- Bauer, K. A. *Hematology Am. Soc. Hematol. Educ. Program* **2013**, *2013*, 464.
- Priestley, E. S. *Drug Discov. Today* **2014**, *19*, 1440.
- (a) Steffel, J.; Lüscher, T. F.; Tanner, F. C. *Circulation* **2006**, *113*, 722; (b) Petrillo, G.; Cirillo, P.; D'Ascoli, G. L.; Maresca, F.; Ziviello, F.; Chiariello, M. *Curr. Cardiol. Rev.* **2010**, *6*, 325;
- (a) Steffel, J.; Lüscher, T. F.; Tanner, F. C. *Circulation* **2006**, *113*, 722; (b) Petrillo, G.; Cirillo, P.; D'Ascoli, G. L.; Maresca, F.; Ziviello, F.; Chiariello, M. *Curr. Cardiol. Rev.* **2010**, *6*, 325; (a) Himber, J.; Kirchhofer, D.; Riederer, M.; Tschopp, T.; Steiner, B.; Roux, S. *Thromb. Haemost.* **1997**, *78*, 1142; (b) Himber, J.; Refino, C. J.; Burcklen, L.; Roux, S.; Kirchhofer, D. *Thromb. Haemost.* **2001**, *85*, 475; (c) Golino, P.; Ragni, M.; Cirillo, P.; D'Andrea, D.; Scognamiglio, A.; Ravera, A.; Buono, C.; Ezban,

- M.; Corcione, N.; Vigorito, F. *Circul. Res.* **1998**, *82*, 39; (d) Szalony, J. A.; Suleymanov, O. D.; Salyers, A. K.; Panzer-Knodle, S. G.; Blom, J. D.; LaChance, R. M.; Case, B. L.; Parlow, J. J.; South, M. S.; Wood, R. S.; Nicholson, N. S. *Thromb. Res.* **2003**, *112*, 167; (e) Suleymanov, O. D.; Szalony, J. A.; Salyers, A. K.; LaChance, R. M.; Parlow, J. J.; South, M. S.; Wood, R. S.; Nicholson, N. S. *J. Pharmacol. Exp. Ther.* **2003**, *306*, 1115. (f) Salyers, A. K.; Szalony, J. A.; Suleymanov, O. D.; Parlow, J. J.; Wood, R. S.; South, M. S.; Nicholson, N. S. *Pharmacology* **2004**, *70*, 100; (g) Olivero, A. G.; Eigenbrot, C.; Goldsmith, R.; Robarge, K.; Artis, D. R.; Flygare, J.; Rawson, T.; Sutherland, D. P.; Kadhodayan, S.; Beresini, M.; Elliott, L. O.; DeGuzman, G. G.; Banner, D. W.; Ultsch, M.; Marzec, U.; Hanson, S. R.; Refino, C.; Bunting, S.; Kirchhofer, D. *J. Biol. Chem.* **2005**, *280*, 9160; (h) Zbinden, K. G.; Banner, D. W.; Hilpert, K.; Himber, J.; Lavé, T.; Ziderer, M. A.; Stahl, M.; Tschopp, T. B.; Obst-Sander, U. *Bioorg. Med. Chem.* **2006**, *14*, 5357; (i) Young, W. B.; Mordenti, J.; Torkelson, S.; Shrader, W. D.; Kolesnikov, A.; Rai, R.; Liu, L.; Hu, H.; Leahy, E. M.; Green, M. J.; Sprengeler, P. A.; Katz, B. A.; Yu, C.; Janc, J. W.; Elrod, K. C.; Marzec, U. M.; Hanson, S. R.; *Bioorg. Med. Chem. Lett.* **2006**, *16*, 2037; (j) Arnold, C. S.; Parker, C.; Upshaw, R.; Prydz, H.; Chand, P.; Kotian, P.; Bantia, S.; Babu, Y. S. *Thromb. Res.* **2006**, *117*, 343; (k) Wong, P. C.; Luettggen, J. M.; Rendina, A. R.; Kettner, C. A.; Xin, B.; Knabb, R. M.; Wexler, R.; Priestley, E. S. *Thromb. Haemost.* **2010**, *104*, 261; (l) Nagakura, T.; Tabata, K.; Kira, K.; Hirota, S.; Clark, R.; Matsuura, F.; Hiyoshi, H., *Thromb. Res.* **2013**, *132*, 271.
9. (a) Liu, Y.; Jiang, P.; Capkova, K.; Xue, D.; Ye, L.; Sinha, S. C.; Mackman, N.; Janda, K. D.; Liu, C. *Cancer Res.* **2011**, *71*, 6492; (b) Sampson, M. T.; Kakkar, A. K. *Biochem. Soc. Trans.* **2002**, *30*, 201. (c) Mueller, B.M.; Ruf, W. *J. Clin. Invest.* **1998**, *101*, 1372.
 10. (a) Taylor, F. B.; Chang, A. C.; Peer, G.; Li, A.; Ezban, M.; Hedner, U. *Blood* **1998**, *91*, 1609; (b) Camerer, E.; Huang, W.; Coughlin, S. R. *Proc. Natl. Acad. Sci.* **2000**, *97*, 5255.
 11. Quan, M. L.; Glunz, P. W.; Smallheer, J. M. *Advances in Anticoagulants (Book Chapter)*, **2016**, accepted for publication.
 12. (a) Schechter, I.; Berger, A., *Biochem. Biophys. Res. Commun.* **1967**, *27*, 157; (b) Mahesh, S. A.; Gabrail, N. Y.; Gandhi, J. G.; Khorana, A. A.; Manges, R.; Shah, S.; Thomas, G. W.; Hamdy, A. M.; Stevens-Brogan, M.; Zhou, C. *J. Clin. Oncol.* **2013**, *31*, e15014; (c) Arnold, C. S.; Parker, C.; Upshaw, R.; Prydz, H.; Chand, P.; Kotian, P.; Bantia, S.; Babu, Y. S. *Thromb. Res.* **2006**, *117*, 343-349; (d) Kotian, P. L.; Krishnan, R.; Rowland, S.; El-Kattan, Y.; Saini, S. K.; Upshaw, R.; Bantia, S.; Arnold, S.; Babu, Y. S.; Chand, P. *Bioorg. Med. Chem.* **2009**, *17*, 3934; (e) Glunz, P. W.; Cheng, X.; Cheney, D. L.; Weigelt, C. A.; Wei, A.; Luettggen, J. M.; Wong, P. C.; Wexler, R. R.; Priestley, E. S. *Bioorg. Med. Chem. Lett.* **2015**, *25*, 2169-2173; (f) Nagakura, T.; Tabata, K.; Kira, K.; Hirota, S.; Clark, R.; Matsuura, F.; Hiyoshi, H. *Thromb. Res.* **2013**, *132*, 271; (g) Hu, H.; Kolesnikov, A.; Riggs, J. R.; Wesson, K. E.; Stephens, R.; Leahy, E. M.; Shrader, W. D.; Sprengeler, P. A.; Green, M. J.; Sanford, E.; Nguyen, M.; Gjerstad, E.; Cabuslay, R.; Young, W. B. *Bioorg. Med. Chem. Lett.* **2006**, *16*, 4567; (h) Miura, M.; Seki, N.; Koike, T.; Ishihara, T.; Niimi, T.; Hirayama, F.; Shigenaga, T.; Sakai-Moritani, Y.; Tagawa, A.; Kawasaki, T.; Sakamoto, S.; Okada, M.; Ohta, M.; Tsukamoto, S. *Bioorg. Med. Chem.* **2007**, *15*, 160; (i) Cheney, D. L.; Bozarth, J. M.; Metzler, W. J.; Morin, P. E.; Mueller, L.; Newitt, J. A.; Nirschl, A. H.; Rendina, A. R.; Tamura, J. K.; Wei, A.; Wen, X.; Wurtz, N. R.; Seiffert, D. A.; Wexler, R. R.; Priestley, E. S. *J. Med. Chem.* **2015**, *58*, 2799; (j) Priestley, E. S.; Cheney, D. L.; DeLucca, I.; Wei, A.; Luettggen, J. M.; Rendina, A. R.; Wong, P. C.; Wexler, R. R. *J. Med. Chem.* **2015**, *58*, 6225.
 13. Wong, P. C.; Luettggen, J. M.; Rendina, A. R.; Kettner, C. A.; Xian, B.; Knabb, R. M.; Wexler, R. R.; Priestley, E. S. *Thromb. Haemostasis* **2010**, *104*, 261.
 14. Glunz, P. W.; Mueller, L.; Cheney, D. L.; Ladziata, V.; Zou, Y.; Wurtz, N. R.; Wei, A.; Wong, P. C.; Wexler, R. R.; Priestley, E. S. *J. Med. Chem.* **2016**, *59*, 4007.
 15. Zhang, X.; Glunz, P. W.; Johnson, J. A.; Jiang, W.; Jacutin-Porte, S.; Ladziata, V.; Zou, Y.; Phillips, M. S.; Wurtz, N. R.; Parkhurst, B.; Rendina, A. R.; Harper, T. M.; Cheney, D. L.; Luettggen, J. M.; Wong, P. C.; Seiffert, D.; Wexler, R. R.; E. Priestley, E. S. *J. Med. Chem.* **2016**, *59*, 7125.
 16. McNaney, C. A.; Dexter, D. M.; Hnatyshyn, S. Y.; Zvyaga, T. A.; Knipe, J. O.; Belcastro, J. V.; Sanders, M. *ASSAY Drug Dev. Techn.* **2008**, *6*, 121.
 17. van Breemen, R. B.; Li, Y. *Expert Opin. Drug. Metab. Toxicol.* **2005**, *1*, 175.
 18. (a) Zhang, X.; Jiang, W.; Jacutin-Porte, S.; Glunz, W. P.; Zou, Y.; Cheng, X.; Nirschl, H. A.; Wurtz, N. R.; Luettggen, M. J.; Rendina, A. R.; Luo, G.; Harper, M. T.; Wei, A.; Anumula, R.; Cheney, L. D.; Knabb, M. R.; Wong, C. P.; Wexler, R. R.; Priestley, E. S. *ACS Med. Chem. Lett.* **2014**, *5*, 188; (b) Glunz, P. W.; Zhang, X.; Zou, Y.; Nirschl, A. H.; Cheng, X.; Weigelt, C. A.; Cheney, D. L.; Wei, A.; Wen, X.; Bozarth, J. M.; Normandin, D. E.; Hartl, K. S.; Rendina, A. R.; Barbera, F. A.; Luettggen, J. M.; Brown, R. L.; Peterson, T.; Zhang, G.; Wong, P. C.; Harpel, M.; Shen, L.; M., K. R.; Wexler, R. R.; Priestley, E. S. *Bioorg. Med. Chem. Lett.* **2013**, *23*, 5244.
 19. Wallace, D. J.; Chen, C-Y. *Tetrahedron Lett.* **2002**, *43*, 6987.
 20. Ma, D.; Cai, Q. *Acc. Chem. Res.* **2008**, *41*, 1450.
 21. (a) Krapcho, A. P.; Weimaster, J. F.; Eldridge, J. M.; Jahng, E. G. E., Jr.; Lovey, A. J.; Stephens, W. P. *J. Org. Chem.* **1978**, *43*, 138; (b) Krapcho, A. P.; Ciganek, E. *Org. React.* **2013**, *81*, 1.
 22. (a) Corey, E. J.; Chaykovsky, M. *J. Am. Chem. Soc.* **1962**, *84*, 867; (b) Corey, E. J.; Chaykovsky, M. *J. Am. Chem. Soc.* **1962**, *84*, 3782; (c) Corey, E. J.; Chaykovsky, M. *J. Am. Chem. Soc.* **1965**, *87*, 1353.
 23. Gao, X.; Fu, H.; Qiao, R.; Jiang, Y.; Zhao, Y. *J. Org. Chem.* **2008**, *73*, 6864.
 24. **53**: MS (ESI) m/z : 612.2 $[M+1]^+$; 1H NMR (400 MHz, CD_3OD) δ ppm 8.09 (dd, $J=9.3$, 2.0 Hz, 1H), 7.65 (dd, $J=8.0$, 1.8 Hz, 1H), 7.45 (d, $J=8.0$ Hz, 1H), 7.42 (d, $J=5.0$ Hz, 1H), 7.28 (dd, $J=9.3$, 2.3 Hz, 1H), 7.21 (s, 1H), 7.16 (d, $J=8.3$ Hz, 1H), 6.97 (s, 1H), 6.67 (dd, $J=8.0$, 2.3 Hz, 1H), 5.97 (d, $J=2.0$ Hz, 1H), 5.77 (s, 1H), 5.54 (d, $J=16.8$ Hz, 1H), 4.64 (t, $J=10.9$ Hz, 1H), 3.89-4.00 (m, 2H), 3.43-3.53 (m, 1H), 3.28 (s, 1H), 2.32 (s, 3H), 1.64 (br. s., 2H), 1.31 (d, $J=7.0$ Hz, 3H), 1.20 (br. s., 2H). **54**: MS (ESI) m/z : 626.3 $[M+1]^+$; 1H NMR (400 MHz, CD_3OD) δ ppm 8.09 (dd, $J=9.2$, 1.9 Hz, 1H), 7.62 (dd, $J=7.9$, 1.6 Hz, 1H), 7.40-7.47 (m, 2H), 7.28 (dd, $J=9.3$, 2.5 Hz, 1H), 7.22 (d, $J=1.5$ Hz, 1H), 7.13 (d, $J=8.3$ Hz, 1H), 6.97 (s, 1H), 6.73 (dd, $J=8.2$, 2.1 Hz, 1H), 5.93 (d, $J=2.0$ Hz, 1H), 5.70-5.79 (m, 1H), 5.36 (d, $J=16.6$ Hz, 1H), 4.61 (t, $J=10.9$ Hz, 1H), 3.97 (dd, $J=10.8$, 4.3 Hz, 1H), 3.81 (d, $J=16.6$ Hz, 1H), 3.42-3.55 (m, 1H), 3.20 (s, 2H), 2.72-2.86 (m, 2H), 2.46-2.61 (m, 2H), 2.10-2.24 (m, 1H), 2.03 (s, 1H), 1.78-1.91 (m, 1H), 1.31 (d, $J=7.0$ Hz, 3H). **55**: MS (ESI) m/z : 640.3 $[M+1]^+$; 1H NMR (400 MHz, CD_3OD) δ ppm 8.86 (s, 1H), 8.08 (dd, $J=9.3$, 1.8 Hz, 1H), 7.62 (dd, $J=8.0$, 1.5 Hz, 1H), 7.38-7.47 (m, 2H), 7.25-7.34 (m, 2H), 7.22 (s, 1H), 6.96 (br. s., 1H), 6.72 (dd, $J=8.3$, 2.3 Hz, 1H), 6.01 (d, $J=2.0$ Hz, 1H), 5.75 (s, 1H), 5.46 (d, $J=16.3$ Hz, 1H), 4.59 (t, $J=10.9$ Hz, 1H), 3.97 (dd, $J=10.8$, 4.3 Hz, 1H), 3.88 (d, $J=16.3$ Hz, 1H), 3.42-3.55 (m, 1H), 3.20 (s, 3H), 2.59 (dt, $J=12.7$, 6.3 Hz, 1H), 2.43-2.53 (m, 1H), 2.33 (s, 3H), 1.90-2.09 (m, 2H), 1.65-1.84 (m, 4H), 1.31 (d, $J=7.0$ Hz, 3H). **56**: MS (ESI) m/z : 654.3 $[M+1]^+$; 1H NMR (400 MHz, CD_3OD) δ ppm 8.10 (dd, $J=9.2$, 1.9 Hz, 1H), 7.60 (dd, $J=7.8$, 1.8 Hz, 1H), 7.36-7.47 (m, 3H), 7.26-7.32 (m, 1H), 7.21 (s, 1H), 6.97 (1 H, s), 6.75 (dd, $J=8.5$, 2.3 Hz, 1H), 6.04 (1 H, s), 5.74 (s, 1H), 5.51 (d, $J=16.3$ Hz, 1H), 4.57 (t, $J=10.9$ Hz, 1H), 3.93-4.03 (m, 2H), 3.44-3.54 (m, 1H), 3.18 (s, 3H), 2.41 (d, $J=15.3$ Hz, 1H), 2.35 (s, 3H), 2.30 (d, $J=13.8$ Hz, 1H), 1.75-1.88 (m, 2H), 1.55-1.74 (m, 5H), 1.34-1.42 (m, 1H), 1.32 (d, $J=7.0$ Hz, 3H). **57**: MS (ESI) m/z : 656.2 $[M+1]^+$; 1H NMR (400 MHz, CD_3OD) δ ppm 8.92 (s, 1H), 8.10 (dd, $J=9.3$, 1.8 Hz, 1H), 7.61 (d, $J=7.8$ Hz, 1H), 7.40-7.48 (m, 2H), 7.37 (d, $J=8.5$ Hz, 1H), 7.29 (dd, $J=9.3$, 2.3 Hz, 1H), 7.19 (1 H, s), 6.96 (1 H, br. s.), 6.77 (dd, $J=8.4$, 2.1 Hz, 1H), 6.09 (s, 1H), 5.75 (s, 1H), 5.50 (d, $J=16.3$ Hz, 1H), 4.58 (t, $J=10.9$ Hz, 1H), 3.93-4.03 (m, 2H), 3.71-3.91 (m, 3H), 3.42-3.54 (m, 1H), 3.20 (s, 3H), 2.46 (d, $J=15.1$ Hz, 1H), 2.34 (s, 3H), 2.03-2.15 (m, 1H), 1.89-2.01 (m, 1H), 1.32 (d, $J=7.0$ Hz, 3H). **59**: MS (ESI) m/z : 636.2 $[M+1]^+$; 1H NMR (400 MHz, CD_3OD) δ ppm 9.03 (s, 1H), 8.09 (dd, $J=9.4$, 1.77 Hz, 1H), 7.65 (dd, $J=8.1$, 1.8 Hz, 1H), 7.47 (d, $J=8.1$ Hz, 1H), 7.43 (d, $J=4.8$ Hz, 1H), 7.36 (d, $J=8.1$ Hz, 1H), 7.29 (dd, $J=9.4$, 2.3 Hz, 1H), 7.14 (s, 1H), 6.96 (s, 1H), 6.77 (dd, $J=8.1$, 2.0 Hz, 1H), 6.09-6.15 (m, 1H), 5.76 (s, 1H), 5.27 (d, $J=16.9$ Hz, 1H), 4.67 (t, $J=11.1$ Hz, 1H), 3.97 (dd, $J=10.7$, 4.4 Hz, 1H), 3.81 (d, $J=16.7$ Hz, 1H), 3.44 - 3.56 (m, 1H), 2.31 (s, 3H), 1.65-1.84 (m, 2H), 1.48-1.61 (m, 2H), 1.32 (d, $J=7.1$ Hz, 3H).
 25. Relative atomic coordinates for the crystallographic structures of **54**, **56**, and the des-F aminoisoquinoline analog of **59** complexed with factor VIIa have been deposited in the PDB; their deposition numbers are 5L2Y, 5L2Z, 5L30, respectively.
 26. For procedures for crystallization, data collection, and structure refinement, please see ref. 12j.
 27. The PyMOL Molecular Graphics System, Version 1.7 Schrödinger, LLC

28. (a) Carini, D. J.; Christ, D. D.; Duncia, J. V.; Pierce, M. E. *Pharm. Biotechnol.* **1998**, *11*, 29. (b) Naylor, E. M.; Chakravarty, P. K.; Costello, C. A.; Chang, R. S.; Chen, T.-B.; Faust, K. A.; Lotti, V. J.; Kivlighn, S. D.; Zingaro, G. J.; Siegl, P. K. S.; Wong, P. C.; Carini, D. J.; Wexler, R. R.; Patchett, A. A.; Greenlee, W. J. *Bioorg. Med. Chem. Lett.* **1993**, *3*, 69.

Graphical Abstract

To create your abstract, type over the instructions in the template box below.
Fonts or abstract dimensions should not be changed or altered.

Synthesis and P1' SAR Exploration of Potent Macrocyclic Tissue Factor-Factor VIIa Inhibitors

Leave this area blank for abstract info.

Vladimir (Uladzimir) Ladziata,^{*} Peter W. Glunz, Yan Zou,[†] Xiaojun Zhang, Wen Jiang, Swanee Jacutin-Porte, Daniel L. Cheney, Anzhi Wei, Joseph M. Luettgen, Timothy M. Harper, Pancras C. Wong, Dietmar Seiffert, Ruth R. Wexler, E. Scott Priestley