Climate teleconnections and indicators of coastal systems response

Ofelia Gutiérrez, Daniel Panario, Gustavo J. Nagy, Mario Bidegain, Carlos Montes

Abstract

This article surveys secular urban sandy beaches erosion – accretion and its relationship with climate teleconnections, e.g., El Niño Southern Oscillation (ENSO), Atlantic Multidecadal Oscillation (AMO), and extreme events, e.g., storm surges, great rivers' floods, and heavy rains in adjacent basins. The paper aims to discuss these issues and the expected coastal retreat as a consequence of manmade climate changes, e.g., sea level rise (SLR) and increased storminess in the coming decades. Several beaches (Buceo, Malvín, Pocitos, Ramírez) and two tidal creek mouths (Carrasco and Pando), with different characteristics but all constrained by coastal linear infrastructure and two tidal creek sandy mouths were studied. These sites are located along the urbanised coast of the middle region of Río de la Plata microtidal river estuary. All of them show a more or less strong retreat trend with alternated fluctuations, e.g., weak retreat likely due to sea level rise, significant erosion very likely due to storm surges, and processes of loss of sediment stock, as well as episodes of sand recovery. Therefore, these beaches require interventions to preserve their beach prism and dry sand surface. In search for answers to better understand why and under what conditions the process of advance and retreat of the coastline occur, we have analysed different teleconnections and carried out reanalyses for wind anomalies during ENSO events from 1951-2010. Both weak and moderate erosion-accretion periods are likely related to atmospheric anomalies, e.g., wind direction changes and the consequent swell and littoral drift changes, related to El Niño and La Niña events. The former associated with accretion and the latter with erosion. In the past most interventions have been reactive. Increased knowledge of climate and weather relationship with the sedimentary balance provides an approach that would allow developing beach risk-management, pro-active strategies and climate adaption measures focused on the generation and recovery planning based on the analysis of the occurrence and prediction of El Niño / La Niña events.

Keywords

Sandy beach; creek mouths; erosion-accretion; El Niño - La Niña events; Río de la Plata estuary.

1. INTRODUCTION

The growing threat that hangs over coastal systems due to global change is induced primarily by forcings such as changing land use and climate change, especially sea level rise (SLR) (Masselink and Gehrels, 2014). Socioeconomic, cultural and environmental importance of beaches make it necessary to create as accurate as possible scenarios of their evolution in the medium and long term (Dias et al., 2013; Ledoux and Turner, 2002), in order to develop mitigation and adaptation plans and measures against these threats (Luisetti et al., 2014).

Ecosystem resilience is a clear objective of modern science (Holling and Gunderson, 2002). For coastal environments this involves analysing the capacity to generate the necessary conditions for maintaining ecosystem functions that support the services provided by beaches (Flood and Schechtman, 2014).

In order to build resilience in an ecosystem subject to fast and slow drivers of change, e.g., extreme events and gradual climate change respectively, there is a need to identify them, particularly the fast ones and their trends in the mid-term (months to years), since they may produce undesirable and irreversible shifts. These threats are very important in urban beaches constrained by linear infrastructure.

World's beaches are threatened not only by the increase in mean sea level (MSL)¹, with a recent trend of +3.26 mm/year (average rate 3.3 ± 0.4 mm/year) measured by high-precision altimetry satellites (Nicholls and Cazenave, 2010), but also by other natural forcings as subsidence, depletion of underwater sand reserves, changes in atmospheric circulation and possible increase in wind speed due to global warming. In addition, attention should be given to local human forcings, such as construction or resizing of harbours, building walls, breakwaters, jetties or breakwaters, sand mining, or urbanisation of exchanging sand dune systems (Gutiérrez and Panario, 2005; Gutiérrez et al., 2015; Marshall and Banks, 2013; Panario and Gutiérrez, 2005; Panario and Piñeiro, 1997; Stevens et al., 2005).

As a result of these actions, most of the Uruguayan beaches are showing evidence of degradation e.g., the loss of the primary dune, and/or the decrease in the area of dry sand and/or the height of the beach prism, which makes them vulnerable. Typically these beaches have their land limit constrained by the presence of coastal linear infrastructure. Several experiences of primary dune reconstruction carried out by our team have been successful while others have failed due to storms that destroyed the installed soft works. The chances of success of these initiatives on these beaches could

---

¹ **Abbreviations:** **AMO:** Atlantic Multidecadal Oscillation, **AMSL:** above mean sea level, **ENSO:** El Niño Southern Oscillation, **MSL:** mean sea level, **NCEP/NCAR:** National Center for Environmental Prediction/National Center for Atmospheric Research, **PDO:** Pacific Decadal Oscillation, **PHTH-WL:** high tide high - water level, **RdIP:** Río de la Plata, **SLR:** sea level rise, **SSTA:** Sea Surface Temperature Anomalies, **WDL-RM:** wet/dry line or run-up maxima, **WSA:** wind speed anomalies.

be fostered if we had better knowledge of both the causality and prognosis of increased storms, and the periods of increased beach prism, hence of the availability of aeolian sand.

From our experience it should be noted that the reconstruction of dunes in long stretches of coastline requires adequate economic, human, and material resources, and logistics to perform the task. Since failures discourage funding sources of these activities, it is of paramount importance to anticipate preparedness for when these activities are most likely to occur and in sufficient time to have the human and material resources for execution.

This article focuses on four beaches Pocitos, Ramírez, Buceo and Malvín of Montevideo city, and two tranches with mouths included (Carrasco and Pando creeks) on the Uruguayan coast of the large Río de la Plata (from now on RdIP) river estuary (38,000 km², 100-230 km wide) (Figure 1). The northern Uruguayan coast of the estuary is characterised by the presence of wave-dominated sandy beaches. This coast and its beaches are subject to the action of tides (< 0.5 m amplitude), wind climate patterns, episodic river floods and storm surges events, and gradual sea level rise (SLR) (Gutiérrez et al., 2015; Nagy et al., 2014a, 2014b; Verocai et al., 2015).

The selected beaches have notable differences, e.g., orientation, bathymetry, exposure length, profile, presence/absence of primary dune, as well as in the mineralogical composition. Human interventions have also contributed to the observed differences and long-term evolution of the studied coastal systems (Gutiérrez, 2016; Gutiérrez et al., 2015). Due to all these differences it is very likely the existence of diverse sources of sediment for the studied beaches and creek mouths.

Among the fast climate stressors on the Uruguayan coast, the wave climate change stands out. This change is associated with: i) atmospheric circulation changes likely attributable to global warming (Young et al., 2011); ii) the Atlantic Multidecadal Oscillation - AMO (Ortega et al., 2013) and the Pacific Decadal Oscillation - PDO (Nagy et al., 2014a, 2014b); and iii) oscillations associated with El Niño Southern Oscillation - ENSO teleconnections (Bidegain et al., 2015, 2014; Nagy et al., 2014a, 2008a, 2008b, 2002). It is not always clear whether these variables are linked to global climate change or natural variability. It is therefore necessary to isolate seasonal, annual, decadal and multi-decadal fluctuations occurring in atmospheric circulation and wind climate over the RdIP basin and estuary (Bidegain et al., 2005; Ortega et al., 2013; Simionato et al., 2005; Verocai et al., 2015).

The aims of this article are: i) to review the background of long-term accretion-erosion processes at urban sandy beaches focused on case studies at the Uruguayan coast of the middle region of the RdIP river estuary; ii) to focus on climate teleconnections, mainly ENSO, extreme storm surges, and their relationship with coastline retreat; iii) to update the analysis of ENSO-related wind anomalies; iv) to give insight on the need of

fostering the analysis, forecast and impacts of fast climate stressors of sandy beaches changes in the light of very likely future sea level rise and likely increased storminess; v) to provide further evidence to develop beach management planning and implement climate adaptation measures.

2. BACKGROUND

The two main observed wind patterns related to the purpose of this article are synthesised as follows:

1. An increasing trend of the occurrence of south-eastern (SE) winds over the RdIP (Bischoff, 2005; Escobar et al., 2004; Nagy et al., 2013; Ortega et al., 2013; Simionato et al., 2005).
2. A relationship between ENSO events and anomalies of atmospheric circulation and winds, e.g., an increase in E-SE-SSE-N and decrease in SW winds (called *Pamperos*) during El Niño, and increase in SW winds during La Niña from 1954-1998 (Gutiérrez, 2010).

For instance, Ortega et al. (2013) have shown in Atlantic ocean beaches near the RdIP, the existence of two variables correlated in time, the anomalies of sea surface temperature (SSTA) and wind speed (WSA) as well as an increase in S, SSE and SE wind directions for the last few decades. Whereas Codignotto et al. (2012) observed an increase in wave energy from the East, which affects the erosion of ocean beaches in the province of Buenos Aires (Argentina). Recent research (Nagy et al., 2015a) on the current (1979-2014) and future climate scenarios (centred in 2030) confirm a slight increase of SE winds along the Uruguayan coast.

During the successive El Niño and La Niña events from 1998-2002 the following observations have been reported:

- The frequency of days greater than 8 m/s during October-March varied from 7.2 days per month from 1977-86 to 11 days in 2001-02 and 17.6 days per month during the moderate to strong El Niño event in 2002-03, an increase of 57 and 150 per cent respectively (Nagy et al., 2008a),
- the wind speed yearly variability varied from 5.8 m/s to 6.2 m/s during El Niño and La Niña respectively (Nagy et al., 2003), and
- the prevailing anomalies' modes were fresh (8.6 m/s) ESE winds and strong (10.9 m/s) W winds (Nagy et al., 2008b).

With regard to waves, Dragani et al. (2010) reported a possible increase in wind wave heights and relatively high inter-annual variability in the south-eastern South American continental shelf between 32°S and 40°S, including the RdIP estuary, based on short-term series of *in situ* (1996-2006) and Topex satellite (1993-2001) data. The authors

suggested that this possible increase would be able to produce changes in the littoral processes and, consequently, in the erosion of the coast. The possible link between this inter-annual variability and ENSO was investigated but no apparent relationship was found.

Pereira and Klumb-Oliveira (2015) found for the coastal zone of Rio de Janeiro, Brazil, a slight reduction in significant wave height during strong El Niño years caused by the attenuation of the frequency of events from the S quadrant. The opposite pattern was observed during strong La Niña years, caused by an increase in wave incidence from the S and SW quadrants, especially in boreal winter from 1979-2013. The decrease could be attributed to the intensification of the South Atlantic High with a corresponding increase in the occurrence of subtropical jets during periods of El Niño. This weather change causes the blocking of southern cold fronts and the consequent reduction in the percentage of waves from the south.

According to our systematic observations of the Uruguayan coast of the RdIP river estuary and Atlantic Ocean since the 1990s, the southern winds play a key role on water and coastal dynamics, but with dissimilar effects depending on the synoptic conditions and fetch associated with wind speed (EcoPlata, 1999; Gutiérrez et al., 2015; López Laborde and Nagy, 1999; MTOP/PNUD/UNESCO, 1979; Nagy et al., 2014a, 2013, 2008a, 2008b, 2007, 2005, 2002; Verocai et al., 2015).

Southerly wind called "Pampero" is associated to the entry over the RdIP of dynamical high pressure system with two different types. The first called "dry Pampero" is formed with air masses from the south-eastern Pacific Ocean, which lose their moisture by passing the Andes and whose entry over the RdIP determines a rapid decrease in cloudiness and rainfall. Its mass air origin is mainly from the W-WSW, and the fetch is weak over shallow RdIP, causing only a slight to moderate increase in water-level along the Uruguayan coast. The other type is called "wet Pampero" characterised by the trajectory over the southern Atlantic Ocean producing rainfall after its entry into the region. Associated winds in the frontal zone can exceed 100 km/hour and its origin can be both S-SSW and SSE. The fetch over the Atlantic Ocean generates a sharp rise in sea level associated with downwelling at beach level. Its prevalence is mostly during winter season (Simionato et al., 2007) including May to September, and its frequency was dominant during dry and cold periods of the Holocene (Iriando et al., 2009; Piovano et al., 2009 and references therein).

The wind rose of tri-hourly average speeds between December 1997 and August 2008 (data from Instituto Nacional de Meteorología-InUMet, Uruguay) at Carrasco airport (**Figure 2**) shows that ESE direction prevailed up to 6 m/s (which is about 7 m/s at the coast), that is to say for most observations, whereas for greater speeds SW and WSW directions prevailed (Panario et al., 2008). The former were mainly related to the austral spring and summer months and the latter to mid-winter (July-August) strong

“pamperos” winds, which is supported by the high frequency of storm surges greater than 1.69 m above mean sea level (AMSL) reported by Verocai et al. (2015).

Wave rose (Dec 2003 - May 2006), available for the studied region (**Figure 3**) generated from the wave sensor data of the Municipal Government of Montevideo closely matches the prevailing ESE to SE and S wind and swell directions: SE 35%; E 24%; S 20%; W and SW <10%; NW, NE and N <5% (Panario et al., 2008).

Eastern and south-eastern winds (E, ESE, and SE) have different impact on the coast. For instance, floodings due to ESE and SE winds varies from + $\approx$ 0.2 m AMSL for average wind speed (5 m/s or 18 km/h) to + $\approx$ 1 m AMSL for strong winds of 20 m/s (72 km/h), whereas MSL (102 cm) matches weak winds speed of 2 m/s (Verocai et al., 2015). Their destructive effects are caused by a strong atmospheric low pressure to the north of Argentinean Mesopotamia (between the great Paraná and Uruguay Rivers), which fetch raises the level of RdIP and adjacent sea. These very destructive events are infrequent (Barros et al., 2005; Bischoff, 2005), only 2 or 3 per year in the Uruguayan coast (Simionato et al., 2007; Verocai et al., 2015) and tend to occur mainly during the summer (Simionato et al., 2007) and mid-winter (Verocai et al., 2015).

ESE winds are largely sea breeze prevailing from October to March. Their intensity is usually weak to moderate therefore produce no significant downwellings or storm surges as they only raise water height a few centimetres AMSL. These winds are associated with rebuilding of the beach prism because they generate waves that reinforce the action of the swell (**Figure 3**), which is the main builder of the beach prism on the Uruguayan coast. Thus, an increase in light ESE and ENE winds, prevailing from October-April and April-September respectively (Nagy et al., 2015a) can result in cushioning the effects of SLR, while an increase in meridional winds (+) may have the opposite effect.

Assuming that current response of the winds to air temperature variations are similar to what happened during the Upper Holocene (Iriondo et al., 2009; Piovano et al. 2009 and references therein), hypotheses can be developed on the basis of trends which forces are known, to the extent that their effects on coastal systems are verified. This approach would allow establishing midterm scenarios. Sedimentological evidences indicates that during the cold periods of the Holocene, including the Little Ice Age (from 1550-1850), there was predominance of dry weather, low temperatures and low rainfall in the pampean region and surrounding areas (Bracco et al., 2011; del Puerto et al., 2013; Iriondo et al., 2011, 2009; Piovano et al., 2009 and references therein). There is also geomorphological evidence, expressed in longitudinal dunes fields, suggesting the dominance of southern winds during the cold period of the Holocene (Iriondo and Garcia, 1993; Iriondo et al., 2011, 2009).

Observations over the last century show a sea level rise (SLR) of 11 cm in Montevideo (Bidegain et al., 2005; Magrín et al., 2007; Nagy et al., 2005) and 17 cm in Buenos Aires

as a result of global climate change and the regional effects of river discharge and winds (Barros et al., 2005; Magrín et al., 2007). Periods of acceleration and stabilisation have been recorded at Montevideo from 1971-2003 and 2004-2013 respectively (Nagy et al., 2014a, 2005; Verocai et al., 2015). The latter was mainly attributed to lower river discharges which could have masked the global trend.

Sea level along the Uruguayan coast fluctuates $\pm 0.1\text{--}0.2$ m (superimposed on SLR) due to ENSO-related anomalies in precipitation, river flow and winds. Wind-induced flooding reaches 1, 2, and 3 m AMSL on yearly, decadal, and historical timescale basis, respectively (Nagy et al., 2015b; Verocai et al., 2015).

While it is recognised that the input of fresh water can produce yearly time-scale differences in MSL of up to 20 cm (Verocai et al., 2015) the effects of atmospheric circulation is a key factor in the behaviour of most estuarine and costal variables (Gutiérrez et al., 2015; Simionato et al., 2007, 2005; Verocai et al., 2015). For instance, the retreat of pocket beaches of Montevideo seems to fit the Bruun Rule, without negative fluctuations associated with the freshwater input (Gutiérrez et al., 2015).

The well-known El Niño/La Niña - Southern Oscillation (ENSO) linked variability at the Uruguayan coast (Nagy et al., 2014a, 2008a, 2008b), and medium-term trends have been analysed, resulting in seasonal and biennial variability (Ortega et al., 2013; Simionato et al., 2005) linkable to Rossby waves (Simionato et al., 2007), and another one with a rate of return of 8-12 years, in addition to long cycles of between 70 and 80 years linked to the AMO (Ortega et al., 2013).

Gutiérrez et al. (2015) have suggested for pocket beaches of Montevideo, a possible relationship between ENSO and fluctuations of beach area. From all the information and assumptions presented here, it is arguable that the trends observed on the beaches of Montevideo and perhaps partly for the whole Uruguayan coast, should be analysed in terms of atmospheric circulation anomalies with effects and variations over time, likely associated with both natural variability and manmade climate change.

3. MATERIALS AND METHODS

This article is based on data compiled and generated by Gutiérrez (2010), as amended by Gutiérrez et al. (2015). A long-term analysis of a dense series of high definition remote sensing was performed since 1927 for four urban beaches (located in Montevideo city), and since 1937 for two tidal creek sandy mouths located few tens of km to the East of downtown.

3.1. Geomorphological analysis

Polygons were plotted on all analysed beach surfaces as proposed by Gutiérrez et al. (2015), Gutiérrez and Panario (2005) to overcome the uncertainty associated with estimating changes by random transect method. This procedure allows reducing uncertainty since the surface is equivalent to infinite transects (method commonly

used for these studies), and ultimately, the beach area is the parameter of greatest management interest. In order to standardise the process, the digitalisation of shorelines was conducted at a scale of 1:3,000, as proposed by Armaroli et al. (2006), Ciavola et al. (2003), and Gutiérrez and Panario (2005). To select indicators of the shoreline, only those identified in the entire time-series of records were used. These were:

i) The *previous high tide high - water level - PHTH-WL* (Boak and Turner, 2005) is used because relatively large differences between tides produce a minimal error in the horizontal cartographic translation of the coastline. This is because the segment of the beach above the high tide level is the one with greater slope in the cross section thereof (Gutiérrez et al., 2015).

ii) Another indicator called *wet/dry line or run-up maxima - WDL-RM* (Boak and Turner, 2005) was used. This line indicates the erosive - constructive limit set within the maximum range of the wave, which is neatly identifiable by the difference in tone on aerial photographs. It is found where the beach changes from a concave profile to a convex one, with a clear break in the slope's angle.

Univariate statistical analyses were developed for the data sets obtained for each study site. The selected statistical indicators were: central tendency and data dispersion: mean, median, mode, standard deviation, variance, coefficient of variation and R^2 . Deviations from the trend line, the R^2 , the existence (or not) of climatic anomalies and their forcing atmospheric circulation were also analysed.

3.2. Climatic and oceanographic analysis

Meteorological, hydrological and oceanographic databases were used to establish possible correlations of the observed changes in the position of the coastline, with series of events that could explain them. This approach seeks to analyse coincident responses in all or most of the analysed shores despite the noticeable differences in the physical parameters of the beaches and the mouths of this study.

Positive changes (accretion) of coastlines in both creeks' mouths were correlated with periods of heavy rainfall. During these periods, the ideal conditions for transporting sandy sediments to RdIP estuary and the ocean are generated. These sediments were introduced into the mouths by the tidal flow or wind where there are dunes next to the mouth, or due to inputs from the watershed, as analysed by Gutiérrez and Panario (2006, 2005).

This analysis was based on maximum rainfall events (> 80 mm/24 hours) in the basins of the Pando and Carrasco creeks using the precipitation rate and precipitation database (data provided by InUMet from the local rainfall network).

The positions of the coastlines, particularly the negative changes (retreats) were compared with the elevations of the RdIP caused by the great floods of the rivers

Paraná and Uruguay, together or separately (Camilloni and Barros, 2003, 2000; Menéndez and Berbery, 2006; Nagy et al., 2002). These floods are able to increase the water height of the RdIP by 10-20 cm or plus (Bidegain et al., 2005; Nagy et al., 2005; Verocai et al., 2015). Therefore, Nagy et al. (2007) raised the hypothesis that river floodings could be a cause of the retreat of the coastline.

The historical series of decadal average and yearly maximum storm surges from 1921-2008 recorded in Montevideo were also analysed focusing on the search for matches from aerial photographic surveys occurred after (from months to a year) SSE and SSW wind-storms producing significant rise in sea levels (storm surges).

In turn, the effect on the response of the coastline (retreat or accretion) was analysed for years with ENSO events through composites of wind anomalies ². Two methods were used: i) atmospheric reanalysis of the NCEP/NCAR (National Center for Environmental Prediction/National Center for Atmospheric Research), NOAA-CIRES Climate Diagnostics Center, Kalnay et al. (1996) based on the NOAA Global Circulation Model to estimate the zonal ³ (from the West: + and from the East: -) and meridional ⁴ (from the South: + and from the North: -) winds from 1948-2012, and ii) the Twentieth Century Reanalysis Project dataset (Version V2) lead by the Physical Sciences Division at the NOAA/ESRL (National Oceanic and Atmospheric Administration, Earth System Research Laboratory), and the CIRES at the University of Colorado, to estimate the wind vector anomaly from 1950-2010 and vector wind components anomalies at surface level from 1948-2011.

Tri-hourly average wind speed data from 36 directions were used, as well as the time-series of significant wave height and direction (Panario et al., 2008) in order to analyse and integrate results from climatic data, models and reanalyses.

These analyses were performed for the period October-March because beaches usually recover after the winter S and SW erosive winds (from July-August) and the ENSO signal is expressed from October onwards over the RdIP (Pisciottano et al., 1994; Ropelewski and Halpert, 1987). For the period 2000-2008 images are available for most of the years, so the possible relationships between the observed coastline fluctuations with Rossby waves were explored, because according to Simionato et al. (2005) they modify the position of the South Atlantic anticyclone (SA High Pressure).

ENSO events, classified as weak, moderate, strong or very strong were determined according to Climate Prediction Center, NOAA (<http://www.cpc.ncep.noaa.gov/>)

---

²/ Climate anomaly - The difference between the average climate over a period of several decades or more, and the climate during a particular month or season. (http://glossary.ametsoc.org/wiki/Climate_anomaly).

³/ Zonal Wind - The wind or component of a wind along a particular parallel of latitude: It is positive if it blows from the West and negative if it blows from the East.

⁴/ Meridional Wind - The wind or component of a wind along a particular meridian: It is positive if it blows from the South and negative if it blows from the North.

products/analysis_monitoring/ensostuff/ensoyears.shtml) and to Severov et al. (2004) for the period before 1950.

4. RESULTS AND DISCUSSION

4.1. Joint analysis of the study cases

Pocitos and Ramírez beaches have almost opposite orientations; however, they have a similar mineralogical composition (originated from rocks of Montevideo Formation). On the other hand, their drifts are divergent, so these are two separate beaches different from each other and the other beaches of Montevideo (Gutiérrez et al., 2015) (see Figure 1, Tables 1 and 2).

These beaches have statistically significant difference in the mineralogy of the sand ($p < 0.01$) with neighbouring Buceo and Malvín beaches, while the mouths of Carrasco and Pando creeks have similar sources of sediment as Malvín beach, but differ from the others by having a recirculation of sediment by wind action and contributions of their watersheds. In turn, these beaches have different gradients of depth, making them change their typology from intermediate beach to dissipative beach. Because of all these factors it is particularly important to analyse the causes of the similarities in their behaviour through time.

Table 1. Characterisation of four beaches.

Beach	Primary dune	Classification	Sand mineralogy	Slope	Coastal tipology	Orientation (normal to the direction tangent)	Exposure length (m)
Ramírez	No	Dissipative	Feldspathic quartz medium sands with amphibole and pyroxene	Smooth (1.9°)	Pocket	N68°E	500
Pocitos	No	Intermediate	Feldspathic quartz medium sands with amphibole and pyroxene as main accessory minerals	Smooth (1.8°)	Pocket	N43°W	1,425
Buceo	Protodune	Dissipative	Feldspar quartz sands, with very little amphibole and pyroxene, very few points garnet points	Smooth (1.9 to 2.2°)	Logarithmic spirals	N28°W	887
Malvín	Partially	Dissipative to intermediate	Medium feldspathic quartz sands, and very little amphibole and pyroxene, calcite, opaque, garnet points, zircon, tourmaline, andalusite and epidote	Smooth (1.9 to 2.2°)	Logarithmic spirals	N3°E	1,072

Table 2. Characterisation of two creek mouths.

Creeks' mouth	Primary dune	Classification	Sand mineralogy	Orientation (normal to the direction tangent)	Trench length (m)
Carrasco	Degraded	Dissipative to intermediate	Feldspar quartz medium sand with very little amphibole and pyroxene, calcite, opaque, garnet points, zircon, tourmaline, andalusite and epidote	N26.3°W and N36°W	1,549
Pando	No	Dissipative to intermediate	Feldspar quartz medium sand with very little amphibole and pyroxene, calcite, opaque, garnet points, zircon and tourmaline.	N24°W and N17.3°W	3,290

4.2. Wind, waves and storm surges

Not including hours without wind (calms), the hourly wind speed data at the Carrasco station (see **Figure 2**) fit well to a Weibull distribution with a tail toward higher-energy events. The so originated positive asymmetry was associated with the fact that the mean was greater than the mode of the distribution, indicating that the average wind conditions were disturbed by strong rare events linked to disturbances such as the passage of cyclone systems and cold fronts (Panario et al., 2008).

The wave rose data (see **Figure 3**) closely matches the prevailing wind and swell directions coinciding with modelled reanalyses. In turn, the exploratory analysis of the wave data in Punta Brava values showed that significant height (H_s , in meters) and period (T , in seconds) had a frequency distribution which fitted a log- Normal function ($p < 0.001$). The main parameters were: Average H_s (m) 0.539, T (s) 5.657427; median H_s (m) 0.46, T (s) 4.4; mode H_s (m) 0.35, T (s) 2.9; standard deviation H_s (m) 0.38121, T (s) 3.983412 (Panario et al., 2008).

Matches between records of aerial photos and previous storm surges were recorded in the years 1943, 1993 and 2005. Although there were no coincidence of storm surges occurrence with photos for all studied sites, the available superimposed trends suggest the existence of storm signs in all cases except Malvín, where the signal was only recorded in the dry sand surface, not in the full beach. This can be attributed to the fact that during strong southern wind storm surges Malvín beach receives sediments passing over the tombolo linking Malvín with Brava beach, since the height of waves exceed its maximum height, offsetting losses related to these extreme events. Three storm events coinciding with elevations of more than 2.11 m AMSL were found coinciding with average daily Southern winds (S) from 24 m/s (85 km/h) to 31 m/s (110

km/h). These events can overcome most of the prism beach in most of the study area. Their occurrence from 1921-2008 was six times (0.068), only once since 1950 (0.017). However, Verocai et al. (2015) found a slight increase of the occurrence of events greater than 1.69 and 1.89 m AMSL over the last decade.

4.3. Exceptional rainfall

There was no evidence of clear trends associated with extreme rainfall in the basins of Pando and Carrasco microtidal creeks and in the beaches near them.

4.4. Floods of the tributaries of the Río de la Plata basin

The comparison between the events of river flooding and the morphodynamics of the studied beaches showed an opposite trend to what might be expected, given that the highest recovery in all the studied sites coincided with major river floods, and relatively high water level values. These results are not consistent with the hypothesis linking high flow and erosion.

4.5. General atmospheric circulation anomalies

The existence of relationships between the evolution of beaches and ENSO events was analysed, showing multiple matches with photogrammetric records with the latter (**Figures 4, 5, and 6**). Several papers (Nagy et al., 2008b; Severov et al., 2004; Simionato et al., 2005), suggest the existence of a signal of ENSO events in the Southwestern Atlantic Ocean at sea level pressure (SLP) and sea surface temperature (SST) with a weakening of the Malvinas Current and strengthening of Brazil current during El Niño, and vice versa during La Niña. It is arguable, in agreement with the findings of Iriondo et al. (2009 and citation therein) for the Holocene, the existence of a higher incidence of strong winds from the South quadrant and more specifically Meridional Wind (+) during La Niña, and increased Zonal wind (-) during El Niño events.

Due to the existence of coincidences between La Niña events with coastline retreats or loss of dry area in all analysed beaches, we decided to verify the hypothesis that ENSO-related circulation anomalies could explain the observed changes. For this, the NCEP/NCAR reanalysis (Kalnay et al., 1996) and the Twentieth Century Reanalysis were used to estimate the meridional and zonal wind anomalies, and the latter was also used to estimate vector wind components at surface level respectively. These reanalyses were applied for moderate and strong ENSO events, and even weak ones, if they occurred in the context of a multi-year event (e.g., from 1973-1975).

Due to the relatively low number of moderate and strong El Niño/La Niña events from 1950-2010 (less than 20) the results of wind anomalies are not conclusive yet but allow stating a working hypothesis to depict the relationship between ENSO-related wind anomalies and beach accretion-erosion processes.

418 The analysis focused on the anomalies of surface winds on the Uruguayan coast,
419 showing a positive anomaly of SW winds associated with retreats of the coastline
420 during La Niña events as suggested by Gutiérrez et al. (2015) for Ramírez and Pocitos.
421 Nevertheless, if all La Niña events were considered, no significant correlation was
422 found as either observed by Simionato et al. (2005), which could be attributed to the
423 fact that the signal of weak ENSO events (± 0.5 °C) is poor. For instance, from
424 October-March ESE-SE winds usually increase and a light ENSO signal can be masked.

425 Wind anomalies were also recorded in neutral years, i.e., in 1961, 1970. In 1970
426 retreats were recorded in all beaches due to the observed increased frequency of
427 southern winds.

428 Thus, despite the strong association between El Niño and increased river discharges,
429 beach area was recovered during several of these events (Gutiérrez et al., 2015).

430 During strong ENSO events yearly sea levels (SL) at Montevideo lies outside the long-
431 term trend line. For instance, during La Niña SL is -5 to -15 cm below MSL and during El
432 Niño it lies +5 to +20 AMSL (Nagy et al., 2005; Verocai et al., 2015).

433 The observed behaviour of the coastline could be interpreted as follows: The SW winds
434 -associated with both winter period and La Niña variability- produce short period
435 waves, change the direction of long shore drift in the analysed beaches (except
436 Ramírez), raise the water level (although to a lesser extent than the S and SSE winds),
437 and also cause the retreat of the coastline. These effects are likely to be compensated
438 during austral spring and summer, or during El Niño events, when winds anomalies are
439 reverted intensifying the spring-summer time usual recovery. Wind climatology show
440 that the strongest wind velocities occur in August associated with SW-WSW winds and
441 October associated with ESE-SE winds (Nagy et al., 1997).

442 The NCEP-NCAR composites of meridional and zonal wind anomalies over the RdIP and
443 the Atlantic coast for moderate and strong ENSO events from 1950-2010 confirmed
444 that the most important anomaly was the meridional component from October to
445 March. During this period an increase of northerly winds (warm, moist and weak
446 winds) expressed by an anomaly of -0.2 m/s (**Figure 7**) was observed. During La Niña
447 southerly winds increased (positive anomaly of +0.2 m/s on the Atlantic coast). The
448 zonal component showed positive anomalies in both phases although somewhat
449 higher during El Niño when the zonal circulation, particularly the easterlies, slightly
450 increased. Results obtained with 20th century reanalysis for the Uruguayan coastal
451 areas were very similar.

452 The reanalysis of the fields of vector wind anomalies (speeds) from October-March for
453 all moderate and strong events showed anomalies from SW of 0.8 m/s during La Niña,
454 and 0.4 m/s from W-WNW during El Niño (**Figure 8**).

Summarising, climate analysis of joint frequency of wind speed and direction indicates higher mean speeds, e.g., in August and October with WSW-SW and ESE-SE winds respectively. The La Niña events increase the frequency of SW to WNW, which typically prevail during fall-winter (April-September). The El Niño increases ESE-SE ones during spring-summer (October-March). Therefore, it is arguable that it is very likely that the effect of the La Niña reinforces climatic winter beach sand losses (July-August), whereas El Niño reinforces and/or extends summer beach sand accretion from October onward.

Results of vector field anomalies' are closely associated with strong southern to western directions (WSW and SW) > 8 and > 10 m/s, whereas the meridional-zonal anomalies are closely associated with the prevailing south-eastern frequencies. The similar field anomalies during La Niña (SW to WNW) for both methods are the resultant of the coincidence of frequency and speed, whereas the differences for El Niño are related to the increased frequency of typically fresh to strong eastern to south-eastern winds (< 8 m/s) and of the far less frequent but stronger (> 8 m/s and > 10 m/s) western winds (see **Figure 2**). Therefore, the increased frequency of fresh SE winds (> 6 m/s) is masked by the small increase of strong W winds (> 8 m/s).

This result coincides with previous observations about ENSO influence on atmospheric circulation anomalies from October-March as well as with the observed and projected increase in eastern to south-eastern winds over the Uruguayan coast. Therefore, it could serve as an analogy to foresee coastal evolution and management scenarios.

The evolution of decadal mean of the maximum yearly storm surges (**Figure 9**) and of the AMO (Ortega et al., 2013) showed a similar development trend which was not reflected in the evolution of the studied beaches, except partially in Ramírez. However, this effect is likely to have occurred and should be explored at lower prism height beaches in some sections of both inner and outer estuary, with cliffs that have actively retreated in the past and which nowadays are stabilised or receding more slowly.

Since 2000 yearly available images show a bi-annual variation in neutral years (see last few years of **Figure 4**). This could be associated with the Rossby wave signal in SST and SLP in the South Atlantic reported by Simionato et al. (2005). Nonetheless, this hypothesis should be analysed using a denser and longer series of images and/or land monitoring.

Therefore, the key climatic teleconnections factor of accretion-erosion processes seem to be the occurrence of moderate to strong La Niña which are pulses of severe events, while El Niño events allow partial or full sand recovery on the short- and mid-term scales. The quantitative balance of this superimposed seasonal and interannual sedimentary budget is difficult to be made due to the parallel effects of gradual sea level rise, storm surges and human interventions.

Fisher et al. (2015) identified three specific challenges in climate change and resilience monitoring and evaluation: i) assessing attribution, ii) creating baselines, and iii) monitoring over long time horizons. This paper addressed the attribution of urban sandy beaches accretion-erosion processes, baselines, long-term series of climate teleconnections and weather extremes, and developed an approach to monitoring and evaluation, aiming to increase beach resilience to SLR and storminess.

Coastal management and adaptation to climate threats have been mostly based on reactive responses to date. In order to achieve practical management implications of ENSO impacts, as well as of other climatic modellers and threats to coastal systems, i.e., storm surges and SLR, it is necessary to foster continuous prognosis, monitoring and modelling of both climatic threats and coastal responses on monthly to seasonal basis. Soft measures such as rebuilding of sandy dunes and planting of psammophyte vegetation among others, require financial resources and logistical planning for their implementation at appropriate times. Any failure due to a lack of foresight often discourage authorities to allocate resources for this purpose. This would allow to transform a threat into an opportunity by taking pro-active actions to increase coastal resilience, particularly in relation to sandy beaches and tidal creeks' mouths.

5. CONCLUSIONS

This article surveyed the behaviour of managed urban sandy beaches at a wave dominated coast of a large and unique microtidal system, the Río de la Plata's northern coast in Uruguay, and climate stressors such as wind extreme events and climate teleconnections, mainly ENSO-linked variability.

The working assumption of this article may be resumed as follows: The shape, size and characteristics of sandy beach and dune systems are the consequence of the particular interaction between the energy of waves and wind, with their grain size, equilibrium profile and roughness. Any change to any of these factors may result in a realignment of the beach. This flexibility and ability to work with pulses is the reason that the sandy beaches are considered the best coastal defense. Therefore, these systems are cyclical, with alternating periods of erosion and accretion, to give a type of position and geometry in the medium and long term.

Main key findings are: i) A relationship was found between the observed changes in urban beaches and teleconnections (ENSO), while it is arguable the existence of an effect of Rossby waves. ii) No correlation was found with water level variations attributable to river discharge. iii) The sand surfaces of the middle estuary beaches analysed in this article have varied due to atmospheric circulation anomalies such as exceptional storm surges associated with more frequent southern winds that erode beaches or zonal winds that enable their recovery. iv) The studied beaches are often more eroded during strong La Niña events due to a higher frequency of typically strong

SW winds than during neutral years, whereas during El Niño a decrease in the frequency of SW winds and a relative stabilisation or even gain in the exposed sand area is observed. This finding is supported by climatic reanalyses of wind anomalies for moderate and strong ENSO events from 1950-2008 besides field and remote sensing observations since 1921. v) The strong surface gain of Ramírez Beach in 1961 could be correlated to a light eastern winds anomaly, which would have strengthened the swell. This was also observed, although to a lesser extent, in Pocitos, Buceo, Malvín beaches and in the Pando and Carrasco creek tidal mouth.

The main conclusion was that the results presented in this article highlight the likely existence of some degree of relationship between ENSO events and beach erosion-accretion processes.

The main questions to be explored in a further study are: i) Which is the response to other teleconnections e.g., Rossby waves and AMO? ii) Will the likely increase in the frequency and/or intensity of El Niño events be a risk or an opportunity for the studied beaches? iii) How was the relationship between ENSO wind-anomalies and beach accretion-erosion processes from 2009/10-2015/16 (when three moderate to strong events occurred).

Since ENSO seems to be a good descriptor of the status of sandy beaches of Montevideo during ENSO years and ENSO +1, there are practical implications in coastal erosion risk-management, climate adaptation, planning of sandy beaches conservation and restoration. The occurrence of El Niño events could become an opportunity to develop pro-active beach management plans to reduce risks provided an improvement of observational, forecasting and modelling tools and capacities are achieved. The implementation of this measure will also serve as an experience to adapt to both storm surges and gradual sea level rise.

Monitoring change within the coast and estuary play an important role in understanding the impacts of climate change.

ACKNOWLEDGMENTS

This research was conducted as part of the “Technical Proposal for the conservation and recovery of Buceo beach and certified beaches of Montevideo”, agreement between UNCIEP, Facultad de Ciencias, Universidad de la República and the Departamento de Desarrollo Ambiental, Intendencia de Montevideo (IdEM). We thanks especially to Gustavo Piñeiro for his collaboration in the mineralogical analysis in the context of this project.

We thank the dataset provided by the NOAA-ESRL Physical Sciences Division, Boulder Colorado from their Web site at <http://www.esrl.noaa.gov/psd/>. Support for the

Twentieth Century Reanalysis Project dataset is provided by the U.S. Department of Energy, Office of Science Innovative and Novel Computational Impact on Theory and Experiment (DOE INCITE) program, and Office of Biological and Environmental Research (BER), and by the National Oceanic and Atmospheric Administration Climate Program Office.

We too sincerely thank José E. Verocai for providing unpublished storm surges data and Daniela Peluffo for her language assistance.

REFERENCES

Armaroli, C., Ciavola, P., Balouin, Y., Gatti, M., 2006. An Integrated Study of Shoreline Variability Using GIS and ARGUS Techniques. *Journal of Coastal Research* (ISSN: 0749-0208) SI 39, 473–477.

Barros, V., Menéndez, A., Nagy, G.J. (Eds.), 2005. El cambio climático en el Río de la Plata. Project assessments of impacts and adaptation to climate change (AIACC), Centro de Investigaciones del Mar y la Atmósfera - CIMA/CONICET, Buenos Aires, Argentina.

Bidegain, M., Caffera, R.M., Blixen, F., Pshennikov, V.A., Lagomarsino, J.J., Forbes, E.A., Nagy, G.J., 2005. Tendencias climáticas, hidrológicas, y oceanográficas en el Río de la Plata y Costa Uruguay, in: Barros, V., Menéndez, Á., Nagy, G. (Eds.), *El Cambio Climático En El Río de La Plata*. Centro de Investigaciones del Mar y la Atmósfera - CIMA/CONICET, Buenos Aires, Argentina, pp. 137–143.

Bidegain, M., Stella, J.L., Betolli, M.L., Quintana, J., 2015. [Regional climates] Southern South America [in ‘State of the Climate in 2014’]. *Bulletin of the American Meteorological Society* 96, S182–S184. doi:10.1175/2015BAMSStateoftheClimate.1

Bidegain, M., Stella, J.L., Penalba, O.C., 2014. [Regional climates] Southern South America [in ‘State of the Climate in 2013’]. *Bulletin of the American Meteorological Society* 95, S172–S173. doi:10.1175/2014BAMSStateoftheClimate.1

Bischoff, S., 2005. Sudestadas, in: Barros, V., Menéndez, Á., Nagy, G. (Eds.), *El Cambio Climático En El Río de La Plata*. Project assessments of impacts and adaptation to climate change (AIACC), CIMA/CONICET, Buenos Aires, pp. 53–67.

Boak, E.H., Turner, I.L., 2005. Shoreline Definition and Detection: A Review. *Journal of Coastal Research* 21, 688–703. doi:10.2112/03-0071.1

Bracco, R., del Puerto, L., Inda, H., Panario, D., Castiñeira, C., García-Rodríguez, F., 2011. The relationship between emergence of mound builders in SE Uruguay and climate change inferred from opal phytolith records. *Quaternary International* 245, 62–73. doi:10.1016/j.quaint.2010.05.025

Camilloni, I., Barros, V., 2000. The Paraná River response to El Niño 1982–83 and 1997–98 events. *Journal of Hydrometeorology* 1, 412–430. doi:10.1175/1525-7541(2000)001<0412:TPRRT>2.0.CO;2

607 Camilloni, I., Barros, V., 2003. Extreme discharge events in the Paraná River and their
608 climate forcing. *Journal of Hydrology* 278, 94–106. doi:10.1016/S0022-
609 1694(03)00133-1

610 Ciavola, P., Corbau, C., Cibir, U., Perini, L., 2003. Mapping of the coastal zone of the
611 Emilia-Romagna region using geographical information systems, in: Ozhan, E.
612 (Ed.), *Proceedings of The Sixth International Conference on the Mediterranean*
613 *Coastal Environment (MEDCOAST)*, Volume III. MEDCOAST, Ravenna, Italy, pp.
614 2363–2374.

615 Codignotto, J.O., Dragani, W.C., Martin, P.B., Simionato, C.G., Medina, R.A., Alonso, G.,
616 2012. Wind-wave climate change and increasing erosion in the outer Río de la
617 Plata, Argentina. *Continental Shelf Research* 38, 110–116.
618 doi:10.1016/j.csr.2012.03.013

619 del Puerto, L., Bracco, R., Inda, H., Gutiérrez, O., Panario, D., García-Rodríguez, F.,
620 2013. Assessing links between late Holocene climate change and
621 paleolimnological development of Peña Lagoon using opal phytoliths, physical,
622 and geochemical proxies. *Quaternary International* 287, 89–100.
623 doi:10.1016/j.quaint.2011.11.026

624 Dias, J.M.A., Cearreta, A., Isla, F.I., de Mahiques, M.M., 2013. Anthropogenic impacts
625 on Iberoamerican coastal areas: Historical processes, present challenges, and
626 consequences for coastal zone management. *Ocean and Coastal Management* 77,
627 80–88. doi:10.1016/j.ocecoaman.2012.07.025

628 Dragani, W.C., Martin, P.B., Simionato, C.G., Campos, M.I., 2010. Are wind wave
629 heights increasing in south-eastern south American continental shelf between
630 32°S and 40°S? *Continental Shelf Research* 30, 481–490.
631 doi:10.1016/j.csr.2010.01.002

632 EcoPlata, 1999. Diagnóstico ambiental y socio-demográfico de la zona costera
633 uruguaya del Río de la Plata. Recopilación de informes técnicos.
634 CIID/PNUD/MVOTMA/UNESCO, Montevideo, Uruguay.

635 Escobar, G., Vargas, W., Bischoff, S., 2004. Wind tides in the Rio de la Plata estuary:
636 Meteorological conditions. *International Journal of Climatology* 24, 1159–1169.
637 doi:10.1002/joc.1026

638 Fisher, S., Dinshaw, A., McGray, H., Rai, N., Schaar, J., 2015. Evaluating climate change
639 adaptation: Learning from methods in international development. *Monitoring and*
640 *evaluation of climate change adaptation: A review of the landscape*. *New*
641 *Directions for Evaluation* 13–35. doi:10.1002/ev.20128

642 Flood, S., Schechtman, J., 2014. The rise of resilience: Evolution of a new concept in
643 coastal planning in Ireland and the US. *Ocean & Coastal Management* 102, 19–31.
644 doi:10.1016/j.ocecoaman.2014.08.015

645 Gutiérrez, O., 2010. Dinámica sedimentaria en la costa uruguaya: evolución y
646 tendencias de playas urbanas en el marco del Cambio Global. (Master's thesis).
647 Universidad de la República, Montevideo, Uruguay.
648 doi:10.13140/RG.2.1.1904.4568

649 Gutiérrez, O., 2016. Dinámica sedimentaria en la costa uruguaya. Un abordaje desde la
650 geografía histórica y los S.I.G. para comprender las tendencias evolutivas
651 vinculadas al Cambio Global. (Unpublished doctoral thesis). Universidad
652 Internacional de Andalucía, Huelva, Spain.

653 Gutiérrez, O., Panario, D., 2005. Dinámica geomorfológica de la desembocadura del
654 Arroyo Pando, Uruguay. Geografía histórica y SIG, análisis de tendencias naturales
655 y efectos antrópicos sobre sistemas dinámicos. Xeográfica, Revista de Xeografía,
656 Territorio e Medio Ambiente (ISSN: 1578-5637) 5, 107–126.

657 Gutiérrez, O., Panario, D., 2006. Evolución de la desembocadura del Arroyo Pando
658 (Canelones, Uruguay): ¿tendencias naturales o efectos antrópicos?, in: Menafrá,
659 R., Rodríguez-Gallego, L., Scarabino, F., Conde, D. (Eds.), Bases Para La
660 Conservación Y El Manejo de La Costa Uruguaya. Vida Silvestre Uruguay,
661 Montevideo, Uruguay, pp. 391–400.

662 Gutiérrez, O., Panario, D., Nagy, G.J., Piñeiro, G., Montes, C., 2015. Long-term
663 morphological evolution of urban pocket beaches in Montevideo (Uruguay):
664 impacts of coastal interventions and links to climate forcing. Journal of Integrated
665 Coastal Zone Management - Revista de Gestão Costeira Integrada 15, (in press).
666 doi:10.5894/rgci553

667 Holling, C.S., Gunderson, L.H., 2002. Resilience and Adaptative Cycles, in: Gunderson,
668 L.H., Holling, C.S. (Eds.), Panarchy, Understanding Transformations in Human and
669 Natural Systems. Island Press, Washington DC, U.S.A., pp. 25–62.

670 Iriondo, M., Brunetto, E., Kröhling, D., 2009. Historical climatic extremes as indicators
671 for typical scenarios of Holocene climatic periods in the Pampean plain.
672 Palaeogeography, Palaeoclimatology, Palaeoecology 283, 107–119.
673 doi:10.1016/j.palaeo.2009.09.005

674 Iriondo, M., García, N.O., 1993. Climatic variations in the Argentine plains during the
675 last 18,000 years. Palaeogeography, Palaeoclimatology, Palaeoecology 101, 209–
676 220. doi:10.1016/0031-0182(93)90013-9

677 Iriondo, M., Kröhling, D., Brunetto, E., 2011. Aridization, dune dissipation and
678 pedogenesis in the quaternary of eastern pampean sand sea, in: Murphy, J.A.
679 (Ed.), Sand Dunes. Nova Science Publishers, Inc., New York, U.S.A., pp. 1–42.

680 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell, M.,
681 Saha, S., White, G., Woollen, J., Zhu, Y., Leetmaa, A., Reynolds, R., Chelliah, M.,
682 Ebisuzaki, W., Higgins, W., Janowiak, J., Mo, K.C., Ropelewski, C., Wang, J., Jenne,
683 R., Joseph, D., 1996. The NCEP/NCAR 40-year reanalysis project. NCEP/NCAR 40-
684 Year reanalysis project. Bulletin of the American Meteorological Society 77, 437–
685 470.

686 Ledoux, L., Turner, R.K., 2002. Valuing ocean and coastal resources: a review of
687 practical examples and issues for further action. Ocean and Coastal Management
688 45, 583–616. doi:10.1016/S0964-5691(02)00088-1

689 López Laborde, J., Nagy, G.J., 1999. Hydrography and Sediment Transport
690 Characteristics of the Río de la Plata: A Review, in: Perillo, G.M.E., Piccolo, M.C.,
691 Pino-Quivira, M. (Eds.), Estuaries of South America. Their Geomorphology and

692 Dynamics. Springer, Berlin, Heidelberg, Germany, pp. 133–159. doi:10.1007/978-
693 3-642-60131-6_7

694 Luisetti, T., Turner, R.K., Jickells, T., Andrews, J.E., Elliott, M., Schaafsma, M.,
695 Beaumont, N., Malcolm, S., Burdon, D., Adams, C.A., Watts, W., 2014. Coastal
696 Zone Ecosystem Services: from science to values and decision making; a case
697 study. *Science of The Total Environment* 493, 682–693.
698 doi:10.1016/j.scitotenv.2014.05.099

699 Magrín, G.O., Gay García, C., Cruz Choque, D., Giménez, J.C., Moreno, A.R., Nagy, G.J.,
700 Nobre, C., Villamizar, A., 2007. Latin America. *Climate Change 2007: impacts,*
701 *adaptation and vulnerability*, in: Parry, M.L., Canziani, O.F., Palutikof, J.P., Linden,
702 P.J. van der, Hanson, C.E. (Eds.), *Contribution of Working Group II to the Fourth*
703 *Assessment Report of the Intergovernmental Panel on Climate Change.*
704 Cambridge University Press, Cambridge, United Kingdom, pp. 581–615.

705 Marshall, F.E., Banks, K., 2013. *Shoreline Habitat: Beaches.*, in: Nettle, W.K., Fletcher,
706 P.J. (Eds.), *Integrated Conceptual Ecosystem Model Development for the*
707 *Southeast Florida Coastal Marine Ecosystem.* NOAA Technical Memorandum,
708 OAR-AOML-103 and NOS-NCCOS-163, Florida, Miami, U.S.A., pp. 94–108.

709 Masselink, G., Gehrels, R. (Eds.), 2014. *Coastal Environments and Global Change.*
710 American Geophysical Union - Wiley-Blackwell, New York, U.S.A.

711 Menéndez, A., Berbery, E.H., 2006. Tendencias hidrológicas en la Cuenca del Plata, in:
712 Barros, V., Clarke, R., Silva Días, P. (Eds.), *El Cambio Climático En La Cuenca Del*
713 *Plata.* CIMA/CONICET, Buenos Aires, Argentina, pp. 81–92.

714 MTOP/PNUD/UNESCO, 1979. *Conservación y mejora de playas - URU. 73.007.* Unesco,
715 Montevideo, Uruguay.

716 Nagy, G.J., Bidegain, M., Caffera, R.M., Norbis, W., Ponce, A., Pshennikov, V., Severov,
717 D.N., 2008a. Fishing strategies for managing climate variability and change in the
718 Estuarine Front of the Río de la Plata, in: Leary, N., Adejuwon, J., Barros, V.,
719 Burton, I., Kulkarni, J. (Eds.), *Climate Change and Adaptation.* Routledge, London,
720 United Kingdom, pp. 353–370.

721 Nagy, G.J., Bidegain, M., de los Santos, B., Verocai, J.E., 2015a. Climate and sea level
722 baselines for the Coastal Areas of Uruguay from 1979-2014. IECA Technical
723 Reports Series: Coastal Areas Science and Management N° 1, December 2015.
724 Facultad de Ciencias, Universidad de la República. Montevideo, Uruguay.
725 doi:10.13140/RG.2.1.3632.9040

726 Nagy, G.J., Gómez-Erache, M., Fernández, V., 2007. El aumento del nivel del mar en la
727 costa uruguaya del Río de la Plata. *Tendencias, vulnerabilidades y medidas para la*
728 *adaptación.* Medio Ambiente y Urbanización (ISSN 0326-7857) 67, 77–93.

729 Nagy, G.J., Gómez-Erache, M., Kay, R., 2015b. A risk-based and participatory approach
730 to assessing climate vulnerability and improving governance in coastal Uruguay,
731 in: Glavovic, B., Kelly, M., Kay, R., Travers, A. (Eds.), *Climate Change and the Coast*
732 *Building Resilient Communities.* CRC Press, Boca Raton, Florida, U.S.A., pp. 357–
733 378. doi:10.1201/b18053-23

734 Nagy, G.J., Gómez-Erache, M., López, C.H., Perdomo, A.C., 2002. Distribution patterns
735 of nutrients and symptoms of eutrophication in the Rio de la Plata River Estuary
736 System. *Hydrobiologia* 475-476, 125–139. doi:10.1023/A:1020300906000

737 Nagy, G.J., Martínez, C.M., Caffera, R.M., Pedrosa, G., Forbes, E.A., Perdomo, A.C.,
738 López Laborde, J., 1997. The Hydrological and Climatic Setting of the Rio de la
739 Plata, in: Wells, P.G., Daborn, G.R. (Eds.), *The Rio de La Plata: An Environmental*
740 *Overview. An EcoPlata Project Background Report.* Dalhousie University, Halifax,
741 Nueva Escocia, Canada, pp. 17–68.

742 Nagy, G.J., Muñoz, N., Verocai, J.E., Bidegain, M., Seijo, L., 2014a. Adjusting to current
743 climate threats and building alternative future scenarios for the Rio de la Plata
744 coast and estuarine front, Uruguay. *Journal of Integrated Coastal Zone*
745 *Management - Revista de Gestão Costeira Integrada* 14, 553–568.
746 doi:10.5894/rgci472

747 Nagy, G.J., Ponce, A., Pshennikov, V.A., Silva, R., Forbes, E.A., Kokot, R.R., 2005.
748 Desarrollo de la capacidad de evaluación de la vulnerabilidad costera al Cambio
749 Climático: Zona Oeste de Montevideo como caso de estudio, in: Barros, V.,
750 Menéndez, Á., Nagy, G. (Eds.), *El Cambio Climático En El Río de La Plata.* Centro de
751 *Investigaciones del Mar y la Atmósfera - CIMA/CONICET,* Buenos Aires, Argentina,
752 pp. 173–180.

753 Nagy, G.J., Pshennikov, V.A., Robatto, P., 2003. Monthly variability of salinity at
754 Montevideo, Rio de la Plata's frontal zone, in response to ENSO fluctuations and
755 to the River Uruguay Flow (1998-2000), in: Vizziano, D., Puig, P., Mesones, C.,
756 Nagy, G.J. (Eds.), *Río de La Plata: Research to Manage the Environment, Fish*
757 *Resources and the Fishery in the Saline Front.* Montevideo, Uruguay, pp. 21–31.

758 Nagy, G.J., Seijo, L., Verocai, J.E., Bidegain, M., 2014b. Stakeholders' climate perception
759 and adaptation in coastal Uruguay. *International Journal of Climate Change*
760 *Strategies and Management* 6, 63–84. doi:10.1108/IJCCSM-03-2013-0035

761 Nagy, G.J., Seijo, L., Verocai, J.E., Brugnoli, E., Bidegain, M., 2013. Enfoque,
762 conocimiento y medidas para enfrentar las amenazas del clima presente en la
763 zona frontal del Río de la Plata, Uruguay. *Costas: revista Iberoamericana de*
764 *manejo costero integrado* (ISSN 2304-0963) 2, 67–87.

765 Nagy, G.J., Severov, D.N., Pshennikov, V.A., De los Santos, M., Lagomarsino, J.J., Sans,
766 K., Morozov, E.G., 2008b. Rio de la Plata estuarine system: Relationship between
767 river flow and frontal variability. *Advances in Space Research* 41, 1876–1881.
768 doi:10.1016/j.asr.2007.11.027

769 Nicholls, R.J., Cazenave, A., 2010. Sea-level rise and its impact on coastal zones. *Science*
770 328, 1517–1520. doi:10.1126/science.1185782

771 Ortega, L., Celentano, E., Finkl, C., Defeo, O., 2013. Effects of climate variability on the
772 morphodynamics of Uruguayan sandy beaches. *Journal of Coastal Research* 29,
773 747–755. doi:10.2112/JCOASTRES-D-13-00003.1

774 Panario, D., Gutiérrez, O., 2005. La vegetación en la evolución de playas arenosas. El
775 caso de la costa uruguaya. *Ecosistemas. Revista de la Asociación Española de*
776 *Ecología Terrestre* 14, 150–161.

777 Panario, D., Gutiérrez, O., Piñeiro, G., 2008. Propuesta técnica para la conservación
778 recuperación de la playa del Buceo y playas certificadas de Montevideo. Informe
779 final. Convenio UNCIEP, Facultad de Ciencias y Departamento de Desarrollo
780 Ambiental, Intendencia Municipal de Montevideo. Setiembre. Montevideo,
781 Uruguay. doi:10.13140/RG.2.1.2508.8725

782 Panario, D., Piñeiro, G., 1997. Vulnerability of oceanic dune systems under wind
783 pattern change scenarios in Uruguay. *Climate Research* 9, 67–72.
784 doi:10.3354/cr009067

785 Pereira, N.E. da S., Klumb-Oliveira, L.A., 2015. Analysis of the influence of ENSO
786 phenomena on wa (Brazil). *Journal of Integrated Coastal Zone Management -*
787 *Revista de Gestão Costeira Integrada* 15, 353–370. doi:10.5894/rgci570

788 Piovano, E.L., Ariztegui, D., Córdoba, F., Cioccale, M., Sylvestre, F., 2009. Hydrological
789 variability in South America below the Tropic of Capricorn (Pampas and eastern
790 Patagonia, Argentina) during the last 13.0 ka, in: Vimeux, F., Sylvestre, F., Khodri,
791 M. (Eds.), *Past Climate Variability from the Last Glacial Maximum to the Holocene*
792 *in South America and Surrounding Regions. From the Last Glacial Maximum to the*
793 *Holocene, Developments in Paleoenvironmental Research*. Springer, Netherlands,
794 pp. 323–351. doi:10.1007/978-90-481-2672-9_14

795 Pisciotto, G., Díaz, A., Cazess, G., Mechoso, C.R., 1994. El Niño-Southern Oscillation
796 Impact on rainfall in Uruguay. *Journal of Climate* 7, 1286–1302. doi:10.1175/1520-
797 0442(1994)007<1286:ENSOIO>2.0.CO;2

798 Ropelewski, C.F., Halpert, M.S., 1987. Global and regional scale precipitation patterns
799 associated with the El Niño/Southern Oscillation. *Monthly Weather Review* 115,
800 1606–1626. doi:10.1175/1520-0493(1987)115<1606:GARSPP>2.0.CO;2

801 Severov, D.N., Mordecki, E., Pshennikov, V.A., 2004. SST anomaly variability in
802 Southwestern Atlantic and El Niño/Southern oscillation. *Advances in Space*
803 *Research* 33, 343–347. doi:10.1016/S0273-1177(03)00479-4

804 Simionato, C.G., Meccia, V.L., Guerrero, R.A., Dragani, W.C., Nuñez, M.N., 2007. Río de
805 la Plata estuary response to wind variability in synoptic to intraseasonal scales: 2.
806 Currents' vertical structure and its implications for the salt wedge structure.
807 *Journal of Geophysical Research* 112, C07005. doi:10.1029/2006JC003815

808 Simionato, C.G., Vera, C.S., Siegmund, F., 2005. Surface wind variability on seasonal
809 and interannual scales over Río de la Plata area. *Journal of Coastal Research* 21,
810 770–783. doi:10.2112/008-NIS.1

811 Stevens, S., Milstead, B., Albert, M., Entsminger, G., 2005. Northern coastal and barrier
812 network vital signs monitoring plan. Technical Report NPS/NER/NRTR-2005/025.
813 U.S. Department of the Interior, National Park Service, Northeast Region, Boston,
814 MA, U.S.A.

815 Verocai, J.E., Gómez-Erache, M., Nagy, G.J., Bidegain, M., 2015. Addressing climate
816 extremes in Coastal Management: The case of the Uruguayan coast of the Río de
817 la Plata System. *Journal of Integrated Coastal Zone Management - Revista de*
818 *Gestão Costeira Integrada* 15, 91–107. doi:10.5894/rgci555

819 Young, I.R., Zieger, S., Babanin, A.V., 2011. Global trends in wind speed and wave
820 height. *Science* (New York, N.Y.) 332, 451–455. doi:10.1126/science.1197219
821

Figure 1

Figure 1. Study area: a) South America, L and H indicate low and high atmospheric pressure. b) Uruguay. c) Beaches: 1) Ramírez, 2) Pocitos, 3) Buceo, 4) Malvín; and Carrasco and Pando creeks. 5) Indicates the location of the Punta Brava wave sensor. 6) Carrasco airport meteorological station. The white arrow indicates the drift direction.

Figure 2

Figure 2. Carrasco wind rose 1998-2008. The frequency of directions is shown for all winds (left), and for those higher than 6 m/s (center left), higher than 8 m/s (center right), and higher than 10 m/s (right) from 1998-2008 (Panario et al., 2008).

Figure 3

Figure 3. Punta Brava wave rose from 2003-2006 (between Pocitos and Ramírez beaches, see Figure 1). The prevailing direction related to swell matches quite well the prevailing E-SE winds, particularly during the austral summer (Panario et al., 2008).

Figure 4

Figure 4. a) Historical development of the beaches Ramírez, Pocitos, Buceo and Malvín, indicated as coastal accretion or retreat process. b) ENSO events, classified as weak, moderate, strong or very strong according to Climate Prediction Center, NOAA (period 1950-2010). c) ENSO events according to Severov et al. (2004) for events before 1950. The vertical gray lines (La Niña) and the vertical dashed black lines (El Niño) suggest the links between the coastal fluctuations and teleconnections.

Figure 5

Figure 5. a) Historical development of the mouth tidal creeks Carrasco and Pando, indicated as coastal accretion or retreat process. b) ENSO events, classified as weak, moderate, strong or very strong according to Climate Prediction Center, NOAA (period 1950-2010). c) ENSO events according to Severov et al. (2004) for events before 1950. The vertical gray lines (La Niña) and the vertical dashed black lines (El Niño) suggest the links between the coastal fluctuations and teleconnections.

Figure 6

Figure 6. Historical trend-line of study sites. Beaches: a) Ramírez, b) Pocitos, c) Buceo, d) Malvín. Creeks' mouths: e) Carrasco, f) Pando. The series of remote sensing observations have a minimum of 17 records and a maximum of 27. Different proxy records were explored as indicators of coastline in order to interpret the responses of the system, wet/dry line or run-up maxima (WDLRM) or previous high tide high water level (PHTH-WL).

Figure 7

Figure 7. Composite of zonal and meridional wind anomalies for moderate and strong ENSO from 1950-2010 (NCEP-NCAR reanalysis). Zonal anomalies: El Niño (above left), La Niña (above right), and meridional anomalies: El Niño (below left) and Niña (below right).

Figure 8

Figure 8. Wind Vector Anomaly (m/s) for ENSO events during Oct-Mar, from 1950-2010 (NOAA 20th Century Reanalysis Project): La Niña years (above) and El Niño years (below).

Figure 9

Figure 9. Evolution of decadal mean of the maximum yearly storm surges for de RldP.