Observations of *in situ* deep-sea marine bioluminescence with a high-speed, high-resolution sCMOS camera

Phillips, BT^a; Gruber, DF^b; Vasan, G^c; Roman, CN^a; Pieribone, VA^{c,d*}; Sparks, JS^{e*}.

^aUniversity of Rhode Island, Graduate School of Oceanography, Narragansett RI 02882 USA

^bCity University of New York, Baruch College, New York, NY USA

^cThe John B. Pierce Laboratory, New Haven, CT USA

^d Cellular and Molecular Physiology & Neurobiology, Yale University School of Medicine, New Haven, CT USA

^eAmerican Museum of Natural History, New York, NY USA

^{*}*contributed equally*

Abstract

Observing and measuring marine bioluminescence *in situ* presents unique challenges, characterized by the difficult task of approaching and imaging weakly illuminated bodies in a three-dimensional environment. To address this problem, a scientific complementary-metal-oxide-semiconductor (sCMOS) microscopy camera was outfitted for deep-sea imaging of marine bioluminescence. This system was deployed on multiple platforms (manned submersible, remotely operated vehicle, and towed body) in three oceanic regions (Western Tropical Pacific, Eastern Equatorial Pacific, and Northwestern Atlantic) to depths up to 2500 m. Using light stimulation, bioluminescent responses were recorded at high frame rates and in high resolution, offering unprecedented low-light imagery of deep-sea bioluminescence *in situ*. The kinematics of light production in several zooplankton groups was observed, and luminescent responses at different depths were quantified as intensity vs. time. These initial results signify a clear advancement in the bioluminescent imaging methods available for observation and experimentation in the deep-sea.

Keywords: Bioluminescence, zooplankton, sCMOS, low-light camera, light stimulation

1. Introduction

Marine bioluminescence has generated increasing interest among the scientific community in the past several decades, spanning the disciplines of comparative biology (e.g. Haddock and Case 1999), biochemistry (e.g. Prasher et al. 1992), physiology (e.g. Contag and Bachmann 2002), neuroscience (e.g. Martin 2008), population dynamics (e.g. Moline et al. 2009), and naval applications (e.g. Fucile 2002). This research began with terrestrial fireflies (Green and McElroy 1956) and later expanded to readily accessible marine organisms, most notably bacteria (Bassler et al. 1993; Surette et al 1999), copepods (Campbell and Herring 1990), and cnidarians (Prasher et al. 1985). Whereas almost all major phyla of deep-sea zooplankton exhibit bioluminescence (Haddock et al. 2010), with new discoveries reported annually, only a handful of studies have been published on *in situ* marine bioluminescence. As a result, the deep-sea offers vast potential for future multidisciplinary discoveries related to bioluminescence.

The greatest limitation to observing marine bioluminescence *in situ* is technology. As early as 1955, photomultiplier tubes (PMTs) were employed to quantitatively measure light in the ocean from a single point (Boden and Kampa 1957), a method still used today (Adrián-Martínez et al. 2014, Johnsen et al. 2014, Craig et al. 2015). In the past three decades, a number of studies utilizing Intensified Silicon Intensifier Target (ISIT) cameras were published (Widder et al. 1989, Robison 1992, Priede et al. 2006). ISIT cameras are very sensitive, but are limited in resolution and retain residual images between frames, offering visually-pleasing images but making them less than ideal quantitative imaging platforms. More recently, image-intensified and electron-multiplied CCD cameras have been employed to measure extremely low light sources in high resolution (e.g. Johnson et al. 2012, Craig et al. 2015). Scientific complementary-metal-oxide-semiconductor (sCMOS) cameras offer high sensitivity, low noise, wide dynamic range, and high-resolution, high-speed capabilities without image amplification (Coates et al. 2009, Fowler et al. 2009).

We hypothesize that applying sCMOS technology *in situ* offers the ability to study the kinematics of luminescent organisms at an unprecedented resolution, and by pairing this camera with light-stimulation methods, it may be possible to quantify the luminescence of the midwater community in both spatial and temporal scales. Here we present the results of cross-platform deployment of a sCMOS camera *in situ* to depths up to 2500 m, utilizing light stimulation to elicit bioluminescent responses in an exploratory fashion.

2. Methods

A Hamamatsu Photonics ORCA-Flash4.0 V2 sCMOS camera was chosen for its high dynamic range (16-bit), high quantum efficiency (72%), 2048 x 2048 resolution, and high speed (up to 100 fps). All recordings in this study were conducted at 30fps using a global shutter. The camera was fitted with a Navitar 17 mm *f*0.95 lens and integrated into a 2500 m-rated housing (Prevco) with a flat acrylic viewport (Figure 1). The entire system was run on 24 VDC provided from surface power, with a nominal power draw of 100 W. Due to the high data output of the camera, a solid-state computer (Copperhead from VersaLogic, Tualatin, OR) with a 1 TB RAID0 SSD drive array was included in the housing along with a water-cooling pump and focus/aperture control. All internal components producing light, such as LEDs, were blacked out inside the housing. The recorded 16-bit intensity images were processed with Matlab v. R2012a. The camera power and focus was controlled by an RS232 serial connection. Remote access via Ethernet was used to control the embedded computer. Images were collected using software either from Stanford Photonics (Palo Alto, CA) or using custom software LabView GUIs. Raw data was uploaded from the embedded SSDs via Ethernet upon recovery of the instrument.

Image processing to count lit ‘blobs’ and calculate Intensity Units (IU) in each frame is described in the following pseudo-code:

for each sequential image in a recorded movie

 Subtract minimum image intensity calculated from entire movie (dark field; average value, 472 AU);

 Calculate maximum image intensity;

 Subtract reference image frame from current image (composite mean of 100 dark field images recorded with no bioluminescent sources);

 High image pixel intensity thresholding to discount strobe-lit frames;

 Further thresholding of low-light image, with threshold defined by the standard deviation of the image added to mean image intensity;

 Morphological opening of thresholded image to isolate and count regions larger than

95
96
97

2 pixels in diameter;
end

98

99
100

Fig. 1. (Top) Block diagram and internal assembly of subsea sCMOS camera system: RC = RS232 power/focus control; FO = fiber optic to Ethernet mux; PC = control computer; WP = cooling water pump; HD = RAID hard drive array; CA = camera head; VP = viewport. (Bottom) Internal view of camera system. The assembled subsea housing measures 75 cm long, 20 cm diameter, and has an approximate in-water weight of 18 kg.

106

Field sites for this study were chosen based on collaborative opportunity with several research expeditions. All recordings were made at night to eliminate the incidence of weak ambient light from the surface. Bioluminescent responses were induced using strobe light stimulation following methods first described by Neshyba (1967). Limited mechanical stimulation was induced when bioluminescence animals randomly came into contact with the viewport.

112

Initial deployment of the camera system was aboard a Triton 3k3 submersible as part of a biodiversity exploration expedition to the Solomon Islands in September 2013 (Figure 2C). The camera was mounted on a forward-facing instrumentation frame so that recorded video matched what observers inside the submersible sphere viewed in real-time. All lights inside and outside the submersible were either turned off or blacked out with opaque tape. Light stimulation was done using a handheld Nikon Speedlight SB-700 strobe, set to produce a sequence of 5 flashes at 2 Hz. Several dives were conducted at night to depths up to 973 m, with two notable dives taking place approximately 22 km SE of Gizo Island and 6 km N of Mborokua Island, respectively. At Mborokua Island, a vertical transect through the water column was conducted via submersible with a depth station every 100 m down to the seafloor at 900 m.

122

A second research cruise, to the New England shelf break, was conducted in November 2014 (Figure 2B). The camera system was attached to a 1-ton towed body equipped with power switching and DSL internet to the surface, run through a standard 0.322" coaxial oceanographic wire. The viewport was oriented sideways to limit transverse water movement against it. Light stimulation was accomplished using a Vivitar Thyristor strobe placed inside a borosilicate instrument sphere, set to

126

produce a sequence of 5 flashes at 2 Hz. Two deployments were conducted at night to 900 m at Veatch Canyon and Atlantis Canyon.

A third deployment took place onboard the ROV HERCULES in the Galapagos Islands region in June/July 2015 (Figure 2A). The camera was mounted vertically pointing downwards into open water. Light stimulation was conducted using a pair of Ocean Imaging Systems 3831 strobes mounted 3 m forward, pointed at an oblique 45° angle towards the camera, and set to produce a sequence of 2 flashes spaced by a system cycle time of 2 s. Recordings were made opportunistically in the water column and on the seafloor, including at a 'black smoker' hydrothermal vent and a lower-temperature vent community dominated by *Riftia* sp. tubeworms.

Fig. 2: Deployment sites (yellow stars) for bioluminescent imaging in the Galapagos Islands (A), the New England shelf break (B) and the Solomon Islands (C). Bathymetry contours are spaced at 500 m.

3. Results

Over 200 15-30 second-length recordings were made at depths ranging from 30 m to 2500 m. Observed bioluminescence responses were most pronounced using repetitive flash stimuli; this was investigated on a submersible dive, when different flash characteristics were compared visually. At depths exhibiting a high bioluminescence response, human observers witnessed an omnidirectional display that momentarily resembled a bright night sky (i.e. the "Milky Way"). In many cases, major zooplankton groups were recognized based on the shape and pattern of the emitted bioluminescence. Siphonophores, ctenophores, and fishes were among the organisms discernable by their light response in otherwise complete blackness. Notably, light-stimulated bioluminescence responses on

153 the seafloor were almost non-existent, and therefore layers of dense zooplankton populations in the
154 water column were targeted for subsequent recordings.

155 Recordings of bioluminescence along a vertical cast in the water column illustrate the system's
156 potential to measure light-stimulated bioluminescence as an assay of zooplankton population density
157 (Figure 3). Spatially, total bioluminescence exhibited an increasing trend with depth to a maximum at
158 400 m, followed by a steady decrease towards the seafloor. Typical bioluminescent responses at each
159 depth were characterized by a sharp increase following the flash stimulation, rising to a maximum
160 within 1 s followed by a complete decline within ~4 s (Figure 4). Figure 4A demonstrates a relatively
161 short-lived bioluminescent response event of approximately 0.5 s that occurred after the fourth
162 strobe event, while the response in Figure 4B shows a discrete response after each strobe with
163 increasing intensity. Figure 4C is characteristic of many light-stimulated responses observed in this
164 study, with a delay in luminescence of ~1 s following an initial strobe event and a gradual decline
165 thereafter. In some cases, the decline in luminescence follows an exponential pattern (Figure 4B) and
166 in others a more linear regression was observed (Figure 4A and C). Low levels of light measured
167 before stimulus (<50 AU) reflect a background signal on the image sensor itself.
168

169 **Fig. 3.** A) Maximum counts of bioluminescent sources, or 'blobs' following light stimulation at 100 m-
170 spaced depth levels at a station located 6 km N of Mborokua Island, Solomon Islands at approx. 22:00-
171 23:30 local time. B) Representative images of maximum bioluminescent response at three depth
172 stations shown in (A).
173
174

Fig. 4. Maximum image intensities, reported in arbitrary units (AU) vs. time, for whole image light-stimulated responses at a) 100 m, b) 200 m, and c) 400 m depths recorded from a manned submersible in the Solomon Islands in September 2013. Vertical red dashed lines indicate strobe events.

Full-frame image sequences recorded at 30fps demonstrate the system's ability to track light production on free-swimming pelagic animals (Figure 5), offering the potential to gain new insight into the pathways that mediate bioluminescence in multicellular organisms. Two recordings of medusa show a swift response from swimming bell to tentacles, indicating the use of eyespots to sense light and a sequential reaction that dissipates quickly. In a spectacular video of a *Tomopteris* sp. pelagic polychaete, ([Video Link](#)), the organism displayed an elongated, >2.5 s luminescent reaction initiating in the head and traveling outward to the parapodia. Several recordings of bioluminescence in *Cestum veneris*, the 'venus girdle' ctenophore, illustrate light traveling through the animal in a single wave (not shown). Many other patterns were observed, but are not presented due to the absence of positive identification of the organism.

Fig. 5: Sample image sequences of *in situ* bioluminescence responses recorded at 30 fps. Time between frames is indicated in milliseconds in the lower left of each image. Top and middle row: tentacles from a hydromedusa show a distinct wave of bioluminescence traveling from the bell outwards and dissipating completely within 0.2 s, recorded at 700 m and 900 m depth (respectively) off the New England shelf. Bottom row: a *Tomopteris* sp. pelagic worm at 850 m in the Galapagos Islands, which exhibited a bright response for a total time of ~2.5 s.

4. Discussion

The three different deployment setups described in this paper allow for a comparison of methodology and recommendations for future research. The manned submersible was by far the most stable platform for observing midwater bioluminescence, and offered the unique opportunity to witness bioluminescent responses with the human eye. Phylum-level identification of several bioluminescent animals was possible, an effort that may be improved upon through repetitive application of this method. The use of a towed body with low-bandwidth telemetry produced poor results that were largely unusable, mostly due to the unstable nature of this setup. The use of a heave-compensated winch would likely absolve this issue, at least in moderate seas. ROV-based observations worked very well due to the stable nature of the decoupled platform and the high-bandwidth telemetry offered by the system. However, in both the manned submersible and ROV, the need to switch off all internal and external lighting hindered all other concurrent observations. While

all of the deployment methods used in this study are viable for further development efforts, the authors envision using an autonomous vehicle to conduct light-stimulated bioluminescence surveys in high spatial and temporal resolution.

The video-based vertical profile of bioluminescence presented in Figure 3 matches previous measurements of deep-sea zooplankton distribution (Banse 1964, Weikert 1982, Sameoto 1986, Kosobokova and Hirche 2000), roughly characterized by a maximum at 300-500 m followed by an exponential decline beyond depths >1000 m. Our observed maxima of bioluminescence at 400m may also represent a vertically migrating population of zooplankton as this dive was conducted in the early evening, just after sunset. Vertically migrating populations of major planktonic groups are known to rise from depths exceeding 700m in the daytime to near-surface waters at night (Ringelberg 2009); repeated vertical casts using this method over a 24-h period might produce results similar to that shown using net tows (Nishikawa and Tsuda 2001), video plankton recorders (Ashjian et al. 2001), and acoustic analysis (Kringel et al. 2003). One other study has observed the vertical migration of plankton using bioluminescence as an indicator (Widder et al. 1992), and Gillibrand et al. (2007) report an extensive bioluminescent layer at ~1500 m occurring seasonally in the NE Atlantic Ocean. Depressed bioluminescence responses approaching the seafloor corroborate similar observations in the Bahamas (Johnsen et al. 2012) and on the Mid-Atlantic Ridge (Craig et al. 2015).

In strict terms, these data represent the light-responsive bioluminescence community that exists in the water column. Because of the widespread prevalence of bioluminescence in deep-sea midwater species (with the exception of one major phylum, Tunicata), we therefore assert that the number of bioluminescent sources may be a proxy of total living biomass, at least among animals that are large enough to be imaged by the camera. The same line of comparison has been drawn with instruments using physical stimulation to elicit a bioluminescence response (e.g. Widder 2002, Herren et al. 2005, Heger et al. 2008, Craig et al. 2015).

Varying patterns of response to individual stimulus events may represent the diversity of the bioluminescent community itself, a corollary also drawn by Dominjon et al. (2012) and Craig et al. (2015). We envision developing a library of bioluminescent response profiles based on intensity vs. time and shape, paired with animal identification, which could allow for automated classification based solely on light production measured passively using a camera system. Applying such a method spatially and temporally would address a major technical challenge in deep-sea biological oceanography that has resulted in a dearth of zooplankton distribution data (Wiebe and Benfield 2003, Webb et al. 2010, Sutton 2013). It may also produce the resolution needed to resolve zooplankton population patchiness at critical time and space scales (Benoit-Bird et al. 2013, Benoit-Bird and McManus 2014).

The utilization of light stimulation to produce a bioluminescent response offers a unique advantage over mechanical, electrical and chemical stimulation techniques. In dinoflagellates, mechanical stimulation results in the deformation of cell membranes, initiating a cascade of reactions within the unicellular organism (Latz and Rohr 1999, Cussatlegras and Le Gal 2005). Larger animals, such as gelatinous zooplankton and fishes, may also exhibit a bioluminescent response based on the location and magnitude of the physical and/or electrical disturbance (e.g. Barnes and Case 1974, Herring and Widder 2004). Chemical stimulation is typically achieved by injecting a solution into laboratory aquaria, resulting in a widespread stimulus unlikely to be encountered in a natural setting. As such, luminescent responses in relation to an organism's neural/sensory network are poorly understood. Light stimulation induces bioluminescence with no mechanical or chemical invasion, and may be the least intrusive method for eliciting a 'natural' response from animals *in situ*.

The high-speed global shutter capability of the sCMOS camera offers a new window into the kinematics of bioluminescence. Our observation of the *Tomopteris* sp. polychaete supports the conclusions of Gouveneaux and Mallefet (2013) who conducted laboratory experiments on *T. helgolandica* and witnessed neural control of bioluminescence under contrived conditions. In several

recordings of siphonophores, most of the organism becomes luminescent at the same time, which correlates with previous work on a siphonophore species that exhibited no coordinated response to mechanical stimulation (Freeman 1987). Without a size scale, velocities cannot be strictly calculated, but the image sequences presented here illustrate the potential of the system for future research in this area. Several methods could be employed to resolve scale such as utilizing stereo cameras (e.g., Rife and Rock 2001), scaling lasers, or simply constraining the volume being imaged to minimize the depth of field. The stereo method is particularly attractive, since it would allow volume-pixel regions or 'voxels' to be constrained and characterized.

A comprehensive review of published literature presenting *in situ* observations of bioluminescence is given in Table 1. While the human eye can still outperform the best digital cameras in terms of sensitivity, dynamic range and resolution, and PMTs have long been able to sense single photons, the inability to record imagery is an obvious drawback to using these methods. The electron-bombarded CMOS camera used by Dominjon et al. (2012), while likely the most sensitive instrument ever used to record bioluminescence *in situ*, does not collect focused imagery and is therefore categorized as similar to a PMT device. The defining feature of sCMOS technology is achieving high quantum efficiency (QE) and low readout noise without electron multiplication or image intensification. This distinction makes direct comparison of sCMOS cameras to electron multiplier CCD cameras (EMCCD) and image intensified CCD cameras (I²CCD) not straightforward, and best quantified by empirical studies (e.g. Fullerton et al. 2012, Beier and Ibey 2014). While contemporary EMCCD cameras still report higher quantum efficiencies (e.g. Andor iXon 2015 series at >90% QE) compared to the most advanced sCMOS cameras (e.g. Hamamatsu ORCA-Flash4.0 2015 model at >82% QE), the electron multiplication process can dramatically reduce the effective QE of EMCCD cameras (Long et al. 2012). Given the upward trajectory in sCMOS QE performance combined with their characteristic high speed and resolution capability, sCMOS cameras have the potential to become the benchmark of bioluminescence imaging technology.

Regardless of the imaging sensor employed, most *in situ* bioluminescence research has relied exclusively on mechanical stimulation to elicit a response. For example, Herren et al. (2005), Moline et al. (2009), and Johnsen et al. (2014) used essentially the same excitation chamber/PMT instrument; Widder (1989), Widder and Johnsen (2000), Widder et al. (2005), and Johnsen et al. (2012) used similar 'splat-screen' approaches; and Priede et al. (2006), Heger et al. (2008), Craig et al. (2010), Craig et al. (2011), and Craig et al. (2015) use the 'splat-screen' method for their vertical profiling work. By far the most comprehensive investigation into light-stimulated bioluminescence was conducted by Neshyba (1967), who measured responses using PMTs. Notably, Buskey and Swift (1985) performed a series of light-stimulation experiments on copepods and euphausiids, but these were conducted in a laboratory environment. Lapota and Losee (1984) conducted a vertical dive in a manned submersible similar to that in this study, but bioluminescence was not recorded beyond written observation. Therefore, we believe these data presented herein represent the only published imagery of the phenomenon *in situ*.

303
304

Table 1
Reference list of studies observing bioluminescence *in situ*

	Sensor	Citation	Depth (m)	Observation platform
<i>Mechanical stimulation</i>	Human observer	Beebe 1934*	0-660	Bathyscape
		Galt et al. 1985	Surface water	SCUBA diver
	Photomultiplier tube	Boden & Kampa 1957	0-300	Vertical profiler
		Hardy & Kay 1964	62	Vertical profiler
		Bradner et al. 1987	4300	Vertical profiler
		Webster et al. 1991	4500	Vertical profiler
		Amram et al. 1999	~2350	Moored array
		Widder et al. 1999	0-250	Vertical profiler
		Herren et al. 2005	0-30	Muiltplatform
		Moline et al. 2009	0-40	AUV
		Tamburini et al. 2013*	~2350	Moored array
		Johnsen et al. 2014	20	Vertical profiler
	ISIT camera	Widder et al. 1989	0-600	Submersible
		Robison 1992	n/a	Submersible
		Widder & Johnsen 2000	0-200	Submersible
		Widder et al. 2005	600	ROV, benthic lander
		Priede et al. 2006	0-4000	Vertical profiler
		Gillibrand et al. 2007	0-4800	Vertical profiler
		Heger et al. 2008	0-3000	Vertical profiler
		Craig et al. 2010	0-5000	Vertical profiler
	I ² CCD (image-intensified CCD)	Craig et al. 2011	1500-2700	Benthic lander, ROV
		Craig et al. 2015	0-2500	Vertical profiler
		Johnsen et al. 2012*	1000	Submersible
	ebCMOS camera (electro-bombarded CMOS)	Dominjon et al. 2012	2500	Moored array
<i>Light stimulation</i>	Human observer	Lapota & Losee 1986	0-600	Submersible
	Photomultiplier tube	Neshyba 1967	700	Profiler

* These studies did not actively stimulate bioluminescence. Their results may result from naturally occurring mechanical stimulation (i.e. currents causing collisions) or may be from animals actively producing bioluminescence with no external stimulation.

5. Conclusions

Marine bioluminescence research has resulted in transformative applications to a wide range of medical, military and industrial disciplines (e.g. Shimomura et al. 1962), and the potential for important new discoveries is high. The diversity of light-producing organisms is remarkable, and observed in every ocean from shallow to deep seas. Despite the ubiquity of bioluminescence among marine taxa, the phenomenon is categorically underexplored and limited by the technology available to observe it *in situ*. To address this, an sCMOS camera was deployed on multiple oceanographic platforms in three different oceanic regions to explore bioluminescence *in situ*. Using light stimulation to elicit responses, counts of bioluminescent sources were compared against water depth, demonstrating a potential new method to assess midwater zooplankton populations. High-resolution, quantitative measurements of light intensity responses were observed to vary greatly among depth zones and individual organisms. The kinematics of bioluminescence was imaged among several animals at a high frame rate and resolution, presenting new visualizations of how marine animals produce light. While this study demonstrates the utility of light stimulation for eliciting bioluminescent responses, it also highlights the current lack of scientific knowledge on the subject; further work is needed to identify which organisms respond to light stimulation and varying light intensities, patterns, and wavelengths. Future work should focus on pairing spatial and temporal observations of light-stimulated bioluminescence with traditional-method biomass assessments, in a true effort to estimate total living biomass. Advanced control software will lead towards support onboard autonomous vehicles and other self-powered platforms, and may allow for bioluminescent surveys to be conducted in high spatial and temporal resolution. We conclude that sCMOS cameras paired with light stimulation can offer a wide range of new applications in bioluminescence research.

Acknowledgements

The authors wish to acknowledge the crew and technical team of the M/Y *Alucia*, the E/V *Nautilus* (leg NA064), and the R/V *Endeavor* (leg EN564). We are grateful for the engineering and fabrication support of The John B. Pierce Laboratory and the generous technical support provided by David Casagrande, Donald Dansereau, John Buckley, Tom Delessandro, Andrew Wilkens and Ron Goodman. This work was funded in part by the Ray and Barbara Dalio Family Foundation, the Rhode Island Endeavor Program, the Ocean Exploration Trust, and NSF grant DEB-1257555 to JSS. We thank Igor Belkin and 6 anonymous reviewers who contributed greatly to this manuscript with critical and constructive commentary.

References

- Adrián-Martínez, S., Albert, A., André, M., Anghinolfi, M., Anton, G., Ardid, M.S., Aubert, J.-J., Baret, B., JBarrios-Martí, J., Basa, S., Bertin, V., Biagi, S., Bogazzi, C., Bormuth, R., Bou-Cabo, M., Bouwhuis, M.C., Bruijn, R., Brunner, J., Busto, J., Capone, A., Caramete, L., Cârloganu, C., Carr, J., Chiarusi, T., Circella, M., Core, L., Costantini, H., Coyle, P., Creusot, A., Curtil, C., De Rosa, G., Dekeyser, I., Deschamps, A., De Bonis, G., Distefano, C., Donzaud, C., Dornic, D., Dorosti, Q., Drouhin, D., Dumas, A., Eberl, T., Elsässer, D., Enzenhöfer, A., Escoffier, S., Fehn, K., Felis, I., Fermani, P., Folger, F., Fusco, L.A., Galatà, S., Gay, P., Geißelsöder, S., Geyer, K., Giordano, V., Gleixner, A., Gómez-González, J.P., Graf, K., Guillard, G., van Haren, H., Heijboer, A.J., Hello, Y., Hernández-Rey, J.J., Herold, B., Herrero, A., Höß, J., Hofestädt, J., James, C.W., de Jong, M., Kadler, M., Kalekin, O., Katz, U., Kießling, D., Kooijman, P., Kouchner, A., Kreykenbohm, I., Kulikovskiy, V., Lahmann, R., Lambard, E., Lambard, G., Lattuada, D., Lefèvre, D., Leonora, E., Loehner, H., Loucatos, S., Mangano, S., Marcelin, M., Margiotta, A., Martínez-Mora, J.A., Martini, S., Mathieu, A., Michael, T., Migliozi, P., Mueller, C., Neff, M., Nezri, E., Palioselitis, D., Păvălaș, G.E., Perrina, C., Piattelli, P., Popa, V., Pradier, T., Racca, C., Riccobene, G., Richter, R., Roensch, K., Rostovtsev, A., Saldaña, M., Samtleben, D.F.E., Sánchez-Losa, A., Sanguineti, M., Sapienza, P., Schmid, J., Schnabel, J., Schulte, S., Schüßler, F., Seitz, T., Sieger, C., Spies, A., Spurio, M., Steijger, J.J.M., Stolarczyk, T., Taiuti, M., Tamburini, C., Tayalati, Y., Trovato, A., Vallage, B., Vallée, C., Van Elewyck, V., Visser, E., Vivolo, D., Wagner, S., Wilms, J., de Wolf, E., Yarkin, K., Yepes, H., Zornoza, J.D., and Zúñiga, J. (2014). Searches for point-like and extended neutrino sources close to the galactic center using the ANTARES neutrino telescope. *The Astrophysical Journal Letters*, 786(1), L5. [doi:10.1088/2041-8205/786/1/L5](https://doi.org/10.1088/2041-8205/786/1/L5)
- Ashjian, C. J., Davis, C. S., Gallager, S. M., Alatalo, P. (2001). Distribution of plankton, particles, and hydrographic features across Georges Bank described using the Video Plankton Recorder. *Deep Sea Research Part II: Topical Studies in Oceanography*, 48(1), 245-282. [doi:10.1016/S0967-0645\(00\)00121-1](https://doi.org/10.1016/S0967-0645(00)00121-1)
- Banse, K. (1964). On the vertical distribution of zooplankton in the sea. *Progress in Oceanography*, 2, 53-125. [doi:10.1016/0079-6611\(64\)90003-5](https://doi.org/10.1016/0079-6611(64)90003-5)
- Barnes, A. T., Case, J. F. (1974). The luminescence of lanternfish (Myctophidae): spontaneous activity and responses to mechanical, electrical, and chemical stimulation. *Journal of Experimental Marine Biology and Ecology*, 15(2), 203-221. [doi:10.1016/0022-0981\(74\)90046-X](https://doi.org/10.1016/0022-0981(74)90046-X)
- Bassler BL, Wright M, Showalter RE, Silverman MR. 1993. Intercellular signalling in *Vibrio harveyi*: sequence and function of genes regulating expression of luminescence. *Molecular Microbiology* 9 (4): 773–786. [doi:10.1111/j.1365-2958.1993.tb01737.x](https://doi.org/10.1111/j.1365-2958.1993.tb01737.x)
- Beier HT, Ibey BL (2014) Experimental Comparison of the High-Speed Imaging Performance of an EM-CCD and sCMOS Camera in a Dynamic Live-Cell Imaging Test Case. *PLoS ONE* 9(1): e84614. [doi:10.1371/journal.pone.0084614](https://doi.org/10.1371/journal.pone.0084614)
- Benoit-Bird, K. J., Shroyer, E. L., & McManus, M. A. (2013). A critical scale in plankton aggregations across coastal ecosystems. *Geophysical Research Letters*, 40(15), 3968-3974. [doi: 10.1002/grl.50747](https://doi.org/10.1002/grl.50747)
- Benoit-Bird, K.J., McManus, M.A. (2014) A critical time window for organismal interactions in a pelagic ecosystem. *PLoS ONE*, 9(5): e97763. [doi:10.1371/journal.pone.0097763](https://doi.org/10.1371/journal.pone.0097763)
- Boden, B.P., Kampa, E.M. (1957) Records of Bioluminescence in the Ocean. *Pacific Science*, Vol. 11(2) <http://hdl.handle.net/10125/8656>
- Bradner, H., Bartlett, M., Blackinton, G., Clem, J., Karl, D., Learned, Lewitus, A., Matsuno, S., O'Connor, D., peatman, W., Reichle, M., Roose, C., Water, J., Webster, M., Yarbrough, M.(1987). Bioluminescence profile in the deep Pacific Ocean. *Deep Sea Research Part A. Oceanographic Research Papers*, 34(11), 1831-1840. [doi:10.1016/0198-0149\(87\)90057-4](https://doi.org/10.1016/0198-0149(87)90057-4)
- Buskey, E. J., & Swift, E. (1985). Behavioral responses of oceanic zooplankton to simulated bioluminescence. *The Biological Bulletin*, 168(2), 263-275. [doi: 10.2307/1541239](https://doi.org/10.2307/1541239)

395 Campbell, A. K., and Herring, P. J. 1990. Imidazolopyrazine bioluminescence in copepods and other
 396 marine organisms. *Mar. Biol.* 104: 219-225. [doi:10.1007/BF01313261](https://doi.org/10.1007/BF01313261)

397 Coates, C., Fowler, B., & Holst, G. (2009). sCMOS-Scientific CMOS technology: a high-performance
 398 imaging breakthrough. White paper, Andor Technology, Fairchild Imaging and PCO AG, 171.

399 Contag, C. H., & Bachmann, M. H. (2002). Advances in in vivo bioluminescence imaging of gene
 400 expression. *Annual review of biomedical engineering*, 4(1), 235-260. [doi:](https://doi.org/10.1146/annurev.bioeng.4.111901.093336)
 401 [10.1146/annurev.bioeng.4.111901.093336](https://doi.org/10.1146/annurev.bioeng.4.111901.093336)

402 Craig, J., Jamieson, A. J., Hutson, R., Zuur, A. F., Priede, I. G. (2010). Factors influencing the abundance
 403 of deep pelagic bioluminescent zooplankton in the Mediterranean Sea. *Deep Sea Research Part I:*
 404 *Oceanographic Research Papers*, 57(11), 1474-1484. [doi:10.1016/j.dsr.2010.08.005](https://doi.org/10.1016/j.dsr.2010.08.005)

405 Craig, J., Jamieson, A. J., Bagley, P. M., Priede, I. G. (2011). Naturally occurring bioluminescence on the
 406 deep-sea floor. *Journal of Marine Systems*, 88(4), 563-567. [doi:10.1016/j.jmarsys.2011.07.006](https://doi.org/10.1016/j.jmarsys.2011.07.006)

407 Craig, J., Youngbluth, M., Jamieson, A. J., Priede, I. G. (2015). Near seafloor bioluminescence,
 408 macrozooplankton and macroparticles at the mid-Atlantic ridge. *Deep Sea Research Part I:*
 409 *Oceanographic Research Papers*. [doi:10.1016/j.dsr.2014.12.011](https://doi.org/10.1016/j.dsr.2014.12.011)

410 Cussatlegras, A. S., & Gal, P. L. (2005). Dinoflagellate bioluminescence in response to mechanical
 411 stimuli in water flows. *Nonlinear Processes in Geophysics*, 12(3), 337-343. [doi:10.5194/npg-12-337-](https://doi.org/10.5194/npg-12-337-2005)
 412 [2005](https://doi.org/10.5194/npg-12-337-2005)

413 Dominjon, A., Ageron, M., Barbier, R., Billault, M., Brunner, J., Cajgfinger, T., ... & Guérin, C. (2012). An
 414 ebCMOS camera system for marine bioluminescence observation: The LuSEApher prototype.
 415 *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers,*
 416 *Detectors and Associated Equipment*, 695, 172-178. [doi:10.1016/j.nima.2011.11.048](https://doi.org/10.1016/j.nima.2011.11.048)

417 Freeman, G. (1987). Localization of bioluminescence in the siphonophore *Nanomia cara*. *Marine*
 418 *Biology*, 93(4), 535-541. [doi: 10.1007/BF00392791](https://doi.org/10.1007/BF00392791)

419 Fowler, B., Liu, C., Mims, S., Balicki, J., Li, W., Do, H., & Vu, P. (2009, May). Low-light-level CMOS image
 420 sensor for digitally fused night vision systems. In *SPIE Defense, Security, and Sensing* (pp. 72981D-
 421 72981D). International Society for Optics and Photonics. [doi:10.1117/12.818053](https://doi.org/10.1117/12.818053)

422 Fucile, P. D. (2002, October). An expendable bioluminescence measuring bathy-photometer. In
 423 *OCEANS'02 MTS/IEEE* (Vol. 3, pp. 1716-1721). IEEE. [doi:10.1109/OCEANS.2002.1191892](https://doi.org/10.1109/OCEANS.2002.1191892)

424 Fullerton, S.; Bennett, K.; Toda, E.; Takahashi, T. (2012) ORCA-flash4.0: Changing the Game.
 425 *Hamamatsu Photonics Whitepaper*. [Permalink to document](#).

426 Galt, C. P., Grober, M. S., & Sykes, P. F. (1985). Taxonomic correlates of bioluminescence among
 427 appendicularians (Urochordata: Larvacea). *The Biological Bulletin*, 168(1), 125-134. [Article link](#)

428 Gillibrand, E. J. V., Jamieson, A. J., Bagley, P. M., Zuur, A. F., & Priede, I. G. (2007). Seasonal development
 429 of a deep pelagic bioluminescent layer in the temperate NE Atlantic Ocean. *Marine Ecology Progress*
 430 *Series*, 341, 37-44. [doi:10.3354/meps341037](https://doi.org/10.3354/meps341037)

431 Gouveneaux, A., Mallefet, J. (2013). Physiological control of bioluminescence in a deep-sea planktonic
 432 worm, *Tomopteris helgolandica*. *The Journal of Experimental biology*, 216(22), 4285-4289.
 433 [doi: 10.1242/jeb.090852](https://doi.org/10.1242/jeb.090852)

434 Green AA, McElroy WD. Crystalline firefly luciferase. *Biochimica et biophysica acta*. 1956 20(1):170-
 435 176. [doi:10.1016/0006-3002\(56\)90275-X](https://doi.org/10.1016/0006-3002(56)90275-X)

436 Haddock, S. H. D., & Case, J. F. (1999). Bioluminescence spectra of shallow and deep-sea gelatinous
 437 zooplankton: ctenophores, medusae and siphonophores. *Marine Biology*, 133(3), 571-582. [doi:](https://doi.org/10.1007/s002270050497)
 438 [10.1007/s002270050497](https://doi.org/10.1007/s002270050497)

439 Haddock, S. H., Moline, M. A., Case, J. F. (2010). Bioluminescence in the sea. *Annual Review of Marine*
 440 *Science*, 2: 443-493 [doi:10.1146/annurev-marine-120308-081028](https://doi.org/10.1146/annurev-marine-120308-081028)

441 Heger, A., Ieno, E. N., King, N. J., Morris, K. J., Bagley, P. M., Priede, I. G. (2008). Deep-sea pelagic
 442 bioluminescence over the Mid-Atlantic Ridge. *Deep Sea Research Part II: Topical Studies in*
 443 *Oceanography*, 55(1), 126-136. [doi:10.1016/j.dsr2.2007.09.014](https://doi.org/10.1016/j.dsr2.2007.09.014)

Herren, C. M., Haddock, S. H., Johnson, C., Orrico, C. M., Moline, M. A., Case, J. F. (2005). A multi-platform bathyphotometer for fine-scale, coastal bioluminescence research. *Limnology and Oceanography: Methods*, 3(5), 247-262. [doi:10.4319/lom.2005.3.247](https://doi.org/10.4319/lom.2005.3.247)

Herring, P. J., Widder, E. A. (2004). Bioluminescence of deep-sea coronate medusae (Cnidaria: Scyphozoa). *Marine Biology*, 146(1), 39-51. [doi:10.1007/s00227-004-1430-7](https://doi.org/10.1007/s00227-004-1430-7)

Johnsen, S., Frank, T. M., Haddock, S. H., Widder, E. A., Messing, C. G. (2012). Light and vision in the deep-sea benthos: I. Bioluminescence at 500–1000 m depth in the Bahamian Islands. *The Journal of Experimental Biology*, 215(19), 3335-3343. [doi: 10.1242/jeb.072009](https://doi.org/10.1242/jeb.072009)

Johnsen, G., Candeloro, M., Berge, J., Moline, M. (2014). Glowing in the dark: discriminating patterns of bioluminescence from different taxa during the Arctic polar night. *Polar Biology*, 37(5), 707-713. [doi: 10.1007/s00300-014-1471-4](https://doi.org/10.1007/s00300-014-1471-4)

Kosobokova, K., Hirche, H. J. (2000). Zooplankton distribution across the Lomonosov Ridge, Arctic Ocean: species inventory, biomass and vertical structure. *Deep Sea Research Part I: Oceanographic Research Papers*, 47(11), 2029-2060. [doi:10.1016/S0967-0637\(00\)00015-7](https://doi.org/10.1016/S0967-0637(00)00015-7)

Kringel, K., Jumars, P. A., & Holliday, D. V. (2003). A shallow scattering layer: High-resolution acoustic analysis of nocturnal vertical migration from the seabed. *Limnology and Oceanography*, 48(3), 1223-1234. [doi:10.4319/lo.2003.48.3.1223](https://doi.org/10.4319/lo.2003.48.3.1223)

Lapota, D., Losee, J. R. (1984). Observations of bioluminescence in marine plankton from the Sea of Cortez. *Journal of Experimental Marine Biology and Ecology*, 77(3), 209-239. [doi:10.1016/0022-0981\(84\)90121-7](https://doi.org/10.1016/0022-0981(84)90121-7)

Latz, M. I., & Rohr, J. (1999). Luminescent response of the red tide dinoflagellates *Lingulodinium polyedrum* to laminar and turbulent flow. *Limnology and Oceanography*, 44(6), 1423-1435. [doi: 10.4319/lo.1999.44.6.1423](https://doi.org/10.4319/lo.1999.44.6.1423)

Long, F., Zeng, S., & Huang, Z. L. (2012). Localization-based super-resolution microscopy with an sCMOS camera Part II: Experimental methodology for comparing sCMOS with EMCCD cameras. *Optics express*, 20(16), 17741-17759. [doi:10.1364/OE.20.017741](https://doi.org/10.1364/OE.20.017741)

Lucas, C. H., Jones, D. O., Hollyhead, C. J., Condon, R. H., Duarte, C. M., Graham, W. M., Robinson, K.L, Pitt, K.A., Schildhauer, M., Regetz, J. (2014). Gelatinous zooplankton biomass in the global oceans: geographic variation and environmental drivers. *Global Ecology and Biogeography*, 23(7), 701-714. [doi: 10.1111/geb.12169](https://doi.org/10.1111/geb.12169)

Neshyba, S. (1967). Pulsed light stimulation of marine bioluminescence *in situ*. *Limnology and Oceanography*, 12(2), 222-235. [doi:10.4319/lo.1967.12.2.0222](https://doi.org/10.4319/lo.1967.12.2.0222)

Nishikawa, J., Tsuda, A. (2001). Diel vertical migration of the tunicate *Salpa thompsoni* in the Southern Ocean during summer. *Polar Biology*, 24(4), 299-302. [doi:10.1007/s003000100227](https://doi.org/10.1007/s003000100227)

Martin, J. R. (2008). In vivo brain imaging: fluorescence or bioluminescence, which to choose?. *Journal of neurogenetics*, 22(3), 285-307. [doi: 10.1080/01677060802298517](https://doi.org/10.1080/01677060802298517)

Moline, M. A., Blackwell, S. M., Case, J. F., Haddock, S. H., Herren, C. M., Orrico, C. M., Terrill, E. (2009). Bioluminescence to reveal structure and interaction of coastal planktonic communities. *Deep Sea Research Part II: Topical Studies in Oceanography*, 56(3), 232-245. [doi:10.1016/j.dsr2.2008.08.002](https://doi.org/10.1016/j.dsr2.2008.08.002)

Prasher, D., McCann, R.O., Cormier, M.J. (1985). Cloning and expression of the cDNA coding for aequorin, a bioluminescent calcium-binding protein. *Biochemical and Biophysical Research Communications*, 126(3):1259-1268. [doi:10.1016/0006-291X\(85\)90321-3](https://doi.org/10.1016/0006-291X(85)90321-3)

Prasher, D. C., Eckenrode, V. K., Ward, W. W., Prendergast, F. G., & Cormier, M. J. (1992). Primary structure of the *Aequorea victoria* green-fluorescent protein. *Gene*, 111(2), 229-233. [doi:10.1016/0378-1119\(92\)90691-H](https://doi.org/10.1016/0378-1119(92)90691-H)

Priede, I. G., Bagley, P. M., Way, S., Herring, P. J., & Partridge, J. C. (2006). Bioluminescence in the deep sea: free-fall lander observations in the Atlantic Ocean off Cape Verde. *Deep Sea Research Part I: Oceanographic Research Papers*, 53(7), 1272-1283. [doi:10.1016/j.dsr.2006.05.004](https://doi.org/10.1016/j.dsr.2006.05.004)

- Rife, J., Rock, S. M. (2001). A pilot-aid for ROV based tracking of gelatinous animals in the midwater. In OCEANS, 2001. MTS/IEEE Conference and Exhibition (Vol. 2, pp. 1137-1144). IEEE. [doi:10.1109/OCEANS.2001.968274](https://doi.org/10.1109/OCEANS.2001.968274)
- Ringelberg, J. (2009). Diel vertical migration of zooplankton in lakes and oceans: causal explanations and adaptive significances. Springer Science & Business Media. [doi:10.1007/978-90-481-3093-1](https://doi.org/10.1007/978-90-481-3093-1)
- Robison, B. H. (1992). Bioluminescence in the benthopelagic holothurian *Enypniasteseximia*. Journal of the Marine Biological Association of the United Kingdom, 72(02), 463-472. [doi:10.1017/S0025315400037826](https://doi.org/10.1017/S0025315400037826)
- Sameoto, D. D. (1986). Influence of the biological and physical environment on the vertical distribution of mesozooplankton and micronekton in the eastern tropical Pacific. Marine Biology, 93(2), 263-279. [doi:10.1007/BF00508264](https://doi.org/10.1007/BF00508264)
- Shimomura O, Johnson FH, Saiga Y. (1962). Extraction, purification and properties of aequorin, a bioluminescent protein from the luminous hydromedusan, Aequorea. Journal of Cellular and Comparative Physiology 59 (3): 223-39. [doi: 10.1002/jcp.1030590302](https://doi.org/10.1002/jcp.1030590302)
- Surette MG, Miller MB, Bassler BL. 1999. Quorum sensing in *Escherichia coli*, *Salmonella typhimurium*, and *Vibrio harveyi*: a new family of genes responsible for autoinducer production. Proceedings of the National Academy of Sciences 96 (4): 1639-44. [doi:10.1073/pnas.96.4.1639](https://doi.org/10.1073/pnas.96.4.1639)
- Sutton, T. T. (2013). Vertical ecology of the pelagic ocean: classical patterns and new perspectives. Journal of Fish Biology, 83: 1508-1527. [doi: 10.1111/jfb.12263](https://doi.org/10.1111/jfb.12263)
- Tamburini, C., Canals, M., Durrieu de Madron, X., Houpert, L., Lefèvre, D., Martini, S., D'Ortenzio, F., Robert, A., Testor, P., Aguilar, J.A., Al Samarai, I., Albert, A., André, M., Anghinolfi, M., Anton, G., Anvar, S., Ardid, M., Assis Jesus, A.C, Astraatmadja, T.L., Aubert, J.-J., Baret, B., Basa, S., Bertin, V., Biagi, S., Bigi, A., Bigongiari, C., Bogazzi, C., Bou-Cabo, M., Bouhou, B., Bouwhuis, M.C., Brunner, J., Busto, J., Camarena, F., Capone, A., Cârloganu, C., Carminati, G., Carr, J., Cecchini, S., Charif, Z., Charvis, P., Chiarusi, T., Circella, M., Coniglione, R., Costantini, H., Coyle, P., Curtil, C., Decowski, P., Dekeyser, I., Deschamps, A., Donzaud, C., Dornic, D., Dorosti, H.Q., Drouhin, D., Eberl, T., Emanuele, U., Ernenwein, J.-P., Escoffier, S., Fermani, P., Ferri, M., Flaminio, V., Folger, F., Fritsch, U., Fuda, J.-L., Galatà, S., Gay, P., Giacomelli, G., Giordano, V., Gómez-González, J.-P., Graf, K., Guillard, G., Halladjian, G., Hallewell, G., Van Haren, H., Hartman, J., Heijboer, A.J., Hello, Y., Hernández-Rey, J.J., Herold, B., Hößl, J., Hsu, C.-C., De Jong, M., Kadler, M., Kalekin, O., Kappes, A., Katz, U., Kavatsyuk, O., Kooijman, P., Kopper, C., Kouchner, A., Kreykenbohm, I., Kulikovskiy, V., Lahmann, R., Lamare, P., Larosa, G., Lattuada, D., Lim, G., Lo Presti, D., Loehner, H., Loucatos, S., Mangano, S., Marcelin, M., Margiotta, A., Martinez-Mora, J.A., Meli, A., Montaruli, T., Moscoso, L., Motz, H., Neff, M., Nezri, E., Palioselitis, D., Păvălaș, G.E., Payet, K., Payre, P., Petrovic, J., Piattelli, P., Picot-Clemente, N., Popa, V., Pradier, T., Presani, E., Racca, C., Reed, C., Riccobene, G., Richardt, C., Richter, R., Rivière, C., Roensch, K., Rostovtsev, A., Ruiz-Rivas, J., Rujoiu, M., Russo, V.G., Salesa, F., Sánchez-Losa, A., Sapienza, P., Schöck, F., Schuller, J.-P., Schussler, F., Shanidze, R., Simeone, F., Spies, A., Spurio, M., Steijger, J.J.M., Stolarczyk, T., Taiuti, M.G.F., Toscano, S., Vallage, B., Van Elewyck, V., Vannoni, G., Vecchi, M., Vernin, P., Wijnker, G., Wilms, J., De Wolf, E., Yepes, H., Zaborov, D., Zornoza, J. De D., Zúñiga, J. (2013). Deep-sea bioluminescence blooms after dense water formation at the ocean surface. PLoS ONE, 8(7), 1-10. [doi: 10.1371/journal.pone.0067523](https://doi.org/10.1371/journal.pone.0067523)
- Webb, T. J., Vanden Berghe, E., O'Dor, R. (2010). Biodiversity's big wet secret: The global distribution of marine biological records reveals chronic under-exploration of the deep pelagic ocean. PLoS One, 5(8), e10223. [doi:10.1371/journal.pone.0010223](https://doi.org/10.1371/journal.pone.0010223)
- Webster, M. S., Roos, C. E., Roberts, A., Okada, A., Ohashi, Y., O'Connor, D., Mitiguy, R., Matsuno, S., March, R., Learned, J.G., Karl, D., Clem, J., Blackinton, G., Bradner, H., Babson, J. (1991). Mechanical stimulation of bioluminescence in the deep Pacific Ocean. Deep Sea Research Part A. Oceanographic Research Papers, 38(2), 201-217. [doi:10.1016/0198-0149\(91\)90080-Y](https://doi.org/10.1016/0198-0149(91)90080-Y)

540 Weikert, H. (1982). The Vertical Distribution of Zooplankton in Relation to Habitat Zones in the Area
 541 of the Atlantis II Deep, Central Red Sea. Marine Ecology Progress Series 8(2), 129-143.
 542 [doi:10.3354/meps00812](https://doi.org/10.3354/meps00812)
 543 Widder, E. A., Bernstein, S. A., Bracher, D. F., Case, J. F., Reisenbichler, K. R., Torres, J. J., Robison, B. H.
 544 (1989). Bioluminescence in the Monterey Submarine Canyon: image analysis of video recordings
 545 from a midwater submersible. Marine Biology, 100(4), 541-551. [doi:10.1007/BF00394831](https://doi.org/10.1007/BF00394831)
 546 Widder, E. A., Greene, C. H., Youngbluth, M. J. (1992). Bioluminescence of sound-scattering layers in
 547 the Gulf of Maine. Journal of plankton research, 14(11), 1607-1624. [doi:10.1093/plankt/14.11.1607](https://doi.org/10.1093/plankt/14.11.1607)
 548 Widder, E. A., Johnsen, S. (2000). 3D spatial point patterns of bioluminescent plankton: a map of the
 549 'minefield'. Journal of Plankton Research, 22(3), 409-420. [doi:10.1093/plankt/22.3.409](https://doi.org/10.1093/plankt/22.3.409)
 550 Widder, E. (2002). Bioluminescence and the pelagic visual environment. Marine and Freshwater
 551 Behaviour and Physiology, 35(1-2), 1-26. [doi:10.1080/10236240290025581](https://doi.org/10.1080/10236240290025581)
 552 Widder, E. A., Robison, B. H., Reisenbichler, K. R., Haddock, S. H. D. (2005). Using red light for *in situ*
 553 observations of deep-sea fishes. Deep Sea Research Part I: Oceanographic Research Papers, 52(11),
 554 2077-2085. [doi:10.1016/j.dsr.2005.06.007](https://doi.org/10.1016/j.dsr.2005.06.007)
 555 Wiebe, P.H., Benfield, M.C. (2003). From the Hensen net toward four-dimensional biological
 556 oceanography. Progress in Oceanography, 56(1), 7-136. [doi:10.1016/S0079-6611\(02\)00140-4](https://doi.org/10.1016/S0079-6611(02)00140-4)

Graphical Abstract