Title: The utility of protein and mRNA correlation

Authors: Samuel H. Payne

Address: Pacific Northwest National Laboratory, 902 Battelle Boulevard, Richland WA 99336

Corresponding author: Payne, S.H. (samuel.payne@pnnl.gov)

Keywords: Bioinformatics, Systems Biology, proteomics, transcriptomics

Abstract

Transcriptomic, proteomic and metabolomic measurements are revolutionizing the way we model and predict cellular behavior, and multi-omic comparisons are being published with increased regularity.

Some have expected a trivial and predictable correlation between mRNA and protein; however the manifest complexity of biological regulation suggests a more nuanced relationship. Indeed, observing this lack of strict correlation provides clues for new research topics, and has the potential for transformative biological insight.

Easy access to global –omic measurements

High throughput transcriptomic, proteomic and metabolomic measurements are revolutionizing the way we model and predict cellular behavior. Disruptive technologies are continually improving measurement speed, coverage, and accuracy of multi-omic data while simultaneously reducing costs. A popular avenue in the post-genomic era is the simultaneous interrogation of global abundance of protein and mRNA [1], [2], [3]. As these studies are becoming more common and the results are being repeated and corroborated on multiple technology platforms, the goals and utilities of such comparisons now need to be evaluated. The purpose of this Forum article is not to exhaustively summarize the literature. Rather, the aim is to highlight the types of experiments that produce scientifically useful conclusions. Specifically, the aim is to highlight the types of experiments which produce scientifically useful conclusions. It is clear from numerous reports that proteome and transcriptome abundances are not sufficiently correlated to act as proxies for each other. The majority of this difference is rooted in fundamental biological regulation, and not measurement bias or platform-specific error. Thus we should not wrest with the differences, but rather leverage them to elucidate the underlying phenomena. With this emerging access to global measurements of multiple –omes, what new biology can be explored?

Biological processes necessarily lead to complexity in abundance measurements

One of the first comparisons of mRNA and protein was performed by the Aebersold group in 1999 using *Saccharomyces cerevisiae* growing at mid-log phase [4]. This study was limited compared to today's technologies, with 106 genes examined. Their major conclusion succinctly stated that “[they] found that the correlation between mRNA and protein levels was insufficient to predict protein expression levels from quantitative mRNA data.” Over the past 15 years, technological innovation has drastically improved the breadth of both mRNA and protein measurements, yet this fundamental observation

continues to be widely, though not universally, supported [5, 6, 7]. An extensive review of the literature of protein/transcript comparisons can be found in Vogel and Marcotte [8].

Fundamental biological processes control the information flow from genome to gene-product to functional output (Figure 1). It is now recognized that biological systems will regulate processes by modification, binding, concentration, and/or localization of nearly any biological molecule. In particular, protein abundance is regulated by a variety of complex mechanisms. By measuring mRNA abundance, only the early steps in a long chain of regulatory events are observed. The following exemplary studies have used multi-omic data to identify and characterize a variety of regulatory mechanisms. Using a cohort of 95 diverse individuals from the HapMap project to identify genetic variation that affects protein abundance, the Snyder group discovered that the loci controlling RNA expression (eQTLs) had only a 50% overlap with the loci controlling protein expression (pQTLs) highlighting distinct genetic regulatory sequences [9]. By coupling high-throughput sequencing of ribosome-protected transcripts to RNA-Seq experiments, Brar et al. teased apart the abundance of a transcript from the use of a transcript. In this genome wide analysis, they show that translational regulation is pervasive [10]. MicroRNAs are an additional specific mechanism used by cells to regulate protein synthesis. A global analysis found that microRNAs can affect protein abundance either through mRNA destabilization, which decreases mRNA abundance, or through translational repression, which does not alter mRNA abundance [11]. Another important factor in the differences between mRNA and protein abundance is the distinct synthesis and decay rates. Not only are these relative rates on different scales (the lifetime for an mRNA is minutes, the lifetime for a protein is hours to years), but the rates of synthesis or decay for mRNA and protein from a single gene are unrelated [12]. Finally, in a detailed study of the *Prochlorococcus* cell cycle, the Chisholm group characterized cycling proteins and transcripts by both phase and amplitude [13]. They concluded that “significant divergence between mRNA and protein levels in the relative timing and/or magnitude of abundance oscillations are the rule rather than the exception.” Given these and numerous

1
2
3
4 other regulatory mechanisms, we should not expect an easy correlation between protein and mRNA
5
6 abundances.
7
8
9

10 **How to use multi-omic data for greatest insight?**

11

12
13 As quantification technologies improve in coverage, accuracy and cost, it will become increasingly
14
15 common to globally profile both protein and transcripts, which has great potential to elucidate novel
16
17 biology. It is clear, however, that transcript measurements do not orthogonally validate proteome
18
19 measurements and vice versa. In the utilitarian perspective, mRNA and protein abundance cannot
20
21 proxy for one another, just as protein abundances of enzymes are not appropriate proxies for their
22
23 enzymatic products. Given this knowledge and perspective, it is essential to consider the purpose for
24
25 multi-omics experiments before one embarks.
26
27
28
29

30
31 For many experiments, collecting both data types will be valuable. In generic experiment, one might ask
32
33 what the effects of a stimulus are. To address this question, proteomics could be used to monitor
34
35 phosphorylation signaling and dynamic cellular localization. Transcriptomics would elucidate the cohort
36
37 of genes up or down-regulated by activated transcription factors. Finally, proteomics would determine
38
39 which transcripts become protein at which time, allowing researchers to see a multi-staged response to
40
41 the stimulus, delineating between a rapid response and a long term adaptation. Carefully considered
42
43 hypotheses and experiments can use multi-omic data to drive biological insight. A large number of
44
45 clinical and synthetic biology experiments are empowered by the recent decrease in cost of data
46
47 generation and pursue genotype-to-phenotype hypotheses. Such experiments ask how changes in a
48
49 gene (e.g. mutation or deletion) affect cellular function. By their nature, these are multi-omic questions.
50
51 For clinical cohorts, these changes could be single nucleotide polymorphisms or more complex genomic
52
53 alterations. In synthetic biology, the focus is often on the insertion of new genes. The measurement of
54
55
56
57
58
59
60
61
62
63
64
65

cellular function varies according to the hypothesis, but can be done with proteomics, metabolomics, or clinical meta-data.

A unique perspective on regulatory networks can be gained from multi-omic integration. Copy number aberrations (CNA) measure the number of DNA segments encode a gene in the genome. The normal copy number for diploid organisms is two; however, in a population there are typically some individuals with more or fewer copies of any given gene. As the copy number of a gene changes, it is expected that the abundance of its cognate protein changes accordingly. In genetics, this is termed a *cis* effect. This change in the cognate protein's abundance may also affect the abundance of other interacting or downstream proteins, or a *trans* effect. In cancer CNAs are often much more common and extreme. In a recent analysis of 90 colon tumors, Zhang et al integrated proteomics data and genomic measurements of CNAs [14]. Through global correlation of these multi-omic data, they identified many CNAs affecting hundreds of proteins in *trans*. From this subset of CNAs, they illuminated new driver mutations for colon cancer phenotypes. This integrative analysis identifies unique relationships in a regulatory network distinct from co-expression analysis, which finds groups of similarly behaving genes, but does not as clearly identify the drivers of the phenotype.

In systems biology, computational modeling methods can derive great insight from multi-omic integration. An exciting new pursuit is whole cell modeling, which accounts for all cellular processes and molecules [15]. This represents a significant expansion from initial metabolism-only models. To properly parameterize the model, proteomics, metabolomics and transcriptome measurements are all utilized. Whole-cell models offer incredible validation of predicted functions and relationships of molecules in all biological processes as well as presenting hypotheses for emergent properties of the cell. This level of characterization currently only exists for the most genome-reduced organisms. However, the recent

1
2
3
4 increase in experimental data generation, particularly elucidation of protein complexes and localization,
5
6 will make such models feasible in the near future for many organisms.
7
8
9

10 As systems biology matures in its ability to both characterize and predict cellular functions, synthetic
11
12 biology emerges as the tool to build and control cellular functions. In the myriad applications of
13
14 synthetic biology, it is essential to understand the complex regulatory relationships that govern
15
16 molecular behavior. This understanding intrinsically comes from multi-omic analyses.
17
18
19

20 Finally, as computational biologists strive to integrate distinct data types, it is essential to develop a
21
22 framework that understands and appreciates the differences between multi-omic data. This same
23
24 appreciation is needed in the biological researchers which utilize the data. The needed task is not to
25
26 determine which of the non-correlating data are ‘correct’; each is correctly measuring a different
27
28 biomolecule. Rather, the task is to foster and utilize analytical paradigms that derive knowledge from
29
30 multiple data types. We need a mindset that is truly integrative, and not simply correlative.
31
32
33
34

35 **Box 1. Outstanding Questions**

- 36 • What is data integration beyond correlation?
- 37
- 38 • How do we educate new scientists to appreciate the diversity of –omics measurements?
- 39
- 40 • How to determine which –omic data type is best to investigate a hypothesis?
- 41
- 42 • When to generate multi-omic data?
- 43
44
45
46
47
48

49 **Acknowledgements**

50
51
52 The author thanks Karin Rodland, Josh Adkins, Jason McDermott and Katrina Waters for critical reading
53
54 of the manuscript, and Grant Fujimoto for assistance with images. SHP is supported by an Early Career
55
56 Award and the Pan-omics project from the US Department of Energy Genome Sciences Program and the
57
58 National Cancer Institute Clinical Proteomic Tumor Analysis Consortium (CPTAC, U24-CA-160019).
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure Legend

Figure 1. Diversity of regulation. The process of obtaining proteins from a genomic template is governed by many modalities of regulation, some of which are shown. Transcription can be regulated by the chromatin state of the DNA region contain the gene. Genetic regulation as inferred by quantitative trait locus (QTL) analysis identifies variable regions of a gene that affect the final abundance of the gene product. Regions that affect protein and transcript levels partially overlap but are not identical. mRNA stability can be affected by both intrinsic factors of the sequence itself, but also by extrinsic regulation such as through microRNAs. Translational efficiency denotes the amount of protein that is made from a transcript, and is affected by ribosome occupancy and other phenomena (e.g. codon usage). Decay rates for proteins are very different from those for mRNA, both as a global average and for a specific gene. Due to the complexities of regulation, it is not currently possible to predict protein abundance from mRNA.

References

- 1 Ghaemmamghami, S., *et al.* (2003) Global analysis of protein expression in yeast. *Nature* 425, 737-741
- 2 Ideker, T., *et al.* (2001) Integrated genomic and proteomic analyses of a systematically perturbed metabolic network. *Science* 292, 929-934
- 3 Marguerat, S., *et al.* (2012) Quantitative analysis of fission yeast transcriptomes and proteomes in proliferating and quiescent cells. *Cell* 151, 671-683
- 4 Gygi, S.P., *et al.* (1999) Correlation between protein and mRNA abundance in yeast. *Molecular and cellular biology* 19, 1720-1730
- 5 Le Roch, K.G., *et al.* (2004) Global analysis of transcript and protein levels across the Plasmodium falciparum life cycle. *Genome research* 14, 2308-2318
- 6 Nagaraj, N., *et al.* (2011) Deep proteome and transcriptome mapping of a human cancer cell line. *Molecular systems biology* 7, 548
- 7 Kislinger, T., *et al.* (2006) Global survey of organ and organelle protein expression in mouse: combined proteomic and transcriptomic profiling. *Cell* 125, 173-186
- 8 Vogel, C. and Marcotte, E.M. (2012) Insights into the regulation of protein abundance from proteomic and transcriptomic analyses. *Nature reviews. Genetics* 13, 227-232
- 9 Wu, L., *et al.* (2013) Variation and genetic control of protein abundance in humans. *Nature* 499, 79-82
- 10 Brar, G.A., *et al.* (2012) High-resolution view of the yeast meiotic program revealed by ribosome profiling. *Science* 335, 552-557
- 11 Baek, D., *et al.* (2008) The impact of microRNAs on protein output. *Nature* 455, 64-71
- 12 Schwanhauser, B., *et al.* (2011) Global quantification of mammalian gene expression control. *Nature* 473, 337-342
- 13 Waldbauer, J.R., *et al.* (2012) Transcriptome and proteome dynamics of a light-dark synchronized bacterial cell cycle. *PloS one* 7, e43432
- 14 Zhang, B., *et al.* (2014) Proteogenomic characterization of human colon and rectal cancer. *Nature* 513, 382-387
- 15 Sanghvi, J.C., *et al.* (2013) Accelerated discovery via a whole-cell model. *Nature methods* 10, 1192-1195

Figure 1

Genome

Types of Regulation

Protein Abundance