Manuscript format: Original article

Title: Polyunsaturated fatty acid interactions and breast cancer incidence: a population-based case-control study on Long Island, New York

Running title: Polyunsaturated fatty acids and breast cancer

Authors: Nikhil K. Khankari^{a*}, Patrick T. Bradshaw^b, Susan E. Steck^c, KaHe^d, Andrew F. Olshan^a, Jing Shen^e, Jiyoung Ahn^{f,g}, Yu Chen^{f,g}, Habibul Ahsan^h, Mary Beth Terryⁱ, Susan L. Teitelbaum^j, Alfred I. Neugut^{i,k}, Regina M. Santella^e, and Marilie D. Gammon^a

Author affiliations: Departments of ^aEpidemiology and ^bNutrition, University of North Carolina, Chapel Hill, NC 27599, USA; ^cDepartment of Epidemiology and Biostatistics, University of South Carolina, Columbia, SC, 29208, USA; ^dDepartment of Epidemiology and Biostatistics, Indiana University, Bloomington, IN 47405, USA; Departments of ^eEnvironmental Health, ⁱEpidemiology, and ^kMedicine, Columbia University, New York, NY, 10032, USA; Departments of ^fPopulation Health and ^gEnvironmental Medicine, New York University, New York, NY 10016, USA; ^hDepartment of Health Studies and Comprehensive Cancer Center, University of Chicago, Chicago, IL, 60637, USA; ^jDepartment of Preventive Medicine, Mount Sinai School of Medicine, New York, NY, 10029, USA

Corresponding author: *Nikhil K. Khankari, Department of Epidemiology, University of North Carolina, CB#7435, Chapel Hill, NC 27599-7435. Email: khankari@email.unc.edu
Corresponding author (present address): Division of Epidemiology, Vanderbilt University, Nashville, TN 37203. Email: nikhil.khankari@vanderbilt.edu

Abstract (word count = 200/200)

Purpose: Experimental studies demonstrate that ω -3 polyunsaturated fatty acids (PUFAs) inhibit inflammatory eicosanoids generated by ω -6 PUFAs. Epidemiologic studies on dietary ω -3 PUFA intake show consistent inverse associations with breast cancer incidence among Asian populations, where ω -3, relative to ω -6, intake is high. In contrast, associations are inconsistent among Western populations, where intake of ω -3, relative to ω -6, intake is low. We hypothesized that examining interactions between ω -3 and ω -6 would help elucidate the PUFA-breast cancer association in the U.S.

Methods: In a Long Island, NY, population-based study of 1463 breast cancer cases and 1500 controls, we estimated multivariable-adjusted odds ratios (ORs) and 95% confidence intervals (CIs) using unconditional logistic regression to examine interactions between ω -3 and ω -6 intake.

Results: We observed a super-additive interaction (Relative Excess Risk Due to Interaction=0.41; 95%CI=0.06,0.76) between ω -3 and ω -6 intake in association with breast cancer incidence, although the CIs for the joint exposure of low ω -3/high ω -6 compared to high ω -3/low ω -6 intake were wide (OR=1.20; 95%CI=0.85,1.69).

Conclusions: Breast cancer risk reduction may be possible for U.S. women with dietary consumption of higher ω -3, which have anti-inflammatory properties, in concert with lower ω -6, which induce inflammation. Replication from future U.S.-based investigations is needed.

MeSHkeywords: polyunsaturated fatty acids, seafood, breast neoplasms

Manuscript word count = 2999/3000

List of abbreviations:

AA	arachidonic acid
ALA	alpha-linolenic acid
Ca	cases
CI(s)	confidence interval(s)
Co	controls
DPA	docosapentaenoic acid
DHA	docosahexaenoic acid
EPA	eicosapentaenoic acid
FFQ	food frequency questionnaire
g	grams
kcal	kilocalories
LA	linoleic acid
LIBCSP	Long Island Breast Cancer Study Project
LRT	likelihood ratio test
N/A	not applicable
NHANES	National Health and Nutrition Examination Survey
N.Y.	New York
NSAID(s)	non-steroidal anti-inflammatory drug(s)
OR(s)	odds ratio(s)
pct	percentile
PUFA(s)	polyunsaturated fatty acid(s)
RERI	relative excess risk due to interaction
SD	standard deviation
U.S.	United States
VITAL	Vitamins and Lifestyle

Introduction

Breast cancer incidence rates are more than two times higher in the United States(U.S.) and European countries than in China or Japan[1,2]. Migration studies conducted among Asian immigrants show that breast cancer incidence patterns begin to reach those of Western countries a few generations after migration [3–6], suggesting environmental factors may play a role in the geographic variation in incidence rates observed in Asian and Western countries.

One potential environmental exposure is polyunsaturated fatty acids (PUFAs), which includes two primary classes, ω -3 and ω -6 fatty acids. Laboratory studies show that ω -3 PUFAs competitively inhibit ω -6 fatty acids, thus lowering levels of inflammatory eicosanoids generated from ω -6 metabolism [7], and that higher ω -3 relative to ω -6 could reduce breast cancer through inflammation, oxidative stress, and estrogen metabolism [7–10]. Asian populations have a substantially higher ratio of ω -3/ ω -6 intake compared to European and U.S. populations [11,12], due to higher fish consumption [13–15], which may partially explain the lower breast cancer risk observed in these populations [16–19]. However, previous U.S. and European epidemiologic studies examining the PUFA and breast cancer association remain inconsistent [20–40]. The biologic influence of PUFAs is unlikely to differ across populations; however, low fish intake and high ω -6 intake [41] in Western countries may mask important risk reductions. We hypothesized that consideration of both ω -3 and ω -6 intake, as an interaction or as the relative balance (ω -3/ ω -6 ratio), may elucidate the potential benefit of ω -3 intake among populations with low fish consumption.

In the study reported here, we examined the interaction between dietary ω -3 and ω -6 PUFA classes in association with breast cancer risk among women on Long Island, New York(NY).

Materials and Methods

We used the population-based case-control component of the Long Island Breast Cancer Study Project (LIBCSP) comprising English-speaking residents of Nassau and Suffolk counties. Details of the LIBCSP have been published previously [42]. Institutional Review Board approval was obtained from all participating institutions.

Study population. Cases were women newly diagnosed with a first primary *in situ* or invasive breast cancer, identified using a “super-rapid” network by contacting hospital pathology departments daily or 2-3 per week. Controls were identified using Waksberg’s method of random digit dialing [43] for women under 65 years of age, and Health Care Finance Administration rosters for women 65 years and older. Controls were frequency matched to the expected age-distribution of cases. There were no age or race restrictions for subject eligibility.

The parent LIBCSP respondents included 1,508 cases and 1,556 controls (82% and 63% response rates, respectively). Respondents ranged in age from 20 to 98 years of age, 67% were postmenopausal, and the majority self-reported their race as white (94%), followed by black or African American (4%), or other (2%), which is consistent with the racial distribution of these two counties at the time of data collection [44]. Most LIBCSP participants were highly educated (>90% graduating from high school), used alcohol (62%), were parous (88%), never used hormone replacement therapy (74%), never used oral contraceptives (55%), and did not have a family history of breast cancer (84%). Among cases, the majority (84%) were diagnosed with first, primary invasive breast cancer [42].

Assessment of PUFAs and other covariates. LIBCSP participants were administered a main risk factor questionnaire by a trained interviewer about 3 months after diagnosis for cases and 5.5 months after identification for controls [42]. Approximately 98% of participants

(1,479 cases and 1,520 controls) also completed the validated [45–47] self-administered 101-item modified Block food frequency questionnaire (FFQ). After excluding participants with implausible total energy intake (± 3 standard deviations from the mean; $n=36$), 1,463 cases and 1,500 controls remained in our analysis.

We estimated PUFA intake by linking responses from the FFQ (i.e., grams per day for each line item) with nutrient values available in the U.S. Department of Agriculture databases for ω -3 and ω -6 PUFAs [48]. The following PUFAs were estimated: (1) ω -3 fatty acids, including α -linolenic acid (ALA), docosapentaenoic acid (DPA), docosahexaenoic acid (DHA), eicosapentaenoic acid (EPA); and (2) ω -6 fatty acids, including linoleic acid (LA) and arachidonic acid (AA). An estimate of total PUFA intake was calculated by combining all individual fatty acids. Additionally, an estimate of total ω -3 and ω -6 fatty acids was obtained by summing each individual fatty acid within category (e.g., total ω -3 = ALA + DPA + DHA + EPA). We also examined fish/seafood intake according to the items recorded in the FFQ: (1) tuna, tuna salad, tuna casserole; (2) shell fish (shrimp, lobster, crab, oysters, etc.); and (3) other fish (either broiled/baked). Total fish intake was calculated by summing each of the fish/seafood items recorded in the FFQ.

Statistical analyses. All analyses were conducted using SAS version 9.2 (SAS Institute, Cary, NC). Unconditional logistic regression was used to estimate odds ratios (ORs) and 95% confidence intervals (95% CI) for the association between PUFA intake and breast cancer risk. PUFA and fish/seafood intake were categorized as quartiles, according to the distribution among controls. Quartiles were selected over other methods of categorization (e.g., tertiles, quintiles, linear, splines) because the shape of the dose-response between exposure and breast cancer incidence was best represented using these cut-points. The relation between any of the

1
2
3
4 PUFA and/or fish and breast cancer incidence was not strictly monotonic [49] thus linear trend
5
6 tests were not conducted.
7

8
9 Interactions between total ω -3 and total ω -6 intake in association with breast
10
11 cancer incidence were assessed on the additive (common referent) and multiplicative scales.
12
13 Additive interaction was evaluated using relative excess risk due to interaction (RERI), with 95%
14
15 CI [50]. Multiplicative interactions were evaluated by comparing nested models using the
16
17 Likelihood Ratio Test (LRT) [49]. Total ω -3, total ω -6, and ω -3/ ω -6 ratio were dichotomized at
18
19 the median for use in the interaction models.
20
21
22

23
24 We also considered effect modification of the association between PUFA intake and
25
26 breast cancer risk by: menopausal status (post- vs. pre-menopausal status); and dietary
27
28 supplement use (yes/no). However, little or no heterogeneity was observed with either of these
29
30 covariates (data not shown). We also considered potential heterogeneity across breast cancer
31
32 subtypes, defined by hormone receptor status (any hormone receptor positive breast cancer vs.
33
34 no hormone receptor positive breast cancer), by constructing polytomous regression models;
35
36 however, no differences were observed across hormone receptor subtype (data not shown).
37
38
39

40
41 All models were adjusted for the frequency matching factor five-year age group. Other
42
43 potential confounders (including total energy intake, non-steroidal anti-inflammatory drugs
44
45 (NSAID), family history of breast cancer, income, body mass index, alcohol use, fruit and
46
47 vegetable intake, and physical activity) were identified using directed acyclic graph [49]. The
48
49 only covariate that changed the estimates by more than 10% was total energy intake for PUFA
50
51 intake. All fish intake models included age, energy intake and NSAID use given the possibility
52
53 that chronic NSAID users experience gastrointestinal problems (e.g., stomach ulcers, reflux)
54
55 which may subsequently influence diet, including fish consumption [51].
56
57
58
59
60
61
62
63
64
65

Results

As presented in **Table 1**, the average intake of total ω -3 fatty acids (1.01 grams per day (SD=0.74)) was lower relative to ω -6 intake [7.66 grams per day (SD=5.68)] among the 1,500 control women in this population-based sample of Long Island residents. The highest contributor to total ω -3 intake was ALA with average intake of 0.86 grams per day (SD=0.71), whereas LA was the highest contributor to total ω -6 intake with an average intake of 7.59 grams per day (SD=5.66). Tuna intake was reported at higher levels [8.80 grams per day (SD=13.98)] in our control population compared to shell fish intake [3.57 grams per day (SD=9.07)]. Fish was a major contributor to high intake of long-chain ω -3 PUFAs, including DPA, DHA, and EPA. In contrast, foods that contributed to high ALA intake were biscuits/muffins and other fried foods, which was similar to what was observed for LA intake. High AA intake appeared to be driven by eggs and meats, including fish, chicken, and ham.

Multivariable-adjusted odds ratios for the associations between all measures of PUFA intake and breast cancer incidence were imprecise (and most were not statistically significant), as presented in **Table 2**. For example, elevated odds ratios were observed for high intake of total PUFA, total ω -3, ALA, ω -6, and LA intakes, but CIs were wide. No associations were observed for the long-chain ω -3 PUFA (DPA, DHA, EPA), AA, or high intake of the ω -3/ ω -6 ratio.

As shown in **Table 3**, we observed an interaction between ω -3 and ω -6 intake, which was statistically significant on the additive scale [adjusted RERI=0.41 (95% CI=0.06, 0.76)]. Risk reductions for breast cancer were modest for women who consumed low levels of both ω -3 and ω -6 [adjusted OR=0.83 (95% CI=0.63, 1.09)], compared to women who consumed high ω -3 and low ω -6. For women who consumed high levels of ω -3 and ω -6, the odds ratios were close to the null value [adjusted OR=0.95 (95% CI=0.72, 1.26)]. In contrast, higher intakes of ω -6 fatty acids in

1
2
3
4 conjunction with lower intake of ω -3 fatty acids was associated with an approximately 20%
5
6 increased breast cancer incidence[adjusted OR=1.20 (95% CI=0.85,1.69)]. The increase
7
8 observed for this group was super-additive (41% greater) compared to the 22% (=5%+17%)
9
10 expected risk reduction, derived from the individual ORs for those consuming either high levels
11
12 ($\geq$ median; 5% risk reduction), or low levels ($<$ median; 17% risk reduction) of both ω -3 and ω -6
13
14 fatty acids. Similar results were observed when we considered interactions between ω -3 and ω -6
15
16 on a multiplicative scale, as shown in **Table 4**.
17
18
19
20

21 Higher intake of total fish, tuna, shell fish, or other fish (broiled/baked) was not
22
23 associated with breast cancer risk in our study, as presented in **Table 5**.
24
25
26
27

28 **Discussion**

29
30

31 We are the first to report an additive interaction between ω -3 intake and ω -6 intake in
32
33 relation to breast cancer risk in a population-based sample of U.S women. Specifically, we
34
35 observed a 20% increase in the odds of breast cancer among consumers of high levels of ω -6 and
36
37 low levels of ω -3 compared to those who consumed low levels of ω -6 and high levels of ω -3.
38
39 The odds ratio for women consuming high ω -6 and low ω -3 was increased, whereas the
40
41 corresponding estimates for intake of high levels or low levels, of either PUFA class, were
42
43 reduced. This interaction underscores the importance of considering intake of ω -3 and ω -6
44
45 simultaneously when examining associations with breast cancer in the U.S.
46
47
48
49

50 No previous U.S. studies have reported the potential interaction between ω -3 and ω -6
51
52 PUFAs and breast cancer risk, and only two examined the ratio of ω -3 and ω -6 intakes [23,30].
53
54 Consideration of an interaction may be preferable, given that a ratio measure permits only one
55
56 type of relation between two exposures, whereas an interaction is more flexible. One previous
57
58
59
60
61
62
63
64
65

Shanghai study reported an interaction between ω -3 and ω -6 intake on breast cancer risk[18], with a significant two-fold increased risk for high ω -6 in combination with low marine-derived ω -3 intake. The LIBCSP results presented here for the interaction between ω -3 and ω -6 intake are in the same direction as those reported in the Shanghai study, but less pronounced. Importantly, daily fish consumption in the Shanghai population was almost five times greater than the frequency reported among our Long Island population, which could partially explain the weaker association observed in our study.

The modest positive association between ω -6 PUFA (total ω -6, LA, AA) and breast cancer incidence observed here is similar to associations reported in other studies conducted among Western populations[16,18,28,30,35]. However, our findings for the association between long-chain ω -3 (DPA, DHA, EPA) PUFAs are not consistent with the risk reductions reported by previous studies conducted among Asian and some European populations[16–19,32]. Additionally, the increased risk for ALA intake we observed is inconsistent with the laboratory evidence for inhibition of breast cancer growth[52,53]. However, in other epidemiologic studies, the association between ALA intake and breast cancer risk remains unclear, with some studies reporting increased risks [18,20,29,32,54], and others reporting risk reductions [24,36]. The variation in results across studies may be due to different dietary assessment methods used, consumption of different food sources of ALA (e.g., biscuits/muffins and fried foods were major contributors in our population), or with potential recall bias that can occur in case-control studies. Or, perhaps, ALA reduces breast cancer growth only after conversion to long-chain ω -3 PUFAs. The *in vivo* conversion of ALA to long-chain ω -3 PUFAs is inefficient in the presence of high ω -6 [55]. Thus, it is possible that in populations with high ω -6 intake, benefits of ALA intake are less evident.

1
2
3
4 The slight breast cancer risk reductions in relation to a high ω -3/ ω -6 ratio observed in our
5
6 study were modest compared to estimates reported in other studies in European [56],
7
8 Mexican [57], and U.S. populations [23,30]. However, this may reflect the relatively low intake of
9
10 ω -3 and ω -6 in our study population. Very low intake of both ω -3 and ω -6 PUFAs could result in
11
12 a high ratio value for ω -3/ ω -6 intake (when ω -6 intake is less than 1). Doses in excess of 2 g/day
13
14 may be required for decreasing prostaglandin E2 production [58,59], a primary inflammatory
15
16 eicosanoid resulting from AA metabolism, which has been implicated in carcinogenesis
17
18 [7]. Thus, a high ratio of ω -3/ ω -6 derived from low intakes of ω -3 and ω -6 PUFAs may not
19
20 represent a sufficient dose for ω -3 to exert a beneficial response *in vivo*. In the U.S., only one
21
22 previous population-based study has considered the PUFA ratio in association with breast cancer
23
24 risk; utilizing data from the Vitamins and Lifestyle Cohort (VITAL) [30], a 16% risk reduction
25
26 was observed in association with high ω -3/ ω -6 intake ratio. This Western Washington-based
27
28 study included marine-derived ω -3 intake from both dietary sources and supplements, and thus
29
30 levels of ω -3 intake were higher than the dietary-derived intake estimates observed in our Long
31
32 Island-based study. Nonetheless, given the modest risk reductions reported for the ω -3/ ω -6 ratio
33
34 in both studies, examining the interaction between ω -3 and ω -6 intake, rather than the ratio, may
35
36 be a more favorable strategy in populations where PUFA intake is relatively low.
37
38
39
40
41
42
43
44

45 Strengths of our population-based, case-control study include observation of an
46
47 interaction between ω -3 and ω -6 on breast cancer development, which has not been previously
48
49 assessed in a U.S. population. We also examined associations with fish intake (a dietary source
50
51 rich in ω -3 fatty acids) among women who reside in a geographic area that is surrounded by
52
53 water, and for whom the variability of fish intake would presumably be greater than for others
54
55 who reside in more land-locked areas of the U.S. [14]. However, we did not observe any notable
56
57
58
59
60
61
62
63
64
65

1
2
3
4 associations with fish intake and breast cancer incidence, even though women living in New
5
6 York City [60] have been reported to consume fish greater than the national estimates from
7
8 NHANES [61].
9

10
11 This study also has limitations. The LIBCSP population includes predominantly
12
13 Caucasian women, which reflects the racial distribution of the residents of the two source
14
15 counties on Long Island. Consequently, examination of racial differences was not possible. Our
16
17 results are therefore generalizable to only Caucasian-American women, for whom breast cancer
18
19 risk remains high [62]. Future studies may also be warranted to evaluate the timing of exposure
20
21 relative to breast cancer development. FFQ responses are assumed to reflect usual adult diet [63],
22
23 although recent changes due to a disease diagnosis or treatment regimens could influence those
24
25 responses. The LIBCSP questionnaires were administered within months of diagnosis, and for
26
27 two-thirds of women this was prior to the onset of chemotherapy [42], which is likely to reduce
28
29 the impact of dietary changes and perhaps recall of diet on the FFQ. However, a single dietary
30
31 assessment via FFQ may not necessarily reflect diet during all etiologically relevant time periods
32
33 for breast cancer development (i.e., adolescence) [64].
34
35
36
37
38
39
40

41 Furthermore, estimating PUFA intake via FFQ linkage with the USDA database could
42
43 result in measurement error. For example, the PUFA content measured in the foods reported in
44
45 the USDA database [48] may differ from those actually consumed by LIBCSP participants due to
46
47 differences in harvesting, storage, processing, and cooking methods [65–67]. Additionally, we
48
49 were unable to assess associations with consumption of different fish varieties. This is important
50
51 given the amount of long-chain ω -3 content found in fish differs by species [61]. However, we
52
53 assessed tuna intake, which is the most frequently consumed fish variety in the U.S. and is also a
54
55 major source of dietary ω -3s [14]. PUFA intake via supplements (i.e., fish oil consumption) was
56
57
58
59
60
61
62
63
64
65

not measured as part of the case-control interview. Given fish oil contributes to intake of long-chain ω -3 this could underestimate DHA and EPA consumption in our population. However, in a follow-up interview, 89% of the LIBCSP cases reported never using fish oil supplements approximately five years after diagnosis [68]. Fish oil supplement use has only recently (after 2002) received attention due to its potential risk reductions for cardiovascular disease [69]. Furthermore, use of fish oil supplements increased post-2002 in two large U.S. cohorts [70]. Thus, prevalence of fish oil supplement use at the time of the LIBCSP case-control interview (during years 1996-1997) was likely low and the potential underestimation of DHA and EPA intake may be negligible. Biomarkers could provide an objective measure of PUFA intake; however, biomarkers may reflect different time periods of exposure, ranging from a few days to one year (depending on the type of biomarker used) [71]. Therefore, use of PUFA biomarker measurements in a case-control study may not reflect the etiologically relevant time period for breast cancer development. Future studies, if feasible, should consider multiple prospective biomarker measurements in order to capture dietary PUFA exposure (including changes in diet) during relevant periods of breast cancer etiology.

In conclusion, we observed among a population-based sample of Long Island residents, women who consume high levels of ω -6 and low levels of ω -3 had an increased risk for breast cancer, compared to women who consume low levels of ω -6 and high levels of ω -3. Our results suggest that high intake of ω -3 PUFA, coupled with low intake of ω -6, and maybe a potential risk reduction strategy for breast cancer among U.S. women. Further confirmation using an objective measurement of PUFA intake via biomarkers may be warranted.

Source of Funding: This study was supported in part with grants from the National Institutes of Health (CA/ES 66572, CA58233, ES10126,ES009089); the Department of Defense (U.S. Army BC972005); and the Breast Cancer Research Foundation.

Acknowledgements: None.

Conflict Disclosure: The authors declare no financial or non-financial conflicts.

References

- [1] Society AC. Cancer Facts & Figures 2014. American Cancer Society; 2014.
- [2] Ferlay J, Shin HR, Bray F, Forman D, Mathers C, Parkin DM. Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008. *Int J Cancer* 2010;127:2893–917.
- [3] Althuis MD, Dozier JM, Anderson WF, Devesa SS, Brinton LA. Global trends in breast cancer incidence and mortality 1973-1997. *Int J Epidemiol* 2005;34:405–12.
- [4] Chie WC, Chen CF, Chen CJ, Chang CL, Liaw YP, Lin RS. Geographic variation of breast cancer in Taiwan: international and migrant comparison. *Anticancer Res* 1995;15:2745–9.
- [5] Stanford JL, Herrinton LJ, Schwartz SM, Weiss NS. Breast cancer incidence in Asian migrants to the United States and their descendants. *Epidemiol Camb Mass* 1995;6:181–3.
- [6] Wu AH, Bernstein L. Breast Cancer among Asian Americans and Pacific Islanders. *Asian Am Pac Isl J Health* 1998;6:327–43.
- [7] Chenais B, Blanckaert V. The janus face of lipids in human breast cancer: how polyunsaturated Fatty acids affect tumor cell hallmarks. *Int J Breast Cancer* 2012;2012:712536.
- [8] Attar E, Tokunaga H, Imir G, Yilmaz MB, Redwine D, Putman M, et al. Prostaglandin E2 via steroidogenic factor-1 coordinately regulates transcription of steroidogenic genes necessary for estrogen synthesis in endometriosis. *J Clin Endocrinol Metab* 2009;94:623–31.
- [9] Bougnoux P, Hajjaji N, Maheo K, Couet C, Chevalier S. Fatty acids and breast cancer: sensitization to treatments and prevention of metastatic re-growth. *Prog Lipid Res* 2010;49:76–86.
- [10] Brueggemeier RW, Richards JA, Petrel TA. Aromatase and cyclooxygenases: enzymes in breast cancer. *J Steroid Biochem Mol Biol* 2003;86:501–7.
- [11] Sekikawa A, Curb JD, Ueshima H, El-Saed A, Kadowaki T, Abbott RD, et al. Marine-derived n-3 fatty acids and atherosclerosis in Japanese, Japanese-American, and white men: a cross-sectional study. *J Am Coll Cardiol* 2008;52:417–24.
- [12] Simopoulos AP. The importance of the omega-6/omega-3 fatty acid ratio in cardiovascular disease and other chronic diseases. *Exp Biol Med Maywood NJ* 2008;233:674–88.
- [13] Kris-Etherton PM, Taylor DS, Yu-Poth S, Huth P, Moriarty K, Fishell V, et al. Polyunsaturated fatty acids in the food chain in the United States. *Am J Clin Nutr* 2000;71:179S – 88S.
- [14] Mahaffey KR, Clickner RP, Jeffries RA. Adult women's blood mercury concentrations vary regionally in the United States: association with patterns of fish consumption (NHANES 1999-2004). *Environ Health Perspect* 2009;117:47–53.
- [15] Tran NL, Barraji LM, Bi X, Schuda LC, Moya J. Estimated long-term fish and shellfish intake—national health and nutrition examination survey. *J Expo Sci Environ Epidemiol* 2013;23:128–36.
- [16] Gago-Dominguez M, Yuan JM, Sun CL, Lee HP, Yu MC. Opposing effects of dietary n-3 and n-6 fatty acids on mammary carcinogenesis: The Singapore Chinese Health Study. *Br J Cancer* 2003;89:1686–92.
- [17] Kim J, Lim S-Y, Shin A, Sung M-K, Ro J, Kang H-S, et al. Fatty fish and fish omega-3 fatty acid intakes decrease the breast cancer risk: a case-control study. *BMC Cancer* 2009;9:216. doi:10.1186/1471-2407-9-216.

- [18] Murff HJ, Shu XO, Li H, Yang G, Wu X, Cai H, et al. Dietary polyunsaturated fatty acids and breast cancer risk in Chinese women: A prospective cohort study. *Int J CancerJournal Int Cancer* 2010.
- [19] Wakai K, Tamakoshi K, Date C, Fukui M, Suzuki S, Lin Y, et al. Dietary intakes of fat and fatty acids and risk of breast cancer: a prospective study in Japan. *Cancer Sci* 2005;96:590–9.
- [20] Barrett-Connor E, Friedlander NJ. Dietary fat, calories, and the risk of breast cancer in postmenopausal women: a prospective population-based study. *J Am Coll Nutr* 1993;12:390–9.
- [21] Byrne C, Rockett H, Holmes MD. Dietary fat, fat subtypes, and breast cancer risk: lack of an association among postmenopausal women with no history of benign breast disease. *Cancer Epidemiol Biomark Prev Publ Am Assoc Cancer Res Cosponsored Am Soc Prev Oncol* 2002;11:261–5.
- [22] Folsom AR, Demissie Z. Fish intake, marine omega-3 fatty acids, and mortality in a cohort of postmenopausal women. *Am J Epidemiol* 2004;160:1005–10.
- [23] Goodstine SL, Zheng T, Holford TR, Ward BA, Carter D, Owens PH, et al. Dietary (n-3)/(n-6) fatty acid ratio: possible relationship to premenopausal but not postmenopausal breast cancer risk in U.S. women. *J Nutr* 2003;133:1409–14.
- [24] Holmes MD, Hunter DJ, Colditz GA, Stampfer MJ, Hankinson SE, Speizer FE, et al. Association of dietary intake of fat and fatty acids with risk of breast cancer. *JAMA J Am Med Assoc* 1999;281:914–20.
- [25] Katsouyanni K, Trichopoulou A, Stuver S, Garas Y, Kritselis A, Kyriakou G, et al. The association of fat and other macronutrients with breast cancer: a case-control study from Greece. *Br J Cancer* 1994;70:537–41.
- [26] Landa MC, Frago N, Tres A. Diet and the risk of breast cancer in Spain. *Eur J Cancer Prev Off J Eur Cancer Prev Organ ECP* 1994;3:313–20.
- [27] Lof M, Sandin S, Laggiou P, Hilakivi-Clarke L, Trichopoulos D, Adami HO, et al. Dietary fat and breast cancer risk in the Swedish women's lifestyle and health cohort. *Br J Cancer* 2007;97:1570–6.
- [28] Martin-Moreno JM, Willett WC, Gorgojo L, Banegas JR, Rodriguez-Artalejo F, Fernandez-Rodriguez JC, et al. Dietary fat, olive oil intake and breast cancer risk. *Int J CancerJournal Int Cancer* 1994;58:774–80.
- [29] Nkondjock A, Shatenstein B, Ghadirian P. A case-control study of breast cancer and dietary intake of individual fatty acids and antioxidants in Montreal, Canada. *Breast Edinb Scotl* 2003;12:128–35.
- [30] Sczaniecka AK, Brasky TM, Lampe JW, Patterson RE, White E. Dietary Intake of Specific Fatty Acids and Breast Cancer Risk Among Postmenopausal Women in the VITAL Cohort. *Nutr Cancer* 2012;64:1131–42.
- [31] Tavani A, Pelucchi C, Parpinel M, Negri E, Franceschi S, Levi F, et al. n-3 polyunsaturated fatty acid intake and cancer risk in Italy and Switzerland. *Int J CancerJournal Int Cancer* 2003;105:113–6.
- [32] Thiebaut AC, Chajes V, Gerber M, Boutron-Ruault MC, Joulin V, Lenoir G, et al. Dietary intakes of omega-6 and omega-3 polyunsaturated fatty acids and the risk of breast cancer. *Int J CancerJournal Int Cancer* 2009;124:924–31.

- [33] Toniolo P, Riboli E, Shore RE, Pasternack BS. Consumption of meat, animal products, protein, and fat and risk of breast cancer: a prospective cohort study in New York. *Epidemiol Camb Mass* 1994;5:391–7.
- [34] Van 't Veer P, van Leer EM, Rietdijk A, Kok FJ, Schouten EG, Hermus RJ, et al. Combination of dietary factors in relation to breast-cancer occurrence. *Int J Cancer J Int Cancer* 1991;47:649–53.
- [35] Velie E, Kulldorff M, Schairer C, Block G, Albanes D, Schatzkin A. Dietary fat, fat subtypes, and breast cancer in postmenopausal women: a prospective cohort study. *J Natl Cancer Inst* 2000;92:833–9.
- [36] Voorrips LE, Brants HA, Kardinaal AF, Hiddink GJ, Brandt PA van den, Goldbohm RA. Intake of conjugated linoleic acid, fat, and other fatty acids in relation to postmenopausal breast cancer: the Netherlands Cohort Study on Diet and Cancer. *Am J Clin Nutr* 2002;76:873–82.
- [37] Wang J, John EM, Ingles SA. 5-Lipoxygenase and 5-Lipoxygenase-Activating Protein Gene Polymorphisms, Dietary Linoleic Acid, and Risk for Breast Cancer. *Cancer Epidemiol Biomark Prev Publ Am Assoc Cancer Res Cosponsored Am Soc Prev Oncol* 2008;17:2748–54.
- [38] Wirfalt E, Mattisson I, Gullberg B, Johansson U, Olsson H, Berglund G. Postmenopausal breast cancer is associated with high intakes of omega6 fatty acids (Sweden). *Cancer Causes Control CCC* 2002;13:883–93.
- [39] Witte JS, Ursin G, Siemiatycki J, Thompson WD, Paganini-Hill A, Haile RW. Diet and premenopausal bilateral breast cancer: a case-control study. *Breast Cancer Res Treat* 1997;42:243–51.
- [40] Zaridze D, Lifanova Y, Maximovitch D, Day NE, Duffy SW. Diet, alcohol consumption and reproductive factors in a case-control study of breast cancer in Moscow. *Int J CancerJournal Int Cancer* 1991;48:493–501.
- [41] Blasbalg TL, Hibbeln JR, Ramsden CE, Majchrzak SF, Rawlings RR. Changes in consumption of omega-3 and omega-6 fatty acids in the United States during the 20th century. *Am J Clin Nutr* 2011;93:950–62.
- [42] Gammon MD, Neugut AI, Santella RM, Teitelbaum SL, Britton JA, Terry MB, et al. The Long Island Breast Cancer Study Project: description of a multi-institutional collaboration to identify environmental risk factors for breast cancer. *Breast Cancer Res Treat* 2002;74:235–54.
- [43] Waksberg J. Sampling methods for random digit dialing. *J Amer Stat Assoc* 1978:40.
- [44] Gammon MD, Santella RM, Neugut AI, Eng SM, Teitelbaum SL, Paykin A, et al. Environmental toxins and breast cancer on Long Island. I. Polycyclic aromatic hydrocarbon DNA adducts. *Cancer Epidemiol Biomark Prev Publ Am Assoc Cancer Res Cosponsored Am Soc Prev Oncol* 2002;11:677–85.
- [45] Block G, Hartman AM, Dresser CM, Carroll MD, Gannon J, Gardner L. A data-based approach to diet questionnaire design and testing. *Am J Epidemiol* 1986;124:453–69.
- [46] Block G, Woods M, Potosky A, Clifford C. Validation of a self-administered diet history questionnaire using multiple diet records. *J Clin Epidemiol* 1990;43:1327–35.
- [47] Potischman N, Swanson CA, Coates RJ, Weiss HA, Brogan DR, Stanford JL, et al. Dietary relationships with early onset (under age 45) breast cancer in a case-control study in the United States: influence of chemotherapy treatment. *Cancer Causes Control CCC* 1997;8:713–21.

- [48] U.S. Department of Agriculture, Agricultural Research Service, USDA National Nutrient Database for Standard Reference, Release 25. Nutrient Data Laboratory Home Page. 2012.
- [49] Rothman KJ, Greenland S, Lash TL. Modern epidemiology. 3 , [thoroughly rev a updat. Philadelphia: Wolters Kluwer Health/Lippincott Williams & Wilkins; 2008.
- [50] Hosmer DW, Lemeshow S. Confidence interval estimation of interaction. *Epidemiol Camb Mass* 1992;3:452–6.
- [51] Sostres C, Gargallo CJ, Lanas A. Nonsteroidal anti-inflammatory drugs and upper and lower gastrointestinal mucosal damage. *Arthritis Res Ther* 2013;15 Suppl 3:S3.
- [52] Chen J, Stavro PM, Thompson LU. Dietary flaxseed inhibits human breast cancer growth and metastasis and downregulates expression of insulin-like growth factor and epidermal growth factor receptor. *Nutr Cancer* 2002;43:187–92. doi:10.1207/S15327914NC432_9.
- [53] Truan JS, Chen JM, Thompson LU. Flaxseed oil reduces the growth of human breast tumors (MCF-7) at high levels of circulating estrogen. *Mol Nutr Food Res* 2010;54:1414–21.
- [54] De Stefani E, Deneo-Pellegrini H, Mendilaharsu M, Ronco A. Essential fatty acids and breast cancer: a case-control study in Uruguay. *Int J Cancer J Int Cancer* 1998;76:491–4.
- [55] Gerster H. Can adults adequately convert alpha-linolenic acid (18:3n-3) to eicosapentaenoic acid (20:5n-3) and docosahexaenoic acid (22:6n-3)? *Int J Vitam Nutr Res Z Vitam-ErnahrungsforschungJournal Int Vitaminol Nutr* 1998;68:159–73.
- [56] Crowe FL, Key TJ, Appleby PN, Travis RC, Overvad K, Jakobsen MU, et al. Dietary fat intake and risk of prostate cancer in the European Prospective Investigation into Cancer and Nutrition. *Am J Clin Nutr* 2008;87:1405–13.
- [57] Chajes V, Torres-Mejia G, Biessy C, Ortega-Olvera C, Angeles-Llerenas A, Ferrari P, et al. omega-3 and omega-6 Polyunsaturated fatty acid intakes and the risk of breast cancer in Mexican women: impact of obesity status. *Cancer Epidemiol Biomark Prev Publ Am Assoc Cancer Res Cosponsored Am Soc Prev Oncol* 2012;21:319–26.
- [58] Calder PC. Omega-3 polyunsaturated fatty acids and inflammatory processes: Nutrition or pharmacology?: Omega-3 fatty acids and inflammation. *Br J Clin Pharmacol* 2012;n/a – n/a. doi:10.1111/j.1365-2125.2012.04374.x.
- [59] Fabian CJ, Kimler BF, Hursting SD. Omega-3 fatty acids for breast cancer prevention and survivorship. *Breast Cancer Res BCR* 2015;17:62. doi:10.1186/s13058-015-0571-6.
- [60] McKelvey W, Gwynn RC, Jeffery N, Kass D, Thorpe LE, Garg RK, et al. A biomonitoring study of lead, cadmium, and mercury in the blood of New York city adults. *Environ Health Perspect* 2007;115:1435–41.
- [61] Mahaffey KR. Fish and shellfish as dietary sources of methylmercury and the omega-3 fatty acids, eicosahexaenoic acid and docosahexaenoic acid: risks and benefits. *Environ Res* 2004;95:414–28.
- [62] Desantis C, Ma J, Bryan L, Jemal A. Breast cancer statistics, 2013. *CA Cancer J Clin* 2014;64:52–62.
- [63] Willett W. Nutritional epidemiology. vol. 30. 2nd ed. New York: Oxford University Press; 1998.
- [64] Biro FM, Deardorff J. Identifying opportunities for cancer prevention during preadolescence and adolescence: puberty as a window of susceptibility. *J Adolesc Health Off Publ Soc Adolesc Med* 2013;52:S15–20. doi:10.1016/j.jadohealth.2012.09.019.

- 1
2
3
4 [65] Beveridge MC, Thilsted SH, Phillips MJ, Metian M, Troell M, Hall SJ. Meeting the food
5 and nutrition needs of the poor: the role of fish and the opportunities and challenges
6 emerging from the rise of aquaculture. *J Fish Biol* 2013;83:1067–84.
7
8 [66] Echarte M, Zulet MA, Astiasaran I. Oxidation process affecting fatty acids and cholesterol
9 in fried and roasted salmon. *J Agric Food Chem* 2001;49:5662–7.
10
11 [67] Li H, Fan YW, Li J, Tang L, Hu JN, Deng ZY. Evaluating and predicting the oxidative
12 stability of vegetable oils with different fatty acid compositions. *J Food Sci* 2013;78:H633–
13 41.
14 [68] Link AR, Gammon MD, Jacobson JS, Abrahamson P, Bradshaw PT, Terry MB, et al. Use
15 of Self-Care and Practitioner-Based Forms of Complementary and Alternative Medicine
16 before and after a Diagnosis of Breast Cancer. *Evid-Based Complement Altern Med ECAM*
17 2013;2013:301549. doi:10.1155/2013/301549.
18
19 [69] Kris-Etherton PM. Fish Consumption, Fish Oil, Omega-3 Fatty Acids, and Cardiovascular
20 Disease. *Circulation* 2002;106:2747–57. doi:10.1161/01.CIR.0000038493.65177.94.
21
22 [70] Kim HJ, Giovannucci E, Rosner B, Willett WC, Cho E. Longitudinal and secular trends in
23 dietary supplement use: Nurses' Health Study and Health Professionals Follow-Up Study,
24 1986-2006. *J Acad Nutr Diet* 2014;114:436–43. doi:10.1016/j.jand.2013.07.039.
25
26 [71] Arab L. Biomarkers of fat and fatty acid intake. *J Nutr* 2003;133 Suppl 3:925S – 932S.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 1. Characteristics of polyunsaturated fatty acid intake (PUFA) and fish intake among the population-based sample of control women (N=1500) in the LIBCSP, 1996-1997

Population-based sample of control women (N=1500) in the LIBCSP, 1998-1999						
Nutrient/Food	Mean	SD	Quartile Cutpoints			Major PUFA-rich foods contributing to high nutrient intake in the LIBCSP
			25 th Pct	50 th Pct	75 th Pct	
Nutrient (g/day)						
Total PUFA ^a	8.67	6.31	4.21	7.27	11.25	Butter, Mayonnaise/salad dressings, safflower/corn oil, margarine, peanuts/peanut butter
Total ω-3 ^b	1.01	0.74	0.49	0.83	1.30	Biscuits/muffins, butter, mayonnaise/salad dressings, fish, safflower/corn oil
ALA	0.86	0.71	0.35	0.68	1.14	Biscuits/muffins, French fries/fried potatoes, butter, cookies, mayonnaise/salad dressings
DPA	0.01	0.01	0.01	0.01	0.02	Tuna, fish, chicken, shellfish, beef
DHA	0.09	0.09	0.03	0.06	0.12	Tuna, fish, eggs, shellfish, chicken
EPA	0.04	0.05	0.01	0.03	0.06	Fish, tuna, shellfish, chicken
Total ω-6 ^c	7.66	5.68	3.68	6.31	10.10	Biscuits/muffins, French fries/fried potatoes, butter, chips/popcorn, mayonnaise/salad dressings
LA	7.59	5.66	3.65	6.23	9.99	Biscuits/muffins, French fries/fried potatoes, butter, chips/popcorn, mayonnaise/salad dressings
AA	0.07	0.06	0.04	0.06	0.09	Eggs, Fish, chicken, ham/lunch meats, shellfish
ω-3/ω-6	0.15	0.14	0.10	0.14	0.17	N/A
Fish (g/day) ^d						
Total fish ^e	19.62	23.83	4.87	13.23	26.86	N/A
Tuna	8.80	13.98	0.00	4.77	12.40	N/A
Shell fish	3.57	9.07	0.00	0.00	4.62	N/A
Other (broiled/baked)	7.25	10.59	0.00	2.80	10.77	N/A

Note:

^a Total PUFA = ALA + DPA + DHA + EPA + LA + AA^b Total ω -3 = ALA + DPA + DHA + EPA^c Total ω -6 = LA + AA^d Controls with null values for total fish(N=161), tuna (N=393), shell fish (N=765), and other (N=592) were included in calculations.^e Total fish = tuna + shell fish + other (broiled/baked)

LIBCSP = Long Island Breast Cancer Study Project

SD = standard deviation

N/A = not applicable

Table 2.Multivariable^a-adjusted ORs and 95% CI for the association between dietary PUFA intake and the risk of breast cancer in the LIBCSP, 1996-1997

PUFA	Quartile 1			Quartile 2				Quartile 3				Quartile 4			
	Co	Ca	OR	Co	Ca	OR	95% CI	Co	Ca	OR	95% CI	Co	Ca	OR	95% CI
Total PUFA	375	342	1.00	375	392	1.23	1.00, 1.52	375	386	1.27	1.02, 1.59	375	343	1.25	0.95, 1.63
Total ω-3	375	340	1.00	375	403	1.25	1.01, 1.54	375	377	1.23	0.98, 1.54	375	343	1.20	0.92, 1.58
ALA	375	335	1.00	375	390	1.23	0.99, 1.51	375	389	1.29	1.04, 1.61	375	349	1.25	0.96, 1.62
DPA	375	365	1.00	375	354	0.99	0.81, 1.22	375	375	1.08	0.88, 1.33	375	369	1.09	0.88, 1.36
DHA	375	372	1.00	375	336	0.91	0.74, 1.13	375	369	1.02	0.83, 1.26	375	386	1.06	0.86, 1.31
EPA	375	350	1.00	375	359	1.02	0.83, 1.25	375	365	1.08	0.88, 1.33	375	389	1.14	0.92, 1.40
Total ω-6	375	347	1.00	375	374	1.15	0.93, 1.42	375	405	1.31	1.05, 1.63	375	337	1.18	0.91, 1.55
LA	375	351	1.00	375	367	1.12	0.91, 1.38	375	407	1.30	1.05, 1.62	375	338	1.18	0.90, 1.54
AA	375	371	1.00	375	378	1.05	0.85, 1.29	375	367	1.03	0.83, 1.27	375	347	1.03	0.81, 1.29
ω-3/ω-6	375	360	1.00	375	384	1.09	0.89, 1.34	375	346	0.95	0.77, 1.17	375	373	0.99	0.80, 1.21

Note:

Co=Controls, Ca=Cases, LIBCSP=Long Island Breast Cancer Study Project

^a Multivariable-adjusted ORs and 95% CI adjusted for matching factor (5-year age group) and total energy intake (kcal/day)

Table 3.Multivariable^a-adjusted ORs and 95% CI for the additive interaction between dietary ω-3 and ω-6 (high and low intake) and the risk of breast cancer in the LIBCSP, 1996-1997

Model	Low ω-6 (< median)			High ω-6 (≥ median)			RERI ^b	95% CI ^c
	N	OR	95% CI	N	OR	95% CI		
High ω-3 (≥ median)	256	1.00		1,214	0.95	0.72, 1.26		
Low ω-3 (< median)	1,215	0.83	0.63, 1.09	278	1.20	0.85, 1.69	0.41	0.06, 0.76

Note:

Co=Controls, Ca=Cases, LIBCSP=Long Island Breast Cancer Study Project

^a Multivariable ORs and 95% CI adjusted for matching factor (5-year age group) and total energy intake (kcal/day)

^b RERI (Relative Excess Risk due to Interaction) = OR₁₁ - OR₁₀ - OR₀₁ +1

^c 95% CI for RERI estimated using Hosmer&Lemeshow[50]

Table 4. Multivariable^a-adjusted ORs and 95% CI for the multiplicative interaction between dietary ω -3 and ω -6 (high and low intake) and the risk of breast cancer in the LIBCSP, 1996-1997

Model	Low ω -6 ($<$ median)			High ω -6 ($\geq$ median)			LRT χ^2 ^b	p value
	N	OR	95% CI	N	OR	95% CI		
High ω -3 ($\geq$ median)	256	1.00		1,214	1.00			
Low ω -3 ($<$ median)	1,215	0.83	0.63, 1.09	278	1.26	0.96, 1.65	4.61	0.03

Note:

Co=Controls, Ca=Cases, LIBCSP=Long Island Breast Cancer Study Project, LRT=likelihood ratio test

^a Multivariable ORs and 95% CI adjusted for matching factor (5-year age group) and total energy intake (kcal/day)

^b LRT χ^2 calculated using nested models for the multiplicative interaction.

Table 5.Multivariable^a-adjusted ORs and 95% CI for the association between fish intake and the risk of breast cancer in the LIBCSP, 1996-1997

Fish	Never			Quartile 1				Quartile 2				Quartile 3				Quartile 4			
	Co	Ca	OR	Co	Ca	OR	95% CI	Co	Ca	OR	95% CI	Co	Ca	OR	95% CI	Co	Ca	OR	95% CI
Total fish intake	161	136	1.00	332	348	1.25	0.94, 1.67	335	323	1.13	0.84, 1.50	336	324	1.12	0.84, 1.50	336	332	1.19	0.89, 1.59
Tuna	393	343	1.00	256	241	1.08	0.85, 1.37	285	320	1.24	0.99, 1.55	141	141	1.12	0.84, 1.49	425	418	1.15	0.93, 1.42
Shell fish	765	750	1.00	126	102	0.78	0.58, 1.05	227	245	1.10	0.89, 1.38	178	169	0.99	0.78, 1.26	204	197	1.09	0.86, 1.38
Other fish (broiled/baked)	592	505	1.00	224	253	1.26	1.00, 1.58	87	104	1.38	1.00, 1.91	346	341	1.08	0.88, 1.32	251	260	1.18	0.94, 1.47

Note:
Co=Controls, Ca=Cases, LIBCSP = Long Island Breast Cancer Study Project
^a Multivariable-adjusted ORs and 95% CI adjusted for matching factor (5-year age group), total energy intake (kcal/day), and NSAID use.