Biological Control

Cold tolerance of *Trissolcus japonicus* and *T. cultratus*, potential biological control agents of *Halyomorpha halys*, the brown marmorated stink bug

Erica Nystrom Santacruz¹, Robert Venette², Christine Dieckhoff³, Kim Hoelmer³ and Robert L. Koch^{1,*}

¹Department of Entomology, University of Minnesota, 1980 Folwell Avenue, St. Paul, MN 55108, USA

²USDA-FS, Northern Research Station, 1990 Folwell Avenue, St. Paul, MN 55108, USA

³USDA-ARS, Beneficial Insect Introduction Research Laboratory, 501 South Chapel St., Newark, DE 19713, USA

*Corresponding author. Address: Department of Entomology, University of Minnesota, 1980 Folwell Avenue, St. Paul, MN 55108, USA. E-mail address: koch0125@umn.edu (R.L. Koch)

ABSTRACT

Halyomorpha halys (Stål) (Hemiptera: Pentatomidae) is native to Asia and has become a severe agricultural and nuisance pest in the U.S. Therefore, foreign exploration was conducted in Asia to identify potential classical biological control agents. Several *Trissolcus* spp. (Hymenoptera: Scelionidae) parasitize *H. halys* eggs in Asia and are being evaluated for potential release in the U.S. Since *H. halys* has invaded regions that experience sub-zero winter temperatures, cold tolerance is important for evaluation of *Trissolcus* spp. Our study compared the cold tolerance of populations of *T. japonicus* and *T. cultratus*, in order to assess relative suitability of the populations for release. We used thermocouple thermometry to determine the supercooling point and lower lethal temperature after brief exposure to cold temperature for each population. In addition, we subjected adult *T. japonicus* to a short photoperiod and low temperature regime, which increases cold tolerance in *H. halys*, to observe whether these conditions cause a change in cold tolerance in the parasitoid. We found that populations of both species froze and survived at colder temperatures than those reported for *H. halys*. In addition, there were no ecologically relevant differences in the temperature at which freezing or survival occurred among populations of either species, indicating that these populations are equally cold tolerant and suitable for introduction. Finally, *T. japonicus* does not acclimate by increasing cold tolerance in response to conditions that increase cold tolerance in *H. halys*, suggesting that the above-mentioned measures of cold tolerance are ecologically relevant.

KEY WORDS

Scelionidae, supercooling point, lower lethal temperature, geographic variation, longevity, acclimation

1. Introduction

Halyomorpha halys (Stål) (Hemiptera: Pentatomidae), the brown marmorated stink bug, is native to China, Japan, Korea, and Taiwan (Rider *et al.*, 2002). In North America, this invasive species was first detected in Pennsylvania in 1996, and by June 2016 had spread to 43 states in the U.S.A., and two Canadian provinces (Hoebeke and Carter, 2003; Northeastern IPM Center, 2016). *Halyomorpha halys* has also invaded several European countries and has been intercepted in New Zealand (Harris, 2010; Haye *et al.*, 2014). This highly polyphagous pest feeds on >100 species of plants in 45 families including fruits, vegetables, field crops, and ornamental plants in both native and invaded habitats (Haye *et al.*, 2014; Hoebeke and Carter, 2003; Lee *et al.*, 2013), and is considered a nuisance when aggregating en masse while overwintering inside man-made structures (Hoebeke and Carter, 2003; Inkley, 2012).

Some native North American insects have been identified as predators or parasitoids of *H. halys*, but are generally inconsistent in suppressing *H. halys* populations (Herlihy *et al.*, 2016; Jones *et al.*, 2014; Ogburn *et al.*, 2016). As part of a classical biological control program for *H. halys*, foreign exploration has been conducted in Asia since 2007 (K. Hoelmer, unpublished data) and four species of egg parasitoids in the genus *Trissolcus* (Hymenoptera: Scelionidae) have been identified as potential biological control agents. These species are being evaluated for potential release against *H. halys* in the U.S.A. (Talamas *et al.*, 2013). One of these species, *T. japonicus* (Ashmead), appears promising due to a short developmental time (10.5 d at 25 °C), multiple generations per year, highly female-biased sex ratio, and parasitism rates of 50-80% on *H. halys* in the native Asian range (Yang *et al.*, 2009). A second candidate species, *T. cultratus* (Mayr), is also under consideration, though parasitism rates are lower than those of *T. japonicus* (Haye *et al.*, 2015).

Low temperature is a limiting factor in the geographic distribution of many parasitoids (Hance *et al.*, 2007); thus, understanding the cold tolerance of a potential biological control agent can help estimate its capacity to survive and establish in novel areas (Daane *et al.*, 2012; Hanson *et al.*, 2013). In fact, successful classical biological control programs ensure that potential biological control agents and the targeted pests share similar responses to temperature (Goolsby *et al.*, 2005; Hoelmer and Kirk, 2005; Ulrichs and Hopper, 2008). Responses to temperature can vary among populations of the same species (Cira *et al.*, 2016; Feng *et al.*, 2014; Rochefort *et al.*, 2011), so it is prudent to compare geographically-distinct populations to identify which population may be better suited for introduction (e.g., Mausel *et al.*, 2011). Furthermore, because classical biological control programs depend on rearing individuals under laboratory settings for release, it is important to ensure that loss of overwintering potential over time due to exposure to long periods of non-diapause laboratory conditions does not occur (e.g., Gariepy *et al.*, 2015).

Commonly-used indices for quantifying cold tolerance in insects include the supercooling point (i.e., the lowest temperature reached prior to the release of latent heat of fusion, observed as an exotherm) and lower lethal temperature (i.e., the temperature at which brief exposure causes a desired level of mortality, often 50 or 90%) (Sinclair *et al.*, 2015). The relationship between freezing and mortality in insects can be used to determine one of three basic cold tolerance categories: chill-intolerance (i.e., death from cold without internal ice formation), freeze intolerance (i.e., death coincides with internal ice formation), or freeze tolerance (i.e. survival of freezing) (Lee, 2010). However, during the onset of fall and through the winter, these strategies may change as insects acclimatize to decreasing temperatures and photoperiods (Hefty, 2016). Acclimatization can result in an increase in cold tolerance over the course of a season (Crosthwaite *et al.*, 2011; Feng *et al.*, 2016; Udaka and Sinclair, 2014). Therefore, it is important

to try to quantify indices of cold tolerance after prolonged exposure to cold temperatures and shorter photoperiods to see if there are any changes over time.

Halyomorpha halys has invaded northern temperate regions (Garipey *et al.*, 2014; Koch, 2014) which experience winter temperatures below 0 °C. In fact, adult *H. halys* collected from Minnesota were found to be chill intolerant and to vary seasonally in their cold tolerance (Cira *et al.*, 2016). In response to colder temperatures and shorter photoperiods, *H. halys* will enter a reproductive diapause (Niva and Takeda, 2002). In contrast, little is known about the cold tolerance or overwintering behavior of *T. japonicus* or *T. cultratus*, except that *T. japonicus* overwinters as an adult (Yang *et al.*, 2009), presumably under bark. Aggregations of adult *Trissolcus* species have been reported during the winter under the bark of various trees in Japan (Y. Hirose, H. Honda, and K. Matsuo, personal communication) and Iran (Safavi, 1968). Furthermore, related parasitoids are known to overwinter as adults (Fathi *et al.*, 2011; Foerster and Nakama, 2002; James, 1988; Ryan *et al.*, 1981) and under bark (Fathi *et al.*, 2011). No information exists on subzero temperature limits for *T. japonicus* and *T. cultratus*, nor the extent of cold tolerance variation among populations of each species.

Adventive populations of *T. japonicus* were recently found in the wild in North America (Milnes *et al.*, 2016; Talamas *et al.*, 2015), so there is a need to determine whether *T. japonicus* can survive cold winter temperatures. To assist in assessing the establishment potential of *T. japonicus* and *T. cultratus* in North America we measured supercooling points and lower lethal temperatures for unacclimated, geographically-distinct populations of these species (e.g. Hanson *et al.*, 2013). In addition, we subjected adult *T. japonicus* to conditions that increase cold tolerance in *H. halys* in order to observe whether this induces a change in cold tolerance in the parasitoid.

2. Materials and Methods

2.1. Parasitoid acquisition and rearing

Trissolcus japonicus and *T. cultratus* populations were collected from four locations in Asia (Beijing and Nanjing, China; Pucheon, South Korea; and Tsukuba, Japan) between 2007 to 2012 (Table 1). Populations were reared at 20 °C, 16:8 h L:D, and 60-65% RH, and provided honey for nutrition and laboratory-reared *H. halys* eggs for reproduction at the U.S. Department of Agriculture (USDA), Agricultural Research Service, Beneficial Insects Introduction Research Unit, in Newark, Delaware. Separate egg masses of *H. halys* were parasitized by four populations of *T. japonicus* or two populations of *T. cultratus* with four parasitized egg masses of each population shipped overnight under USDA PPQ permit P526-14-01139 to a Biosecurity-Level-2 containment facility in Saint Paul, Minnesota during July 2014, and February and May 2015 for experiments (Table 1). At the Saint Paul location, some individuals of each species were provided with laboratory-reared *H. halys* eggs masses and reared under the conditions described above unless otherwise specified.

2.2. Cold tolerance experiments

The cold tolerance of adult *T. japonicus* and *T. cultratus* was evaluated separately in August 2014, and March and July 2015. The supercooling points and lower lethal temperatures of *T. japonicus*, and the lower lethal temperatures of *T. cultratus* were examined. Due to the limited number of individuals, supercooling points for *T. cultratus* were not measured. In order to facilitate comparison to the host *H. halys*, the cold tolerance testing methods generally followed those employed by Cira *et al.* (2016) for *H. halys*, with the substitution of a smaller thermocouple (e.g. Stephens *et al.*, 2015). Recently emerged (i.e., 12-36 h old) adult parasitoids

were used in all experiments, and the sex of each individual was determined after death by examination of the antennae (Yang *et al.*, 2009). All statistics were performed using R (R Core Development Team, 2015; RStudio Team, 2015), and all commands originate from the base package unless otherwise specified.

2.2.1. Supercooling point determination of *T. japonicus*

Methods for contact thermocouple thermometry followed Stephens *et al.* (2015). Adult *T. japonicus* were placed individually inside 1.5-ml microcentrifuge tubes. Each tube was sealed with a milled plastic dowel that had been fitted with an o-ring. A 0.127-mm copper-constantan thermocouple was threaded through a hole in the center of the dowel. Thermocouples with insects were placed inside $20 \times 20 \times 20$ cm polystyrene cubes calibrated to cool at a rate of 1 °C per minute when placed inside an ultralow freezer at -80 °C (Carrillo *et al.*, 2004). Cubes were cooled in groups (i.e., blocks) of 4 to 16 cubes and the temperatures of the insects were recorded every second by use of an analog data acquisition unit (USB-TC, Measurement Computing Core, Norton, MA), and visualized in real time on a computer using TracerDAQ Pro (Measurement Computing Core, Norton MA). For each individual, the supercooling point was recorded as the lowest temperature reached before the release of latent heat of fusion (i.e., an exotherm) (Carrillo *et al.*, 2004). For each population, 14-22 exotherms were detected (Table 1).

Regression analysis was used to compare the distributions of supercooling points among populations of *T. japonicus* because such distributions are often not normal (Sinclair *et al.*, 2015). Logistic regression was used to compare the cumulative proportion of individuals that began to freeze (i.e., an event) by a particular temperature among populations of *T. japonicus* (e.g., Stephens *et al.*, 2015). For this analysis a generalized linear model was created using the

glm() command with the probability of observing freezing as the response variable, and population, temperature, sex, and their interaction as explanatory variables. There was no effect of sex ($Z = -0.15$, $P = 0.88$), so this variable was dropped, and the data were pooled in the reduced model. To test for differences among populations in the temperature at which freezing occurred, intercepts (b_0) and slopes (b_1) among the four populations of *T. japonicus* were compared by changing the model reference population, and a Bonferroni adjustment (critical $\alpha = 0.008$ per comparison) was used to maintain an overall $\alpha = 0.05$. The dose.p() command from the MASS package (Ripley, 2015) followed the general equation: $P(\text{Insect freezes}) = 1/[1 + e^{-(b_0 + b_1 * \text{temperature})}]$, to identify the 50th percentile freezing temperature (i.e., $P(\text{Insect freezes}) = 0.5$) ($\pm$ SE) for each population.

2.2.2. Lower lethal temperature determination of *T. japonicus* and *T. cultratus*

To determine the lower lethal temperature of *T. japonicus* and *T. cultratus*, two experiments were performed (one for each species), and both experiments followed a randomized block design. For each population of a given species, individual adults were cooled to an assigned exposure temperature following the same protocol as the supercooling point experiments, except that individuals were removed from the freezer and cubes upon reaching the assigned exposure temperatures. Individuals were assigned randomly to exposure temperatures of -23, -21, -19, -17 °C, or a control of 25 °C. Previously-collected data suggested that mortality varied from 0-100% in this range. The number of individuals randomly assigned to each exposure temperature ranged from 12-22 individuals for each population of both species (Table 1). Up to 20 individuals were cooled in a block, for a total of 18 blocks for *T. japonicus* and 12 blocks for *T. cultratus*. Due to the limited supply of individuals and the inability to synchronize

emergence, it was not possible to represent each population and exposure temperature equally within a block; however, each population represented in a block included one control. Survivorship of each individual was visually determined 30 min after returning to room temperature and survival was defined as the parasitoid walking within the centrifuge tube. To determine the effect of low temperature exposure on longevity, individuals from the lower lethal temperature experiments were transferred in their centrifuge tubes to a growth chamber at 23 °C, 16:8 h L:D, and 60-65% RH, and were starved and monitored every 1-3 d until death.

To compare cumulative mortality as a function of exposure temperature among populations of each species, generalized linear models were prepared as described for the supercooling point analysis. However, for lower lethal temperature, an event was defined as whether an insect died by the exposure temperature. As before, sex was dropped from both models because the effect was not significant (*T. japonicus*: $Z = -1.21$, $P = 0.22$, and *T. cultratus*: $Z = -0.74$, $P = 0.46$), and the data were pooled in the reduced models. Testing for differences in the temperature at which mortality occurs among populations and identification of the 50th percentile mortality temperature ($\pm$ SE) for each population were determined as outlined for the supercooling point analysis.

To compare the effect of low temperature exposure on longevity among populations of each species, a parametric survival regression model was created for each species by using the `survreg()` command from the Survival package (Therneau, 2015). For these models, the response variable was the number of days an individual was alive, and individuals that died between monitored dates were censored at the last day that they were seen alive. The explanatory variables for each model consisted of population, temperature, and their interaction. Models were reduced with a maximum likelihood ratio test (χ^2) through use of the `anova()` command. The

predict() command was used to identify the median number of days until death ($\pm$ 95% CI) (i.e. longevity) for each combination of population and temperature. Tukey pairwise comparisons within a population were made using the glht() command from the multcomp package (Hothorn *et al.*, 2015). All reported p-values were adjusted at $\alpha = 0.05$.

2.2.3. Cold-tolerance strategy analysis of *T. japonicus*

To determine cold-tolerance strategies for the four populations of *T. japonicus*, cumulative mortality and cumulative freezing as a function of temperature were compared within a population as per Stephens *et al.* (2015). The cumulative mortality curve describes the total proportion of individuals that would be expected to die after brief exposure to a specified temperature; death may have occurred at a warmer temperature than the specified temperature. Similarly, the cumulative supercooling point curve describes the total proportion of individuals that would be expected to start freezing at a temperature greater than or equal to the specified temperature. This comparison was performed by creating generalized linear models as described for the supercooling point analysis for each population, but in this case the probability of observing an event (i.e., freezing or dying) as the response variable, and treatment (i.e., supercooling point or lower lethal temperature), temperature, and their interaction, as explanatory variables.

2.2.4. Cold acclimation experiment for *T. japonicus*

An experiment was performed to determine if temperature and photoperiod conditions cooler than those known to increase cold tolerance in *H. halys* (e.g., 20 °C, 12:12 h L:D, T. Cira, personal communication) could enhance the cold tolerance of *T. japonicus*. Adult parasitoids

from the Beijing, China population were received from Newark in May 2015 and maintained at summer conditions (23 °C, 16:8 h L:D, and 60-65% RH), for two generations. Recently emerged (i.e., 12-36 h old) adult males (n=1-3) and females (n=1-3) were transferred into eight separate (9 dram) plastic vials provisioned with honey and were allowed to mate for 24 h in the summer conditions. After 24 h, males were removed and the vials were evenly split between two sets of rearing conditions: summer or fall (18 °C, 12:12 h L:D, and 60-65% RH). Each vial was provisioned with a thawed *H. halys* egg mass (~25-28 eggs per egg mass) prepared following Haye *et al.* (2015). Six-24 h old *H. halys* egg masses were collected from a University of Minnesota (St. Paul, MN) laboratory colony (maintained at 23 °C, 16:8 h L:D, and 60-65% RH), and stored in a -80 °C freezer for up to four weeks until needed. The egg masses were thawed for 30 min at room temperature before use. After 48 h, females were removed from the vials and the egg masses (first generation, or F₁) were maintained in their respective rearing conditions until emergence. Upon emergence, the mating and oviposition process outlined above was repeated for two additional generations (F₂ and F₃), though individuals remained in their respective rearing conditions instead of completing these steps in the summer conditions.

Each generation of adult female parasitoids produced under the two rearing conditions were used in lower lethal temperature experiments similar to those described for the lower lethal temperature determination. However, exposure temperatures for these lower lethal temperature measurements spanned every degree between -25 to -14 °C, and a control of 25 °C and the number of individuals randomly assigned to each exposure temperature ranged from 4-6 individuals for each generation of both rearing conditions (Table 1). Again, up to 20 individuals were cooled in a block, for a total of 40 blocks.

Like the lower lethal temperature analyses, generalized linear models were created to compare cumulative mortality as a function of exposure temperatures between rearing conditions of the same generation, and among generations of a rearing condition. Three models were created to compare rearing conditions (summer or fall) within a generation (F₁, F₂, or F₃), where the explanatory variables included temperature, rearing condition, and their interaction. Two models were created to compare generations (P, F₁, F₂, and F₃) within each rearing condition (summer or fall), where the explanatory variables included temperature, generation, and the interaction. For the latter two models, testing for differences in the temperature at which mortality occurs among generations and identification of the 50th percentile mortality temperature ($\pm$ SE) for each generation was determined as outlined for the supercooling point analysis.

3. Results

3.1. Supercooling point of *T. japonicus*

Among all populations of *T. japonicus*, supercooling points ranged from -23.67 to -16.73 °C. The temperatures that caused 50% of the population to begin to freeze ($\pm$ SE) ranged from -22.04 $\pm$ 0.07 (Puceon) to -21.43 $\pm$ 0.19 °C (Nanjing) (Table 2). The distributions of supercooling points were not the same for each population. From the logistic regressions of cumulative percent of individuals freezing (i.e., supercooling) as a function of temperature for *T. japonicus*, the intercept (b_0) and slope (b_1) for Nanjing were significantly higher than the intercepts and slopes of Beijing and Puceon, but not significantly different from those of Tsukuba (critical α = 0.008) (Table 2, Supplemental Table 1). In general, more individuals from Nanjing began to freeze at warmer temperatures than individuals from Beijing or Puceon, but

not Tsukuba, and the change in the rate at which individuals began to freeze as temperatures declined was significantly greater for individuals from Nanjing than from Beijing or Pucheon, but not Tsukuba (Figure 1A-D).

3.2. Lower lethal temperature of *T. japonicus* and *T. cultratus*

For *T. japonicus*, the 50th percentile mortality temperature ($\pm$ SE) was lowest for the southernmost population of Nanjing (-20.03 ± 0.61 °C) and highest for the second northernmost population of Pucheon (-17.36 ± 0.84 °C) (Table 2). The 50th percentile mortality temperature ($\pm$ SE) for Tsukuba 2007 and Tsukuba 2012 of *T. cultratus* was -19.88 ± 0.46 °C and -19.86 ± 0.45 °C, respectively. In general, while mortality increased as individuals were exposed to decreasing temperature, the likelihood of mortality did not differ significantly among populations within either species (Figure 1A-F). Logistic regressions describing mortality as a function of temperature for *T. japonicus* did not have significantly different intercepts or slopes among populations (Table 2, Supplemental Table 1). Similarly, the intercepts (-11.28 ± 2.67 and -13.98 ± 3.39) and the slopes (-0.57 ± 0.13 , and -0.70 ± 0.17) for the Tsukuba 2007 and Tsukuba 2012 populations of *T. cultratus* did not differ significantly between collection years ($Z = -0.63$, $P = 0.53$ and $Z = -0.63$, $P = 0.53$, respectively).

For *T. japonicus* and *T. cultratus*, longevity for each population was generally significantly lower after exposure to low temperature (i.e., -23 and -21 °C), than to the control temperature (i.e., 25 °C) (Figure 2). Among all populations of *T. japonicus* and *T. cultratus*, median longevity after exposure to -23 and 25 °C ranged from 0.78 - 1.33 , and 4.28 - 6.44 d, respectively (Figure 2). Maximum likelihood ratio tests revealed that the interaction of population and temperature in the survival regressions comparing the effect of low temperature

exposure on longevity within a species was not significant in either model for *T. japonicus* ($\chi^2_3 = 3.32$, $P = 0.34$) nor *T. cultratus* ($\chi^2_1 < 0.002$, $P = 0.97$). Therefore, the relative differences in longevity among populations within a species were constant across all exposure temperatures. In fact, in the reduced model for *T. japonicus* where population was significant, the median longevity ($\pm$ 95% CI) for Nanjing was 1.50 (1.12, 2.00) days longer than Beijing ($Z = 2.75$, $P = 0.03$), 1.73 (1.32, 2.27) days longer than Pucheon ($Z = 3.92$, $P < 0.001$), and 1.98 (1.46, 2.69) days longer than Tsukuba ($Z = 4.37$, $P < 0.001$). A maximum likelihood ratio test revealed that population was not significant and therefore dropped from the reduced model for *T. cultratus* ($\chi^2_1 < 0.001$, $P = 0.97$), thus there were no differences in longevity between the two populations. In the final model for each species, temperature was significant for both *T. japonicus* ($Z = 6.98$, $P < 0.001$) and *T. cultratus* ($Z = 6.48$, $P < 0.001$), indicating that longevity within a population generally increased with increasing (i.e., warmer) exposure temperatures (Figure 2).

3.3. Cold-tolerance strategy of *T. japonicus*

From the comparisons of curves that described cumulative freezing and cumulative mortality within each population of *T. japonicus*, mortality appeared to precede freezing as temperature declined (Figure 1A-D). For each population, the slope and intercept of the logistic regression model characterized the cumulative frequency of supercooling points. In the logistic regression models, intercepts for supercooling points intercepts and slopes were all significantly lower than the lower lethal temperature intercepts and slopes (Table 2).

3.4. Cold acclimation of *T. japonicus*

From the logistic regressions of cumulative percent of individuals dying as a function of temperature, there were no significant differences among the intercepts and slopes of generations within rearing conditions or rearing conditions within a generation (Table 3, Supplemental Table 2). The 50th percentile mortality temperature ($\pm$ SE) for the summer and fall conditions were similar, spanning from -17.95 ± 0.50 to -15.44 ± 0.91 °C (Table 3). Therefore, as temperature decreased, mortality increased at the same rate among all generations within a rearing condition, as well as between rearing conditions for a given generation (Figure 3A-D). Therefore, exposure over three generations to conditions known to induce reproductive diapause in *H. halys* did not affect the likelihood of survival of *T. japonicus*.

4. Discussion

Classical biological control by Asian *Trissolcus* spp. may provide an additional tool for managing *H. halys* beyond North American natural enemies (Herlihy *et al.*, 2016; Ogburn *et al.*, 2016) and the current use of pesticides (Leskey *et al.*, 2012). Recently, populations of *T. japonicus* were detected in both the eastern and western United States (Milnes *et al.*, 2016; Talamas *et al.*, 2015). However, these areas currently represent a fraction of the range of *H. halys* in North America (Northeastern IPM Center, 2016), and do not reflect the northern extent of the invaded range of this pest, which seasonally experiences cold, winter temperatures (USDA, 2012). The present research is the first assessment of cold tolerance for *T. japonicus* and *T. cultratus*, and will facilitate evaluation of classical biological control for *H. halys*.

Many insects (Crosthwaite *et al.*, 2011; Kim and Song, 2000; Régnière and Bentz, 2007), including parasitic Hymenoptera (Hanson *et al.*, 2013), are known to seasonally acclimatize to cold conditions by increasing cold tolerance as a response to environmental conditions such as

photoperiod or temperature (Denlinger, 1991). Such seasonal acclimatization could affect ecological relevance of cold tolerance estimates attained from unacclimated, laboratory populations. Whether *T. japonicus* seasonally acclimatizes to colder temperatures or shorter photoperiod was unknown to date. Therefore, we began our investigation with unacclimated individuals that were reared under standard conditions (23 ± 1.5 °C, 16:8 h L: D, 65% RH) to determine a reference point for comparing populations reared under cooler temperature or shorter photoperiod conditions.

Interestingly, median supercooling point and lower lethal temperature results for unacclimated populations of *T. japonicus* and *T. cultratus* were all below the winter mean supercooling point of *H. halys* from Minnesota (-17.06 °C), an area that experiences severe winter temperatures (Cira *et al.*, 2016). In addition, prolonged exposure to laboratory conditions for *T. cultratus* (e.g. Tsukuba 2007) did not appear to affect the baseline cold tolerance when compared to a population collected five years later (e.g. Tsukuba 2012). This suggests that both parasitoid species, when unacclimated, are capable of surviving colder temperatures than their acclimated host, and that this unacclimated cold tolerance is retained during laboratory rearing. Such results are promising for potential long-term laboratory populations reared for classical biological control. Nevertheless, we evaluated lower lethal temperature of *T. japonicus* under more relevant conditions by performing an experiment that investigated the ability of *T. japonicus* to modify its cold tolerance in response to conditions known to increase in cold tolerance in *H. halys*. Our results suggest that unlike *H. halys*, exposure to conditions cooler than 20 °C and a photoperiod of 12:12 h does not increase the cold tolerance *T. japonicus*. Parasitic wasps are dependent on their hosts and for most species synchrony in behavior to that of their host is crucial to their survivorship (Hance *et al.*, 2007). *Trissolcus japonicus* has been recorded

from several other Asian pentatomid species (Matsuo *et al.*, 2016; Yang *et al.*, 2009), and it overwinters as an adult (versus inside of the host) (Yang *et al.*, 2009). This lack of dependency on a single host species and behavior of overwintering outside of the host may in part explain why *T. japonicus* did not respond to the same photoperiod cues as *H. halys*. However, our study does not rule out the possibility that *T. japonicus* can further adjust its cold tolerance if exposed to other conditions. Further work should explore how *T. japonicus* responds to temperature or photoperiod regimes beyond those used in this study.

Supercooling points of *T. japonicus* were not the same for all source populations. Populations originating from the southernmost population (i.e., Nanjing) began to freeze at significantly higher temperatures than the northernmost populations (i.e., Beijing and Pucheon). This pattern aligns with the latitude and 30-year average lowest temperature from January for the source population areas in Asia (Table 1). Similar correlations have also been found with other studies that demonstrate that the mean supercooling points of geographically-distinct populations of a single insect species increase in temperature as latitude decreases (Jing and Kang, 2003; Zhou *et al.*, 2011). However, the small magnitude of difference in median freezing values among the populations tested here is not likely to confer an ecologically-relevant advantage for the populations that supercooled at statistically lower temperatures. Therefore, from the standpoint of using supercooling point as an index of cold tolerance, these four populations of *T. japonicus* should have similar likelihoods for freezing at cold winter temperatures.

We did not find significant differences in mortality as a function of exposure temperature among any of the populations of *T. japonicus* or *T. cultratus*. This suggests that all populations should have similar likelihoods for surviving acute exposures to cold and that all of the populations are equally suitable for introduction with respect to winter survival. Because

freezing and mortality can occur at significantly different temperatures (Sømme, 1982), it is important to assess the relationship between these two phenomena in order to determine whether supercooling point and lower lethal temperature are interchangeable cold-tolerance indices for mortality (Renault *et al.*, 2002). Our results demonstrate that each of the populations of *T. japonicus* appeared to be dying before freezing. Thus, lower lethal temperature is a more ecologically-relevant source of information than the supercooling point for assessing the cold tolerance of *T. japonicus* (e.g. Baust and Rojas, 1985), and should be used for more accurate comparisons of *T. japonicus* cold tolerance to that of *H. halys*. Although informative, our study is limited in that we investigated survivorship after instantaneous exposure to low temperatures (e.g. lower lethal temperature). This deliberate measurement was used to decouple low temperature and exposure time, as duration of exposure to a low temperature can differentially affect survivorship (Renault *et al.*, 2002). Further cold tolerance studies with *T. japonicus* should expand on these initial measures of lower lethal temperature by incorporating the element of time (e.g. lower lethal time). This could help assess how survivorship of this species is affected under conditions that more closely simulate what the insects may encounter in nature.

Comparison of the temperatures at which insects begin to freeze or die also can be used to infer cold tolerance strategy (e.g. Hanson and Venette, 2013), and those that die from cold temperatures before body fluids begin to freeze are considered to be chill-intolerant (Lee, 2010). Our results demonstrate that *T. japonicus* is chill-intolerant, which is a cold-tolerance strategy shared by other parasitoids (Carrillo *et al.*, 2005; Hanson *et al.*, 2013; Rivers *et al.*, 2000), as well as by *H. halys* (Cira *et al.*, 2016). Therefore, similar to *H. halys*, populations of *T. japonicus* are freezing at temperatures significantly lower than what they could survive.

Interestingly, while we did not find differences among populations of either parasitoid in lower lethal temperature, we did find that the southernmost population, Nanjing, lived longer without food after brief exposure to all low temperatures in comparison with the other *T. japonicus* populations. This increase of approximately two days in longevity was relatively small, but it could provide the individuals from this population with additional time to seek nutrition during the spring, when food is not yet available or scarce. Little is known about the natural feeding habits of *T. japonicus* except that it does not host-feed; however, if it is similar to that of other scelionids, this species likely also feeds on a flower nectar and insect honeydew (Orr, 1988; Safavi, 1968). Other studies have documented that longevity decreases as a result of prolonged exposure to decreasing temperatures (Colinet *et al.*, 2006; Langer and Hance, 2000; Lysyk, 2004). Our results instead provide evidence of an effect on longevity after brief cold exposure, where individuals of *T. japonicus* and *T. cultratus* that are able to survive brief exposure to subzero temperatures do so at the expense of their longevity. Such sublethal trade-offs could have negative implications for establishment success of *T. japonicus* or *T. cultratus* in areas that experience even brief subzero temperatures (e.g. Bale, 2002).

Localized detections of *T. japonicus* in the United States over consecutive summers provide evidence that the parasitoid is able to successfully overwinter or re-colonize here in the United States. In Winchester, Virginia, the coldest area in the United States where *T. japonicus* was found for multiple summers (2015 and 2016, K. Hoelmer, unpublished data), the average annual extreme minimum temperature ranges from -20.6 to -17.8°C (USDA, 2012). Based on our laboratory experiments, in which the insects were in direct contact with low temperature, this range of temperatures caused mortality in approximately 25-91% of the laboratory populations of *T. japonicus*. However, some insects seek overwintering habitats that buffer extreme cold as

measured by ambient temperature (Inkley, 2012; Koch *et al.*, 2004; Vermunt *et al.*, 2012). Therefore, it is possible that our predictions underestimate survivorship in nature because our methodology does not account for potential thermal buffering provided by an overwintering habitat. *Trissolcus japonicus* may overwinter as an adult below *Zelkova* spp. tree bark (Matsuo, H. Honda and Y. Hirose, unpublished data), which is a similar behavior to that of its host, *H. halys* (Lee *et al.*, 2014), and other *Trissolcus* spp. (Safavi, 1968). It is likely that during this overwintering period *T. japonicus* adults experience some degree of thermal buffering to ambient air temperatures, which could improve their ability to survive in these areas (e.g. Turnock and Fields, 2005). Field studies targeting the adventive populations of *T. japonicus* in the United States could help investigate the overwintering behavior of the wasps and the degree of thermal buffering provided by bark. In addition, the adventive populations provide a unique opportunity to compare cold tolerance measurements between laboratory and field populations.

As *H. halys* has established in areas that experience cold temperatures, effective biological control agents must be able to withstand similar conditions. Based on our comparison of cold tolerance indices for geographically-distinct populations of two *Trissolcus* spp., it appears that the populations are equally suited for survival in these areas. In regards to assessing the potential for classical biological control of *H. halys*, these *Trissolcus* spp. appear more cold tolerant than their host and therefore may not be limited by cold in areas that *H. halys* has invaded. However, if microclimate of specific overwintering habitat is not accounted for, resulting estimates of potential geographic range will be conservative. Therefore, to refine estimates of the geographic range of these species, further work is needed to determine the overwintering habitats of these *Trissolcus* spp. and to quantify cold exposure in such habitats.

Finally, arthropod overwintering success is contingent on a number of factors beyond temperature stress. In addition, arthropods employ a number of mechanisms to survive cold exposure including (but not limited to), cryoprotectant and antifreeze protein synthesis, membrane restructuring, adjustments to ion transports, upregulation of heat shock proteins, etc. (Teets and Denlinger, 2013). Further studies with *T. japonicus* or *T. cultratus* should examine additional factors affecting overwintering potential (e.g., Colinet *et al.*, 2007), or the mechanisms that drive that success (e.g., Foray *et al.*, 2013).

Acknowledgements

We would like to thank Theresa Cira, Paul Botch, and Trisha Samota for guidance with maintaining populations of *Trissolcus* spp, and *H. halys*; and Kathleen Tatman, Jaana Iverson, and James Menger for technical support. We thank Stephanie Dahl, Jeanne Ciborowski, and Nik Prenevost for facilitating research in the Minnesota Agricultural Experiment Station - Biosecurity Facility, and Amy Morey, Mark Asplen, and Luke Skinner for comments on a previous version of this manuscript. This work was supported by the Minnesota Environment and Natural Resources Trust Fund, the Minnesota Soybean Research and Promotion Council, and the U.S. Department of Agriculture – National Institute of Food and Agriculture – Specialty Crop Research Initiative (USDA–NIFA–SCRI) #2011-51181-30937. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the USDA. USDA is an equal opportunity provider and employer.

References

- Bale, J.S., 2002. Insects and low temperatures: from molecular biology to distributions and abundance. *Philos. Trans. R. Soc. London B* 357, 849–62. doi:10.1098/rstb.2002.1074
- Baust, J.G., Rojas, R.R., 1985. Review—Insect cold hardiness: Facts and fancy. *J. Insect Physiol.* 31, 755–759. doi:10.1016/0022-1910(85)90067-8
- Carrillo, M.A., Heimpel, G.E., Moon, R.D., Cannon, C.A., Hutchison, W.D., 2005. Cold hardiness of *Habrobracon hebetor* (Say) (Hymenoptera: Braconidae), a parasitoid of pyralid moths. *J. Insect Physiol.* 51, 759–768. doi:10.1016/j.jinsphys.2005.03.006
- Carrillo, M.A., Kaliyan, N., Cannon, C.A., Morey, R.V., Wilcke, W.F., 2004. A simple method to adjust cooling rates for supercooling point determination. *CryoLetters* 25, 155–160.
- Cira, T.M., Venette, R.C., Aigner, J., Kuhar, T., Mullins, D.E., Gabbert, S.E., Hutchison, W.D., 2016. Cold tolerance of *Halyomorpha halys* (Hemiptera: Pentatomidae) across geographic and temporal scales. *Environ. Entomol.* 45, 484–491. doi:10.1093/ee/nvv220
- Colinet, H., Hance, T., Vernon, P., 2006. Water relations, fat reserves, survival, and longevity of a cold-exposed parasitic wasp *Aphidius colemani* (Hymenoptera: Aphidiinae). *Environmental Entomol.* 35, 228–236. doi:10.1603/0046-225X-35.2.228
- Colinet, H., Vernon, P., Hance, T., 2007. Does thermal-related plasticity in size and fat reserves influence supercooling abilities and cold-tolerance in *Aphidius colemani* (Hymenoptera: Aphidiinae) mummies? *J. Therm. Biol.* 32, 374–382. doi:10.1016/j.jtherbio.2007.03.005
- Crosthwaite, J.C., Sobek, S., Lyons, D.B., Bernards, M.A., Sinclair, B.J., 2011. The overwintering physiology of the emerald ash borer, *Agrilus planipennis* Fairmaire (Coleoptera: Buprestidae). *J. Insect Physiol.* 57, 166–73. doi:10.1016/j.jinsphys.2010.11.003
- Daane, K.M., Wang, X., Johnson, M.W., Cooper, M.L., 2012. Low temperature storage effects on two olive fruit fly parasitoids. *BioControl* 58, 175–185. doi:10.1007/s10526-012-9481-z
- Denlinger, D.L., 1991. Relationship between cold hardiness and diapause, in: Lee, R.E., Denlinger, D.L. (Eds.), *Insects at Low Temperature*. Chapman & Hall, New York, NY, pp. 174–198. doi:10.1007/978-1-4757-0190-6
- Fathi, S.A.A., Bakhshizadeh, N., Michaud, J.P., 2011. Spatial and temporal correspondence of sunn pest oviposition and egg parasitism by *Trissolcus* spp. in Iran. *J. Appl. Entomol.* 135, 666–672. doi:10.1111/j.1439-0418.2010.01607.x
- Feng, Y., Xu, L., Li, W., Xu, Z., Cao, M., Wang, J., Tao, J., Zong, S., 2016. Seasonal changes in supercooling capacity and major cryoprotectants of overwintering Asian longhorned beetle (*Anoplophora glabripennis*) larvae. *Agric. For. Entomol.* 18, 302–312. doi:10.1111/afe.12162
- Feng, Y., Xu, L., Tian, B., Tao, J., Wang, J., Zong, S., 2014. Cold hardiness of Asian longhorned beetle (Coleoptera: Cerambycidae) larvae in different populations. *Environ. Entomol.* 43, 1419–1426. doi:10.1603/EN13352
- Foerster, L.A., Nakama, P.A., 2002. Efeito da estocagem em baixa temperatura na capacidade reprodutiva e longevidade de *Trissolcus basalis* (Wollaston) e *Telenomus podisi* Ashmead (Hymenoptera: Scelionidae). *Neotrop. Entomol.* 31, 115–120. doi:10.1590/S1519-566X2002000100016
- Foray, V., Desouhant, E., Voituron, Y., Larvor, V., Renault, D., Colinet, H., Gibert, P., 2013. Does cold tolerance plasticity correlate with the thermal environment and metabolic profiles of a parasitoid wasp? *Comp. Biochem. Physiol. - A Mol. Integr. Physiol.* 164, 77–83.

- doi:10.1016/j.cbpa.2012.10.018
- Garipey, T.D., Fraser, H., Scott-Dupree, C.D., 2014. Brown marmorated stink bug (Hemiptera: Pentatomidae) in Canada: recent establishment, occurrence, and pest status in southern Ontario. *Can. Entomol.* 146, 579–582. doi:10.4039/tce.2014.4
- Garipey, V., Boivin, G., Brodeur, J., 2015. Why two species of parasitoids showed promise in the laboratory but failed to control the soybean aphid under field conditions. *Biol. Control* 80, 1–7. doi:10.1016/j.biocontrol.2014.09.006
- Goolsby, J.A., Debarro, P.J., Kirk, A.A., Sutherst, R.W., Canas, L., Ciomperlik, M.A., Ellsworth, P.C., Gould, J.R., Hartley, D.M., Hoelmer, K.A., Naranjo, S.E., Rose, M., Roltsch, W.J., Ruiz, R.A., Pickett, C.H., Vacek, D.C., 2005. Post-release evaluation of biological control of *Bemisia tabaci* biotype “B” in the USA and the development of predictive tools to guide introductions for other countries. *Biol. Control* 32, 70–77. doi:10.1016/j.biocontrol.2004.07.012
- Hance, T., van Baaren, J., Vernon, P., Boivin, G., 2007. Impact of extreme temperatures on parasitoids in a climate change perspective. *Annu. Rev. Entomol.* 52, 107–26. doi:10.1146/annurev.ento.52.110405.091333
- Hanson, A.A., Venette, R.C., 2013. Thermocouple Design for Measuring Temperatures of Small Insects. *CryoLetters* 34, 261–266.
- Hanson, A.A., Venette, R.C., Lelito, J.P., 2013. Cold tolerance of Chinese emerald ash borer parasitoids: *Spathius agrili* Yang (Hymenoptera: Braconidae), *Tetrastichus planipennisi* Yang (Hymenoptera: Eulophidae), and *Oobius agrili* Zhang and Huang (Hymenoptera: Encyrtidae). *Biol. Control* 67, 516–529. doi:10.1016/j.biocontrol.2013.08.015
- Harris, A.C., 2010. *Halyomorpha halys* (Hemiptera: Pentatomidae) and *Protaetia brevitarsis* (Coleoptera: Scarabaeidae: Cetoniinae) intercepted in Dunedin. *Weta* 40, 42–44.
- Haye, T., Fischer, S., Zhang, J., Garipey, T., 2015. Can native egg parasitoids adopt the invasive brown marmorated stink bug, *Halyomorpha halys* (Heteroptera: Pentatomidae), in Europe? *J. Pest Sci.* (2004). 88, 693–705. doi:10.1007/s10340-015-0671-1
- Haye, T., Wyniger, D., Garipey, T., 2014. Recent range expansion of brown marmorated stink bug in Europe, in: Muller, G., Pospischi, R., Robinson, W.H. (Eds.), *Proceedings of the 8th International Conference on Urban Pests*, 20–23 July. Zurich, Switzerland, pp. 309–314.
- Hefty, A.R., 2016. Risk of invasion by walnut twig beetle throughout eastern North America. University of Minnesota.
- Herlihy, M. V., Talamas, E.J., Weber, D.C., 2016. Attack and Success of Native and Exotic Parasitoids on Eggs of *Halyomorpha halys* in Three Maryland Habitats. *PLoS One* 11, e0150275. doi:10.1371/journal.pone.0150275
- Hoebeke, E.R., Carter, M.E., 2003. *Halyomorpha halys* (Stål) (Heteroptera: Pentatomidae): a polyphagous plant pest from Asia newly detected in North America. *Am. Proc. Entomol. Soc. Washingt.* 105, 225–237.
- Hoelmer, K.A., Kirk, A.A., 2005. Selecting arthropod biological control agents against arthropod pests: Can the science be improved to decrease the risk of releasing ineffective agents? *Biol. Control* 34, 255–264. doi:10.1016/j.biocontrol.2005.05.001
- Hothorn, T., Bretz, F., Westfall, P., 2015. Multcomp: Simultaneous inference in general parametric models [WWW Document]. URL <https://cran.r-project.org/web/packages/multcomp/index.html> (accessed 7.7.16).
- Inkley, D.B., 2012. Characteristics of home invasion by the brown marmorated stink bug (Hemiptera: Pentatomidae). *J. Entomol. Sci.* 47, 125–130.

- doi:<http://dx.doi.org/10.18474/0749-8004-47.2.125>
- James, D.G., 1988. Fecundity, longevity and overwintering of *Trissolcus biproruli* Girault (Hymenoptera: Scelionidae) a parasitoid of *Biprorulus bibax* Breddin (Hemiptera: Pentatomidae). *Aust. J. Entomol.* 27, 297–301. doi:10.1111/j.1440-6055.1988.tb01177.x
- Jing, X.-H., Kang, L., 2003. Geographical variation in egg cold hardiness: a study on the adaptation strategies of the migratory locust *Locusta migratoria* L. *Ecol. Entomol.* 28, 151–158. doi:10.1046/j.1365-2311.2003.00497.x
- Jones, A.L., Jennings, D.E., Hooks, C.R.R., Shrewsbury, P.M., 2014. Sentinel eggs underestimate rates of parasitism of the exotic brown marmorated stink bug, *Halyomorpha halys*. *Biol. Control* 78, 61–66. doi:10.1016/j.biocontrol.2014.07.011
- Kim, Y., Song, W., 2000. Effect of thermoperiod and photoperiod on cold tolerance of *Spodoptera exigua* (Lepidoptera: Noctuidae). *Environ. Entomol.* 29, 868–873. doi:10.1603/0046-225X-29.5.868
- Koch, R.L., 2014. Detections of the brown marmorated stink bug (Hemiptera: Pentatomidae) in Minnesota. *J. Entomol. Sci.* 49, 313–317. doi:10.18474/0749-8004-49.3.313
- Koch, R.L., Carrillo, M.A., Venette, R.C., Cannon, C.A., Hutchison, W.D., 2004. Cold hardiness of the multicolored Asian lady beetle (Coleoptera: Coccinellidae). *Environ. Entomol.* 33, 815–822. doi:10.1603/0046-225X-33.4.815
- Langer, A., Hance, T., 2000. Overwintering strategies and cold hardiness of two aphid parasitoid species (Hymenoptera: Braconidae: Aphidiinae). *J. Insect Physiol.* 46, 671–676. doi:10.1016/S0022-1910(99)00155-9
- Lee, D., Short, B.D., Joseph, S. V., Bergh, J.C., Leskey, T.C., 2013. Review of the biology, ecology, and management of *Halyomorpha halys* (Hemiptera: Pentatomidae) in China, Japan, and the Republic of Korea. *Environ. Entomol.* 42, 627–641. doi:10.1603/EN13006
- Lee, D.-H., Cullum, J.P., Anderson, J.L., Daugherty, J.L., Beckett, L.M., Leskey, T.C., 2014. Characterization of overwintering sites of the invasive brown marmorated stink bug in natural landscapes using human surveyors and detector canines. *PLoS One* 9, 1–9. doi:10.1371/journal.pone.0091575
- Lee, R.E., 2010. A primer on insect cold-tolerance, in: Denlinger, D.J., Lee, R.E. (Eds.), *Low Temperature Biology of Insects*. Cambridge University Press, New York, NY, pp. 3–34.
- Leskey, T.C., Short, B.D., Butler, B.R., Wright, S.E., 2012. Impact of the invasive brown marmorated stink bug, *Halyomorpha halys* (Stål), in mid-Atlantic tree fruit orchards in the United States: case studies of commercial management. *Psyche A J. Entomol.* 2012, 1–14. doi:10.1155/2012/535062
- Lysyk, T.J., 2004. Effects of cold storage on development and survival of three species of parasitoids (Hymenoptera: Pteromalidae) of house fly, *Musca domestica* L. *Environ. Entomol.* 33, 823–831. doi:10.1603/0046-225X-33.4.823
- Matsuo, K., Honda, T., Itoyama, K., Toyama, M., Hirose, Y., 2016. Discovery of three egg parasitoid species attacking the shield bug *Glaucias subpunctatus* (Hemiptera: Pentatomidae). *Japanese J. Appl. Entomol. Zool.* 60, 43–45. doi:10.1303/jjaez.2016.43
- Mausel, D.L., Van Driesche, R.G., Elkinton, J.S., 2011. Comparative cold tolerance and climate matching of coastal and inland *Laricobius nigrinus* (Coleoptera: Derodontidae), a biological control agent of hemlock woolly adelgid. *Biol. Control* 58, 96–102. doi:10.1016/j.biocontrol.2011.04.004
- Milnes, J.M., Wiman, N.G., Talamas, E.J., Brunner, J.F., Hoelmer, K.A., Buffington, M.L., Beers, E.H., 2016. Discovery of an exotic egg parasitoid of the brown marmorated stink

- bug, *Halyomorpha halys* (Stål) in the Pacific Northwest. *Proc. Entomol. Soc. Washingt.* 118.
- Niva, C.C., Takeda, M., 2002. Color changes in *Halyomorpha brevis* (Heteroptera: Pentatomidae) correlated with distribution of pteridines: regulation by environmental and physiological factors. *Comp. Biochem. Physiol. Part B Biochem. Mol. Biol.* 132, 653–660. doi:10.1016/S1096-4959(02)00081-7
- Northeastern IPM Center, 2016. State-by-State [WWW Document]. URL <http://www.stopbmsb.org/where-is-bmsb/state-by-state/> (accessed 7.7.16).
- Ogburn, E., Bessin, R., Dieckhoff, C., Dobson, R., Grieshop, M., Hoelmer, K., Matthews, C., Moore, J., Nielsen, A.L., Poley, K., Pote, J., Rogers, M., Welty, C., Walgenbach, J., 2016. Natural enemy impact on the invasive brown marmorated stink bug, *Halyomorpha halys* (Stål) (Hemiptera: Pentatomidae), in organic agroecosystems: A regional assessment. *Biol. Control* 101, 39–51. doi:10.1016/j.biocontrol.2016.06.002
- Orr, D.B., 1988. Scelionid wasps as biological control agents: A review. *Florida Entomol.* 71, 506–528. doi:10.2307/3495011
- R Core Development Team, 2015. R: A language and environment for statistical computing [WWW Document]. R Found. Stat. Comput. URL <http://www.r-project.org> (accessed 7.7.16).
- Régnière, J., Bentz, B., 2007. Modeling cold tolerance in the mountain pine beetle, *Dendroctonus ponderosae*. *J. Insect Physiol.* 53, 559–72. doi:10.1016/j.jinsphys.2007.02.007
- Renault, D., Salin, C., Vannier, G., Vernon, P., 2002. Survival at low temperatures in insects: What is the ecological significance of the supercooling point? *CryoLetters* 23, 217–228.
- Rider, D.A., Zheng, L.Y., Kerzhner, I.M., 2002. Checklist and nomenclatural notes on the Chinese Pentatomidae (Heteroptera). II. Pentatominae. *Zoosystematica Ross.* 11, 135–153.
- Ripley, B., 2015. MASS: Support functions and datasets for venables and Ripley's MASS [WWW Document]. URL <https://cran.r-project.org/web/packages/MASS/index.html> (accessed 7.7.16).
- Rivers, D.B., Lee, R.E., Denlinger, D.L., 2000. Cold hardiness of the fly pupal parasitoid *Nasonia vitripennis* is enhanced by its host *Sarcophaga crassipalpis*. *J. Insect Physiol.* 46, 99–106. doi:10.1016/S0022-1910(99)00106-7
- Rocheft, S., Berthiaume, R., Hébert, C., Charest, M., Bause, É., 2011. Effect of temperature and host tree on cold hardiness of hemlock looper eggs along a latitudinal gradient. *J. Insect Physiol.* 57, 751–759. doi:10.1016/j.jinsphys.2011.02.013
- RStudio Team, 2015. RStudio: Integrated development environment for R [WWW Document]. RStudio, Inc. URL <http://www.rstudio.com> (accessed 7.7.16).
- Ryan, R.B., Mortensen, R.W., Torgersen, T.R., 1981. Reproductive biology of *Telenomus californicus* Ashmead, an egg parasite of the Douglas-fir tussock moth: laboratory studies. *Ann. Entomol. Soc. Am.* 74, 213–216. doi:10.1093/aesa/74.2.213
- Safavi, M., 1968. Chapitre 4. Biologie et écologie des *Trissolcus* in Iran. A. Développement des *Trissolcus*, in: *Etude Biologique et Ecologique Des Hymenopteres Parasites Des Oeufs Des Punaises Des Cereales*. Entomophaga, pp. 381–495.
- Sinclair, B.J., Coello Alvarado, L.E., Ferguson, L. V., 2015. An invitation to measure insect cold tolerance: Methods, approaches, and workflow. *J. Therm. Biol.* 53, 180–197. doi:10.1016/j.jtherbio.2015.11.003
- Sømme, L., 1982. Supercooling and winter survival in terrestrial arthropods. *Comp. Biochem.*

- Physiol. Part A Physiol. 73, 519–543. doi:10.1016/0300-9629(82)90260-2
- Stephens, A.R., Asplen, M.K., Hutchison, W.D., Venette, R.C., 2015. Cold hardiness of winter-acclimated *Drosophila suzukii* (Diptera: Drosophilidae) adults. Environ. Entomol. 1–8. doi:10.1093/ee/nvv134
- Talamas, E.J., Buffington, M., Hoelmer, K., 2013. New synonymy of *Trissolcus halyomorphae* Yang. J. Hymenopt. Res. 33, 113–117. doi:10.3897/JHR.33.5627
- Talamas, E.J., Herlihy, M. V., Dieckhoff, C., Hoelmer, K.A., Buffington, M., Bon, M.-C., Weber, D.C., 2015. *Trissolcus japonicus* (Ashmead) (Hymenoptera, Scelionidae) emerges in North America. J. Hymenopt. Res. 43, 119–128. doi:10.3897/JHR.43.4661
- Teets, N.M., Denlinger, D.L., 2013. Physiological mechanisms of seasonal and rapid cold-hardening in insects. Physiol. Entomol. 38, 105–116. doi:10.1111/phen.12019
- Therneau, T.M., 2015. Survival: Survival analysis [WWW Document]. URL <https://cran.r-project.org/web/packages/survival/index.html> (accessed 7.7.16).
- Turnock, W.J., Fields, P.G., 2005. Winter climates and coldhardiness in terrestrial insects. Eur. J. Entomol. 102, 561–576. doi:10.14411/eje.2005.081
- Udaka, H., Sinclair, B.J., 2014. The overwintering biology of the acorn weevil, *Curculio glandium* in southwestern Ontario. J. Therm. Biol. 44, 103–109. doi:10.1016/j.jtherbio.2014.02.019
- Ulrichs, C., Hopper, K.R., 2008. Predicting insect distributions from climate and habitat data. BioControl 53, 881–894. doi:10.1007/s10526-007-9143-8
- USDA, 2012. Plant hardiness zone map [WWW Document]. URL <http://planthardiness.ars.usda.gov/PHZMWeb/> (accessed 7.7.16).
- Vermunt, B., Cuddington, K., Sobek-Swant, S., Crosthwaite, J., 2012. Cold temperature and emerald ash borer: Modelling the minimum under-bark temperature of ash trees in Canada. Ecol. Modell. 235–236, 19–25. doi:10.1016/j.ecolmodel.2012.03.033
- Yang, Z., Yao, Y., Qiu, L., Li, Z., 2009. A new species of *Trissolcus* (Hymenoptera: Scelionidae) parasitizing eggs of *Halyomorpha halys* (Heteroptera: Pentatomidae) in China with comments on its biology. Ann. Entomol. Soc. Am. 102, 39–47. doi:10.1603/008.102.0104
- Zhou, Z.S., Guo, J.Y., Michaud, J.P., Li, M., Wan, F.H., 2011. Variation in cold hardiness among geographic populations of the ragweed beetle, *Ophraella communa* LeSage (Coleoptera: Chrysomelidae), a biological control agent of *Ambrosia artemisiifolia* L. (Asterales: Asteraceae), in China. Biol. Invasions 13, 659–667. doi:10.1007/s10530-010-9857-x

Table 1

Source information for populations of *T. japonicus* and *T. cultratus* used to measure lower lethal temperatures (LLTemp) and supercooling points (SCP).

Table 2

Coefficients ($\pm$ SE) for logistic regression models to describe the cumulative frequency of freezing (from supercooling point measures) and the cumulative extent of mortality from brief cold exposure (from lower lethal temperature treatments) intercepts or slope estimates for populations of *Trissolcus japonicus*.

Table 3

Coefficients ($\pm$ SE) for logistic regression models to describe the extent of mortality after acute cold exposure among *Trissolcus japonicus* adults that had been reared under summer or fall conditions for up to three generations.

Fig. 1. Observed and predicted ($\pm$ SE band) cumulative freezing and/or mortality for populations of *Trissolcus japonicus* (A: Beijing, China; B: Pucheon, South Korea; C: Tsukuba, Japan; D: Nanjing, China) and *T. cultratus* (E: Tsukuba 2007, Japan; F: Tsukuba 2012, Japan).

Fig. 2. Median longevity ($\pm$ 95% CI), following brief exposure to low temperatures for populations of *Trissolcus japonicus* (A: Beijing, China; B: Pucheon, South Korea; C: Tsukuba, Japan; D: Nanjing, China) and *T. cultratus* (E: Tsukuba 2007, Japan; F: Tsukuba 2012, Japan).

Fig. 3. Observed and predicted mortality ($\pm$ SE band) of *Trissolcus japonicus* reared over multiple generations (A: parental (P) generation; B: first (F₁) generation; C: second (F₂) generation; D: third (F₃) generation) under summer and fall rearing conditions.

Table 1

Experiment	Population or generation	Experimental date	Latitude (°N)	Longitude (°E)	JALT † (°C)	Collection year	Shipment date	No. SCP tested	No. LLTemp tested (n/exposure temperature)
Supercooling point and lower lethal temperature	<i>T. japonicus</i>								
	Beijing, China	Aug.-Oct. 2014	39.91	116.39	-9.4	2009	Jul. 2014	22	88 (15-20)
	Pucheon, South Korea	Aug.-Oct. 2014	37.89	127.20	-6.1‡	2009	Jul. 2014	22	111 (20-25)
	Tsukuba, Japan	Aug.-Oct. 2014	36.03	140.07	2.5‡	2012	Jul. 2014	18	68 (12-15)
	Nanjing, China	Aug.-Oct. 2014	32.05	118.77	-1.6	2009	Jul. 2014	14	74 (14-15)
Lower lethal temperature	<i>T. cultratus</i>								
	Tsukuba 2007, Japan	Feb.-Mar. 2015	36.03	140.07	2.5‡	2007	Feb. 2015	-	80 (16)
	Tsukuba 2012, Japan	Feb.-Mar, 2015	36.03	140.07	2.5‡	2012	Feb. 2015	-	100 (20)
Cold acclimation	<i>T. japonicus</i> *								
	Summer, P	Jul.-Oct. 2015	39.91	116.39	-9.4	2009	May 2015	-	78 (6)
	Summer, F ₁	Jul.-Oct. 2015	39.91	116.39	-9.4	2009	May 2015	-	78 (6)
	Summer, F ₂	Jul.-Oct. 2015	39.91	116.39	-9.4	2009	May 2015	-	65 (5)
	Summer, F ₃	Jul.-Oct. 2015	39.91	116.39	-9.4	2009	May 2015	-	78 (6)
	Fall, F ₁	Jul.-Oct. 2015	39.91	116.39	-9.4	2009	May 2015	-	56 (4-5)
	Fall, F ₂	Jul.-Oct. 2015	39.91	116.39	-9.4	2009	May 2015	-	65 (5)
	Fall, F ₃	Jul.-Oct. 2015	39.91	116.39	-9.4	2009	May 2015	-	65 (5)

1

* Experiment included individuals from Beijing, China population.

2

† JALT January average lowest temperature (30-year period). Data from the World Meteorological Organization, 2016.

3

‡ Due to the lack of climatological information available for Pucheon, South Korea, and Tsukuba, Japan, a substitution was made with information

4

from the nearest cities: Seoul, South Korea (20km) and Tokyo, Japan (70km).

Table 2

Population	df	Intercept (b_0)				Slope (b_1)				50 th percentile	
		Freezing	Mortality	Z	P	Freezing	Mortality	Z	P	Freezing	Mortality
Beijing	106	-37.88 ± 3.85 _a	-7.71 ± 2.64 _a	-6.46	<0.001	-1.76 ± 0.18 _a	-0.42 ± 0.13 _a	-6.02	<0.001	-21.50 ± 0.06	-18.56 ± 0.72
Pucheon	129	-35.71 ± 3.41 _a	-7.26 ± 2.50 _a	-6.74	<0.001	-1.62 ± 0.15 _a	-0.42 ± 0.13 _a	-5.99	<0.001	-22.04 ± 0.07	-17.36 ± 0.84
Tsukuba	82	-33.54 ± 4.40 _{ab}	-12.34 ± 4.32 _a	-3.44	<0.001	-1.52 ± 0.20 _{ab}	-0.71 ± 0.23 _a	-2.66	0.01	-22.00 ± 0.09	-17.41 ± 0.64
Nanjing	84	-19.71 ± 3.27 _b	-9.67 ± 2.92 _a	.2.60	0.02	-0.92 ± 0.15 _b	-0.48 ± 0.15 _a	-2.09	0.04	-21.43 ± 0.19	-20.03 ± 0.61

- 1 Models follow the general equation: $P(\text{Insect freezes or dies}) = 1/[1 + e^{-(b_0 + b_1 * temperature)}]$.
- 2 Within rows (i.e., within populations), intercepts and slopes from the models for freezing were all significantly different from the models for
- 3 mortality at $\alpha = 0.05$.
- 4 Within a column, different letters indicate significant differences for intercepts or slopes among populations freezing ($df = 68$) or mortality ($df =$
- 5 333) treatments at an adjusted $\alpha = 0.008$ to maintain an overall $\alpha = 0.05$.

Generation	df	Intercept (b_0)				Slope (b_1)				50 th percentile (°C)	
		Summer	Fall	Z	P	Summer	Fall	Z	P	Summer	Fall
P*	76	-12.60 ± 2.89 _a	-	-	-	-0.70 ± 0.16 _a	-	-	-	-17.95 ± 0.50	-
F ₁	130	-8.19 ± 2.29 _a	-7.68 ± 2.79 _a	0.06	0.95	-0.50 ± 0.13 _a	-0.50 ± 0.16 _a	0.14	0.89	-16.24 ± 0.71	-15.44 ± 0.91
F ₂	126	-7.65 ± 2.25 _a	-8.52 ± 2.51 _a	0.26	0.80	-0.45 ± 0.12 _a	-0.51 ± 0.14 _a	0.33	0.74	-16.92 ± 0.79	-16.56 ± 0.74
F ₃	139	-8.73 ± 2.84 _a	-14.05 ± 4.31 _a	1.03	0.30	-0.58 ± 0.17 _a	-0.88 ± 0.26 _a	0.96	0.33	-15.09 ± 0.76	-15.98 ± 0.52

- 1 Models followed the general equation: $P(Insect\ dies) = 1/[1 + e^{-(b_0 + b_1 * temperature)}]$.
- 2 Within rows (i.e. within generations), intercepts and slopes from the summer conditions were not significantly different from those of the fall
- 3 conditions at $\alpha = 0.05$.
- 4 Within columns, different letters indicate significant differences for intercepts or slopes among generations for summer ($df = 291$) or fall ($df =$
- 5 256) conditions at an adjusted $\alpha = 0.008$. The summer P generation was also included in comparison among those reared under fall conditions.
- 6 *The P generation only reared under summer conditions, hence no comparison for this generation to a fall counterpart, nor a predicted 50th
- 7 percentile.

Figure 1

Figure 2

Figure 3

• Source locations of *Trissolcus* spp.

Freezing and mortality temperatures of *T. japonicus*