The Effect of Ochratoxin A on Antimicrobial Polypeptide Expression and Resistance to Water Mold Infection in Channel Catfish (*Ictalurus punctatus*)

Eman Zahran^{1,3,6}, Bruce Manning², Jung-Kil Seo^{1,4}, and Edward J. Noga^{1,5}

¹Department of Clinical Sciences, College of Veterinary Medicine, North Carolina State University, Raleigh, NC, USA 27606

²National Warmwater Aquaculture Center, Mississippi State University, Stoneville, MS 38776 USA

³Present address: Department of Internal Medicine, Infections and Fish Diseases, College of Veterinary Medicine, 1 Ahmed Mahar, Street, Mansoura, EGYPT

⁴Present address: Department of Food Science and Biotechnology, Kunsan National University, Kunsan, Jeonbuk 54150, REPUBLIC OF KOREA

⁵Present address: South Eastern Aquatechnologies, Inc. 250 NE 20th Street, #523, Boca Raton, FL 33431 USA

⁶Corresponding author: Tel: +20121100560 Fax: +20-50-2200696

E-mail address: eman_zahran2000@yahoo.com

1

Abstract

² Deceased

Mycotoxin contamination of agricultural commodities poses a serious risk to animal health, including aquaculture species. Ochratoxin A (OA) is the most immunotoxic ochratoxin, yet little is known about its effect on immune function in fish. Antimicrobial polypeptides (AMPPs) are one of the most potent, innate, host defense factors, yet very little is known about what types of chronic stressors affect their expression. Among the most prevalent and potent AMPPs in fish are histone-like proteins (HLPs). In this study, fish were fed 2, 4, or 8 mg OA/kg diet. Skin antibacterial activity and HLP-1 levels were measured on Days 0, 28 and 56. Feeding 2, 4 or 8 mg OA/kg diet resulted in significant growth depression, but higher levels (4 or 8 mg OA/kg diet) resulted in lowering feed intake (FI) and impaired feed conversion ratio. In addition, feeding 8 mg OA/kg diet increased susceptibility to experimental water mold (*Saprolegnia*) challenge, suggesting that OA toxicity might contribute to some saprolegniosis outbreaks. However, there were no changes in AMPP expression in any treatment group. Our data suggests that the increased disease susceptibility of channel catfish due to OA is probably due to mechanisms other than a direct effect on antimicrobial polypeptide expression.

Keywords: Antimicrobial peptides, Ochratoxin A, Water mold

1. Introduction

Mycotoxins are secondary fungal metabolites, which contaminate various types of feed commodities such as corn, wheat, cottonseed meal, peanuts, and soybean meal [1, 2]. Ochratoxin A (OA), produced by *Penicillium* and *Aspergillus* species, especially *Aspergillus ochraceus*, is one of the most important mycotoxins [3] and is associated with many signs such as reduced growth rate, feed efficiency ratio, reproductive performance, resistance to infectious diseases. It also causes damage to liver and other organs of fish and farm animals [4, 5]

OA mediates its toxic effect by acting on cellular respiratory enzymes through competitive inhibition of ATPase, succinate dehydrogenase, and cytochrome C oxidase in mitochondria. Moreover, because of its phenylalanine moiety, it competitively inhibits phenylalanyl-tRNA synthase, thus disrupting protein synthesis. In addition, cellular damage is caused by hydroxyl radical formation and lipid peroxidation [6]. OA induces oxidative damage in vivo [7] and in vitro [8] that eventually, lead to mitochondrial dysfunction, apoptosis and DNA damage [9].

OA is toxic to fish, but susceptibility varies considerably among species. Sea bass (*Dicentrarchus labrax* L.), are highly sensitive, having a 96 h LC₅₀ of 9.23 mg OA/kg diet [10]. The LD₅₀ for rainbow trout (*Oncorhynchus mykiss*) by injection is 4.76 mg OA/kg [11]. Channel catfish are much more resistant, tolerating as high as 4 mg OA/kg diet for at least 8 weeks without mortalities, and 8 mg OA/kg diet with 80% survival [12].

However, the sublethal effects of OA intoxication are much more economically damaging [13]. OA is well-known to increase disease susceptibility in homeotherms [14]. Pigs fed 3 mg OA/kg diet for up to three weeks spontaneously contracted salmonellosis, with *Salmonella choleraesuis* found in the faeces and liver, and deaths between days 15 and 17. Fewer pigs were affected at a lower dose (1 mg OA/kg diet) and none was affected in the control group [15]. Broiler chickens fed 2 mg OA/kg diet and challenged after 14 days with *Escherichia coli* O₇₈ had increased mortality (by 21.4%) compared with chicks inoculated with *E. coli* alone [16]. Eleven-day-old chickens orally challenged with *Salmonella typhimurium* for 2 consecutive days (3 mg OA/kg) had increased *S. typhimurium* in both their duodenal and cecal contents [17].

OA also increases disease susceptibility to *Edwardsiella ictaluri*, one of the most important pathogens affecting channel catfish (*Ictalurus punctatus*) [18]. Channel catfish fed 2.0 or 4.0 mg OA/kg diet for 6 weeks and then challenged by immersion with a highly virulent isolate of *Edwardsiella ictaluri* had significantly greater mortality (80%) than the control (68%). This result together with the less weight gain in both doses compared to control, indicate that feeding intoxicated diet could be implicated in higher susceptibility to the infection.

One of the most prevalent innate defenses in animals is antimicrobial polypeptides (AMPPs) [19]. Studies that have shown that depressed AMPP levels can greatly increase disease susceptibility. For example, recurrent bacterial infections occur in cases where there is a deficiency of α -defensins in neutrophils. Morbus Kostmann

syndrome, which is a severe congenital neutropenia in humans that is typified by low concentrations of AMPPs in the mouth, results in recurrent oral infections [20]. Depressed levels of histatins, AMPPs in human oral mucosa, have been associated with an increased risk of developing human immunodeficiency virus (HIV) infection [21]. Furthermore, Furci et al. [22] showed that α -defensin-5 (HD5) is a potent and broad-spectrum inhibitor of biologically diverse HIV-1 strains, through their interaction with the major HIV-1 envelope glycoprotein, gp120, and with its primary cellular receptor, CD4 interfering with their reciprocal binding. In the same context, immunomodulatory peptide (IDR-1002) enhanced in vitro chemokine induction activities and stronger in vivo anti-infective properties were confirmed. Additionally, it showed protection against the Gram-negative bacterial pathogen *Escherichia coli* [23].

Among the most common AMPPs in fish are histone-like proteins (HLPs), which have high homology to core nuclear histones. Originally isolated from the skin of channel catfish (*Ictalurus punctatus*) [24], HLPs were subsequently identified in skin, gill and/or spleen of hybrid striped bass (*Morone saxatilis* male x *M. chrysops* female) and rainbow trout (*Oncorhynchus mykiss*) [25, 26]. HLP-1 is the most prevalent and potent HLP, with broad-spectrum activity against bacteria, parasites and water molds [24, 26]. HLPs are also the predominant AMPP in normal healthy channel catfish [24].

Water molds (Oomycetes), one of the most important pathogens of freshwater fish [27], are classical opportunists that typically cause outbreaks when fish are exposed to some type of stress, such as adverse water temperature, poor water quality, handling or crowding. These factors may compromise immunity, increasing susceptibility to

infection [28, 29]. Members of the genus *Saprolegnia* are the most common and important water molds and thus the disease is often termed saprolegniosis. Saprolegniosis is associated with “winter kill” a syndrome in channel catfish in the southeastern United States that is associated with rapid temperature drop (below 15°C) in fall and winter [30, 31].

There have been no studies that have examined the effect of OA or any other mycotoxin on the expression of AMPPs in any animal. Thus, to better understand possible mechanisms underlying OA immunotoxicity, we assessed the effect of OA on the expression of AMPPs in channel catfish. We also determined if there was any relationship between AMPP tissue levels and resistance to challenge with the water mold *Saprolegnia*.

2. Materials and Methods

2.1. Experimental Fish

Channel catfish were obtained from a local producer and transported to North Carolina State University. Fish were acclimated in a 380 liter fiberglass aquarium at 24°C for 60 days prior to beginning the experiment. During that time, fish were fed ad lib with a 2 mm pellet (40% crude protein, 10% crude fat, and 4% crude fiber, Zeigler Bros, Inc., Gardners, PA). Ammonia, nitrite, and pH were monitored weekly. Two weeks before initiating the experiment, fish were sedated with 60 mgL⁻¹ buffered tricane and transferred to the experimental aquaria.

2.2. *Experimental Design*

Fifteen fish (7 month old and 17-21cm length) were placed in each of twelve, 60-liter freshwater aquaria (total N = 180). All aquaria were connected to a central filtration system having a conditioned biofilter (biocubes and bead filter [Aquadyne, Koi Camp Aquariology, Loganville, GA]) and a titanium heater (Process Technology Co., Mentor, OH). Fish began feeding normally almost immediately and were fed close to apparent satiation twice daily. After 7 days, the temperature was increased from 24°C to 29°C over 7 days. After 14 d at 29°C (day 0), all fish were weighed, the mean weight of all fish on day 0 was 55 ± 15 g and the day 0 sampling was performed. At day 1, all fish were switched to semi-purified diets (Table 1) prepared as described previously (Manning et al. 2003), and having one of four concentrations of OA (0, 2.0, 4.0, or 8.0 mg OA/ kg diet). “The experiment was conducted with a protocol approved by the North Carolina State University Institutional Animal Care and Use Committee. Triplicate aquaria were assigned to one of four treatments. During the experiment, 75% water changes were performed thrice weekly, and water quality was measured (ammonia, nitrite, nitrate, pH, and dissolved oxygen) twice weekly via water quality test kits (Aquarium Pharmaceuticals, Inc.).

2.3. *Sample Collection*

Three fish from each aquarium were sampled at days 0, 28 and 56. Each aquarium was sampled one at a time; the three fish were sedated with a low dose of buffered tricane; (**MS-222, ARGENT**), (30 mg/L tricane + 60 mg/L sodium

bicarbonate) and each fish was then euthanized one at a time in a separate container having a high dose of buffered tricane (200 mg/L tricane + 400 mg/L sodium bicarbonate). Each fish was weighed and measured to **calculate the condition factor, feed intake (FI), and feed conversion ratio (FCR) according to the following formulae:**

$$K (K = W (g) \times 10^2 / L^3 (cm))$$

$$\text{Feed intake (FI) (g /fish) = Dry feed intake (g)/Number of fish}$$

$$\text{Feed conversion ratio (FCR) (\%) = Feed intake (g)/ Gain in body mass (g)}$$

A skin scraping and a gill clip were then taken and examined immediately for pathogens under a light microscope. A skin sample was then collected for AMPP levels. Skin was processed for measuring AMPPs as previously described [25, 32]. Briefly, 50 μ L of tissue was collected to yield a total volume of 200 μ L in 150 μ L of 1% acetic acid (1:4 dilution of tissue). After boiling for 5 min, the sample was homogenized and then centrifuged at 14,000 x g for 10 min at 4°C. The resulting extract (supernatant) was used to measure antibacterial activity and HLP-1.

2.4. Bacterial Culture

Blood from representative fish in each treatment (2 fish per aquarium) was collected from the caudal vasculature with a 1 cc syringe having a 23 GA needle. A drop was placed on a Columbia blood agar plate and then spread with a sterile swab (Mini-tip Culturette, Becton Dickinson, Franklin Lakes, NJ). Culture plates were incubated at room temperature and observed daily for 14 d.

2.5. Total Antibacterial Activity

Total antibacterial activity was measured using the radial diffusion assay as described previously [26]. Briefly, *Escherichia coli* D31 was propagated overnight in trypticase soy broth with 1% NaCl and then washed in PBS (pH 7) and diluted until to an OD₅₇₀ of 0.10 with a Vitek colorimeter (Hach, Loveland, CO). One mL of the bacterial suspension and 1 mL of streptomycin sulfate (10 mg/mL stock) was added to an autoclaved agarose medium (1.57 g of low electroendosmosis agarose, 0.5 g of NaCl, 20.7 mL of 5X Luria-Bertani broth, 20.7 mL of 1 M phosphate buffer pH 6.7, and 58.6 ml of deionized water) that was cooled in a water bath to 45-48°C. Ten milliliters of the suspension was poured into each sterile Petri dish, which was used immediately or stored at 4°C until use. Samples were pipetted into 2.5 mm diameter wells punched in the agarose plate. After diffusion of the sample into the agarose, the plate was incubated at 37°C for 18 h, at which time clearing zone diameters were measured by calipers to the nearest 0.1 mm. The radial diffusion assay clearing zone diameters were then converted to Units of activity by reference to a standard curve prepared by serially diluting calf histone H2B (Roche Diagnostics Co, Basel, Switzerland). The 1000 µg/mL concentration of calf histone H2B was considered to be 100 Units of activity. All values were multiplied by 10 to report the tissue concentrations (as µg/mL⁻¹ equivalents of calf histone H2B).

2.6. Anti-*Edwardsiella ictaluri* Activity

We have previously discovered that other AMPPs are present in channel catfish skin besides HLPs [33]. These AMPPs appear to be mainly small polypeptides (<10 kDa). We have recently discovered that some of these AMPPs have potent activity against *Edwardsiella ictaluri* (A Ullal and E Noga, North Carolina State University, Unpublished Data). We subsequently developed a method for selectively detecting this activity (see below), which involves concentrating the activity via reverse-phase chromatography (using a Sep-Pak [Waters]) followed by measuring the activity using a pathogenic isolate of *Edwardsiella ictaluri*. *Edwardsiella ictaluri* is not sensitive to HLPs [24]; thus, using this bacterium in the assay allows one to exclude measurement of antibacterial activity due to HLPs.

Pooled skin extracts were tested from fish sampled at Day 0 and Day 56 in the 0 and 8 mg OA/ kg diet treatments. Each sample (25 μ L) was loaded onto a Sep-Pak Plus C₈ cartridge (Waters #036775), which was previously washed with 3 mL of acidified water (0.1% trifluoroacetic acid); the activity was then eluted with 2 mL of 80% acetonitrile (ACN) in 0.1% trifluoroacetic acid. The effluents was lyophilized and resuspended in 10 μ L of 0.01% HAc, and screened for antibacterial activity against *Edwardsiella ictaluri* using the ultrasensitive assay as described in Seo et al. [34]. Clearing zone diameters were then converted to Units of activity by reference to a standard curve prepared by serially diluting synthetic piscidin 1, an AMPP which is known to have potent activity against *E. ictaluri* (J Stephenson and E Noga, Unpublished Data). All values were then divided by 2.5 to account for the 10-fold concentration of the extract; the values were then multiplied by 10 to report the tissue concentrations as μ g/ml equivalents of piscidin 1.

2.7. Quantification of HLP-1

HLP-1 was measured via ELISA according to Noga et al. [35]. Briefly, microtiter plates were coated for one h at room temperature using primary antibodies that were homologous to the N-terminus and C-terminus of HLP-1 (anti-PDPA and anti-VSEG antibodies). After a washing step, the plates were incubated with Stabilcoat (Surmodics, Inc., Eden Prairie, MN) for 20 min to eliminate nonspecific binding and then stored at 4°C until use. Immediately before use, plates were washed with 0.05 M tris-buffered saline with 0.05% Tween 20. The pooled skin tissue extracts were diluted to the desired concentration (1:100) in 0.05 M tris-buffered saline with 0.05% Tween 20 and 1% bovine serum albumin (BSA) (Sigma #T6789, St. Louis, MO). All samples were added in duplicate to the ELISA plate and incubated for 1 h. The plate was washed again before the secondary antibody (affinity-purified, peroxidase-labeled antibody to calf histone H2B) was added and incubated for 1 h. The plate was then washed and TMB peroxidase substrate was added and incubated for 15 min. The reaction was terminated with 3 M phosphoric acid. Absorbance was measured at 450 nm with an ELISA plate reader (KCjunior, Bio-Tek Instruments, Inc., Winooski, VT). A standard curve was prepared using a serial dilution of calf histone H2B, and the positive control was prepared from a synthetic antigen created by linking the N-terminus and C-terminus peptide antigens (used to prepare the coating antibodies) to BSA.

Tissue concentration of HLP-1 was calculated by the software (KCjunior, Bio-Tek Instruments, Inc., Winooski, VT) at 450 nm absorbance based on the optical density (OD) of the 4 parameter curve-fitting standards. Samples with an $OD \geq 3$ were

diluted twofold and re-ran. All values were multiplied by 4 to account for the 1:4 dilution of tissue in 1% acetic acid (50 μ L of tissue in 150 μ L of 1% acetic acid) during sample collection. Tissue concentrations were reported as μ g/mL equivalents of calf histone H2B.

2.8. Water Mold Challenge

Zoospores for challenge were prepared according to Udomkusionsri and Noga [28]. Briefly, *Saprolegnia* sp. (Isolate #97-2010A) from a diseased channel catfish was cultured on glucose yeast extract (GY) agar [36] at 19°C for 2 days. Agar with mycelia was then aseptically cut into 8×8 mm squares and placed into a petri-dish having 30 mL GY broth. After 2 days, the agar remnants were removed and the mycelium was washed three times with sterile sporulation medium [MSM (modified Griffin's sporulation medium) [37]]. The mycelium was then incubated in MSM at 19°C for 24 h to induce zoospore formation. **About 300 mL of zoospore suspension was collected from about 10 petri dishes containing sterile MSM and placed in a plastic aquarium bag for inoculation into the challenge aquarium.**

On day 58 of the feeding trial, the remaining fish were challenged with zoospores. Nine fish from each group (0, 2.0, 4.0, and 8.0 mg OA/ kg diet) were placed in separate plastic cages (48 cm in diameter, 38 cm deep) in a 1100 liter closed system fiberglass aquarium at 29°C; over ~12 hr, the temperature was dropped to 11°C by unplugging the heaters and placing an external chiller (Aqua Logic, Inc, San Diego, CA) in the water. When the water temperature reached 11°C, the plastic bag containing the

zoospore suspension was placed in the aquarium for ~30 min to acclimate the zoospore suspension to the aquarium temperature. The suspension was then mixed with 25% aquarium water for 25 min, followed by pouring the suspension into the aquarium. The challenged fish were examined thrice daily for typical signs of water mold infection (e.g., cottony white growth on skin and fins). Moribund or dead fish were immediately removed. Skin scrapings and gill and fin biopsies of all removed fish were examined; water mold infection was confirmed via identification of broad, aseptate hyphae, sporangia and encysted zoospores with light microscopy.

2.9. Statistical Analyses

All data were analyzed using SPSS version 17.0 (SPSS Inc, Chicago, Illinois). The effect of experimental variables (OA feed rate, aquarium replicate, sampling day) was tested by analysis of variance (ANOVA). A significant difference was accepted if $P < 0.05$. The Kruskal-Wallis test, the nonparametric version of ANOVA, was used to analyze the the effect of treatment on survival time after water mold challenge. In addition, a Wilcoxon test was used for each pair of treatments, adjusting for multiple comparisons using Tukey's method for pairwise comparisons. Values were considered significant if $P < 0.05$.

3. Results

3.1. Necropsy Examinations

There were no gross external lesions in any fish during the entire period of OA exposure (8 weeks); also, no pathogens were detected in skin or gill biopsies at any time. Blood cultures had a small number of bacterial colonies in some fish of each of the four treatment groups (ranging from 1 to 26 colonies). No predominant colony type was observed and in the sample with the largest number of colonies, three colony types were present.

3.2. Growth Parameters

At Day 28, the 8 mg OA/kg diet treatment group weighed significantly less than all other groups (Fig. 1A). By Day 56, all treatment groups (2, 4 and 8 mg OA/ kg diet) weighed significantly less than the control. Changes in condition factor (K) were less consistent; the 8 mg OA/kg diet treatment was significantly lower than the control at day 28, but was not different at day 56 (Fig. 1B). Feed intake (FI) was significantly lower in treatment groups compared to the control, with an apparent dose-related trend with increasing OA dose. This same trend was also evident with feed conversion ratio (FCR) (Fig. 2) suggesting poor feed utilization.

3.3. AMPP Levels

There were no differences in the total antibacterial activity among any treatment groups at any of the sampling times (Fig. 3). This was also reflected in the lack of obvious changes measured in pooled extracts when measuring HLP-1 levels (Table 2).

There was no detectable activity in these same samples against *E. ictaluri* (data not shown).

3.4. Susceptibility to Water Mold Challenge

Water mold infection was first grossly visible on Day 4 post-challenge, where, in the 8 mg OA/ kg diet treatment, one fish was dead and all others were heavily infected. Affected fish had typical signs of water mold infection, with cotton-like growths on the body and fins. By Day 8, fish in all groups were dead except one in the control. **There was a significant effect of treatment on survival time (Kruskal-Wallis test, $p=0.0074$). The 8 mg OA/kg diet group had a significantly shorter survival time than all three other treatments ($p<0.05$) as determined using Wilcoxon test for each pair of treatments (Fig. 4).**

4. Discussion

OA is well-known to be immunosuppressive to vertebrates. OA reduces the size of immune organs, such as thymus, spleen, and lymph nodes, and the number of immune cells within tissues. It also causes profound alterations in the function of mononuclear leukocytes [38], affecting production of reactive oxygen species, chemotaxis, phagocytosis and cytokine synthesis [39].

Feeding 4 mg OA/kg for 10 weeks caused lymphoid depletion in the thymus of turkeys and was accompanied by reduced T-lymphocyte-mediated delayed cutaneous hypersensitivity responses [40]. In other studies, OA inhibited natural killer (NK) cell activity and increased the growth of tumor cells in mice. Ochratoxin had little direct effect on NK cells, but did impair the induction of interferon. Because interferon is

necessary for NK cell activity, the suppression of NK cell activity was attributed to a decrease of endogenous interferon levels in the ochratoxin-treated mice [41]. The effect of 0.5 or 2.0 mg OA/kg feed on broiler chicks revealed significant reductions in cell-mediated immunity, as indicated by diminished skin sensitivity, graft versus host reactions and T-lymphocyte counts. Total lymphocyte counts, total serum protein, serum albumin and serum globulin levels were significantly depressed on day 21. Phagocytosis was inhibited, and the number of splenic macrophages was drastically reduced in both intoxicated groups, as well as the weights of the thymus, bursa of Fabricius and spleen [42]. Feeding Nile tilapia on AFB1 contaminated diet (0.2 mg/kg diet) showed up-regulation of CYP1A and down-regulation of SOD, IL-1 β and TGF- β after 16 weeks exposure [43]. OA (0.4 and 0.6 mg/kg diet) fed to Nile tilapia for 90 days, revealed reduction in RBCs, Hb and Hct values. Besides, a significant decrease in plasma total protein, total lipids, AST and lactate dehydrogenase along with significant increase in the glucose concentration and ALT levels [44]. Furthermore, Zheng et al. [45] confirmed the association between the OA induced toxicity and zinc, they showed that OA exposure was associated with inhibition of cell proliferation, decrease in the intracellular zinc concentration, induction of ROS production, decreases of SOD activity and DNA strand breaks, DNA oxidation and hypomethylation. But, zinc supplementation alleviated OA induced such toxicity.

To our knowledge, we are the first to examine the effect of any mycotoxin on expression of any AMPP. However, Fujiwara et al. [46] reported that another food-borne toxin, 2-amino-1-methyl-6 phenylimidazo[4,5-b]pyridine (PhIP), one of the most abundant heterocyclic amines produced while cooking meat and fish,

induces an overexpression of α -defensin genes by Paneth cells in rats developing colon cancers. A synthetic lytic peptide (D4E1) gene, were expressed in transgenic hybrid poplar (*Populus tremula* L. $\times$ *Populus alba* L.) against bacterial disease. One transgenic poplar line, Tr23, was the highest transcript accumulated for the D4E1 gene, and exhibited a significant relief in symptoms caused by *A. tumefaciens* and *X. populi* [47]. Recently, Badea et al. [48] cloned antimicrobial peptides genes, MsrA2 and 10R and showed that T3 generation genetically modified (GM) plants had a 53% reduction in *Fusarium* damaged kernels and some lines also had a 59% reduction in powdery mildew susceptibility compared with the non-GM control.

HLP-1 levels at Day 0 for both control and treatment groups in the pooled skin extracts were similar (12 and 10.8 $\mu\text{g/mL}$, respectively; Table 2); at Day 56, both groups showed nominally higher HLP-1 levels (15.6 and 25.6 $\mu\text{g/mL}$, respectively; Table. 2). The latter levels are within concentrations that are inhibitory or lethal to some important pathogens such as the protozoan ectoparasite, *Amyloodinium ocellatum* (lethal at 12.5 $\mu\text{g/mL}$) [25] and the bacterium *Aeromonas hydrophila* (inhibitory at 25 $\mu\text{g/mL}$) [24]. However, our resting levels of antibacterial activity for skin extracts at both Day 0 and 56 for both groups were low ($\leq 0.5 \text{ U} = 5 \mu\text{g/mL}$, Fig. 3). This is in contrast to our previous study [33], where there was much higher antibacterial activity in control channel catfish (5.7 U = 57 $\mu\text{g/mL}$). However, some of the activity measured in the Ullal et al. [33] study might have been due to the upregulation of certain AMPPs related to hemoglobin, due to the possible presence of a subclinical ectoparasite infestation on some fish. Thus, the resting levels that we measured in the present study might more closely reflect the true baseline levels in channel catfish skin.

Nonetheless, our data did not show any effect of OA on antimicrobial polypeptide expression. Total antibacterial activity was not significantly different among treatment groups. This was supported by the measurements of HLP-1 in pooled tissue extracts, which did not reveal any apparent trends among OA feed levels. HLP-1 is the major AMPP in unstimulated channel catfish skin and gills [24, 33]. Similar results were seen in measurements of anti-*E. ictaluri* activity. These data suggest that OA does not affect AMPP levels in channel catfish. Our data also suggests that the increased susceptibility to edwardsiellosis observed with 8 mg OA/ kg diet in the feed [18] is not due to depressed AMPP levels but rather to some other form of immunosuppression. OA has been found to affect intestinal epithelial cells and alter both intestinal barriers and selected transport functions. Moreover, OA selectively affects the protein content of plasma membrane microdomains, inducing perturbation of microdomain composition, which may account for the decrease of transepithelial resistance and thus destabilizing the tight junctions of intestinal epithelial cells, facilitating its own absorption [49]. **Thus, OA might primarily target an intrinsic component of intestinal immunity (its barrier function) rather than having a direct effect on extrinsic components of immunity such as AMPPs. However, the immunosuppressant effect of OA was still evidenced by increased fish susceptibility to *Saprolegnia* challenge.**

Manning et al. [50] found that in channel catfish fed OA for 8 weeks, survival was high for fish fed diets containing 0–4 mg OA/kg feed, but fish fed the diet containing 8.0 mg OA/kg had significantly lower survival. At 8 weeks, histopathological examination showed a significant increase in the severity of liver and

posterior kidney lesions among catfish fed diets containing 2.0 mg OA/kg or above. These lesions were characterized as enlarged melanomacrophage centers, which replaced normal hepatopancreatic or posterior kidney cells. Exocrine pancreatic cells, that normally surround the hepatic portal veins, were decreased or absent in fish fed 1.0 mg OA/kg diet or greater. Hematocrit was significantly lower for catfish fed 8 mg OA/kg of feed, but there were no significant differences in white blood cell (WBC) counts at any dietary levels of OA.

We found that OA significantly reduced weight gain and feed efficiency in a dose-dependent manner. After 56 days (8 weeks), body weight was significantly lower in fish fed 2, 4, or 8 mg OAk⁻¹ diet. This is consistent with the findings of Manning et al. [50], where, after 8 weeks, there was a significant reduction in weight gain in channel catfish fed diets containing 1, 2, 4, or 8 mg OA/kg. In a separate study, catfish fed 2.0 or 4.0 mg OA/kg diet for 6 weeks also weighed significantly less than the control fish [18].

In the current study, feed intake (FI) and FCR were impaired particularly, in fish fed 4 or 8 mg OA/kg of feed after 28 days and 56 days. This is in line with the findings of Saad [51], who reported impaired FCR of *O.niloticus* treated with OA. Similar results were seen by Srour [5], who found reduction of feed and nutrient utilization of *O. niloticus* fed OA contaminated diet. Manning et al. [50] also found poorer FCR in fish fed 4 or 8 mg OA/kg of feed. In same context, Srour [5], found a reduction in growth performance parameters in Nile tilapia exposed to OA at 4.8, 9.6 and 14.4 mg/kg diet. Similarly, Farag [52] found depression in weight, gain and specific growth rate (SGR)

of *O. niloticus* and *Cyprinus carpio* fed on diet contaminated with 7 mg of OA/kg diet. Feeding Nile tilapia on OA contaminated diet at low and high doses (0.8 and 0.16 mg/kg fish), respectively, led to a significant reduction in final body weight, body gain, average daily gain (ADG) and SGR in a dose dependent manner [53]. This may be attributed to different mechanism of OA such as, inhibition of protein synthesis [54], induction of ROS with consequent damage of nucleic acids, protein and lipids [55]. Thus, it is reasonable to expect that OA would cause growth retardation.

The condition factor (K) of the 8 mg OA/kg diet treatment was significantly lower than the control at day 28, but K was not different at day 56, and that of the control fish was similar to what has been reported for channel catfish [56]. Our results were corroborated with Abdallah et al. [53] who reported that K was not affected with OA treatment in Nile tilapia at lower (0.8 mg/kg fish) and higher (0.16 mg/kg fish) doses. Thus, condition factor changes were less consistent and did not reflect the lower innate immune status of the fish group fed the highest tested dose 8 mg OA/ kg diet.

Despite the lack of effect on AMPP levels, exposure of channel catfish to *Saprolegnia* significantly increased mortality in fish that were fed a diet containing the highest level of OA (8 mg OA/ kg diet). OA exerts suppression of the immune response renders animals more susceptible to pathogen with subsequent increasing in the severity of infection and aggravating the OA toxicity [57]. Our results are consistent with previous studies showing that 4 mg OA/kg feed increases susceptibility of channel catfish to *Edwardsiella ictaluri* challenge causing significant higher mortalities (80%) versus the control (68%) [18]. Previous studies conducted in homeotherms, showed that

OA increased susceptibility of rabbits to *Pasteurella multocida* associated with higher mortalities [58]. OA increased chicks' susceptibility to *Eimeria acervulina* leading to severe duodenal coccidiosis [59]. In the present study, impairment of the protein metabolism one of the possible causes to lowering animal resistance, along with suppression of the immune response [60] can ultimately enhance the susceptibility to various infections. Recent surveys of the occurrence of mycotoxin-contaminated feed or feed ingredients are not extensive; however, an evaluation of OA contamination of feed grains or animal feeds conducted in the USA indicates that levels of OA of up to 16 mg/kg were found in surveys and up to 90 mg/kg in feeds from OA case studies [61]; thus, high doses can be found naturally in fish feed and pose a risk for animals and humans as well. Thus, OA might be a contributing factor to some saprolegniosis outbreaks in channel catfish.

Conclusions

In summary, channel catfish exposed to moderate to high dietary levels of ochratoxin A (2-8 mg OA/ kg diet) exhibited a significant decrease in growth rate and significant increase in feed conversion ratio, supporting findings in prior studies. However, only very high levels (8 mg OA/ kg diet) had any effect on the susceptibility to water mold infection and this effect appeared relatively small. This increased susceptibility did not appear to be associated with depressed AMPP levels, suggesting that other aspects of impaired immune function are responsible for this greater susceptibility.

Acknowledgements

This work was supported by Grant NA16-RG-2251 (R/AF-45) from the National Sea Grant College Program, National Oceanic and Atmospheric Administration, to the North Carolina Sea Grant College Program, North Carolina Fishery Research Grant Project #06-AM-04, and the National Research Initiative of the USDA Cooperative State Research, Education and Extension Service, Grant number #2006-35204-17508 to E.J.N. This work was also supported as a joint supervision programme, Mansoura University, mission sector for 2006/2007 plan. We thank S Salger, S Christian and especially J. Stephenson for technical assistance, and S Wood for assistance with statistical analysis. J. Corrales and A. Ulall provided helpful discussions.

References

1. Zinedine A, Brera C, Elakhdari S, Catano C, Debegnach F, Angelini S, et al. Natural occurrence of mycotoxins in cereals and spices commercialized in Morocco. *Food Control*. 2006 17:868-74.
2. Sangare-Tigori B, Dem A, Kouadio H, Betbeder A, Dano D, Moukha S, et al. Preliminary survey of ochratoxin A in millet, maize, rice and peanuts in Cote d'Ivoire from 1998 to 2002. *Human & experimental toxicology*. 2006 25:211-6.
3. Dietrich DR, Heussner AH, O'Brien E. Ochratoxin A: comparative pharmacokinetics and toxicological implications (experimental and domestic animals and humans). *Food additives and contaminants*. 2005 22 Suppl 1:45-52.

4. Tuan NA, Manning BB, Lovell RT, Rottinghaus GE. Responses of Nile tilapia (*Oreochromis niloticus*) fed diets containing different concentrations of moniliformin or fumonisin B 1. *Aquaculture*. 2003 217:515-28.
5. Srour T. Effect of ochratoxin-A with or without Biogen® on growth performance, feed utilization and carcass composition of Nile tilapia (*Oreochromis niloticus*) fingerlings. *J Agric Sci Mansoura Univ*. 2004 29:51-61.
6. Hussein HS, Brasel JM. Toxicity, metabolism, and impact of mycotoxins on humans and animals. *Toxicology*. 2001 167:101-34.
7. Schaaf G, Nijmeijer S, Maas R, Roestenberg P, De Groene E, Fink-Gremmels J. The role of oxidative stress in the ochratoxin A-mediated toxicity in proximal tubular cells. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease*. 2002 1588:149-58.
8. Petrik J, Žanić-Grubišić T, Barišić K, Pepeljnjak S, Radić B, Ferencić Ž, et al. Apoptosis and oxidative stress induced by ochratoxin A in rat kidney. *Archives of toxicology*. 2003 77:685-93.
9. Le Bras M, Clement M, Pervaiz S, Brenner C. Reactive oxygen species and the mitochondrial signaling pathway of cell death. 2005.
10. El-Sayed YS, Khalil RH, Saad TT. Acute toxicity of ochratoxin-A in marine water-reared sea bass (*Dicentrarchus labrax* L.). *Chemosphere*. 2009 75:878-82.
11. Doster R, Sinnhuber R, Wales J. Acute intraperitoneal toxicity of ochratoxins A and B in rainbow trout (*Salmo gairdneri*). *Food and Cosmetics Toxicology*. 1972 10:85-92.

12. Manning BB, Ulloa RM, Li MH, Robinson EH, Rottinghaus GE. Ochratoxin A fed to channel catfish (*Ictalurus punctatus*) causes reduced growth and lesions of hepatopancreatic tissue. *Aquaculture*. 2003 219:739-50.
13. Kuiper-Goodman T, Magan N, Olsen M. Risk assessment and risk management of mycotoxins in food. *Mycotoxins in food: detection and control*. 2004:3-31.
14. Pfohl-Leszkowicz A, Manderville RA. Ochratoxin A: An overview on toxicity and carcinogenicity in animals and humans. *Molecular nutrition & food research*. 2007 51:61-99.
15. Stoev S, Goundasheva D, Mirtcheva T, Mantle P. Susceptibility to secondary bacterial infections in growing pigs as an early response in ochratoxicosis. *Experimental and toxicologic pathology*. 2000 52:287-96.
16. Kumar A, Jindal N, Shukla C, Pal Y, Ledoux D, Rottinghaus G. Effect of ochratoxin A on *Escherichia coli*-challenged broiler chicks. *Avian diseases*. 2003 47:415-24.
17. Fukata T, Sasai K, Baba E, Arakawa A. Effect of ochratoxin A on *Salmonella typhimurium*-challenged layer chickens. *Avian diseases*. 1996:924-6.
18. Manning BB, Terhune JS, Li MH, Robinson EH, Wise DJ, Rottinghaus GE. Exposure to feedborne mycotoxins T-2 toxin or ochratoxin A causes increased mortality of channel catfish challenged with *Edwardsiella ictaluri*. *Journal of aquatic animal health*. 2005 17:147-52.
19. Zasloff M. Antimicrobial peptides of multicellular organisms. *nature*. 2002 415:389-95.

20. Pütsep K, Carlsson G, Boman HG, Andersson M. Deficiency of antibacterial peptides in patients with morbus Kostmann: an observation study. *The Lancet*. 2002 360:1144-9.
21. Lal K, Pollock JJ, Santarpia III RP, Heller HM, Kaufman HW, Fuhrer J, et al. Pilot study comparing the salivary cationic protein concentrations in healthy adults and AIDS patients: correlation with antifungal activity. *JAIDS Journal of Acquired Immune Deficiency Syndromes*. 1992 5:904-14.
22. Furci L, Tolazzi M, Sironi F, Vassena L, Lusso P. Inhibition of HIV-1 infection by human α -defensin-5, a natural antimicrobial peptide expressed in the genital and intestinal mucosae. *PLoS ONE* 2012 7:e45208.
23. Nijnik A, Madera L, Ma S, Waldbrook M, Elliott MR, Easton DM, et al. Synthetic cationic peptide IDR-1002 provides protection against bacterial infections through chemokine induction and enhanced leukocyte recruitment. *The Journal of Immunology*. 2010 184:2539-50.
24. Robinette D, Wada S, Arroll T, Levy M, Miller W, Noga E. Antimicrobial activity in the skin of the channel catfish *Ictalurus punctatus*: characterization of broad-spectrum histone-like antimicrobial proteins. *Cellular and Molecular Life Sciences CMLS*. 1998 54:467-75.
25. Noga EJ, Fan Z, Silphaduang U. Histone-like proteins from fish are lethal to the parasitic dinoflagellate *Amyloodinium ocellatum*. *Parasitology*. 2001 123.
26. Noga EJ, Fan Z, Silphaduang U. Host site of activity and cytological effects of histone-like proteins on the parasitic dinoflagellate *Amyloodinium ocellatum*. *Diseases of aquatic organisms*. 2002 52:207-15.

27. El Ashram AMM, Abd El-Rhman AM, Sakr SF. A contribution to saprolegniosis in cultured Nile tilapia (*Oreochromis niloticus*) with special reference to its control. Egypt J Aquat Biol & fish. 2007 11:943-55.
28. Udomkusionsri P, Noga E. The acute ulceration response (AUR): A potentially widespread and serious cause of skin infection in fish. Aquaculture. 2005 246:63-77.
29. Roberge C, Paez DJ, Rossignol O, Guderley H, Dodson J, Bernatchez L. Genome-wide survey of the gene expression response to saprolegniasis in Atlantic salmon. Molecular immunology. 2007 44:1374-83.
30. Ali EH, Hashem M, Al-Salahy MB. Pathogenicity and oxidative stress in Nile tilapia caused by *Aphanomyces laevis* and *Phoma herbarum* isolated from farmed fish. Diseases of Aquatic Organisms. 2011 94:17-28.
31. Thoen E, Evensen O, Skaar I. Pathogenicity of *Saprolegnia* spp. to Atlantic salmon, *Salmo salar* L., eggs. Journal of Fish Diseases. 2011 34:601-8.
32. Robinette DW, Noga EJ. Histone-like protein: a novel method for measuring stress in fish. Diseases of aquatic organisms. 2001 44:97-107.
33. Ullal AJ, Litaker RW, Noga EJ. Antimicrobial peptides derived from hemoglobin are expressed in epithelium of channel catfish (*Ictalurus punctatus*, Rafinesque). Developmental and comparative immunology. 2008 32:1301-12.
34. Seo JK, Crawford JM, Stone KL, Noga EJ. Purification of a novel arthropod defensin from the American oyster, *Crassostrea virginica*. Biochemical and biophysical research communications. 2005 338:1998-2004.
35. Noga EJ, Borron PJ, Hinshaw J, Gordon WC, Gordon LJ, Seo J-K. Identification of histones as endogenous antibiotics in fish and quantification in rainbow trout

- (*Oncorhynchus mykiss*) skin and gill. Fish physiology and biochemistry. 2011 37:135-52.
36. Hatai K, Takahashi S, Egusa S, Oe K. Study on the pathogenic fungus of mycotic granulomatosis, 1: Isolation and pathogenicity of the fungus from cultured-Ayu infected with the disease. Fish Pathology. 1977.
 37. Griffin D. *Achlya bisexualis*. Lower fungi in the laboratory: University of Georgia Press Athens, GA; 1978, p. 67-8.
 38. AL-ANATI L, Petzinger E. Immunotoxic activity of ochratoxin A. Journal of veterinary pharmacology and therapeutics. 2006 29:79-90.
 39. Alvarez RM, Morales AE, Sanz A. Antioxidant defenses in fish: biotic and abiotic factors. Rev Fish Biol Fisher. 2005 15:75-88.
 40. Dwivedi P, Burns R. Immunosuppressive effects of Ochratoxin a in young Turkeys 1. Avian Pathology. 1985 14:213-25.
 41. Luster MI, Germolec DR, Burleson GR, Jameson CW, Ackermann MF, Lamm KR, et al. Selective immunosuppression in mice of natural killer cell activity by ochratoxin A. Cancer research. 1987 47:2259-63.
 42. Singh G, Chauhan H, Jha G, Singh K. Immunosuppression due to chronic ochratoxicosis in broiler chicks. Journal of comparative pathology. 1990 103:399-410.
 43. Mahfouz M. Ameliorative effect of curcumin on aflatoxin B1-induced changes in liver gene expression of *Oreochromis niloticus*. Molecular Biology. 2015 49:275-86.
 44. Shalaby A. The opposing effect of ascorbic acid (vitamin C) on ochratoxin toxicity in Nile tilapia (*Oreochromis niloticus*). Proceedings of the 6th

- International Symposium on Tilapia in Aquaculture (RB Remedios, GC Mair and K Fitzsimmons, eds); 2004, p. 209-21.
45. Zheng J, Zhang Y, Xu W, Luo Y, Hao J, Shen XL, et al. Zinc protects HepG2 cells against the oxidative damage and DNA damage induced by ochratoxin A. *Toxicology and applied pharmacology*. 2013 268:123-31.
 46. Fujiwara K, Ochiai M, Ohta T, Ohki M, Aburatani H, Nagao M, et al. Global gene expression analysis of rat colon cancers induced by a food-borne carcinogen, 2-amino-1-methyl-6-phenylimidazo [4, 5-b] pyridine. *Carcinogenesis*. 2004 25:1495-505.
 47. Mentag R, Luckevich M, Morency M-J, Seguin A. Bacterial disease resistance of transgenic hybrid poplar expressing the synthetic antimicrobial peptide D4E1. *Tree Physiology*. 2003 23:405-11.
 48. Badea A, Eudes F, Laroche A, Graf R, Doshi K, Amundsen E, et al. Antimicrobial peptides expressed in wheat reduce susceptibility to Fusarium head blight and powdery mildew. *Canadian Journal of Plant Science*. 2013 93:199-208.
 49. Maresca M, Mahfoud R, Pfohl-Leszkowicz A, Fantini J. The mycotoxin ochratoxin A alters intestinal barrier and absorption functions but has no effect on chloride secretion. *Toxicology and applied pharmacology*. 2001 176:54-63.
 50. Manning B. Ochratoxin A fed to channel catfish (*Ictalurus punctatus*) causes reduced growth and lesions of hepatopancreatic tissue. *Aquaculture*. 2003 219:739-50.
 51. Saad TT. Some studies on the effects of ochratoxin-A on cultured *oreochromis niloticus* and carp species. [M.V.SC]: Alexandria University; 2002.

52. Farag M. PLASMA PROTEINS PROFILE IN NILE TILAPIA (*OREOCHROMIS NILOTICUS*) AND COMMON CARP (*CYPRINUS CARPIO*) EXPOSED TO OCHRATOXIN-A AND TREATED WITH SOME ANTI-TOXINS. EGYPTIAN JOURNAL OF AGRICULTURAL RESEARCH. 2005 83:1397.
53. Abdallah T, Eglal A, Magdy K, Tarek M, Abdelaziz M. The Antagonistic Effect of Whey on Ochratoxin A toxicity on the Growth Performance, Feed Utilization, Liver and Kidney Functions of Nile Tilapia (*Oreochromis niloticus*). Middle East Journal of Applied Sciences. 2015 5:176-83.
54. Ringot D, Chango A, Schneider YJ, Larondelle Y. Toxicokinetics and toxicodynamics of ochratoxin A, an update. Chemico-biological interactions. 2006 159:18-46.
55. Pietsch C, Schulz C, Rovira P, Kloas W, Burkhardt-Holm P. Organ damage and hepatic lipid accumulation in carp (*Cyprinus carpio* L.) after feed-borne exposure to the mycotoxin, deoxynivalenol (DON). Toxins. 2014 6:756-78.
56. Perkins E, Griffin B, Hobbs M, Gollon J, Wolford L, Schlenk D. Sexual differences in mortality and sublethal stress in channel catfish following a 10 week exposure to copper sulfate. Aquatic toxicology. 1997 37:327-39.
57. Gupta S, Jindal N, Khokhar RS, Asrani RK, Ledoux DR, Rottinghaus GE. Individual and combined effects of ochratoxin A and *Salmonella enterica* serovar *Gallinarum* infection on pathological changes in broiler chickens. Avian Pathology. 2008 37:265-72.
58. Mir M, Dwivedi P. Immunopathology of ochratoxicosis-A in New Zealand white rabbits (*Oryctolagus cuniculus*). feedback. 2010 5:51.

59. Koynarski V, Stoev S, Grozeva N, Mirtcheva T, Daskalov H, Mitev J, et al. Experimental coccidiosis provoked by *Eimeria acervulina* in chicks simultaneously fed on ochratoxin A contaminated diet. *Research in veterinary science*. 2007 82:225-31.
60. Duarte SC, Lino CM, Pena A. Ochratoxin A in feed of food-producing animals: An undesirable mycotoxin with health and performance effects. *Veterinary microbiology*. 2011 154:1-13.
61. Kuiper-Goodman T, Scott P. Risk assessment of the mycotoxin ochratoxin A. *Biomedical and environmental sciences: BES*. 1989 2:179-248.

Table 1. Ingredient composition of the semipurified basal diet^a.

Ingredient	Amount (g/kg dry mixture)
Casein, vitamin free (USB #12866) ^b	290.0 g
Gelatin	80.0 g
Dextrin (USB #9004-53-9)	360.0 g
Cellulose	106.0g
Fish meal, menhaden	20.0 g
Carboxymethyl cellulose	30.0 g
Corn oil	30.0 g
Menhaden oil (USB #8002-50-4)	30.0 g
Mineral premix ^c	40 g
Vitamin premix ^d	12.5 g
Vitamin C ^e	1.5 g

^a This prepares 1000 g (1 kg) of diet with a calculated crude protein concentration of 32.7 %. Prepare feed in 2-4 kg batches.

^b USB: USB Corp., Cleveland, OH

^c Williams and Briggs mineral premix (Purina Mills Test Diets, Richmond, IN).

^d Vitamin premix supplies per kg of diet the following: vitamin A, 5500 IU; vitamin D3, 1835 IU; vitamin E, 110 IU; vitamin K, 7.3 mg; thiamin, 8.4 mg; riboflavin, 22 mg; pyridoxine, 18.4 mg; pantothenic acid, 58.7 mg; niacin, 36.7 mg; biotin, 2 mg; folic acid, 3.7 mg; vitamin B12, 0.018 mg; choline, 2327 mg; selenium, 0.1 mg.

^e L-ascorbyl 2-polyposphate (Stay-C, Hoffman La Roche, Nutley, NJ, USA, 25% ascorbic acid active)(Manning et al. 2003)

Table 2. HLP-1 levels in pooled skin extracts of channel catfish fed 0 or 8 mg OA/ kg diet. HLP-1 levels were measured at Day 0 and Day 56 of the experiment. Values for both the diluted acidic extract and concentration present in the undiluted tissue are reported; both are expressed in $\mu\text{g/ml}$ equivalents of calf histone H2B.

Time	Treatment group			
	0 (mg OA/ kg diet)		8 (mg OA/ kg diet)	
	Extract concentration	Tissue concentration	Extract concentration	Tissue concentration
Day 0	3.0	12.0	2.7	10.8
Day 56	3.9	15.6	6.4	25.6

Figure Legend

Figure 1. Weight (A) and condition factor (B) of channel catfish fed either 0, 2, 4, or 8 mg OA/ kg diet ochratoxin A. **Data is expressed as the mean of nine fish $\pm$ SEM. Values with a different letter superscript within a sampling day are significantly different between treatment groups (ANOVA, $p < 0.05$). N = 9 fish per treatment group per sampling time.**

Figure 2. Feed intake (FI) and feed conversion ratio (FCR) of channel catfish fed either 0, 2, 4, or 8 mg OA/ kg diet ochratoxin A. **Data is expressed as the mean of nine fish $\pm$ SEM. Values with a different letter superscript within a sampling day are significantly different between treatment groups (ANOVA, $p < 0.05$). N = 9 fish per treatment group per sampling time.**

Figure 3. Antibacterial activity Units (mean $\pm$ SE) in the skin of channel catfish fed either 0, 2, 4, or 8 mg OA/ kg diet ochratoxin A. N = 9 fish per treatment group per sampling time.

Fig. 4. Mortality curves of fish that had been fed various doses of ochratoxin A and challenged with water mold (*Saprolegnia*). **Curves with different letters were significantly different between treatment groups ($p < 0.05$) using the Wilcoxon test for each comparison prior to multiple testing correction. Performing Tukey's method for pairwise comparisons yielded identical results.**

Fig. 1

Fig. 2

Fig. 3

Fig. 4