The Green Clade grows: a phylogenetic analysis of *Aplastodiscus* (Anura; Hylidae).

Bianca V. M. Berneck^a; Célio F. B. Haddad^a; Mariana L. Lyra^a; Carlos A. G. da Cruz^b; and Julián Faivovich^{c,d*}.

^aDepartamento de Zoologia, Instituto de Biociências, UNESP - Universidade Estadual Paulista, Campus de Rio Claro, São Paulo, Brazil.

^bDepartamento de Vertebrados, Museu Nacional/UFRJ, Quinta da Boa Vista, Rio de Janeiro, Rio de Janeiro, Brazil.

^cDivisión Herpetología, Museo Argentino de Ciencias Naturales “Bernardino Rivadavia” CONICET, Buenos Aires, Argentina.

^dDepartamento de Biodiversidad y Biología Experimental, Facultad de Ciencias Exactas y Naturales, Universidad de Buenos Aires, Argentina.

*Corresponding author: jfaivovich@gmail.com

Short running title: A phylogenetic analysis of *Aplastodiscus*.

Abstract

Green tree frogs of the genus *Aplastodiscus* occur in the Atlantic Forest and Cerrado biomes of South America. The genus comprises 15 medium-sized species placed in three species groups diagnosed mainly by cloacal morphology. A phylogenetic analysis was conducted to: (1) test the monophyly of these species groups; (2) explore the phylogenetic relationships among putative species; and (3) investigate species boundaries. The dataset included eight mitochondrial and nuclear gene fragments for up to 6642 bp per specimen. Data strongly support the monophyly of *Aplastodiscus* and of the *A. albofrenatus* and *A. perviridis* groups. *Aplastodiscus sibilatus* is the sister taxon of all other species of *Aplastodiscus*, making the *A. albosignatus* Group non-monophyletic as currently defined. At least six unnamed species are recognized for *Aplastodiscus*, increasing the diversity of the genus by 40%. A fourth species group, the *A. sibilatus* Group is recognized. *Aplastodiscus musicus* is transferred from the *A. albofrenatus* Group to the *A. albosignatus* Group, and *A. callipygius* is considered a junior synonym of *A. albosignatus*. Characters related to external cloacal morphology reveal an interesting evolutionary pattern of parallelisms and reversions, suggesting an undocumented level of complexity. We analyze, in light of our phylogenetic results, the evolution of reproductive biology and chromosome morphology in *Aplastodiscus*.

Key words: Tree frogs, Hylidae, South America, molecular phylogeny, taxonomy.

1. Introduction

The hylid tribe Cophomantini includes *Aplastodiscus* Lutz, 1950; *Bokermannohyla* Faivovich, Haddad, Garcia, Frost, Campbell, and Wheeler, 2005; *Hyloscirtus* Peters, 1882; *Hypsiboas* Wagler, 1830; and *Myersiohyla* Faivovich, Haddad, Garcia, Frost, Campbell, and Wheeler, 2005. The largest genus is *Hypsiboas* with 90 species, followed by *Bokermannohyla* and *Hyloscirtus* with 33 species each, *Aplastodiscus* with 15 species, and *Myersiohyla* with six species forming the sister group to the remainder of the tribe (Faivovich *et al.*, 2005, 2013; Frost, 2014).

Aplastodiscus is distributed mainly in the Atlantic Forest of northeastern, southeastern, and southern Brazil, and adjacent Argentina, with one species reaching gallery forests in the Cerrado biome in central-eastern Brazil (Frost, 2014). All but one species are green, with usually colorful eyes with hues of copper, orange, pink, red, violet, and white (Cruz and Peixoto, 1985, 1987; Garcia *et al.*, 2001; Orrico *et al.*,

2006). *Aplastodiscus* formerly was included in the genus *Hyla* Laurenti, 1768; because of the coloration of the frogs, they have traditionally been called “the green species” of *Hyla* of the Brazilian Atlantic Forest.

Aplastodiscus was erected as a monotypic genus by A. Lutz (in B. Lutz, 1950) to include *A. perviridis*. The genus was considered valid or as a synonym of *Hyla* by different authors (Garcia *et al.*, 2001, and Faivovich *et al.*, 2005, for a review). Several authors (Lutz, 1950; Haddad and Sawaya, 2000; Garcia *et al.*, 2001; Hartmann *et al.*, 2004; Haddad *et al.*, 2005) presented phylogenetic grouping evidence for *Aplastodiscus* and some green species formerly included in *Hyla* (*i.e.*, *H. albosignata* and *H. albofrenata* complexes of the *H. albomarginata* Group of Cochran, 1955); this relationship was corroborated by Faivovich *et al.* (2005) in a comprehensive phylogenetic analysis.

In their review of hylid systematics, Faivovich *et al.* (2005) included 10 species of the complexes previously included in the former *H. albomarginata* Group in their taxon sampling—*viz.*, the *H. albofrenata* Complex (3 of 6 species included at that time), the *H. albosignata* Complex (4 of the 7 species included at that time), and the *H. albomarginata* Complex (3 of the 4 species included at that time). This was the first assessment of the relationships of *Aplastodiscus* and the former *Hyla albofrenata* and *H. albosignata* complexes. Their results showed that the former *Hyla albomarginata* Group was polyphyletic and that the exemplars of the *H. albofrenata* and *H. albosignata* complexes were individually monophyletic, and that *Aplastodiscus* was the sister taxon of the *H. albosignata* Complex.

Faivovich *et al.* (2005) redefined *Aplastodiscus* to include the former *Hyla albofrenata* and *H. albosignata* complexes, thereby increasing the number of species in the genus from two to fifteen. Furthermore, on the basis of the monophyly of their

exemplar species, they recognized each complex as a species group (*A. albofrenatus* and *A. albosignatus* groups), as well as a third group (the *A. perviridis* Group) for the two original species included in *Aplastodiscus*. Based on published information, Faivovich *et al.* (2005) suggested that the reproductive mode, advertisement call, development of metacarpal and metatarsal tubercles, and white parietal peritoneum are putative synapomorphies for *Aplastodiscus*. Putative phenotypic synapomorphies for the *A. perviridis* and *A. albosignatus* species groups also were discussed.

Wiens *et al.* (2005) presented a phylogenetic analysis of Hylidae with a smaller taxon sampling than Faivovich *et al.* (2005), including only *A. leucopygius* and *A. arildae* (as *H. leucopygia* and *H. albofrenata*, respectively). Their results recovered a large clade composed of the current genera *Aplastodiscus*, *Hypsiboas*, and *Bokermannohyla*, to which they applied collectively the generic name *Boana* Gray, 1825. Aside from *Boana* being a *nomem nudum* (Faivovich *et al.*, 2005), for reasons of priority, these combinations never gained acceptance. Subsequent reanalyses of the DNA sequences produced by Faivovich *et al.* (2005) yielded a topology for *Aplastodiscus* congruent with that obtained by these authors (Wiens *et al.*, 2005; Wiens *et al.*, 2006: supp. data; 2010; Pyron and Wiens, 2011; Pyron, 2014: supp. data).

Currently, three species groups are recognized in *Aplastodiscus*. The *A. perviridis* Group includes two species, *A. perviridis* and *A. cochranae* (Mertens, 1952) that share the lack of webbing between Toes I and II, reduction of webbing among the remaining toes, and a bicolored iris (Garcia *et al.*, 2001). The *A. albofrenatus* Group includes six species: *A. albofrenatus* (Lutz, 1924), *A. arildae* (Cruz and Peixoto, 1987), *A. ehrhardti* (Müller, 1924), *A. eugenioi* (Carvalho-e-Silva and Carvalho-e-Silva, 2005), *A. musicus* (Lutz, 1949), and *A. weygoldti* (Cruz and Peixoto, 1987). The monophyly of this group is supported only by molecular data (Faivovich *et al.*, 2005). The *A.*

albosignatus Group includes seven species: *A. albosignatus* (Lutz and Lutz, 1938), *A. cavicola* (Cruz and Peixoto, 1985), *A. callipygius* (Cruz and Peixoto, 1985), *A. flumineus* (Cruz and Peixoto, 1985), *A. ibirapitanga* (Cruz, Pimenta, and Silvano, 2003), *A. leucopygius* (Cruz and Peixoto, 1985), and *A. sibilatus* (Cruz, Pimenta, and Silvano, 2003). Species of this group share elaborated tubercles in the cloacal region (Cruz and Peixoto, 1985; Cruz *et al.*, 2003; Faivovich *et al.*, 2005).

Although the monophyly of the three groups has been tested and corroborated with several species, there are six species that have been unavailable for phylogenetic analyses, and the relationships of these with the groups have been inferred on the basis of putative phenotypic synapomorphies: *Aplastodiscus albofrenatus*, *A. ehrhardti*, *A. flumineus*, *A. musicus*, *A. ibirapitanga*, and *A. sibilatus*.

The taxonomy of *Aplastodiscus* was last reviewed by Cruz and Peixoto (1985, 1987), who described four species related to the former *Hyla albosignata* and three related to the former *Hyla albofrenata* (One of them was considered a junior synonym of the former *H. ehrhardti*; see Faivovich *et al.*, 2002.) Cruz *et al.* (2003) and Carvalho-e-Silva and Carvalho-e-Silva (2005) described three more species. Thus, the last taxonomic review of the genus was nearly 30 years ago, and several new populations have been discovered since then.

Our goal in this paper is to present a phylogenetic analysis of *Aplastodiscus* to test the monophyly of each species group, as well as to discuss the phylogenetic relationships among all included terminal taxa. On the basis of our results, we discuss the current taxonomy of the genus, putative synapomorphies, and the evolution of cloacal ornamentation, reproductive biology, and chromosome morphology.

2. Materials and Methods

2.1. Taxon sampling

We included 48 samples of 14 of the 15 currently recognized species of *Aplastodiscus*. If available, up to five specimens for each species from different localities were included, as well as topotypes for 10 of the species (complete list in Appendix A). *Aplastodiscus musicus* is the only species for which samples were unavailable. This species is known only from its type locality, where it was last collected in 1995 by A. Carvalho-e-Silva, S. Carvalho-e-Silva, and M. T. Rodrigues; despite multiple visits to the type locality, it has never been found again (Berneck, pers. obs.).

Outgroup terminals were selected on the basis of the current phylogenetic hypothesis of Cophomantini (Faivovich *et al.*, 2005) corroborated by several reanalyses (Wiens *et al.*, 2006: supp. data, 2010; Pyron and Wiens, 2011; Faivovich *et al.*, 2013). Thus, we included one member of each species group of *Hypsiboas* (the sister taxon of *Aplastodiscus*), *Bokermannohyla*, *Hyloscirtus*, and *Myersiohyla*; the latter was used to root the trees.

2.2. Character sampling and laboratory protocols

Our analyses included three fragments of mitochondrial genes: the almost complete ribosomal RNA gene *12S*, the intervening *tRNA-Val*, and partial sequence of *rRNA16S*, a fragment including the downstream region of the *rRNA 16S*, the intervening *tRNA-Leu*, the complete sequence of the *NADH dehydrogenase subunit 1* (*ND1*), *tRNAIle*, and a fragment of the gene *Cytochrome c oxidase I* (*COI*). The nuclear gene sequences analyzed include portions of *rhodopsin exon 1* (*RHOD*), *tyrosinase* (*TYR*), *Recombinase-Activation 1* (*RAG-1*), *Tensin 3* (*TNS3*), and *Seventh in Absentia homolog 1* (*SIAH*). DNA extraction, amplification, and sequencing methods are those

described by Blotto *et al.* (2013). The *COI* primers are those of Jungfer *et al.* (2013), the *TNS3* primers are of Smith *et al.* (2007), and the remaining primers used are those of Faivovich *et al.* (2013). See Appendix A for voucher data and the GenBank access numbers of sequences employed in this study. We included some sequences produced by Faivovich *et al.* (2005). For all samples we calculated uncorrected pair-wise distance for the *16S* fragment delimited by the primers *16sAR* (Palumbi *et al.*, 1991) and *Wilk2* (Wilkinson *et al.* 1996) and *COI* using PAUP (Swofford, 2002).

2.3. Phylogenetic analysis

The phylogenetic analysis under direct optimization was performed with POY5.1.1 (Wheeler *et al.*, 2014), using equal weights for all transformations (substitutions and insertion/deletion events). Sequences of *12S*, *16S*, *tRNAVal*, and *tRNALeu* were initially delimited in sections of putative homology (Wheeler *et al.*, 2006), and equal-length sequences of protein-coding genes were used as static alignments to accelerate the searches.

Searches were performed using the command "Search." This command implements a driven search, building Wagner trees with random addition sequences (RAS), Tree Bisection and Reconnection (TBR) branch swapping followed by Ratchet (Nixon, 1999), and Tree Fusing (Goloboff, 1999). The command (Search) stores the shortest trees of each independent run and computes the final fused tree using the pooled trees as a source of topological diversity. The resulting topologies were submitted to a final round of TBR using iterative pass optimization (Wheeler, 2003).

Phylogenetic analyses using POY were executed in parallel using the Museu de Zoologia da Universidade de São Paulo's high-performance computing cluster Ace, which consists of 12 quad-socket AMD Opteron 6376 16-core 2.3-GHz CPU, 16MB

cache, 6.4 GT/s compute nodes (= 768 cores total), eight with 128 GB RAM DDR3 1600 MHz (16 x 8GB), two with 256 GB (16 X 16GB), and two with 512 GB (32 X 16GB), and QDR 4X InfiniBand (32 GB/s) networking.

We performed a multiple alignment with MAFFT Version 7 (Kato and Standley, 2013). For the regions of *12S*, *tRNA-Val*, and the *16S*, *tRNA-Leu*, *ND1* and *tRNA-Ile*, we employed the alignments generated with Q-INS-i strategy (secondary structure of RNA is considered), whereas the alignments for the remaining genes were generated with G-INS-i (global homology considered). For the phylogenetic analysis using parsimony, we used T.N.T Willi Hennig Society Edition (Goloboff *et al.*, 2008). Searches were conducted using the new technology search under Search Level 50, which included sectorial searches, tree drift, and tree fusing (Goloboff, 1999), and requesting the driven search to hit the best length 100 times. Parsimony Jackknife absolute frequencies (Farris *et al.*, 1996) were estimated using new technology, as well requesting 10 hits with driven searches, for a total of 1000 replicates. Trees were edited with FigTree (Rambaut, 2014).

For the Bayesian analysis, models for each partition were chosen with jModelTest v0.1.1 (Posada, 2008). First, second, and third codon positions were treated as separate partitions for each protein-coding gene. The regions of *12S*, *tRNAVal*, *16S*, *tRNALeu*, and *tRNAIle* were treated as a single partition for model selection. The Akaike Information Criterion (AIC) was used to select the best fitting model for each gene (Pol, 2004; Posada and Buckley, 2004). Bayesian analyses were performed in MrBayes 3.2 (Ronquist *et al.*, 2012) in the CIPRES web cluster (Miller *et al.*, 2010). Analyses consisted of four runs, each consisting of two replicate Monte-Carlo Markov Chains. Each run used four chains and default settings of priors (Dirichlet for substitution rates and state frequencies, uniform for the gamma shape

parameter and proportion of invariable sites, all topologies equally likely *a priori*, and branch lengths unconstrained: exponential). Two analyses running 60 million generations were performed (with a burn-in fraction of 0.20). Stabilization of resulting parameters was evaluated using Tracer (Rambaut *et al.*, 2014).

3. Results

Direct Optimization in POY produced four most parsimonious trees (MPTs) with 11,054 steps (Fig. 1). The static parsimony analysis in T.N.T also yielded in four MPTs with 11,656 steps; Direct Optimization in POY, static parsimony in T.N.T., and Bayesian analysis revealed similar topologies (Appendix B). The main differences among alternate topologies for each optimality criterion are the internal relationships among terminals of *A. eugenioi* (a polytomy in Direct Optimization but not in static parsimony), and *A. leucopygius* (a polytomy in static parsimony but not in Direct Optimization). For the three analyses, see Figure 1 and Appendix B.

The monophyly of *Aplastodiscus* is supported with 100% Parsimony Jackknife Frequency (hereafter PJF) (Fig. 1). Six unnamed species were recognized for *Aplastodiscus* based on topology and genetic p-distances (Fig. 1). See Table 1 for species and clades distances and Appendix C for pair-wise comparisons among all samples; the discussion contains further details and evidence for each unnamed species. A clade formed by *A. sibilatus* and *Aplastodiscus* sp. 1 (aff. *sibilatus*) is the sister taxon of all other species of *Aplastodiscus*, making the *A. albosignatus* Group non-monophyletic as currently recognized. The remaining 13 described species are recovered as a well-supported clade with 95% PJF.

Figure 1: One of the four most parsimonious trees recovered by direct optimization. Black dots indicate nodes that collapse in strict consensus; an asterisk at nodes indicates 100% Parsimony Jackknife absolute frequency. The terminals included in Faivovich et al. (2005) are with an asterisk with names given by these authors in parenthesis. See Appendix A for details and Appendix B for results of the T.N.T. and Bayesian analyses.

All species of the *A. albofrenatus* Group are monophyletic and recovered with 100% PJF. This clade is the sister taxon of a clade composed of the monophyletic *A. perviridis* Group plus a well-supported clade that includes the remaining six species of the *A. albosignatus* Group, both with 100% PJF support. *Aplastodiscus ehrhardti* is the sister taxon of all other species of the *A. albofrenatus* Group. These are grouped in two clades; one includes *A. arildae* with 100% PJF, and the other has 77% PJF support, and includes *A. weygoldti*, *Aplastodiscus* sp. 2, and *A. albofrenatus* + *A. eugenioi*. The *A. perviridis* Group is monophyletic, with *A. perviridis* being the sister taxon of *A. cochranæ* + *Aplastodiscus* sp. 3.

Most species of the *A. albosignatus* Group are monophyletic, with 100% PJF, and with internal relationships having support values of 52–100%. *Aplastodiscus ibirapitanga* and *Aplastodiscus* sp. 4 form the sister taxon of the remaining species in the clade. These are further grouped in two clades. One includes *A. cavicola*, *A. leucopygius*, and *Aplastodiscus* sp. 6. The other clade includes *Aplastodiscus* sp. 5, *A. flumineus*, and a clade including topotypes of *A. albosignatus* and exemplars of *A. callipygius* (including topotypes).

Table 1: Ranges of uncorrected pair-wise sequence distances (p-distances) of the *I6s* fragment (below the diagonal) delimited by primers *Ar* and *Wilk2* and *Cytochrome c oxidase subunit 1* (above the diagonal). For individual values of all samples see Appendix C. Values are percentages.

Species	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20
1 <i>A. albofrenatus</i>	—	19.3	12.1–12.4	19.7	19.9	20.9–21.2	16.4–16.9	7.5–7.8	20.2	19.9	19–19.5	19.9–20.9	22.1	14.1–14.4	22.1	11.9	20.1	19.8–21.8	20.2	18.9
2 <i>A. albosignatus</i>	10.45	—	19.0–19.6	3.2	11.8	14.6–14.7	19.8–20.1	19–19.36	5.1	13.8	10.9–11.8	13.5–17.1	21	18.1–19.1	21	18.4	16.1	13.7–15.3	11	11.0
3 <i>A. arildae</i>	6.33–6.87	10.83–11.94	—	19.1–20.3	19.5–19.6	17.5–18.8	14.7–15.6	11.7–13.5	19.6–20.8	18.1–19	18–19.2	17.3–20.2	21.0–21.5	12–14.1	21–21.2	11.9–12.9	17.5–17.9	17.4–20	18.8–19	18.4
4 <i>A. callipygius</i>	10.8–11.1	0.3–0.7	10.8–12.3	—	12.6	14.6–14.8	19.7–19.7	19.5–19.6	4	13.1	10.4–11.4	13.2–16.5	20.7	18.3–18.6	20.7	19.1	15.8	13.2–14.9	11.2	9.7
5 <i>A. cavicola</i>	11.3–11.7	4.77–4.78	11.75–13.4	4.5–4.7	—	12.6	18.1–18.5	20.4–20.7	11.9	14.4	8.3–9.8	13.8–15.5	20.8	19.9–20.2	20.8	18.7	13.8	13–15.3	12.4	9.8
6 <i>A. cochranæ</i>	11.2–11.4	5.32–5.67	12.72–13.82	5.1–5.6	6.9–7.2	—	19.9–20.4	20.4–20.9	14.8–15.6	16.2–16.5	13.2–15.2	9–10.3	20.3–21.3	17.9–18.9	20.3–21.3	19.5–19.6	9–9.4	14.3–16.3	13.6–13.7	11.8–12.3
7 <i>A. ehrhardti</i>	8.4–11.4	11.93–11.98	8.18–10.49	12.1–12.4	12.1–13.4	12–12.7	—	15.4–16.4	19.6–19.9	19.8	18.9–19.9	18.1–20.4	22.5–22.7	16.9–17.4	22.5–22.7	15.6–15.8	18.4–19	18.7–20.8	19.2–19.5	19.3–19.8
8 <i>A. eugenioi</i>	2.4–3.3	9.54–10.12	5.70–7.01	9.9–10.4	11.2–11.4	11.1–11.5	7.8–9.5	—	20.6–21.0	19.8–21	18.8–19.9	18.3–20.8	21.9–22.3	12.3–14.7	21.9–22.3	11.4–12.7	19.6–19.9	19.1–21.2	18.8–19.2	18.2–19.5
9 <i>A. flumineus</i>	11.71	2.6	11.9–12.8	2.2–2.6	5.4–5.6	5.1–5.5	12.3–12.4	10–10.4	—	13.1	10.2–11.4	13.6–16.8	19.5	18.7–19	19.5	18.5	15.9	12.8–14.4	10.5	10.2
10 <i>A. ibirapitanga</i>	11.2–11.7	4.9–5.0	12.1–12.7	4.7–5.2	5.8–6.5	6.7–8.4	12.5–13.5	10.3–11.4	4.4–5.5	—	12.9–13.6	14.5–17.8	20.1	18.2–18.2	20.1	19	16.1	9.13–10.4	14.5	13
11 <i>A. leucopygius</i>	11–11.2	3.1–3.3	11.2–12.1	3–3.3	4–5.1	5.8–6.5	12–12.7	10.1–10.6	3.7–3.9	4.1–6.3	—	13–16.4	20.3–20.5	17.5–19.4	20.3–20.5	18.6–19.3	13.3–14.5	12.8–14.7	11.8–13.5	6.2–7.2
12 <i>A. perviridis</i>	11.2–11.8	6.2–6.7	12.1–13.6	6.0–6.9	6–7.6	2.1–3	12.2–13.1	11.4–11.8	6.3–7	7.1–8	6–6.4	—	20.1–21.6	17–19.3	20.1–21.6	18.2–19.9	9.2–10.4	13.3–17.3	14.4–16.2	12.5–13.8
13 <i>A. sibilatus</i>	10.1–10.8	12.7–13.2	12.8–14.1	12.7–13.6	12.2–13.9	12.9–13.1	11.4–12.3	10–11.8	12.7–13.6	12.4–13.3	13.6–14.3	13.1–14	—	21.9–23.4	—	20.8	21.2	20.3–21.6	21.8	20.9
14 <i>A. weygoldti</i>	7.5	11.1	6.7–7.6	11.5–11.5	11.5–12.3	12.6–12.8	9.8–10.2	7–7.2	12.2	11.2–11.6	11.2–11.4	12.4–12.4	11.4	—	21.9–23.4	14.5–15.6	18.4–19.0	17.4–19.6	18.7–19.2	18.4–19.5
15 <i>A. sp. 1</i>	10.1	12.7	12.6–12.8	12.7–13.1	13.1–13.3	13.1–13.3	11.4–12.1	10–10.1	12.7	12.1–12.8	13.6–13.8	13.1–14	3.18	11.4	—	20.8	21.2	20.3–21.6	21.8	20.9
16 <i>A. sp. 2</i>	6.7–6.8	10.4–10.5	6.1–7.4	10.4–10.6	11–11.2	11–11.2	9.1–9.3	6–6.2	11	11.8–12	10.1–10.9	10.9–11	11.4–11.6	6.9–7.1	11.4–11.6%	—	19.6	18.9–20.7	17.9	18.9
17 <i>A. sp. 3</i>	12.3	7	13.2–14.1	6.9–7	7.2–8.8	4	12.2–12.7	11.4–12	6.2	7.9–9.5	7.1–7.2	4.6–5.5	14.4	13.5	14.4	11.8	—	15.4–17	15.6	12.9
18 <i>A. sp. 4</i>	11.2	5.1	12.7–13.4	4.6–5.1	6.2–6.3	7.6–8	12.7–13.1	11–11.2	5.4	2.6–3.1	5.1–5.7	7.4–8.3	12.9	11.4	12.9	11.2	9	—	12.3–13.5	13–14.2
19 <i>A. sp. 5</i>	10.1–11.2	2.6–3.7	10.6–13	2.6–3.8	4.9–6	4.9–5.8	11.4–12	9.4–10.4	2.8–3.3	3.9–5.2	3.2–4.4	5.3–6	12.2–12.5	11.2–11.5	12.2–12.5	9.9–10.4	6–6.3	4.8–5.4	—	11.6
20 <i>A. sp. 6</i>	11.2–11.5	3.5–3.7	11.4–12.7	3.5–3.8	3.8–4.7	6.3–6.9	12–12.5	10.6–11	3.53–4.23	4.2–6.1	2.3–3.7	6.5–7.4	13.1–13.3	11.5–11.9	13.1–13.3	10.3–11	6.7–7.4	5.1–5.5	3.3–4.7	—

4. Discussion

4.1. Outgroups

As in previous phylogenetic studies, *Hyloscirtus* is the sister taxon of a clade comprising *Hypsiboas*, *Aplastodiscus*, and *Bokermannohyla*, and these genera are monophyletic (Faivovich *et al.*, 2005; Pyron and Wiens, 2011; Faivovich *et al.*, 2013). The placement of *Hypsiboas* as the sister group of *Aplastodiscus* is corroborated by our analysis. The internal relationships in *Hypsiboas* differ from those recently recovered by Faivovich *et al.* (2013). However, our study has a much reduced taxon sampling for the study of the relationships within *Hypsiboas*, and thus, does not constitute a valid test of previous phylogenetic hypotheses within this genus.

4.2. *Aplastodiscus*

Our analysis corroborates the monophyly of the *A. albofrenatus* and *A. perviridis* groups. However, the position of *A. sibilatus* as the sister taxon of all other species of *Aplastodiscus* makes the *A. albosignatus* Group not monophyletic as currently recognized. The relationships recovered among species groups of *Aplastodiscus* (in which the *A. albofrenatus* Group is sister taxon of the *A. albosignatus* + *A. perviridis* groups) corroborate previous results (Faivovich *et al.*, 2005; Pyron and Wiens, 2011; Faivovich *et al.*, 2013).

Our results indicate the possible existence of at least six unnamed species of *Aplastodiscus*, which represents 40% of the known diversity of this clade. This inference stems from the analysis of our topological results, vouchers, and current taxonomic knowledge of the group. All these putative new species have been confused in collections, something usual with the admittedly complex taxonomy of *Aplastodiscus*.

4.3. The position of *A. sibilatus*

In its original description, *Aplastodiscus sibilatus* was associated with *A. albosignatus*, *A. callipygius*, and *A. flumineus* on the basis of the absence of a cloacal flap (Cruz *et al.*, 2003). In the context of our results, this character state is plesiomorphic for *Aplastodiscus* because the taxonomic distribution of the cloacal flap is restricted to some species of the *A. albosignatus* Group (per discussion below); thus, there is no conflict with our finding that *A. sibilatus* is the sister taxon of all other species of *Aplastodiscus*.

Aplastodiscus sibilatus is known from Atlantic Forest patches in southern and central Bahia, and in southwestern Alagoas (Cruz *et al.*, 2003; Lima *et al.*, 2006). Two samples of this clade were included, one from southern Bahia and the other from Alagoas (Appendix A). These two samples have a p-distance of 3.18% for *16S* (Table 1 and Appendix C). A parallel morphological study by Cruz *et al.* (in prep.) concludes that the voucher from Alagoas belongs to an unnamed species, morphologically distinct from *A. sibilatus*. The p-distances corroborate the specific identity of the population from Alagoas, here referred to as *Aplastodiscus* sp. 1 (Fig. 1).

The topology of *A. sibilatus* and the unnamed species necessitates the recognition of a fourth group of species in *Aplastodiscus*, the *A. sibilatus* Group. Some character states, in contrast with other species of *Aplastodiscus* and the outgroups, uniquely characterize *A. sibilatus* and *Aplastodiscus* sp. 1 and are considered putative phenotypic synapomorphies of the *A. sibilatus* Group—*viz.*, subcloacal dermal fold (Cruz *et al.*, 2003; Appendix D) and white gastrointestinal peritoneum (Berneck, pers. obs.). According to Mercês and Juncá (2010), tadpoles of *A. sibilatus* differ from other tadpoles described for *Aplastodiscus*; however, further studies on Cophomantini are

needed to confirm any putative synapomorphies for the *A. sibilatus* Group or other clades of *Aplastodiscus*.

4.4. The *Aplastodiscus albofrenatus* Group and the putative position of *A. musicus*.

The erection of the *Hyla albofrenata* Complex (now the *A. albofrenatus* Group) was a consequence of the recognition that the name *Aplastodiscus albofrenatus* actually was applied to several different species (*A. arildae*, *A. ehrhardti*, and *A. weygoldti*; Cruz and Peixoto, 1987; Faivovich *et al.*, 2002). Faivovich *et al.* (2005) reported molecular support for the monophyly of this group. (Cochran, 1955; Lutz, 1973; Cruz and Peixoto, 1985; among others). Some authors suggested a red- orange iris as a possible synapomorphy of the group (Carvalho-e-Silva and Carvalho-e-Silva, 2005; Orrico *et al.*, 2006). However, we find that iris color varies among individuals of the same population; at the type locality of *A. arildae*, we recorded two individuals with copper-colored irises, whereas all other individuals had red irises. We have similar observations for *A. calipygius* (of the *A. albosignatus* Group), which is known to have a reddish iris; however, we recorded individuals from Campos do Jordão (State of São Paulo) with white irises. A similar situation was reported by Rivera-Correa and Faivovich (2013) for *Hyloscirtus larinyption*. Whereas there is notable variation in the pattern and coloration of irises within *Aplastodiscus*, we think it premature to hypothesize homologies until intraspecific variation and the nature of iris coloration are better known.

The only species of *Aplastodiscus* not available for our study is *A. musicus*, a species that was assigned to the *A. albofrenatus* Group by Cruz and Peixoto (1987) without specific discussion. In her description of *A. musicus*, Lutz (1949) noted some unusual characters, such as an unpigmented nuptial pad in males and a strong peculiar

odor of a slimy secretion—characters absent in all species now included in *Aplastodiscus*. A strong odor was reported for *Myersiohyla* by Faivovich *et al.*, (2013), and for *Hyloscirtus* by Rivera-Correa and Faivovich, (2013). Lutz (1949) also noted the musical call of *A. musicus*, reminiscent of “an old fashioned glockenspiel,” after which the species was named. She described “a green tree-frog closely akin to *H. albofrenata* and especially to *H. albosignata*” and “though nearer to *H. albosignata*, it is less heavy in build, has a shorter and wider snout and lacks the rows of yellow, milium-like, post-anal glands.” The contents of the original description and the study of external morphology of six topotypes of *Aplastodiscus musicus* (Appendix E) suggest the association of *A. musicus* with the *A. albosignatus* Group rather than the *A. albofrenatus* Group.

Our results indicate the existence of one new species (*Aplastodiscus* sp. 2) of the *A. albofrenatus* Group. This species is the sister taxon of *A. albofrenatus* + *A. eugenioi*, and was confused with *A. eugenioi* by previous authors (Salles *et al.*, 2012). Although this species is morphologically difficult to diagnose from *A. albofrenatus* and *A. eugenioi*, its specific distinctiveness is supported by a unique advertisement call (Berneck, pers. obs.). Uncorrected p-distances of *16S* between *A. albofrenatus* and *A. eugenioi* are 2.4–3.3%. The latter has been distinguished from *A. albofrenatus* by a smaller snout–vent length, larger calcar tubercle, femur and tibia size, iris color, morphology and coloration of the tadpole (Carvalho-e-Silva and Carvalho-e-Silva, 2005), and was considered *A. aff. ehrhardti* by Hartmann *et al.* (2004). Our data suggest that the diagnostic characters of *A. albofrenatus* and *A. eugenioi* require further study.

Only four species of this group were included in previous analyses. The voucher specimen identified as *A. weygoldti* by Faivovich *et al.* (2005; and subsequent analyses) has been re-identified as *A. arildae* in our analysis. The specimen is from Domingos

Martins (State of Espírito Santo). Apparently, both species occur at this locality (Silva-Soares *et al.*, 2011; Silva *et al.*, 2012). Orrico *et al.* (2006) described the call of *A. weygoldti* from Domingos Martins and compared it with topotypic *A. arildae*, finding them to differ significantly in duration (details in Orrico *et al.*, 2006).

Wiens *et al.* (2005) included sequences of *12S* and *16S-tRNA^{Leu}-ND1* fragment of a voucher specimen identified as *Aplastodiscus albofrenatus* (USNM 208734), collected in 1975 by W. R. Heyer in Boraceia (Salesópolis, São Paulo). So far, only *A. arildae* is known from that locality (Cruz and Peixoto, 1987; Berneck, unpubl.). Without further comments, Wiens *et al.* (2006) replaced the *12S* fragment with the sequence DQ380346, produced from a voucher identified as *A. albofrenatus* (USNM 303022) from the same locality. The latter sequence is identical to our sequences of *A. arildae* from Boraceia. The position of the terminal USNM 208734 in the analysis of Pyron and Wiens (2011), as the sister taxon of the remaining exemplar species is unusual. Our study of the voucher specimen USNM 208734 indicates that it actually is *A. arildae*. A study of the available *12S* sequences shows striking differences even in conserved regions, when compared with the sequences of other specimens of *A. arildae* and also all other species of the genus. This pattern might indicate problems with sequence quality (Lyra, pers. obs.), but the unavailability of the chromatograms hinders the confirmation of this hypothesis. We did not include the sequences from Wiens *et al.* (2005, 2006) because they are nearly identical to the samples we have from the same locality (*12S*, Boraceia); moreover, apparently there are sequencing-quality issues.

The *A. albofrenatus* Group is distributed in the Atlantic Forest of Brazil from Bahia (15°25'S 39°34'W) to Santa Catarina (27°46'S; 49°45'W). The distributions of most members of this group are associated with coastal mountain systems (Serra do Mar, Serra Bonita, Serra da Bocaina), with the notable exception of *A. arildae*, the

distribution of which extends westward toward the Serra da Mantiqueira Range and central regions of Minas Gerais (19°56'S; 43°54'W). The first putative phenotypic synapomorphies for the *A. albofrenatus* Group are discussed below in “Evolution of external morphology of the cloacal region.”

4.5. The *Aplastodiscus perviridis* Group

Based on observations by Garcia *et al.* (2001), Faivovich *et al.* (2005) suggested the following putative synapomorphies for the *A. perviridis* Group—*viz.*, the bicolored iris (the superior third white silver and inferior two thirds reddish copper); absence of an interdigital membrane between Toes I and II (with instances of homoplasy as identified by Faivovich *et al.*, 2005); and reduction of webbing between the other toes. In addition, we consider the following features to be putative synapomorphies—the absence of white parietal peritoneum (present in all other species of *Aplastodiscus*) and absence of any cloacal ornamentation as described for the *A. albosignatus*, *A. albofrenatus*, and *A. sibilatus* groups. (In *A. cochranæ*, the cloaca is not glandular; only a pigmentary white line is present).

The *A. perviridis* Group includes the nominal species, *Aplastodiscus* sp. 3 a species identified as *A. perviridis* in previous works (*e.g.*, Bastos *et al.*, 2003; Valdujo *et al.*, 2012), and its sister species, *A. cochranæ*. The specific status of *Aplastodiscus* sp. 3 is supported by a smaller snout–vent length and longer advertisement call than those of *A. perviridis*. The uncorrected p-distances in *16S* of *Aplastodiscus* sp. 3 are 4.6–5.5% with *A. perviridis* and 4.8% with *A. cochranæ* (Table 1). The *A. perviridis* Group is distributed from the states of Minas Gerais and São Paulo, southward to northeastern Argentina and Rio Grande do Sul, and westward to reach the gallery forests in the Cerrado of central-eastern Brazil.

4.6. The *Aplastodiscus albosignatus* Group

Our results indicate that the *A. albosignatus* Group comprises seven described species (given the exclusion of *A. sibilatus* as discussed above and the inclusion of *A. musicus*), as well as three unnamed species—viz., *Aplastodiscus* sp. 4, *Aplastodiscus* sp. 5, and *Aplastodiscus* sp. 6.

Cruz and Peixoto (1985) distinguished some species of this group by the presence of a white dermal ridge above the cloacal opening (“flap”). Faivovich *et al.* (2005) suggested that the combination of elaborate tubercles and ornamentation around the cloacal region is a putative morphological synapomorphy of this group. All characters related to the pericloacal region are discussed below in “Evolution of external morphology of the cloacal region.” The presence of a white submandibular dermal fold is also a putative morphological synapomorphy of this group, because it is absent in all species of *Aplastodiscus* from the other three groups and outgroups studied so far.

Abrunhosa *et al.* (2005) studied nine call parameters of five species of the *Aplastodiscus albosignatus* Group and *A. sibilatus* (then in the *A. albosignatus* Group). These authors proposed a phenetic grouping of species based on their observation that the most important parameter is the dominant frequency, which can be found in different harmonics. However, Zina and Haddad (2006b) reported that the dominant frequency varies, depending on social context for *A. leucopygius* and *A. arildae*; thus, it probably does not adequately characterize calls of *Aplastodiscus*.

The topology of the clade including exemplars *Aplastodiscus albosignatus* and *A. callipygius* indicates that topotypes of the latter are nested within topotypes of the former. The 16S distances among exemplars of this clade vary from 0.35–0.70%

(Appendix C). *Aplastodiscus callipygius* and *A. albosignatus* also are morphologically similar. Both have a supratympanic fold, calcar tubercle, and cloacal ornamentation. These species were diagnosed originally by snout shape (rounded in *A. albosignatus* and pointed in *A. callipygius*), size of vocal sac (medium in *A. albosignatus* and large in *A. callipygius*), and size of calcar tubercle (small in *A. albosignatus* and large in *A. callipygius*). However, these differences are not consistent among the many specimens available for study after the description of *A. callipygius*. Abrunhosa *et al.* (2005) described the calls of *A. albosignatus* and *A. callipygius* as differing somewhat from each other; however, the population on which the call descriptions were based for *A. albosignatus* is, in fact, *Aplastodiscus* sp. 5 herein (Ribeirão Branco, São Paulo, Brazil). Thus, we consider *Aplastodiscus callipygius* (Cruz and Peixoto, 1985) a junior synonym of *A. albosignatus* (Lutz and Lutz, 1938) based on the position of the topotypes of *A. albosignatus*, nested among the samples of *A. callipygius*, their morphological resemblance (Appendix F), and low genetic distances.

Our results reveal three previously unrecognized species in the *A. albosignatus* Group. *Aplastodiscus* sp. 4, the sister species of *A. ibirapitanga*, differs from that species by having a distinct advertisement call (Berneck, pers. obs.) and smaller size (Pimenta *et al.*, 2014; Cruz *et al.*, 2003). Furthermore, the *16S* distances are 2.66–3.16%. *Aplastodiscus* sp. 5 was previously identified as *A. albosignatus* (e.g., Cruz and Peixoto, 1985; Abrunhosa *et al.*, 2005), but our best hypotheses indicate that it is only distantly related to *A. albosignatus*. Furthermore, it differs in the morphology of the cloacal region. *Aplastodiscus* sp. 6, a species of the clade of *A. leucopygius*, was misidentified initially as *A. leucopygius* or *A. cavicola*. The vouchers are medium-sized and match both species. *Aplastodiscus* sp. 6 is diagnosed by morphology of the pericloacal region, and the *16S* genetic distances are 3.89–4.78% with *A. cavicola* and

2.3–3.72% with *A. leucopygius*. The same clade includes the sample of Faivovich *et al.* (2005) from Domingos Martins (Brazil, Espírito Santo) identified by these authors as *A. cavicola* owing to proximity to the type locality. Samples of topotypes of *A. cavicola* were not available; therefore, we assigned the terminals belonging to this species on the basis of the vouchers and the holotype.

4.7. Evolution of external morphology of the cloacal region

The presence of calcar tubercles and dermal folds on the limbs (as in most species of *Aplastodiscus*) has been considered as mimetic and disruptive morphology, especially in resting position (Duellman and Trueb, 1986). However, little attention has been given to cloacal ornamentation in anurans and its possible role in disruptive morphology. Several groups of arboreal frogs (*e.g.*, Duellman, 1970; Cisneros-Heredia and McDiarmid, 2007; Faivovich, pers. obs.) have pericloacal ornamentation. Because the variation of pericloacal ornamentation in *Aplastodiscus* is taxonomically informative, it is well described for many species. Information available in the literature allowed us to study the evolution of some characters related to the morphology of the pericloacal region of *Aplastodiscus* and to standardize the terms suggested by different authors (Appendix D).

Our study revealed four hypotheses of character evolution for *Aplastodiscus* and outgroups (Fig. 2). Faivovich *et al.* (2005) suggested that the presence of ornamentation around the cloaca and elaborated tubercles are putative synapomorphies of this group. We redefined this character with the understanding that the tubercles in the subcloacal region are independent of the pericloacal ornamentation (optimizations in outgroups, character descriptions and figures in Appendix D). The presence of pericloacal ornamentation is a synapomorphy of the *A. albosignatus* Group (present in all species of this group including *A. musicus*, with one instance of homoplasy in *A. albofrenatus*

and *A. eugenioi*). The occurrence of an elliptical area of pericloacal ornamentation around the cloacal opening is a synapomorphy for the clade comprising *A.*

albosignatus, *A. flumineus*, and *A. sp. 5*.

Our study indicates a putative morphological synapomorphy of the *A. albofrenatus* Group, the presence of iridophores in the outer cloacal epithelium (with an instance of homoplasy in *A. sp. 6*). The subcloacal dermal fold is a putative synapomorphy for the *A. sibilatus* Group.

The optimization of the characters from pericloacal morphology shows an interesting pattern of parallelisms and reversions, suggesting a level of complexity undocumented thus far (Appendix D). It is unclear why only *Aplastodiscus* among the Cophomantini has this diversity of external cloacal morphology. The morphology described in the *Hyloscirtus bogotensis* and *H. larinopygion* groups possibly differs histologically and anatomically (Faivovich pers. obs.). Members of the *A. perviridis* Group have the same arboreal habits as their congeners; however, cloacal and tarsal ornamentation are absent in all three species, which brings more complexity to the evolution of characters of the cloacal region.

Figure 2. Optimizations of the four pericloacal characters studied. The characters are (a) Pericloacal ornamentation (yellow: present, restricted to the supracloacal region, blue: present and elliptical around the cloacal opening; black: absent). (b) Iridophores in the outer cloacal

epithelium (blue: present; black: absent). (c) Concentration of heterogeneous granules below the cloacal opening (blue: present; black: absent). (d) Subcloacal dermal fold (blue: present; black: absent). See Appendix D for figures of characters and more details.

4.8. Reproductive biology

The reproductive biology of *Aplastodiscus* was first reported in detail for *A. leucopygius* (Haddad and Sawaya, 2000), although there were earlier anecdotal observations (e.g., Lutz and Lutz, 1938; Cochran, 1955; Cruz and Peixoto, 1985; 1987). In this species, the male excavates a burrow with an inconspicuous entrance in the mud near, or in, the stream edge. The male then finds a spot to call for females, usually perched on vegetation distant from the ground. When a female approaches a calling male, both exchange a complex sequence of tactile stimuli that includes contact of the hind limbs and snout with hands, snout with snout, and chin on head (Haddad and Sawaya, 2000; Hartmann *et al.*, 2004; Haddad *et al.*, 2005; Carvalho-Jr. *et al.*, 2006; Zina and Haddad, 2006; 2007), where sexually dimorphic skin glands have recently been described in males (Brunetti *et al.*, 2015). The transit from the calling and meeting point to the burrow can last from 40 min to 9 h (Hartmann *et al.*, 2004; Zina and Haddad, 2007); during this period, the male emits a courtship call (similar to the advertisement call but less intense) while the female follows him to the burrow. Still, after inspection, the female may reject the burrow and leave (Hartmann *et al.*, 2004; Zina and Haddad, 2007; Carvalho *et al.*, 2006). Clutches of *Aplastodiscus* have about 230 unpigmented eggs, which are deposited in a floating layer inside the burrow where they spend the first stages of development (Haddad *et al.*, 2005). After hatching, the exotrophic tadpoles complete metamorphosis in lotic water (Haddad and Prado, 2005).

Haddad *et al.* (2005) described the reproductive biology of *Aplastodiscus perviridis*. For the *A. albofrenatus* Group, data are available for *A. arildae* (Zina and Haddad, 2006a,b; 2007; Carvalho-Jr. *et al.*, 2006) and *A. eugenioi* (Hartmann *et al.*,

2004). The descriptions are quite similar to that of *A. leucopygius*. In the *A. albosignatus* Group, data on reproductive biology have been reported for *A. albosignatus* (as *A. callipygius*; Gomes and Peixoto, 1997), *A. leucopygius* (Haddad and Sawaya, 2000; Zina and Haddad, 2006a,b; 2007), and *A. cavicola* (Cruz and Peixoto, 1985). *Aplastodiscus cavicola* commonly calls from inside the burrow (Cruz and Peixoto, 1985). Lutz and Lutz (1938) and Cochran (1955) reported the same behavior for a species identified as *H. albosignata*: it is likely that the species is the one called *A. leucopygius* today. Gomes and Peixoto (1997) and Carvalho-Jr. *et al.* (2006) reported male *A. albosignatus* (as *A. callipygius*) and *A. arildae*, respectively, calling from rock crevices. Calling sites are intra- and interspecifically variable (Cruz *et al.*, 2003); on one occasion we found on the same night and along the same stream, a male of *A. ibirapitanga* calling from inside a burrow, whereas others called from shrubs. Males probably call from inside the nest while excavating it and, after the construction is finished, climb the vegetation then used as a calling site (Haddad, pers. obs.). The reproductive biology of *A. sibilatus* is unknown. Cruz *et al.* (2003) reported that male *A. sibilatus* call from leaf litter, shrubs, and trees, along streams, always above the water level.

Despite occasional observations of males of other species of the *A. albosignatus* Group calling from inside burrows, *A. cavicola* is the only species in which this behavior is common (Cruz and Peixoto, 1985; Cruz *et al.*, 2003; Berneck, pers. obs.). It is unknown if the courtship behavior described for other species of *Aplastodiscus* occurs in this species as well, or if the behaviour is modified with respect to the females encountering males directly in the burrow, without the pair moving from an outside meeting point to the nest.

Hyliid frogs usually do not call from burrows. Wells (2007) noted that calling from inside burrows, subterranean nests, and rock crevices can interfere with sound transmission, but in some species, the burrow facilitates resonance. In *Aplastodiscus*, only males of the *A. albosignatus* Group call from burrows, and these species have the lowest frequency of advertisement calls of all members of the genus (calls descriptions in Haddad and Sawaya, 2000; Garcia *et al.*, 2001; Hartmann *et al.*, 2004; Abrunhosa *et al.*, 2005; Conte *et al.*, 2005; Haddad *et al.*, 2005; Orrico *et al.*, 2006; Zina and Haddad, 2006b). Lower frequency advertisement calls usually are associated with sound propagation over long distances, and the call of *A. cavicola* has the lowest frequency of all species in the *A. albosignatus* group (Abrunhosa *et al.*, 2005). The taxonomic distribution of calling from burrows may represent a synapomorphy for one of the internal clades of the *A. albosignatus* Group.

4.9. Chromosome evolution in *Aplastodiscus*

With few exceptions, the diploid number in the subfamily Hyliinae is 24 chromosomes (Catroli and Kassahara, 2009; Catroli *et al.*, 2011). So far as is known, among hylids only *Aplastodiscus* has reductions in the diploid chromosome number involving the small-sized pairs (Bogart, 1973; Carvalho *et al.*, 2009 a,b; Gruber *et al.*, 2012). Members of the *A. perviridis* Group (*A. cochranae* and *A. perviridis*) have a $2n = 24$ complement, whereas those of the *A. albofrenatus* group have a $2n = 22$ complement (known in *A. albofrenatus*, *A. arildae*, *A. ehrhardti*, and *A. eugenioi*), and those of the *A. albosignatus* Group have a $2n = 20$ or $2n = 18$ complement (*A. albosignatus* with $2n = 20$; *A. leucopygius* with $2n = 18$; Carvalho *et al.* 2009a, b; Gruber *et al.*, 2012). The karyotype of *A. sibilatus* is unknown.

Figure 3: Optimization of chromosome complement showing reductions in *Aplastodiscus*. Note that the synapomorphies are the events of fusion, which causes the reduction in chromosome number. Data are from Bogart (1973), Carvalho *et al.* (2009a, b), and Gruber *et al.* (2012). See text for more details.

On the basis of the plesiomorphic $2n = 24$ complement inferred for Hylini (Faivovich *et al.*, 2005), Gruber *et al.* (2012) hypothesized homeologies for several chromosomes in *Aplastodiscus*, and suggested a plesiomorphic condition of $2n = 24$ for

the genus. These authors also suggested two independent events of fusion—one giving rise to large-sized pairs for species with karyotypes of $2n = 20$ and $2n = 18$ (*A. albosignatus* Group) from small chromosomes (Pairs 7–10), and another involving fusion of the small Pair 12 chromosomes and the large Pair 3 chromosomes in species of $2n = 22$ (*A. albofrenatus* Group.)

Although our results support the occurrence of independent reductions in chromosome complements in the *Aplastodiscus albofrenatus* and *A. albosignatus* groups, our increased taxon sampling includes many species with unknown karyotypes; thus, there are several ambiguities regarding the nodes at which these chromosome reductions actually occurred in the *A. albosignatus* Group (Fig. 3). It is most parsimonious to infer that the fusion of Pairs 12 and 3, leading to a $2n = 22$ karyotype in four species of the *A. albofrenatus* Group (*A. albofrenatus*, *A. arildae*, *A. ehrhardti*, *A. eugenioi*) also occurred in *A. weygoldti* and *Aplastodiscus* sp. 2. The karyotype of *A. sibilatus* is unknown, but putative fusions of Pairs 3 and 12 occur neither in the *A. perviridis* and *A. albosignatus* groups, nor in any other known clade of Cophomantini (e.g., Catroli and Kasahar, 2010; Catroli et al., 2011). It is more parsimonious to infer a $2n = 24$ karyotype for *A. sibilatus* and to consider this fusion a synapomorphy of the *A. albofrenatus* Group. (Note that the synapomorphy is the fusion, with the reduction from $2n = 24$ to $2n = 22$ being a consequence of the fusion.)

Transformations in species of the *Aplastodiscus albosignatus* Group leading to the reductions to $2n = 20$ and $2n = 18$ are ambiguous, because karyotypes are known for only three of the nine species included in the group. In the context of our hypothesis, it is most parsimonious to interpret the rearrangement leading to the reduction to $2n = 20$ in *A. albosignatus* to be a synapomorphy at least at the level of the

common ancestor of *A. albosignatus*, *A. cavicola*, *A. flumineus*, *A. leucopygius*, *Aplastodiscus* sp. 5, and *Aplastodiscus* sp. 6. Thus far, the additional rearrangement leading to the $2n = 18$ complement has been inferred only in *A. leucopygius*; given that the karyotype is unknown in the closely related *A. cavicola* and *Aplastodiscus* sp. 6, the node at which these rearrangements occurred is ambiguous. When the karyotype of *A. ibirapitanga* is determined, it will be possible to ascertain whether the chromosomal reduction events (*i.e.*, $2n = 24$ to $2n = 20$) are a synapomorphy (or synapomorphies if more than one event can be inferred) of the *A. albosignatus* Group (Fig. 3).

5. Final remarks

Our phylogenetic analyses yielded a stable phylogenetic hypothesis with high support for most nodes. We highlight four major clades of *Aplastodiscus* as named species groups, each of which contains at least one newly revealed species. Our phylogenetic results revise major hypotheses of morphological, behavioral, and chromosomal evolution in *Aplastodiscus*, and identify the data needed for critical testing of these hypotheses.

Acknowledgments

This work was part of the PhD of BVMB, who is indebted to many colleagues who helped collect samples and conduct field and laboratory work. We thank the following curators and persons who allowed access to the material included in this work: Felipe Leite (UFV), Hussam Zaher (MZUSP), José P. Pombal-Jr. (MNRJ), Miguel T. U. Rodrigues (MTR), Mirco Solé/Iuri Dias (MZUESC), Paulo C. A. Garcia (UFMG), Renato N. Feio (UFV), and William R. Heyer (USNM). We thank the Centro de Estudos de Insetos Sociais (CEIS), UNESP, Rio Claro, Brazil, for allowing us to use its

molecular facilities. We thank A. F. Sabbag, T. Grant, B. Blotto, and M. Pereyra for help with some of the analyses; R. Murta (*A. sp. 4*), L. Malagoli (*A. sp. 5*), C. Conte (*A. ehrhardti*), P. Peloso (*A. weygoldti* and *A. sp. 6*), and M. Segalla (all other spp.) kindly provided photos. The Instituto Chico Mendes de Conservação da Biodiversidade provided the license (#25033). For comments and suggestions that improved the manuscript, we thank two anonymous reviewers and A. Larson. BVMB particularly thanks M. Segalla for his support, as well as that of the CFBH and MACN staff from 2008–2015, especially V. G. D. Orrico and N. Pupin; H. R. da Silva; M. Martins (Labvert, USP); T. Grant; R. Heyer and the Smithsonian Institution for the STV award (2011); and the São Paulo Research Foundation (FAPESP) for Grants #2008/55235–4 and #2013/18807–8. CFBH thanks FAPESP for Grants #2008/50928–1 and #2013/50741–7 and CNPq for the financial support of a research fellowship. MLL thanks FAPESP for grant #2010/50124–0 and Capes for postdoctoral fellowships. JF is grateful to ANPCyT PICTs 2011–1895, 2013–404, CONICET PIP 1112008010 2422, and FAPESP for Grants #2005/56756–0, #2006/52088–5, #2012/10000–5, #2013/50741–7, #2013/20423–3.

References

- Abrunhosa, P. A., Pimenta, B. V. S., Cruz, C. A., Haddad, C. F. B. 2005. Advertisement calls of species of the *Hyla albosignata* group (Amphibia, Anura, Hylidae) Arq. Mus. Nac. Rio de Janeiro. 63, 2, 275–282.
- Bastos, R. P., Motta, J. A. O., Lima, L. P., Guimarães, L. D. A. 2003. Anfíbios da Floresta Nacional de Silvânia, Estado de Goiás. Goiânia, Brazil.
- Blotto, B. L., Nunez, J. J., Basso, N. G., Ubeda, C. A., Wheeler, W. C., Faivovich, J. 2013. Phylogenetic relationships of a Patagonian frog radiation, the *Alsodes*+*Eupsophus* clade (Anura: Alsodidae), with comments on the supposed paraphyly of *Eupsophus*. Cladistics. 29, 2, 113–131.
- Bogart, J. P. 1973. Evolution of anuran karyotypes. In: Vial, J. L. (Ed). Evolutionary

- Biology of Anurans. University of Missouri Press, Columbia, pp. 337–349.
- Brunetti, A.E., Hermida, G.N., Luna, M.C., Barsotti, A.M.G., Jared, C., Antoniazzi, M.M., Rivera- Correa, M., Berneck, B. V. M., and Faivovich, J. 2015. Diversity and evolution of sexually dimorphic mental and lateral glands in Cophomantini treefrogs (Amphibia: Anura: Hylidae). *Biological Journal of the Linnean Society*. 114, 12–34.
- Carvalho, K.A., Garcia, P.C.A., Recco-Pimentel, S.M. 2009a. NOR Dispersion, telomeric sequence detection in centromeric regions and meiotic multivalent configurations in species of the *Aplastodiscus albofrenatus* Group (Anura, Hylidae). *Cytogenet. Genome Res.* 126, 359–367.
- Carvalho, K. A., Garcia, P. C. A., Recco-Pimentel, S. M. 2009b. Cytogenetic comparison of tree frogs of the genus *Aplastodiscus* and the *Hypsiboas faber* group (Anura, Hylidae). *Genet. and Molec. Res.* 8, 1498–1508.
- Carvalho-e-Silva, A. M. P. T., Carvalho-e-Silva, S. P. 2005. New species of the *Hyla albofrenata* Group, from the States of Rio de Janeiro and São Paulo, Brasil (Anura, Hylidae). *J. Herpetol.* 39, 79–81.
- Carvalho-Jr, R. R., Galdino, C. A. B., Nascimento, L. B. 2006. Notes on the courtship behavior of *Aplastodiscus arildae* (Cruz & Peixoto, 1985) at an urban forest fragment in southeastern Brazil (Amphibia, Anura, Hylidae). *Arq. Mus. Nac. Rio de Janeiro*. 64, 247–254.
- Catrolí, G. F., and Kasahara, S. 2009. Cytogenetic data on species of the family Hylidae (Amphibia, Anura): results and perspectives. *Publ. Cienc. Biol. Saúde*. 15, 67–86.
- Catrolí, G. F., Faivovich, J., Haddad, C. F. B., Kasahara, S. 2011. Conserved karyotypes in Cophomantini: cytogenetic analysis of 12 species from 3 species groups of *Bokermannohyla* (Amphibia: Anura: Hylidae). *J. Herpetol.* 45, 120–128.
- Cisneros-Heredia, D.F., McDiarmid, R.W. 2007. Revision of the characters of Centrolenidae (Amphibia: Anura: Athespatanura), with comments on its taxonomy and the description of new taxa of glass frogs. *Zootaxa* 1572: 1–82.
- Cochran, D. M. 1955. Frogs of southeastern Brazil. *US Natl. Mus. Bull.* 206, 1–423.
- Conte, C.E., Lingnau, R., Kwet, A. 2005. Description of the advertisement call of *Hyla ehrhardti* Müller, 1924 and new distribution records (Anura: Hylidae). *Salamandra*, 41, 3, 147–151.

- Cruz, C. A., and Peixoto, O. L. 1985 “1984”. Espécies verdes de *Hyla*: o complexo “*albosignata*” (Amphibia, Anura, Hylidae). Arq. Univ. Fed. Rur. Rio de J. 7, 37–47.
- Cruz, C. A., and Peixoto, O. L. 1987 “1985”. Espécies verdes de *Hyla*: o complexo “*albofrenata*” (Amphibia, Anura, Hylidae). Arq. Univ. Fed. Rur. Rio de J. 8, 59–70.
- Cruz, C.A.G., Pimenta, B.V., Silvano, D.L. 2003. Duas novas espécies pertencentes ao complexo de *Hyla albosignata* Lutz & Lutz, 1938, do leste do Brasil (Amphibia, Anura, Hylidae). Boletim do Museu Nacional, Rio de Janeiro, 503, 1–13.
- Duellman, W. E. 1970. The hylid frogs of Middle America. Monograph of the Museum of Natural History, University of Kansas. 1–2, 1–753.
- Duellman, W. E. 1986. Diversidad y evolucion adaptativa de los hílidos Neotropicales (Amphibia: Anura: Hylidae) An. Mus. Hist. Nat. Valparaiso, 17, 143–150.
- Duellman, W. E. 1988. Patterns of species diversity in anuran amphibians in the American tropics. Ann. Mo. Bot. Gard. 75, 79–104.
- Duellman, W.E., and Trueb, L. 1986. Biology of Amphibians. McGraw-Hill, New York, 670 pp.
- Faivovich, J., Peixoto O., Cruz, C.A.G. 2002. The identity of *Hyla ehrhardti* Müller, 1924 (Anura, Hylidae). Journal of Herpetology, 36, 2, 325–327.
- Faivovich, F., Haddad, C. F. B., Garcia, P. C. A., Frost, D. R., Campbell, J. A., and Wheeler, W. C. 2005. Systematic review of the frog family Hylidae, with special reference to Hyline: phylogenetic analysis and taxonomic revision. Bull. Am. Mus. Nat. Hist. 294, 240 pp.
- Faivovich, J., McDiarmid, R. W. Myers, C. W. 2013. Two new species of *Myersiophyla* (Anura: Hylidae) from Cerro de la Neblina, Venezuela, with comments on other species of the genus. Am. Mus. Novit. 3792, 1–63.
- Farris, J. S., Albert, V. A., Källersjö, M., Lipscomb, D., Kluge, A. G. 1996. Parsimony jackknifing outperforms neighbor-joining. Cladistics. 12, 99–124.
- Frost, D. R. 2014. Amphibian Species of the World: an Online Reference. Version 6.0 (2015, May 29). Electronic Database accessible at <http://research.amnh.org/herpetology/amphibia/index.html>. American Museum of Natural History, New York, USA.
- Garcia, P. C. A., Caramaschi, U., Kwet, A. 2001. O status taxonômico de *Hyla*

- cochranae* e recaracterização de *Aplastodiscus* A. Lutz (Anura, Hylidae) Rev. Bras. Zool. 18, 1197–1218.
- Goloboff, P.A. 1999. Analyzing large datasets in reasonable times: solutions for composite optima. Cladistics. 15, 415–428
- Goloboff, P.A., Farris, J.S., Nixon, K.C. 2008. TNT, a free program for phylogenetic analysis. Cladistics. 24, 1–13.
- Gomes, M. R., and Peixoto, O. L. 1997. *Hyla callipygia*: geographic distribution. Herpetological Review. 28, 3, 110–111.
- Gruber, S. L., Zina, J., Narimatsu, H., Haddad, C. F. B., Kasahara, S. 2012. Comparative karyotype analysis and chromosome evolution in the genus *Aplastodiscus* (Cophomantini, Hylineae, Hylidae). BMC Genetics. 13, 28.
- Haddad, C. F. B., and Sawaya, R. J. 2000. Reproductive modes of Atlantic Forest hyliid frogs: a general overview and the description of a new mode. Biotropica. 32, 862–871.
- Haddad, C. F. B., and Prado, C. 2005. Reproductive modes in frogs and their unexpected diversity in the Atlantic Forest of Brazil. BioScience. 55, 3 207–217.
- Haddad, C. F. B., Faivovich, J., Garcia, P. C. A. 2005. The specialized reproductive mode of the treefrog *Aplastodiscus perviridis* (Anura: Hylidae). Amphibia-Reptilia. 26, 87–92.
- Hartmann, M. T.; Hartmann, P. A.; Haddad, C. F. B. 2004. Visual signaling and reproductive biology in the nocturnal treefrog, genus *Hyla*. Anura: Hylidae). Amphibia-Reptilia, 25, 395–406.
- Heyer, W. R., Rand A. S., Cruz C. A. G., Peixoto O. L., Nelson C. E. 1990. Frogs of Boracéia. Arquivos de Zoologia, Museu de Zoologia da Universidade de São Paulo. 31, 4.
- Jungfer, K. H., Faivovich, J., Padial, J. M., Castroviejo-Fisher, S., Lyra, M.L., Berneck, B.V.M., Iglesias, P.P., Kok, P. J. R., MacCulloch, R. D., Rodrigues, M. T., Verdade, V. K., Torres Gastello, C. P., Chaparro, J. C., Valdujo, P. H., Reichle, S., Moravec, J., Gvoždík, V., Gagliardi-Urrutia, G., Ernst, R., De la Riva, I., Means, D. B., Lima, A. P., Señaris, J. C., Wheeler, W. C., Haddad, C. F. B. 2013. Systematics of spiny-backed treefrogs (Hylidae: *Osteocephalus*): an Amazonian puzzle. Zoologica Scripta. 42, 351–380.
- Katoh, K., Standley, D. M. 2013. MAFFT multiple sequence alignment software

- version 7: improvements in performance and usability. *Mol Biol Evol.* 30,772–780.
- Lima, M. G. D., Skuk, G. O., Silva, M. C. C. D. 2006. Geographic distribution: *Aplastodiscus* (=Hyla) *sibilatus*. *Herpetological Review.* 37, 485.
- Lutz, B. 1949 "1948". Anfíbios anuros da coleção Adolpho Lutz. II. Espécies verdes do gênero *Hyla* do Leste-Meridional do Brasil. *Mem. Inst. Oswaldo Cruz.* 46, 551–577.
- Lutz, B. 1950. Anfíbios anuros da coleção Adolpho Lutz do Instituto Oswaldo Cruz. V. *Mem. Inst. Oswaldo Cruz.* 48, 599–637.
- Lutz, B. 1973. *Brazilian Species of Hyla*. University of Texas Press, 260pp.
- Lutz, A., and B. Lutz. 1938. Two new hylae: *H. albosignata* n. sp. & *H. pickeli*. *Anais da Academia Brasileira de Ciências.* Rio de Janeiro 10, 175–194.
- Mercês, E. A., and Juncá, F. A. 2010. Girinos de três espécies de *Aplastodiscus* Lutz, 1950 (Anura - Hylidae) ocorrentes no Estado da Bahia, Brasil. *Biota Neotrop.* 10, 167–172.
- Miller, M. A., Pfeiffer, W., Schwartz, T. 2010. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. *Proceedings of the Gateway Computing Environments Workshop (GCE)*, New Orleans, 1–8.
- Nixon, K.C. 1999. The parsimony ratchet, a new method for rapid parsimony analysis. *Cladistics* 15, 407–414.
- Orrico, V. G. D., Carvalho-e-Silva, A. M. P. T., Carvalho-e-Silva, S. P. 2006. Redescription of advertisement call of *Aplastodiscus arildae* and description of the call of *Aplastodiscus weygoldti* with general notes about the genus in Southeastern Brazil (Anura, Hylidae). *Rev. Bras. de Zool.* 23, 994–1001.
- Palumbi, S.R., Martin, A., McMillan, W.O., Stice, L., Grabowski, G. 1991. The simple fool's guide to PCR, Version 2.0. Available in <http://palumbi.stanford.edu/SimpleFoolsMaster.pdf>.
- Pimenta, B., Costa, D., Murta-Fonseca, R., Pezzuti, T. 2014. Anfíbios: Alvorada de Minas, Conceição do Mato Dentro, Dom Joaquim - Minas Gerais. Bicho do Mato Press. Belo Horizonte, Brazil. 196pp.
- Pol, D. 2004. Empirical problems of the hierarchical likelihood ratio test for model selection. *Syst. Biol.* 53, 949–962.
- Posada, D. 2008. jModelTest: phylogenetic model averaging. *Mol. Biol. Evol.* 25, 1253–1256.

- Posada, D., and Buckley, T.R. 2004. Model selection and model averaging in phylogenetics: advantages of Akaike Information Criterion and Bayesian approaches over likelihood ratio tests. *Syst. Biol.* 53, 793–808.
- Pyron, R. A. 2014. Biogeographic analysis of amphibians reveals both ancient continental vicariance and recent oceanic dispersal. *Systematic Biology*. 63, 5, 779–797.
- Pyron, R. A., and Wiens, J. J. 2011. A large-scale phylogeny of Amphibia including over 2800 species, and a revised classification of advanced frogs, salamanders, and caecilians. *Mol. Phyl. Evol.* 61, 543–583.
- Rambaut, A. 2014. FigTree, ver. 1.4.2. (Online 2015, May 29). Available: <http://tree.bio.ed.ac.uk/software/figtree/>.
- Rivera-Correa, M., and Faivovich, J. 2013. A new species of *Hyloscirtus* (Anura: Hylidae) from Colombia with a rediagnosis of *Hyloscirtus larinopygion* (Duellman, 1973). *Herpetologica* 69, 298–313.
- Ronquist, F., Teslenko, M., van der Mark, P., Ayres, D. L., Darling, A., Höhna, S., Larget, B., Liu, L., Suchard, Marc A., Huelsenbeck, J. P. 2012. MrBayes 3.2: Efficient bayesian phylogenetic inference and model choice across a large model space. *Syst. Biol.* 61, 3, 539–542.
- Salles, R. O. L., Pontes, R. C., Silva-Soares, T. 2012. New records and geographic distribution of *Aplastodiscus eugenioi* (Anura: Hylidae) in southeastern Brazil. *Herpetology Notes*. 5, 431–433.
- Silva, G. R., Luna-Dias, C., Hepp, F. S. F. S., Carvalho-e-Silva, A. M. P. T., Carvalho-e-Silva S. P. 2012. New record of *Aplastodiscus weygoldti* (Cruz & Peixoto, 1987) in the municipality of Mimoso do Sul, Espírito Santo State, southeastern Brazil (Anura, Hylidae). *Herpetology Notes*. 5, 371–373.
- Silva-Soares, T., Valadares, A. P., Koski, D. A., Ferreira, R. B., Cruz, C. A. G. 2011. New records and distribution of *Aplastodiscus arildae* (Cruz & Peixoto, 1985) (Amphibia, Anura, Hylidae) in the southeast of Brazil. *Herpetology Notes*. 4, 255–258.
- Smith, S. A., Arif, S. A., Nieto, M. O., Wiens, J. J. 2007. A phylogenetic hotspot for 1045 evolutionary novelty in Middle American treefrogs. *Evolution*. 61, 2075–2085.
- Swofford, D. L. 2002. PAUP: Phylogenetic analysis using parsimony, version 4.0b10. Sinauer 1052 Associates, Sunderland, Massachusetts.

- Valdujo, P. H., Silvano, D. L., Colli, G., Martins, M. 2012. Anuran species composition and distribution patterns in Brazilian Cerrado, a Neotropical hotspot. *South American Journal of Herpetology*. 7, 63–78.
- Wells, K. D. 2007. *The Ecology and Behavior of Amphibians*. The University of Chicago Press. Chicago. 1400pp.
- Wheeler, W.C. 2003. Iterative pass optimization of sequence data. *Cladistics*. 19, 254–260.
- Wheeler, W.C., Aagesen, L., Arango, C., Faivovich, J., Grant, T., D’Haese, C., Janies, D., Smith, W.L., Varón, A., Giribet, G. 2006. *Dynamic Homology and Phylogenetic Systematics: A Unified Approach using POY*. The American Museum of Natural History, New York.
- Wheeler, W. C., Lucaroni, N., Hong, L., Crowley, L. M., Varón, A. 2014. POY version 5: phylogenetic analysis using dynamic homologies under multiple optimality criteria. *Cladistics*. 31, 189–196.
- Wilkinson, J.A., Matsui, M., Terachi, T. 1996. Geographic variation in a Japanese tree frog (*Rhacophorus arboreus*) revealed by PCR-aided restriction site analysis of mtDNA. *Journal of Herpetology*. 30, 418–423.
- Wiens, J. J., Fetzner, J. W., Parkinson, C. L., Reeder, T. W. 2005. Hylid frog phylogeny and sampling strategies for speciose clades. *Systematic. Biol.* 54, 719–748.
- Wiens, J. J., Graham, C. H., Moen, D. S., Smith, S. A., Reeder, T. W. 2006. Evolutionary and ecological causes of the latitudinal diversity gradient in hylid frogs: treefrog trees unearth the roots of high tropical diversity. *Am. Nat.*, 168, 1–27.
- Wiens, J. J., Kuczynski, C. A., Xia Hua, Moen, D. S. 2010. An expanded phylogeny of treefrogs (Hylidae) based on nuclear and mitochondrial sequence data. *Mol. Phyl. Evol.* 55, 871–882.
- Zina, J., and Haddad, C. F. B. 2006a. Ecology and reproductive biology of two species of *Aplastodiscus* (Anura: Hylidae) in the Atlantic forest, Brazil. *Journal of Natural History*. 40, 1831–1840.
- Zina, J., and Haddad, C. F. B. 2006b. Acoustic repertoire of *Aplastodiscus arildae* and *A. leucopygius* (Anura: Hylidae) in Serra do Japi, Brazil. *South American Journal of Herpetology*. 1, 227–236.
- Zina, J., and Haddad, C. F. B. 2007. Courtship behavior of two treefrog species,

Aplastodiscus arildae and *A. leucopygius* (Anura: Hylidae), from the Atlantic Rainforest, Southeastern Brazil. *Herpetological Review*. 38, 282–285.

Supplementary data associated with this article can be found in the online version at:

Appendix A. Vouchers, localities, and Genbank numbers of all samples included.

Appendix B. Results of Parsimony and Bayesian analyses using the static alignment.

Appendix C. Genetic p-distances of 16S and COI fragments among all samples employed.

Appendix D. Cloacal characters studied.

Appendix E. Additional specimens studied.

Appendix F. Morphological variation of diagnostic characters in *A. albosignatus* and *A. callipygius*.