Phylogenetics and biogeography of the endemic Madagascan millipede assassin bugs (Hemiptera:
Reduviidae: Ectrichodiinae)

MICHAEL FORTHMAN and CHRISTIANE WEIRAUCH

University of California, Department of Entomology, 900 University Avenue, Riverside, CA 92521, USA

Corresponding author: M. Forthman, mfort001@ucr.edu, 1-951-827-5708

Abstract

For at least the past 80 my, Madagascar, a major biodiversity hotspot, has been isolated from all other landmasses. This long-term isolation, along with geologic and climatic factors within Madagascar and throughout the Indian Ocean, has undoubtedly influenced the evolution of the island's biota. However, few systematic analyses incorporating modern divergence dating and biogeographic analyses have focused on Madagascan insects. The diverse Madagascan millipede assassin bugs (Heteroptera: Reduviidae: Ectrichodiinae) offer an opportunity to contribute to a limited body of insect-related research that explores Madagascar's historical biogeography. A molecular dataset (COI mtDNA and 18S, 28S D2 and D3-D5 rDNAs) for 56 taxa (39 ingroup) and a combined morphological (145 characters) and molecular dataset for 110 taxa (93 ingroup) are analyzed with maximum likelihood (ML) and parsimony approaches. Based on the molecular ML phylogeny, divergence times were estimated using fossil and secondary calibrations and biogeographic analyses performed using DIVA, DEC, and DEC+j models to determine the role and patterns of vicariance and dispersal in the origin of Madagascan Ectrichodiinae. Results indicate that Ectrichodiinae in Madagascar do not form a monophyletic group, different clades are closely related to Afrotropical and Oriental lineages, and have colonized the island via transoceanic dispersal at least twice from the Oriental region and once from the Afrotropical region in the last ~68 my. Additionally, the DEC+j and DIVA models infer a single out-of-Madagascar dispersal event to the Afrotropical region. Oceanic and geologic factors that may have facilitated dispersal between these three regions are discussed. Results of the combined analyses are used to explore character support for Madagascan taxa and inform taxonomic diagnoses. Our results are congruent with the small but growing body of biogeographic research supporting Cenozoic transoceanic dispersal for Madagascan invertebrates to and from Oriental and Afrotropical regions.

Keywords: Ectrichodiinae, Madagascar, phylogeny, divergence dating, biogeography, dispersal

1. Introduction

Madagascar, the fourth largest island on Earth, is one of the world's biodiversity hotspots with more than 13,000 plant, 900 vertebrate, and 5,800 invertebrate species known (Goodman and Benstead, 2005; Phillipson et al., 2006). For non-invasive species, ~86% of invertebrates, ~50% of birds, more than 95% of other vertebrates, and more than 90% of vascular plants are endemic to the island (Goodman and Benstead, 2005; Yoder and Nowak, 2006; Phillipson et al., 2006; Buerki et al., 2013). Numerous phylogenetic studies have provided evidence that a large proportion of Madagascar's biota is closely related to African lineages (~37% of plants, ~30% of vertebrates, ~58% of invertebrates [see Yoder and Nowak, 2006]). However, a significant proportion of the biota is closely related to Oriental taxa (see Vences, 2004; Yoder and Nowak, 2006; Warren et al., 2010); about 32% of plants, 32% of vertebrates, and 17% of invertebrates are sister to Oriental lineages (Yoder and Nowak, 2006). This extant, unique biodiversity is, in the words of Ganzhorn et al. (2014), "rooted in the plate tectonics of the Mesozoic" (i.e., long isolation period), as well as the island's geography, climate, and late colonization by humans (Goodman and Benstead, 2003; Scales, 2014).

Madagascar was once part of Gondwana, but ~160 mya, the Gondwanan landmass began to split with Indo-Madagascar separating from Africa (Plummer and Belle, 1995; Gnos et al., 1997). By at least 80 mya, Madagascar separated from India and has been isolated ~400 km from Africa's southeastern coast since (Storey et al., 1995; Torsvik et al., 2000; Seward et al., 2004; Ali and Aitchison, 2008). Many plant and vertebrate studies have investigated historical biogeographic patterns for various Madagascan taxa. Although some systematic studies support influences of Gondwanan vicariance for the presence of some Madagascan lineages (e.g., ranid frogs [Bossuyt et al., 2006], boid snakes [Noonan and Chippindale, 2006], chameleons [Okajima and Kumazawa, 2010]), many recent studies based on divergence dating analyses support post-Gondwanan transoceanic dispersal from Afrotropical and Oriental regions (e.g., endemic mammalian lineages [Poux et al., 2005], Chrysophylloideae [Bartish et al., 2011], gerrhosaurid lizards [Raselimanana et al., 2009]). Comparatively fewer studies have investigated biogeographic histories of Madagascan insects. Several Madagascan insects have been hypothesized to be relicts of Gondwanan vicariance, e.g., Diplatyidae (Dermaptera) (Popham, 2000), Notonemourinae

(Plecoptera) (Illies, 1965; Paulian and Viette, 2003), some Blephariceridae (Diptera) (Paulian and Viette, 2003), and Sialidae (Liu et al., 2015), but there is an increasing number of systematic studies that support Cenozoic transoceanic dispersal from the Afrotropical and Oriental regions. For example, allodapine bees (Fuller et al., 2005; Schwarz et al., 2006; Chenoweth and Schwarz, 2011) and scarabid beetles (Wirta et al., 2008; Wirta et al., 2010; Sole et al., 2011) colonized Madagascar multiple times in the last 65 my. Fungus-growing termites in Madagascar originated from a single colonization event ~7–11 mya (Nobre et al., 2010). Other studies on carpenter bees (Rehan et al., 2010), mayflies (Monaghan et al., 2005), pierid butterflies (Nazari et al., 2011), and dytiscid beetles (Bukontaite et al., 2015) have also supported Cenozoic colonization.

Millipede assassin bugs (Heteroptera: Reduviidae: Ectrichodiinae) are a diverse group of specialized millipede predators with 736 species in 121 genera that are distributed in circumtropical and some temperate regions (Maldonado, 1990; Dougherty, 1995; Carpintero and Maldonado, 1996; Forthman and Weirauch, 2012; Forthman et al., in press). Recently, Forthman et al. (in press) published a monograph of Madagascan millipede assassin bugs (excluding the genus *Distirogaster* Horváth) in which 63 new species and three new genera were described, increasing the known Madagascan millipede assassin bug fauna by six-fold. All species and six of the eight genera found on Madagascar are endemic; of the 37 species of *Glymmatophora* Stål and 39 species of *Maraenaspis* Karsch, two and one, respectively, are endemic to Madagascar with the remainder distributed in Africa. This diverse Madagascan millipede assassin bug fauna thus presents an opportunity to investigate and contribute to a limited yet rapidly growing body of research exploring historical influences that have shaped Madagascar's terrestrial insect fauna. However, such an evolutionary investigation has been hindered by a lack of phylogenetic hypotheses; relationships between Ectrichodiinae genera have not been tested cladistically beyond a genus-level phylogeny of the New World fauna based on morphological data (Dougherty, 1995). A recent systematic study of the Reduviidae indicates that Ectrichodiinae originated 57–80 mya and, therefore, after the Madagascar-India split (Hwang and Weirauch, 2012). Thus, dispersal is likely responsible for the current distribution of millipede assassin bugs worldwide, but it is unknown what factors (e.g., ocean

currents, land bridges, and stepping-stone islands) may have facilitated dispersal between Madagascar and other biogeographic regions.

Given the recent taxonomic treatment, Upper Cretaceous age, and worldwide distribution of the subfamily, as well as the general restriction of genera to particular biogeographic regions, millipede assassin bugs are an excellent model for biogeographic studies of the Madagascan fauna. Here, we present molecular phylogenetic results that include representatives of six of the eight Madagascan Ectrichodiinae genera and provide a framework for testing the number, timing, and origin of Madagascan lineages. Based on these results, we perform divergence dating and biogeographic analyses to investigate the geographic and oceanic influences that could have facilitated long-distance dispersal between Madagascar and other biogeographic areas. We also present results of a combined morphological and molecular phylogenetic analysis on a large sample of Madagascan Ectrichodiinae species that is used to formalize taxonomic decisions and inform diagnoses in Forthman et al.'s (in press) Madagascan Ectrichodiinae monograph.

2. Material and Methods

2.1. Taxon sampling, vouchers, and databasing

A total of 110 terminal taxa were examined, comprising 93 ingroup (Ectrichodiinae) and 17 outgroup taxa (seven Reduviidae subfamilies). Ingroup sampling comprised all eight genera and 67 species of Madagascan Ectrichodiinae. Due to the lack of DNA quality material, only 56 terminal taxa were sequenced (39 ingroups, including 14 Madagascan species in six genera; sequence data for the Madagascan genus *Toliarus* Forthman, Chłond, and Weirauch and the species *Maraenaspis bidens* [Reuter] are unavailable). Each taxon is represented by a mounted primary specimen voucher (molecular vouchers are listed in Table 1). For specimens sampled for molecular data, a hind leg was removed for DNA extraction; once extraction was completed, the leg was card mounted and associated with the pinned specimen. Each molecular voucher is associated with a unique specimen identifier (USI) label to connect it with specimen information and, where possible, images in the Planetary Biodiversity Inventory

database (<http://www.research.amnh.org/pbi/locality/index.php>) and Heteroptera Species Pages (<http://research.amnh.org/pbi/heteropterasespeciespage/>) maintained by the American Museum of Natural History. All USI labels are comprised of the prefix AMNH_PBI or UCR_ENT followed by an 8-digit number. Sequenced specimens are also associated with an RCW number, which is the ethanol specimen collection number.

Specimens examined for this study are deposited in the following institutions: **AMNH**, American Museum of Natural History, New York, USA; **BMNH**, Natural History Museum, London, United Kingdom; **CAS**, California Academy of Sciences, California, USA; **FCAP**, Universidade Federal do Pará, Pará, Brazil; **MNHN**, Muséum National d'Histoire Naturelle, Paris, France; **MRAC**, Musée Royal de l'Afrique Centrale, Tervuren, Belgium; **MTEC**, Montana State University, Montana, USA; **SU**, Department of Zoology, University of Silesia, Poland; **UCR**, University of California, Riverside Entomological Research Museum, California, USA; **USNM**, National Museum of Natural History, Washington D.C., USA.

2.2. Morphological methods, terminology, and abbreviations

Extreme sexual dimorphism (Fig. 1) is common among millipede assassin bugs and poses a significant problem when analyzing evolutionary relationships. As such, male ectrichodiine specimens were targeted for morphological character coding. However, two terminal taxa, *Maraenaspis coccinea* Horváth and *Racelda* sp., are represented by apterous females. The male of *Maraenaspis coccinea* has, to our knowledge, yet to be discovered and available specimens of *Racelda* were restricted to the one female at the time of this project. Despite the possibility of extreme dimorphism, some morphological features (e.g., antennal segmentation, relative lengths of labial segments, scutellar apical processes, etc.) remain similar among males and females within ectrichodiine species and genera. As such, these sexually static morphological features are inferred and coded in the analysis for taxa only represented by females.

External morphology and genitalic characters were examined using Nikon SMZ1000 and SMZ1500 dissecting microscopes. For males, genitalia (abdominal segment 8, pygophore, and phallus) were

dissected, cleared in heated 10% potassium hydroxide (KOH) for 3–8 minutes, washed with distilled water and subsequently with 100% ethanol (EtOH), stained with Chlorazol Black E in 70% EtOH solution to give contrast to membranous areas, examined in glycerol, and permanently stored in genitalic capsules pinned to the specimen. A total of 145 morphological characters were coded in the Descriptive Language for Taxonomy (DELTA) program (Dallwitz, 1980; Dallwitz et al., 1999) using a modified morphological matrix from Weirauch (2008) and Weirauch (2010). In general, terminology follows a subset of terms used by Dougherty (1995), Weirauch (2008), and Forero and Weirauch (2012). Terminology for wing venation follows Hill (2014) and Weirauch (2008), although homology concepts are currently being reviewed across Paraneoptera (Dávid Rédei, pers. comm.). The morphological characters and character states used in this study are given in Table 2. The morphological matrix is provided in Supplementary Material (SM) 1 in nexus format.

2.3. Molecular markers, PCR, sequencing, and alignment

One mitochondrial (COI) and three nuclear (18S, 28S D2, and 28S D3–D5 rDNAs) gene regions were sampled. Sequences from Weirauch and Munro (2009) and Hwang and Weirauch (2012) were used for outgroup terminals and some Ectrichodiinae taxa (Table 1). To sequence remaining ectrichodiine taxa, primer sequences for 18S (18SF, 18SR) and 28S D3–D5 (D3Fa, D5Ra) were obtained from Weirauch and Munro (2009); 28S D2 (D2Fa, D2Ra) from Forero et al. (2013); and COI (C1-J-2183F) from Simon et al. (1994) and (C1-N-2609R) from Damgaard and Sperling (2001). DNA was extracted from a hind leg for each specimen using QIAGEN DNeasy Blood and Tissue Kit protocols. PCR was performed using either GE Healthcare Life Sciences PuReTaq-Ready-To-Go-PCR-Beads or EmeraldAmp GT PCR Master Mix and Fisher Scientific or BioRad T100 Thermal Cyclers. Gene regions were amplified following Weirauch and Munro (2009), except the initial denaturation was set at 5 minutes, and for the 28S gene regions, the annealing temperature was set to 50°C. Gel electrophoresis with SyberSafe gel stain and a UV illuminator were used to check amplification results. PCR products were cleaned using Bioline SureClean and sequenced on an Applied Biosystems 3730xl DNA Sequencer at UCR's Institute for Integrative Genome

Biology. Sequences were assembled and edited using Sequencher v4.8 and are available on GenBank (Table 1). Each gene region was treated as a separate partition and aligned in MAFFT (Katoh and Standley, 2013) using the G-INS-i algorithm. Gene regions were concatenated in SequenceMatrix v1.7.8. (Vaidya et al., 2011) and exported with external gaps coded as question marks. The concatenated, aligned molecular matrix is provided in nexus format in SM2.

2.4. Phylogenetic analyses

2.4.1. Molecular phylogenetic analyses

A maximum likelihood (ML) analysis was performed using the molecular dataset and RAxML-HPC2 on XSEDE v8.0.24 (Stamatakis, 2014) on the Cyberinfrastructure for Phylogenetic Research Science Gateway v3.1 (CIPRES; http://www.phylo.org/sub_sections/portal/). Each gene partition was analyzed under a GTR+ Γ +I model, which is the best fit model for each partition as determined by MEGA v6.06 (Tamura et al., 2013). Rapid bootstrap analysis (BS) was performed for 1,000 iterations and followed by a ML best tree search. Aside from estimating the proportion of invariable sites for each partition, default settings were used.

The molecular matrix was also subjected to parsimony analyses with New Technology search and equal (EW) and implied weights (IW) in TNT v.1.1 (Goloboff et al., 2008). Uninformative characters (1,990) were inactivated prior to analysis. For IW, three concavity constant values (k) were used: 3, 6, and 10. Internal gaps were treated as a fifth state, while external gaps were treated as missing. Default settings were used for sectorial search, drift, and tree fusing. The initial driven search level was set at 100 with level checked every 3 hits, initial addition sequences = 14, find minimum length 100 times, and random seed = 4325. Standard BS resampling with absolute frequencies was performed using 500 replicates and New Technology search (initial driven search = 38, check level every 3 hits, initial addition sequences = 7, find minimum length 10 times).

2.4.2. Combined morphological and molecular phylogenetic analyses

Morphological and molecular datasets were concatenated using Mesquite v2.75 (Maddison and Maddison, 2011) and subjected to ML (RAxML-HPC v.8.1.15; Stamatakis, 2014) and EW parsimony analyses (New Technology search in TNT). For the ML analysis, each gene partition was analyzed under a GTR+ Γ model, while the morphological partition was analyzed with a Mkv model (Lewis, 2001). Rapid BS was performed for 1,000 iterations and followed by a ML best tree search. Our equal weights parsimony analysis used the same search and bootstrap approaches applied to our molecular dataset (see section 2.4.1). Unambiguous character optimizations were examined on the strict consensus tree in WinClada v1.00.08 (Nixon, 2002) (SM3).

2.5. Divergence dating estimation

Divergence dating estimates were generated on the molecular ML best tree using the CIPRES web server's BEAST2 on XSEDE (Bouckaert et al., 2014). Each gene region was assigned an unlinked site model (GTR+ Γ +I) and linked to the same tree model (Yule). The number of gamma rate categories for each site model was set to 4 and the proportion of invariant sites set to 0.5, with the option to estimate the value of both parameters and the gamma shape. All other site model parameters and estimation options remained at default settings. Following Hwang and Weirauch (2012), a linked clock model was assigned to 28S D2 and 28S D3–D5 to reflect their single identity. As such, three unlinked relaxed lognormal clock models were assigned for 18S, 28S, and COI. Initially, all non-calibration priors were not modified from default settings, however low posterior, prior, and mutation rate effective sample size (ESS) values were extremely low (<50) after performing four independent runs in BEAST. To improve ESS values for these parameters, the default Jeffrey's prior was replaced with a log normal prior for the mutation rates (A. Rambaut & A. Drummond, <https://groups.google.com/forum/#!forum/beast-users>).

Given that there are no described Ectrichodiinae fossils, two outgroup fossils (*Triatoma dominicana* Poinar and *Apicrenus fossilis* Maldonado, Santiago-Blay, and Poinar) and two secondary calibrations (root node and Ectrichodiinae+Tribelocephalinae node) were used from a recent cladistic analysis of Reduviidae (Hwang and Weirauch, 2012). Fossils were assigned to nodes following Hwang and

Weirauch (2012), using an apomorphy-based approach (Parham et al., 2011). For fossil-based node calibrations, a prior lognormal distribution was assigned with the minimum age corresponding to the minimum age of the fossil and with the 95% confidence interval encompassing the range of estimated dates recovered by Hwang and Weirauch (2012). For secondary calibrations, a prior normal distribution was assigned with the 95% confidence interval corresponding to the estimated 95% highest posterior density (HPD) credibility interval from Hwang and Weirauch (2012). Calibration nodes, means, standard deviations or sigma values, and offsets (in real space) are given in SM4. Four Markov Chain Monte Carlo (MCMC) chains were ran for 200 million generations, with parameters and trees logged every 20,000 generations. Each chain was assessed for convergence in Tracer v1.6 and combined after discarding 5% burn-in. Combined ESS values for each parameter were ≥ 200 . LogCombiner v2.1.3 was used to discard 5% of trees as burn-in and combine the remaining trees from each run. Median heights were annotated using TreeAnnotator v.2.1.2.

2.6. *Biogeographic analyses*

Ancestral ranges were reconstructed on the maximum clade credibility tree from the BEAST analysis using DIVA (Dispersal-Vicariance Analysis) in RASP v3.02 (Ronquist, 1997; Yu et al., 2010, 2015) and DEC (dispersal-extinction-cladogenesis) (Ree et al., 2005; Ree and Smith, 2008) and DEC+j (Matzke, 2014) models in the BioGeoBEARS v0.2.1 package (Matzke, 2013) in R v3.2.0 (R Core Team, 2015). Distribution ranges of Ectrichodiinae were divided into four areas: Neotropical, Afrotropical, Madagascar, and Oriental. Outgroup taxa were trimmed due to limited sampling of other assassin bug subfamilies.

The parsimony, event-based DIVA method was developed by Ronquist (1997) and reconstructs ancestral distributions based on a simple biogeographic model and a three-dimensional cost matrix. No penalty is allotted when speciation is the result of vicariance, but dispersal and extinction have a penalty of one per unit area added to or deleted from a distribution (Ronquist, 1997). To reduce the acquisition of multiple areas at nodes near the root, the maximum number of areas for each node was two.

The DEC models geographic range evolution by stochastic dispersal and local extinction events along branches and estimates the likelihood of ancestral ranges at cladogenesis (Ree et al, 2005; Ree and Smith, 2008). This model has two free parameters specifying the dispersal rate (d ; range expansion) and extinction rate (e ; range contraction) along branches (Ree et al., 2005; Ree and Smith, 2008; Matzke, 2014). The DEC model also has a fixed cladogenesis model that assumes that one daughter lineage will always inherit one subarea of a widespread ancestral range (Matzke, 2014). The DEC+j model has an additional free parameter j that enables one daughter lineage to disperse to a new range outside the ancestral range, thus modeling founder-event speciation events (Matzke, 2014). For both DEC and DEC+j models, the range constraint matrix was set at default with the maximum number of areas for each node set to two. Five time scales were constructed in the dispersal matrix to reflect changes in ocean currents in the Mozambique channel and the emergence of Indian Ocean islands: 1) 0–15 mya, 2) 15–23 mya, 3) 23–34 mya, 4) 34–56 mya, and 5) 56–66 mya. Dispersal rates between areas were assigned as 0.1 (very low; oppositely directed ocean circulation, lack of islands and land bridges, and areas not connected), 0.25 (low; ocean circulation in dispersal direction or presence of islands or land bridges), 0.5 (moderate; ocean circulation in dispersal direction and presence of islands or land bridges), 0.75 (high; areas closely adjacent or directly connected), and 1.0 (same area) based on paleoceanographic and paleogeographic reconstructions (Sanmartín et al., 2001; Heydt and Dijkstra, 2006; Ali and Huber, 2010; Lomolino, 2010; Pyron, 2014). Dispersal rates assigned for each time scale are given in SM5. The DEC model is nested within the DEC+j model, and, as a result, we compare the two models using a Likelihood Ratio Test (LTR).

3. Results

3.1 Molecular phylogenetic analyses

The first published molecular ML best tree (Ln score = -25426.19528) of millipede assassin bugs is shown in Fig. 2. Many nodes received moderate (BS = 70–89) to high support (BS = 90–100), but several deeper nodes were poorly supported (BS < 70). Ectrichiinae is monophyletic with high support and

recovered as the sister group to Tribelocephalinae, which is congruent with previous morphological and molecular cladistic analyses (Weirauch, 2008; Weirauch and Munro, 2009; Hwang and Weirauch, 2012). Two major ectrichodiine clades are recovered with low support: a predominantly Neotropical clade (node 1), which includes the Afrotropical genus *Santosia* Stål, and an Old World clade (node 2) that includes Afrotropical, Oriental, and Madagascan taxa.

Four Madagascan Ectrichodiinae lineages are recovered with low to high support. A representative of the Madagascan genus *Gibbosella* Chłond is nested within a clade of Oriental taxa with moderate support, but relationships within this clade, especially the *Gibbosella* and *Caecina* Stål sister group relationship, are poorly supported. The genera *Tanindrazanus* Forthman, Chłond, and Weirauch, *Toxopus* Bergroth, and *Marojejycoris* Forthman, Chłond, and Weirauch is a poorly supported clade. Relationships within this clade are largely weakly supported. *Tanindrazanus* is recovered as the sister group of *Marojejycoris*, and together are closely related to *Toxopus*. *Distirogaster* also forms a clade, but is weakly supported as the sister group to a clade comprising Afrotropical, Madagascan, and Oriental taxa. The Madagascan species *Glymmatophora crassipes* Horváth is highly supported as the sister to the Afrotropical genus *Ectrichodia* Lepeletier and Serville.

A number of unresolved relationships are recovered throughout the EW parsimony phylogeny (SM6), e.g., near the root of the phylogeny (i.e., most outgroups and Ectrichodiinae+Tribelocephalinae), between *Tanindrazanus*+*Marojejycoris* and two other clades (*Toxopus* and *Toxopus*+*Distirogaster*), among others. Ectrichodiinae is also rendered paraphyletic with respect to Tribelocephalinae. The ectrichodiine genus *Vilius* Stål is sister to Tribelocephalinae genera, which is a questionable result. Due to the lack of resolution in our EW phylogeny, we used IW under three concavity constant values that resulted in one ($k = 6$ or 10) or three ($k = 3$) most parsimonious trees (SM6). Some of the relationships recovered from these IW analyses are similar with the molecular ML phylogeny. However, differences are observed (e.g., non-monophyly of *Toxopus* and Ectrichodiinae), some of which are questionable and unlikely based on morphology (e.g., *Vilius* nested with Tribelocephalinae, *Maraenaspis coccinea* sister to *Caecina* sp.). Some IW results consistently differed between weighting schemes, e.g., the phylogenetic

positions of *Marojejycoris*, *Toxopus toamasina*, and *Toxopus griswoldi*. Despite topological differences between EW and IW parsimony trees, these results provide evidence for the non-monophyly of Madagascan Ectrichodiinae.

3.2. Combined morphological and molecular phylogenetic analyses

The TNT consensus tree of 938 parsimonious trees is shown in Fig. 3 (outgroups not shown but same as in Fig. 2) and was used to inform taxonomic decisions in Forthman et al. (in press). As in the molecular (SM2) and combined ML analyses, Ectrichodiinae are monophyletic and recovered as the sister to Tribelocephalinae with high support (BS = 99). A predominantly Neotropical clade (node 1) is also recovered with low support, but unlike the ML analysis, this clade includes the Oriental genus *Vilius* along with *Santosia*. An Old World clade (node 2) is recovered with several topological differences from the ML tree. *Mendis* Stål+*Bannania* Hsiao+*Neolibavius* Miller is weakly supported as the sister to the remaining taxa, followed by a weakly supported *Microstemmaoides* Putschkov+*Synavecoris* Villiers+nr *Bannania* clade. A clade (node 3) comprised of *Ectrichodia*, *Glymmatophora*, *Cleptria* Stål, *Maraenaspis*, and *Ectrychotes* Burmeister is recovered as sister to a predominantly Madagascan clade (node 4) with low support. Relationships within the latter clade are supported with BS<50, with *Marojejycoris* sister to *Gibbosella*+*Caecina*, *Marojejycoris*+*Gibbosella*+*Caecina* sister to *Distirogaster*+*Toliarus*+*Tanindrazanus*+*Toxopus* (node 6), *Distirogaster* sister to *Toliarus*+*Tanindrazanus*+*Toxopus* (node 7), and *Toliarus* sister to *Tanindrazanus*+*Toxopus*. Character optimization results for the parsimony tree are given in SM3.

Our combined ML result (SM7) is largely congruent with the combined parsimony topology with the following exceptions: 1) *Maraenaspis bidens* is sister to *Racelda* sp., 2) *Gibbosella* is the sister to the *Caecina*+*Mendis*+*Neolibavius*+*Bannania*+nr *Bannania* clade, and 3) *Distirogaster* is the sister group to the *Glymmatophora*+*Ectrichodia*+*Centraspis*+*Ectrychotes*+*Microstemmaoides*+*Cleptria*+*Maraenaspis coccinea* clade. The ML result supports the taxonomic decisions of Forthman et al. (in press) and

produces minor character optimization differences in comparison to the parsimony tree for taxa and clades of interest (data not shown).

3.3. Divergence dating estimation

Results of the BEAST molecular divergence dating analysis are shown in Fig. 4, with HPD intervals shown in SM8. Based on the analysis, Ectrichodiinae diverged from Tribelocephalinae approximately 68.53 mya (95% HPD = 57.61–79.39 mya), which is congruent with Hwang and Weirauch (2012). Within Ectrichodiinae, the Neotropical Ectrichodiinae+*Santosia* clade diverged from the Old World clade about 63.51 mya (95% HPD = 52.39–74.44 mya). Within the Old World clade, all Madagascan Ectrichodiinae lineages and genera diverged from their respective sister groups within the last 46 my: 1) *Gibbosella quadocris* Forthman, Chłond, and Weirauch (median = 27.33 mya, 95% HPD = 12.22–42.12 mya), 2) *Marojejycoris*+*Tanindrazanus*+*Toxopus* (median = 35.55 mya, 95% HPD = 25.34–45.60 mya), 3) *Distirogaster* (median = 32.86 mya, 95% HPD = 23.48–42.59 mya), and 4) *Glymmatophora crassipes* (median = 19.24 mya, 95% HPD = 11.46–27.85 mya).

3.4. Biogeographic analyses

Ancestral range reconstruction results based on the DEC and DEC+j models are reported and shown in Fig. 4. Results of the ancestral area reconstruction using the DIVA model are shown in SM9 and are most similar to the highest probability ranges reconstructed by DEC+j. The likelihood, *d*, *e*, and *j* parameter values for the DEC and DEC+j models are as follows: (1) DEC: LnL = -39.09, *d* = 0.0056, *e* = 0.0006, *j* = 0 and (2) DEC+j: LnL = -27.10 *d* = $1e^{-12}$, *e* = $1e^{-12}$, *j* = 0.1238. The DEC+j model performed significantly better than the DEC (LRT $X^2 = 23.97$, df = 1, *p* = $9.80e^{-7}$), with the *j* parameter (or founder speciation effect) by far the most significant contributor to the current distribution.

In analyses with all three biogeographic models, a Neotropical and Oriental distribution is recovered for the ancestral node of Ectrichodiinae. The ancestral areas for the Neotropical Ectrichodiinae+*Santosia* clade and the Old World clade are inferred as Neotropical and Oriental, respectively, in all analyses. In

the DEC+j analysis, the ancestral range of *Gibbosella quadocris* and *Caecina* sp. is recovered as Oriental with high probability, indicating the ancestor of *Gibbosella* dispersed to Madagascar from this region. An Oriental distribution is also reconstructed for the node that includes *Gibbosella* and *Ectrychotes* with subsequent dispersal to Madagascar for the node containing *Marojejycoris* and *Ectrychotes*. The ancestor of the clade containing *Centraspis* Schaum and *Ectrychotes* is reconstructed to have dispersed to the Afrotropical region from Madagascar, with a re-colonization from Africa to Madagascar by *Glymmatophora crassipes*. Results from DIVA are similar to DEC+j highest probability results, but differ in the following with respect to the Madagascan fauna: 1) the node containing *Gibbosella* and *Ectrychotes* is inferred as Madagascar+Oriental, 2) the ancestral range of *Gibbosella quadocris* and *Caecina* sp. is Madagascar+Oriental, 3) an Afrotropical+Madagascar distribution is recovered for the clade containing *Distirogaster* and *Ectrychotes*, and 4) the ancestral range of *Glymmatophora crassipes*+*Ectrichodia* is Afrotropical+Madagascar.

Reconstructions from the DEC analyses differ from DEC+j results for some nodes and by the presence of more widespread ancestral ranges. In the DEC analysis, a Madagascar+Oriental ancestral distribution is recovered for *Gibbosella quadocris*+*Caecina* sp. with high probability, as well as for the node containing *Marojejycoris* and *Ectrychotes* and the node containing *Distirogaster* and *Ectrychotes*. An Afrotropical+Oriental ancestral range is recovered for the clade including *Centraspis* and *Ectrychotes*, with the ancestral node of *Glymmatophora crassipes*+*Ectrichodia* having an Afrotropical+Madagascar distribution.

4. Discussion

4.1. Phylogeny of Madagascan Ectrichodiinae

Results of our molecular ML phylogeny are used to investigate the number of Madagascan Ectrichodiinae lineages, as well as to estimate temporal divergences and biogeographic history. Our combined morphological and molecular phylogeny was used by Forthman et al. (in press) to formalize taxonomic decisions and inform diagnostic features for Madagascan millipede assassin bugs. Both

molecular ML and combined parsimony phylogenetic analyses recovered some similar relationships. Both phylogenetic hypotheses support a paraphyletic New World fauna, with respect to *Santosia* in the molecular ML phylogeny or *Santosia+Vilius* in the combined parsimony result. This predominately New World clade is recovered as sister to a larger clade of Afrotropical, Oriental, and Madagascan Ectrichodiinae with relatively weak support. Madagascan lineages are recovered with close relationships to Oriental and Afrotropical taxa in both analyses, which is congruent with Vences' (2004) and Yoder and Nowak's (2006) conclusions that a large proportion of the Madagascan biota exhibits relationships with Afrotropical and Oriental taxa. Species of *Glymmatophora* share a close relationship with *Ectrichodia* species. *Gibbosella* species are recovered as sister to *Caecina*, albeit with low support in both phylogenies. Although the molecular and combined phylogenies differ in their phylogenetic placement of *Marojejycoris*, the genera *Tanindrazanus* and *Toxopus* retain a close relationship in both analyses.

Despite recovering similar relationships, the two phylogenies show some degree of discordance in higher-level relationships within Ectrichodiinae, a result that may be due to the increase in missing molecular data in the combined analysis. Unlike the molecular phylogeny, most Madagascan taxa form a clade in the combined phylogeny (node 4, including *Caecina*) with low support. The phylogenetic placement of *Marojejycoris* and *Distirogaster* differ; in the ML phylogeny, *Marojejycoris* is recovered as sister to *Tanindrazanus* and *Distirogaster* as sister to a clade containing *Ectrichodia* and *Ectrychotes*, whereas the two genera are included within node 4 of the combined analysis. In general, more nodes are recovered with $BS \geq 50$ in the molecular phylogeny compared to the combined phylogeny. Regardless of the discordance between the two phylogenetic hypotheses, the combined phylogeny has permitted exploration of potential character support for taxa and been useful in determining diagnostic features for Forthman et al.'s (in press) taxonomic monograph.

4.2 Temporal divergence and biogeography

Based on our temporal divergence estimates and ancestral range reconstructions, millipede assassin bugs colonized Madagascar via transoceanic dispersal more than once within the last ~68 my. The DEC+j

model outperformed the DEC model based on LRT and infers two colonization events from the Oriental region to Madagascar, once between 25–57 mya and once within the last 42 my. Subsequently, a single colonization event from Madagascar to the Afrotropical region is inferred to have occurred around the Eocene-Oligocene boundary (21–43 mya) with a back-colonization event (ancestral node of *Glymmatophora*+*Ectrichodia*) from the Afrotropical region to Madagascar within the last ~28 my. These results are congruent with the DIVA results (SM9). Although the DEC results differ from the DEC+j in terms of the ancestral ranges reconstructed for several nodes, in general we find similar patterns: two dispersal events into Madagascar from the Oriental region and one dispersal from Africa to Madagascar. However, an out-of-Madagascar dispersal event to the Afrotropical region is not recovered in the DEC analysis; upon diverging from *Distirogaster*, the branch leading to the node containing *Centraspis* and *Ectrychotes* has an Oriental distribution, with a range expansion into the Afrotropical region at that node.

Several hypotheses have been proposed for dispersal events from the Oriental region to Madagascar and may explain the two colonization events reconstructed from the DEC+j model. While many researchers reconstruct an isolated India in the Late Cretaceous (~65 mya) (see Yoder and Nowak, 2006; Ali and Aitchison, 2008), Ali and Aitchison (2008) proposed a paleogeographic model in which India became progressively isolated but remained connected to Madagascar via the Seychelles-Mascarene Plateau. This connection was extended in the Palaeogene by the development of volcanic islands (Ali and Aitchison, 2008; Zhou et al., 2012) and would have facilitated dispersal as India moved north. Warren et al. (2010) reviewed geologic evidence and also proposed that the rise and fall of sea levels exposed islands between India and Madagascar over the last 65 my, thus reducing the distance of open ocean to traverse for potential colonizers. Warren et al. (2010) further proposed climatic influences that would facilitate long-distance dispersal (e.g., winter monsoon winds blowing from India towards Madagascar facilitating aerial dispersal).

Several studies have supported “out-of-Madagascar” dispersal events to surrounding islands and continents for plants and vertebrates (e.g., Jansa et al., 1999; Wikström et al., 2010; Harmon et al., 2008), as well as for several groups of insects. Torres et al. (2001) concluded that satyrine butterflies in the

subtribe Mycalesina have a complex biogeographical history that involves dispersal from Madagascar to Africa. The mayfly genus *Cloeodes* Traver is inferred to have a Madagascan origin followed by dispersal into Africa (Monaghan et al., 2005). Zakharov et al. (2004) inferred out-of-Madagascar dispersal for species of the butterfly genus *Papilio* Linnaeus to Oriental and Afrotropical regions within the last 10 my. Recently, the out-of-Madagascar biogeographical hypothesis has been supported by systematic analyses of Madagascan diving beetles (Dytiscidae), with results indicating dispersal to Oriental and Afrotropical regions around the Oligocene and Miocene (Bukontaite et al., 2015).

Our DEC+j results support an out-of-Madagascar dispersal event to Afrotropical regions about 21–43 mya. Westward dispersal could not have been facilitated by ocean currents as these currents had an eastward movement during the Eocene and reversed direction only ~15 mya (Ali and Huber, 2010). McCall (1997) hypothesized that a land bridge known as the Davie Ridge once connected Africa and Madagascar during the mid-Eocene to early Miocene (i.e., 45–26 mya). However, this land bridge has not been supported by some studies (e.g., Leclaire et al., 1989; Bassias, 1992; Rogers et al., 2000), although some suggest that isolated islands may have been present (Rogers et al., 2000; de Wit, 2003). Thus, it is possible that islands in the Mozambique Channel have facilitated westward aerial dispersal during this time. Female specimens for extant Madagascan species of *Distirogaster* (5 spp.), *Toxopus* (2 spp.), and *Glymmatophora* (1 sp.) are apterous, which makes present-day aerial dispersal unlikely. However, it is unknown if the ancestors of these taxa were apterous or capable of flight, thus enabling long-distance aerial dispersal. Female specimens for the majority of Madagascan species, as well as many Afrotropical and Oriental taxa, remain unknown and hinder our ability to reconstruct ancestral states. Future discoveries of female specimens will allow us to investigate the evolution of winglessness for each sex and to subsequently refine hypotheses on how ancestral species may have dispersed in and out of Madagascar.

Based on our DEC+j analysis, the presence of islands and the eastward ocean currents of the Mozambique Channel within the last ~28 my may have facilitated the re-colonization of Madagascar from Africa. The divergence time estimate of *Glymmatophora crassipes* (95% HPD = 11.46–27.85 mya)

is consistent with this hypothesis. However, aerial dispersal cannot be ruled out as ocean currents reversed direction ~15 mya, which would hinder eastward dispersal via rafting. *Glymmatophora crassipes* is apterous in both sexes, but some African species in this genus are macropterous, and it is uncertain when and where the ancestor of *G. crassipes* has lost the capability of flight. Apterous and macropterous *Glymmatophora* species will need to be targeted for future analyses to determine if ocean currents or aerial dispersal are responsible for extant Madagascan *Glymmatophora*. Ancestral state reconstructions on the evolution of winglessness across *Glymmatophora* may provide further insight into possible avenues for the colonization of Madagascar.

Five Ectrichodiinae species in three genera — *Mascaregnasa* Distant, *Quinssyana* Distant, and *Rochonia* Distant — are known from another Indian Ocean island, the Seychelles (Maldonado, 1990). Each species is only known from the male holotype and, thus, were not included in our molecular or morphological analyses. We hypothesize, based on morphological similarities examined in habitus images of the holotypes, that *Quinssyana* is closely related to the Madagascan genus *Gibbosella* and the African genus *Synavecoris*, while *Toxopus* is closely related to *Rochonia*. *Mascaregnasa* is very different from other millipede assassin bug genera, and we are unable to hypothesize affinities with other taxa. Future inclusion of these taxa in systematic analyses will be critical for testing the role of the Seychelles as a stepping-stone island, which has been postulated for other taxa (e.g., *Tachycinemis* frogs [Vences et al., 2003a, b], *Nephilengys* hermit spiders [Kuntner and Agnarsson, 2011], and baetid mayflies [Monaghan et al., 2005]). Millipede assassin bugs from other Indian Ocean islands are currently unknown, but future taxonomic surveys on these islands may result in material that would also benefit future biogeographic investigations for Madagascan Ectrichodiinae.

4.3. Combined parsimony analysis to inform taxonomic diagnoses of Madagascan Ectrichodiinae

Character optimizations of the combined parsimony analysis were used as the basis for diagnoses in the taxonomic monograph of Madagascan Ectrichodiinae (Forthman et al., in press). A detailed discussion of important morphological characters for selected nodes shown in Fig. 3 is provided in SM10. Here, we

only discuss some examples of male diagnostic features for endemic Madagascan genera that are also included in Forthman and et al. (in press). Species of *Marojejycoris* are distinguished by features such as a complete pronotal transverse suture (57–1; Fig. 3H), the laterally visible MGE (71–1; Fig. 3K), and the fore wing lacking the distal part of M beyond the M+Cu junction (104–1; Fig. 3M). Diagnostic features for *Gibbosella* include the 8-segmented antennae (38–3; except 6-segmented in *G. pallidalata* Forthman, Chłond, and Weirauch), dorsally directed paramedian scutellar processes (62–1; except in *G. planiscutum* Forthman, Chłond, and Weirauch) (Fig. 3I), and transversely bicolored dorsal laterotergites (105–1) with posterior protuberances (107–1, 108–1, 109–1, 110–1; shared with *Caecina*). A number of features are diagnostic for *Distirogaster*, e.g., the metallic coloration (2–1), 8-segmented antennae (38–3), posterior tubercles on dorsal laterotergites III–VI (107–1, 108–1, 109–1, 110–1), and paramedian carinae on abdominal sternites (120–1; Fig. 3N). The small (1–0) red and black (2–2) body and a punctate posterior pronotal lobe (51–2) distinguish *Toliarus*. Although *Tanindrazanus* is not supported by any synapomorphic characters in our analysis, it is consistently recovered as a clade in molecular and combined analyses. An incomplete transverse suture between the meso- and metasterna (66–1), a laterally expanded dorsal laterotergite II (106–1), and the transversely bicolor dorsal laterotergites (105–1) are treated as diagnostic characters for *Toxopus*.

5. Conclusion

Madagascar's unique biodiversity is, at least partially, the result of long-term geologic isolation, transoceanic dispersal events followed by speciation, local geography, and climate change. The recently revised and diverse millipede assassin bug fauna of Madagascar have presented an opportunity to contribute to this growing knowledge of Madagascan invertebrates by investigating its historical biogeography. Phylogenetic hypotheses generated from molecular and combined morphological and molecular cladistic analyses are similar in some respects but discordant at higher-level relationships. Overall, the molecular phylogeny received higher BS support compared to the combined phylogeny.

Regardless, results from the combined phylogeny have been important for informing taxonomic diagnoses in Forthman et al. (in press).

Based on our molecular dataset, millipede assassin bugs are shown to have colonized Madagascar within the last ~68 my. Given this relatively recent age, transoceanic dispersal rather than vicariance is responsible for the Madagascan Ectrichodiinae fauna we see today. Millipede assassin bugs colonized Madagascar twice from the Oriental region and once from the Afrotropical region. However, DEC+j and DIVA models reconstruct an out-of-Madagascan dispersal event to the Afrotropical region, whereas the DEC model does not. Temporal divergence estimates and biogeographic results indicate that dispersal from the Oriental region to Madagascar may have been facilitated by the Seychelles-Mascarene Plateau and volcanic islands in the Indian Ocean over the last ~65 my. Factors facilitating dispersal from the Afrotropical region to Madagascar are more difficult to determine, but aerial dispersal via stepping-stone islands in the Mozambique Channel is the most probable hypothesis when accounting for ancient oceanographic reconstructions (i.e., ocean currents had an eastward direction). Further testing of the geologic and oceanic factors facilitating dispersal will require a larger sample of Afrotropical millipede assassin bugs, as well as Seychellois species. Furthermore, apterous females are known for several Madagascan species, making aerial dispersal impossible. However, it remains to be investigated if the ancestors of these taxa possessed an apterous condition or were capable of flight. Our results are congruent with a small, yet, largely growing body of biogeographic studies for Madagascan invertebrates; Madagascan millipede assassin bugs have a complex biogeographic history, with Cenozoic transoceanic dispersal between Oriental, Afrotropical, and Madagascan regions solely responsible for the endemic fauna we currently find present on the island.

Acknowledgements

We extend our appreciation to Dominik Chłond, Eliane de Coninck, Eric Guilbert, Tom Henry, Michael Ivie, Norm Penny, Ruth Salas, and Mick Webb for loaning specimens, as well as Jason Cryan, Rudolf Meier, Joshua Oyelade, Michael Sharkey, and David Wagner for donating specimens for study.

We thank John Heraty, Mark Springer, Brad Mullens, Elizabeth Murray, Jason Mottern, and members of the Weirauch lab for feedback on the project and manuscript. We acknowledge the National Science Foundation (NSF) PEET DEB 093333853 grant (awarded to CW), the Entomological Society of America Systematic, Evolution, and Biodiversity Travel Award (awarded to MF), and funding provided by UCR. Specimens used for this research were also acquired under NSF DEB 0542864 (awarded to Michael Sharkey and Brian Brown) and NSF BSI 0072713 (awarded to Brian Fisher and Charles Griswold).

References

- Ali, J.R., Aitchison, J.C. 2008. Gondwana to Asia: plate tectonics, paleogeography and the biological connectivity of the Indian sub-continent from the Middle Jurassic through latest Eocene (166–35 Ma). *Earth-Sci. Rev.* 88, 145–166.
- Ali, J.R., Huber, M. 2010. Mammalian biodiversity on Madagascar controlled by ocean currents. *Nature* 463, 653–656.
- Álvarez-Padilla, F., Ubick, D., Griswold, C.E. 2012. *Noideattella* and *Tolegnaro*, two new genera of goblin spiders from Madagascar, with comments on the gamasomorphoid and silhouettelloid oonopids (Araneae, Oonopidae). *Am. Mus. Novit.* 3745, 1–76.
- Bartish, I.V., Antonelli, A., Richardson, J.E., Swenson, U. 2011. Vicariance or long-distance dispersal: historical biogeography of the pantropical subfamily Chrysophylloideae (Sapotaceae). *J. Biogeogr.* 38, 177–190.
- Bassias, Y. 1992. Petrological and geochemical investigation of rocks from the Davie Fracture Zone (Mozambique Channel) and some tectonic implications. *J. Afr. Earth Sci.* 15, 321–339.
- Bossuyt, F., Brown, R.M., Hillis, D.M., Cannatella, D.C., Milinkovitch, M.C. 2006. Phylogeny and biogeography of a cosmopolitan frog radiation: Late Cretaceous diversification resulted in continent-scale endemism in the family Ranidae. *Syst. Biol.* 55, 579–594.
- Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., Suchard, M.A., Rambaut, A., Drummond, A.J. 2014. BEAST 2: a software platform for Bayesian evolutionary analysis. *PLoS Comput. Biol.* 10: e1003537.

- Buerki S., Devey, D.S., Callmander, M.W., Phillipson, P.B., Forest, F. 2013. Spatio-temporal history of the endemic genera of Madagascar. *Bot. J. Linn. Soc.* 171, 304–329.
- Bukontaite, R., Ranarilalaitiana, T., Randriamihaja, J.H., Bergsten, J. 2015. In or out-of-Madagascar?—colonization patterns for large-bodied diving beetles (Coleoptera: Dytiscidae). *PLoS ONE* 10: e0120777
- Carpintero, D.J., Maldonado, J. 1996. Diagnostic characters and key to the genera of American Ectrichodiinae (Heteroptera, Reduviidae). *Caribb. J. Sci.* 32, 125–141.
- Chenoweth, L.B., Schwarz, M.P. 2011. Biogeographical origins and diversification of the exoneurine allodapine bees of Australia (Hymenoptera, Apidae). *J. Biogeogr.* 38, 1471–1483.
- Dallwitz, M.J. 1980. A general system for coding taxonomic descriptions. *Taxon* 29, 41–46.
- Dallwitz, M.J., Paine, T.A., Zurcher, E.J. 1999. User's guide to the DELTA Editor. Accessible at <http://delta-intkey.com>.
- Damgaard, J., Andersen, N.M., Cheng, L., Sperling, F.A.H. 2000. Phylogeny of sea skaters, *Halobates* Eschscholtz (Hemiptera, Gerridae), based on mtDNA sequence and morphology. *Zool. J. Linn. Soc., Lond.* 130, 511–526.
- Dougherty, V. 1995. A review of the New World Ectrichodiinae genera (Hemiptera: Reduviidae). *T. Am. Entomol. Soc.* 121, 173–225.
- Forero, D., Berniker, L., Weirauch, C., 2013. Phylogeny and character evolution in the bee-assassins (Insecta, Heteroptera: Reduviidae). *Mol. Phylogenet. Evol.* 66, 283–302.
- Forero, D., Weirauch, C. 2012. Comparative genitalic morphology in the New World resin bugs Apiomerini (Hemiptera, Heteroptera, Reduviidae, Harpactorinae). *Dtsch. Entomol. Z.* 59, 5–41.
- Forthman, M., Chlond, D., Weirauch, C. (in press). Taxonomic monograph of the endemic millipede assassin bug fauna of Madagascar (Hemiptera: Reduviidae: Ectrichodiinae). *B. Am. Mus. Nat. Hist.*
- Forthman, M., Weirauch, C. 2012. Toxic associations: a review of the predatory behaviors of millipede assassin bugs (Hemiptera: Reduviidae: Ectrichodiinae). *Eur. J. Entomol.* 109, 147–153.
- Fuller, S., Schwarz, M., Tierney, S. 2005. Phylogenetics of the allodapine bee genus *Braunsapis*: historical biogeography and long-range dispersal over water. *J. Biogeogr.* 32, 2135–2144.

- Ganzhorn, J.U., Wilmé, L., Mercier, J.-L. 2014. Explaining Madagascar's biodiversity. In: Scales, I.R. (Ed.), Conservation and Environmental Management in Madagascar. Routledge, New York, pp. 17–43.
- Gnos, E., Immenhauser, A., Peters, T. 1997. Late Cretaceous/early Tertiary convergence between the Indian and Arabian plates recorded in ophiolites and related sediments. *Tectonophysics* 271, 1–19.
- Goloboff, P.A., Farris, J.S., Nixon, K.C. 2008. TNT, a free program for phylogenetic analysis. *Cladistics* 24, 774–786.
- Goodman, S.M., Benstead, J.P. (Eds.) 2003. The Natural History of Madagascar. University of Chicago Press, Illinois.
- Goodman, S.M., Benstead, J.P. 2005. Updated estimates of biotic diversity and endemism for Madagascar. *Oryx* 39, 73–77.
- Harmon, L.J., Melville, J., Larson, A., Losos, J.B. 2008. The role of geography and ecological opportunity in the diversification of day geckos (*Phelsuma*). *Syst. Biol.* 57, 562–573.
- von der Heydt, A., Dijkstra, H.A. 2006. Effect of ocean gateways on the global ocean circulation in the late Oligocene and early Miocene. *Paleoceanography* 21: PA1011.
- Hill, L. 2014. Revision of *Silhouettanus* with description of nine new species (Hemiptera: Heteroptera: Schizopteridae). *Zootaxa* 3815, 353–385.
- Hwang, W.S., Weirauch, C. 2012. Evolutionary history of assassin bugs (Insecta: Hemiptera: Reduviidae): insights from divergence dating and ancestral state reconstruction. *PLoS ONE* 7: e45523.
- Illies, J. 1965. Phylogeny and zoogeography of the Plecoptera. *Annu. Rev. Entomol.* 10, 117–140.
- Jansa, S.A., Goodman, S.M., Tucker, P.K. 1999. Molecular phylogeny and biogeography of the native rodents of Madagascar (Muridae: Nesomyinae): a test of the single-origin hypothesis. *Cladistics* 270, 253–270.
- Katoh, K., Standley, D.M. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* 30, 772–780.
- Krishnankutty, S.M., Dietrich, C.H. 2011. Taxonomic revision and phylogeny of an endemic leafhopper genus

- Nesocerus* (Hemiptera: Cicadellidae: Idiocerinae) from Madagascar. Zool. J. Linn. Soc., Lond. 162, 499–543.
- Kuntner, M., Agnarsson, I. 2011. Biogeography and diversification of hermit spiders on Indian Ocean islands (Nephilidae: *Nephilengys*). Mol. Phylogenet. Evol. 59, 477–488.
- Leclaire, L., Bassias, Y., Clocchiatti, M., Ségoufin, J. 1898. La Ridge de Davie dans le Canal de Mozambique: approche stratigraphique et géodynamique. C. R. Acad. Sci. Paris Ser. II 308, 1077–1082.
- Lewis, P.O. 2001. A likelihood approach to estimating phylogeny from discrete morphological character data. Syst. Biol. 50, 913–925.
- Liu, X., Hayashi, F., Yang, D. 2015. Phylogeny of the family Sialidae (Insecta: Megaloptera) inferred from morphological data, with implications for generic classification and historical biogeography. Cladistics 31, 18–49.
- Lomolino, M.V., Riddle, B.R., Whittaker, R.J., Brown, J.H. (Eds.) 2010. Biogeography..Sinauer Associates, Maryland.
- Maddison, W.P., Maddison, D.R., 2011. Mesquite: a modular system for evolutionary analysis. Version 2.75. Available: <http://mesquiteproject.org>.
- Maldonado, J. 1990. Systematic catalogue of the Reduviidae of the world (Insecta: Heteroptera). Caribb. J. Sci. Special Edition, 1–694.
- Matzke, N.J. 2013. Probabilistic historical biogeography: new models for founder-event speciation, imperfect detection, and fossils allow improved accuracy and model-testing. Frontiers Biogeogr. 5, 242–248.
- Matzke, N.J. 2014. Model selection in historical biogeography reveals that founder-event speciation is a crucial process in island clades. Syst. Biol. Available online August 14, 2014, doi: 10.1093/sysbio/syu056.
- McCall, R.A. 1997. Implications of recent geological investigations of the Mozambique Channel for the mammalian colonization of Madagascar. Proc. R. Soc. Lond. B 264. 663–665.
- Monaghan, M.T., Gattolliat, J.-L., Sartori, M., Elouard, J.-M., James, H., Derleth, P., Glaizot, O., de Moor, F., Vogler, A.P. 2005. Trans-oceanic and endemic origins of the small minnow mayflies (Ephemeroptera,

- Beatidae) of Madagascar. *Proc. R. Soc. B* 272, 1829–1836.
- Mugrabi, D.F., Azevedo, C.O. 2010. Insecta, Hymenoptera, Bethyridae: range extension and filling gaps in Madagascar. *Check List* 6, 062–063.
- Nazari, V., Larsen, T.B., Lees, D.C., Brattström, O., Bouyer, T., van de Poel, G., Hebert, P.D.N. 2011. Phylogenetic systematics of *Colotis* and associated genera (Lepidoptera: Pieridae): evolutionary and taxonomic implications. *J. Zool. Syst. Evol. Res.* 49, 204–215.
- Nixon, K.C. 2002. WinClada ver. 1.00.08. Published by the author, Ithaca, New York.
- Nobre, T., Eggleton, P., Aanen, D.K. 2010. Vertical transmission as the key to the colonization of Madagascar by fungus-growing termites? *Proc. R. Soc. B* 277, 359–365.
- Noonan, B.P., Chippindale, P.T. 2006. Vicariant origin of Malagasy reptiles supports Late Cretaceous Antarctic land bridge. *Am. Nat.* 168, 730–741.
- Okajima, Y., Kumazawa, Y. 2010. Mitochondrial genomes of acrodont lizards: timing of gene rearrangements and phylogenetic and biogeographic implications. *BMC Evol. Biol.* 10: 141, 1–15.
- Parham, J.F., Donoghue, P.C.J., Bell, C.J., Calway, T.D., Head, J.J., Holroyd, P.A., Inoue, J.G., Irmis, R.B., Joyce, W.G., Ksepka, D.T., Patané, J.S.L., Smith, N.D., Tarver, J.E., van Tuinen, M., Yang, Z., Angielczyk, K.D., Greenwood, J.M., Hipsley, C.A., Jacobs, L., Makovicky, P.J., Müller, J., Smith, K.T., Theodor, J.M., Warnock, R.C.M. 2011. Best practices for justifying fossil calibrations. *Syst. Biol.* 61: 1–14.
- Paulian, R., Viette, P. 2003. An introduction to terrestrial and freshwater invertebrates. In: Goodman, S.M., Benstead, J.P. (Eds.), *The Natural History of Madagascar*. University of Chicago Press, Illinois, pp. 503–511.
- Phillipson, P.B., Schatz, G.E., Lowry II, P.P., Labat, J.-N. 2006. A catalogue of the vascular plants of Madagascar. In: Ghazanfar, S.A., Beentje, H.J. (Eds.), *Taxonomy and Ecology of African Plants: Their Conservation and Sustainable Use*, Proceedings XVIIth AETFAT Congress, Royal Botanic Gardens, Kew. pp. 613–627.
- Plummer, P.S., Belle, E.R. Mesozoic tectono-stratigraphic evolution of the Seychelles microcontinent. *Sediment.*

- Geol. 96, 73–91.
- Popham, E.J. 2000. The geographical distribution of the Dermaptera (Insecta) with reference to continental drift. *J. Nat. Hist.* 34, 2007–2027.
- Poux, C., Madsen, O., Marquard, E., Vieites, D.R., de Jong, W.W., Vences, M. 2005. Asynchronous colonization of Madagascar by the four endemic clades of primates, tenrecs, carnivores, and rodents as inferred from nuclear genes. *Syst. Biol.* 54, 719–730.
- Pyron, R.A. 2014. Biogeographic analysis reveals ancient continental vicariance and recent oceanic dispersal in Amphibians. *Syst. Biol.* Available online July 17, 2014, doi: 10.1093/sysbio/syu042.
- R Core Team. 2015. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Raselimanana, A.P., Noonan, B., Karanth, K.P., Gauthier, J., Yoder, A.D. 2009. Phylogeny and evolution of Malagasy plated lizards. *Mol. Phylogenet. Evol.* 50, 336–344.
- Ree, R.H., Moore, B.R., Webb, C.O., Donoghue, M.J. 2005. A likelihood framework for inferring the evolution of geographic range on phylogenetic trees. *Evolution* 59, 2299–2311.
- Ree, R.H., Smith, S.A. 2008. Maximum likelihood inference of geographic range evolution by dispersal, local extinction, and cladogenesis. *Syst. Biol.* 57, 4–14.
- Rehan, S.M., Chapman, T.W., Craigie, A.I., Richards, M.H., Cooper, S.J.B., Schwarz, M.P. 2010. Molecular phylogeny of the small carpenter bees (Hymenoptera: Apidae: Ceratinini) indicates early and rapid global dispersal. *Mol. Phylogenet. Evol.* 55, 1042–1054.
- Rogers, R.R., Hartman, J.H., Krause, D.W. 2000. Stratigraphic analysis of Upper Cretaceous Rocks in the Mahajanga Basin, Northwestern Madagascar: implications for ancient and modern faunas. *J. Geol.* 108, 275–301.
- Ronquist, F. 1997. Dispersal-vicariance analysis: a new approach to the quantification of historical biogeography. *Syst. Biol.* 46, 195–203.
- Sanmartín, I., Enghoff, H., Ronquist, F. 2001. Patterns of animal dispersal, vicariance and diversification in the Holarctic. *Biol. J. Linn. Soc.* 73, 345–390.

- Scales, I.R. 2014. Conservation as the crossroads: biological diversity, environmental change and natural resource use in Madagascar. In: Scales, I.R. (Ed.), *Conservation and Environmental Management in Madagascar*. Routledge, New York, pp. 1–13.
- Schwarz, M.P., Fuller, S., Tierney, S.M., Cooper, S.J.B. 2006. Molecular phylogenetics of the exoneurine allodapine bees reveal an ancient and puzzling dispersal from Africa to Australia. *Syst. Biol.* 55, 31–45.
- Seward, D., Grujic, D., Schreurs, G. 2004. An insight into the breakup of Gondwana: identifying events through low-temperature thermochronology from the basement rocks of Madagascar. *Tectonics* 23: TC3007.
- Simon, C., Frati, F., Beckenbach, A., Crespi, B., Liu, H., Flook, P. 1994. Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Ann. Entomol. Soc. Am.* 87, 651–701.
- Sole, C.L., Wirta, H., Forgie, S.A., Scholtz, C.H. 2011. Origin of Madagascan Scarabaeini dung beetles (Coleoptera: Scarabaeidae): dispersal from Africa. *Insect Syst. Evol.* 42, 29–40.
- Stamatakis, A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30, 1312–1313.
- Storey, M., Mahoney, J.J., Saunders, A.D., Duncan, R.A., Kelly, S.P., Coffin, M.F. 1995. Timing of hot spot-related volcanism and the breakup of Madagascar and India. *Science* 267, 852–855.
- Tamura, K., Stecher, G., Peterson, D., Filipski, A., Kumar, S. 2013. MEGA6: Molecular Evolutionary Genetics Analysis version 6.0. *Mol. Biol. Evol.* 30, 2725–2729.
- Torres, E., Lees, D.C., Vane-Wright, R.I., Kremen, C., Leonard, J.A., Wayne, R.K. 2001. Examining monophyly in a large radiation of Madagascan butterflies (Lepidoptera: Satyrinae: Mycalesina) based on mitochondrial DNA data. *Mol. Phylogenet. Evol.* 20, 460–473.
- Torsvik, T.H., Tucker, R.D., Ashwal, L.D., Carter, L.M., Jamtveit, B., Vidyadharan, K.T., Venkataramana, P. 2000. Late Cretaceous India-Madagascar fit and timing of break-up related magmatism. *Terra Nova* 12, 220–224.
- Vaidya, G., Lohman, D.J., Meier, R. 2011. SequenceMatrix: concatenation software for the fast assembly of multi-gene datasets with character set and codon information. *Cladistics* 27, 171–180.

- Vences, M., Kosuch, J., Glaw, F., Bohme, W., Veith, M. 2003a. Molecular phylogeny of hyperoliid treefrogs: biogeographic origin of Malagasy and Seychellean taxa and re-analysis of familial paraphyly. *J. Zool. Syst. Evol. Res.* 41, 205–215.
- Vences, M., Vieites, D.R., Glaw, F., Brinkmann, H., Kosuch, J., Veith, M., Meyer, A. 2003b. Multiple overseas dispersal in amphibians. *P. R. Soc. B* 270, 2435–2442.
- Vences, M. 2004. Origin of Madagascar's extant fauna: a perspective from amphibians, reptiles and other non-flying vertebrates. *Ital. J. Zool.* 71: S2, 217–228.
- Warren, B.H., Strasberg, D., Bruggemann, J.H., Prys-Jones, R.P., Thébaud, C. 2010. Why does the biota of the Madagascar region have such a strong Asiatic flavour? *Cladistics* 26, 526–538.
- Weirauch, C. 2008. Cladistic analysis of Reduviidae (Heteroptera: Cimicomorpha) based on morphological characters. *Syst. Entomol.* 33, 229–274.
- Weirauch, C. 2010. *Tribelocodia ashei*, new genus and new species of Reduviidae (Insecta: Hemiptera), has implications on character evolution in Ectrichodiinae and Tribelocephalinae. *Insect Syst. Evol.* 41, 103–122.
- Weirauch, C., Munro, J.B. 2009. Molecular phylogeny of the assassin bugs (Hemiptera: Reduviidae), based on mitochondrial and nuclear ribosomal genes. *Mol. Phylogenet. Evol.* 53, 287–299.
- Wikström, N., Avino, M., Razafimandimbison, S.G., Bremer, B. 2010. Historical biogeography of the coffee family (Rubiaceae, Gentianales) in Madagascar: case studies from the tribes Knoxieae, Naucleaeae, Paederieae and Vanguerieae. *J. Biogeogr.* 37, 1094–1113.
- Wirta, H., Orsini, L., Hanski, I. 2008. An old adaptive radiation of forest dung beetles in Madagascar. *Mol. Phylogenet. Evol.* 47, 1076–1089.
- Wirta, H., Viljanen, H., Orsini, L., Montreuil, O., Hanski, I. 2010. Three parallel radiations of Canthonini dung beetles in Madagascar. *Mol. Phylogenet. Evol.* 57, 710–727.
- de Wit, M.J. 2003. Madagascar: heads it's a continent, tails it's an island. *Annu. Rev. Earth Planet. Sci.* 31, 213–248.

- Yoder, A.D., Nowak, M.D. 2006. Has vicariance or dispersal been the predominant biogeographic force in Madagascar? Only time will tell. *Annu. Rev. Ecol. Evol. Syst.* 37, 405–431.
- Yu, Y., Harris, A.J., He, X. 2010. S-DIVA (Statistical Dispersal-Vicariance Analysis): a tool for inferring biogeographic histories. *Mol. Phylogenet. Evol.* 56, 848–850.
- Yu, Y., Harris, A.J., Blair, C., He, X. 2015. RASP (Reconstruct Ancestral State in Phylogenies): a tool for historical biogeography. *Mol. Phylogenet. Evol.* 87, 46–49.
- Zakharov, E.V., Smith, C.R., Lees, D.C., Cameron, A., Vane-Wright, R.I., Sperling, F.A.H. 2004. Independent gene phylogenies and morphology demonstrate a Malagasy origin for a wide-ranging group of swallowtail butterflies. *Evolution* 58, 2763–2782.
- Zhou, L., Su, Y.C.F., Thomas, D.C., Saunders, R.M.K. 2012. ‘Out-of-Africa’ dispersal of tropical floras during the Miocene climatic optimum: evidence from *Uvaria* (Annonaceae). *J. Biogeogr.* 39, 322–335.

Figure Legends

Fig. 1. Examples and criteria of limited (top) and extreme (bottom) sexual dimorphism in Ectrichodiinae.

Arrows in top panel indicate distal margin of wings.

Fig. 2. Best tree based on ML analysis of 56 taxa, four G-INS-I aligned gene partitions (COI, 18S, 28S D2, 28S D3–D5), and GTR+ Γ +I model of sequence evolution (Final Ln = -25426.19528). Terminal branches are colored by occurrence of a taxon in a given biogeographic region, which corresponds to the map legend. Numbers in gray circles refer to nodes discussed in the text. Bootstrap (BS) values ≥ 50 are reported below branches, with the exception that BS values are reported next to the node of *Microstematoides atrocyanea* and *Centraspis ducalis*, as well as the node of the clade containing *Microstematoides atrocyanea* and *Ectrichodia crux*.

Fig. 3. Strict consensus tree from 938 parsimonious trees from a TNT analysis of 110 taxa (outgroups removed from figure) and a combined morphological and molecular dataset (length = 7250 steps; RI = 0.587; CI = 0.356). Terminal branches are colored by taxon distribution, which corresponds to the map

legend. Numbers in gray circles refer to nodes discussed in the text and SM10. Bootstrap values ≥ 50 are reported below branches. Examples of morphological characters that are discussed in the text are given: **A.–D.** Dorsal (**A.**, **B.**) and lateral (**C.**, **D.**) head morphology: **A.** *Glymmatophora crassipes*, **B.** *Toxopus tibialis*, **C.** *Maraenaspis bidens*, **D.** *Tanindrazanus kathrynae*; **E.** Lateral antennal morphology of *Toxopus insignis*; Dorsal (**F.–H.**), lateral (**I.**), and ventral (**J.**): **F.** *Maraenaspis bidens*, **G.** *Toxopus italaviana*, **H.** *Marojejycoris notadichroa*, **I.** *Gibbosella conisimilis*, **J.** *Tanindrazanus irwini*; **K.** Metathoracic gland evaporatorium of *Tanindrazanus varicolor* in lateral view; **L.** Ventral mid leg morphology of *Glymmatophora crassipes*. **M.** Hemelytral morphology of *Gibbosella planiscutum*; **N.** Ventral abdominal morphology of *Distirogaster tarsalis*. **A.**, **C.–G.**, and **I.–M.** modified from Forthman et al. (in press). Character numbers and character state codings are listed in Table 2. Abbreviations used in figures: **1A**, first anal vein; **aa**, antennal articulation; **ain**, antennal insertion site; **ap**, antennal pseudoarticulation; **apl**, anterior pronotal lobe; **app**, anterolateral pronotal projection; **asp**, anterior femoral subapical protuberance; **bf**, basiflagellomere; **cl**, clypeus; **co**, corium; **cp**, corial pterostigma; **Cu**, cubitus; **df**, distiflagellomere; **e**, compound eye; **exM**, extension of M beyond M+Cu distal junction; **fmp**, femoral medial protuberance; **g**, gula; **is**, interocular sulcus; **L2**, labial segment II (first visible segment); **L3**, labial segment III (second visible segment); **M**, media; **mc**, metacoxa; **mf**, mid femur; **mge**, metathoracic gland evaporatorium; **mms**, transverse suture between meso- and metasterna; **mss**, mesosternum; **mt**, mid trochanter; **mts**, metasternum; **p**, pedicel; **pa**, papillae; **pc**, sternal paramedian carinae; **pcd**, postclypeal depression; **ppl**, posterior pronotal lobe; **psp**, posterior femoral subapical protuberance; **pts**, pronotal transverse suture; **R**, radius; **s**, scape; **s2**, sternite II (fused sternites I and II); **s3**, sternite III; **sc**, scutellum; **sis**, sternal intersegmental suture; **sl**, synthlipsis; **sld**, sternal medial longitudinal depression.

Fig. 4. Ancestral range reconstructions from DEC (left) and DEC+j (right) models for select nodes. Pie charts indicate the relative likelihoods of each reconstructed distribution, with geographical distributions color coded according to the legend at the top of the figure. The DEC+j model performed significantly better than the DEC: DEC+j LnL = -27.10; DEC LnL = -39.09; LRT p-value = $9.80e^{-7}$. Select dispersal

events contributing to Madagascar's Ectrichodiinae diversity is shown for each model. Dispersal events unique to each model are indicated by dashed arrows.

Table Legends

Table 1. Taxon sampling, USI codes, ethanol molecular specimen collection numbers (RCW), depositories, and GenBank accession numbers for sequenced specimens. Asterisks indicate sequences retrieved from GenBank.

Table 2. Morphological characters and character state codings.

Supplementary Material Legends

Supplementary Material 1. Morphological character matrix in nexus format for cladistic analysis.

Supplementary Material 2. Molecular character matrix in nexus format for cladistic analysis.

Supplementary Material 3. Unambiguous character optimizations on the strict consensus tree from parsimony analysis of 110 taxa and a combined morphological and molecular dataset.

Supplementary Material 4. Fossil information and calibration nodes, means, standard deviations (SD) or sigma values, and offsets (in real space).

Supplementary Material 5. DEC and DEC+j time-stratified dispersal rate matrix. N = Neotropical, A = Afrotropical, M = Madagascar, O = Oriental.

Supplementary Material 6. Equal (EW) and implied weights (IW) molecular parsimony results for 56 taxa. **A.** Strict consensus tree from 147 parsimonious trees from an EW analysis (length = 6232 steps; RI = 0.563; CI = 0.336). **B.** Strict consensus tree from 3 parsimonious trees from an IW analysis with $k = 3$ (length = 470.59899 steps; RI = 0.560; CI = 0.335). **C. & D.** Single parsimonious trees from IW analyses when $k = 6$ (**C.**) (length = 340.36981 steps; RI = 0.560; CI = 0.335) and $k = 10$ (**D.**) (length = 252.28554 steps; RI = 0.561; CI = 0.336). For **A.–D.**, bootstrap values ≥ 50 are reported below branches.

Supplementary Material 7. Best tree based on ML analysis of 110 taxa, GTR+ Γ +I model of sequence evolution for each G-INS-I aligned gene partitions (COI, 18S, 28S D2, 28S D3–D5), and Mkv model of

morphological evolution (Final Ln = -31224.182191). Outgroups not shown. Bootstrap values ≥ 50 are reported below branches.

Supplementary Material 8. Divergence time estimates based on a BEAST analysis using the ML best tree (Fig. 2) and the same molecular dataset (COI, 18S, 28S D2, 28S D3–D5). Each gene partition was assigned an unlinked site model (GTR+ Γ +I) and were linked to the same tree model. A linked clock model was assigned to 28S D2 and 28S D3–D5 to reflect their single identity. Three unlinked relaxed lognormal clock models were assigned for 18S, 28S, and COI. Placement of two outgroup fossil and two secondary calibrations are indicated by red diamonds. Purple node bars indicated 95% HPD credibility intervals.

Supplementary Material 9. Ancestral range reconstructions from the DIVA model for select nodes. Geographical distributions are color coded according to the legend.

Supplementary Material 10. Discussion of morphological support for major lineages recovered in the combined morphological and molecular phylogenetic analysis.

Figure 1

Fig. 1. Examples and criteria of limited (top) and extreme (bottom) sexual dimorphism in Ectrichodiinae. Arrows in top panel indicate distal margin of wings.

Figure 2

Fig. 2. Best tree based on ML analysis of 56 taxa, four G-INS-I aligned gene partitions (COI, 18S, 28S D2, 28S D3–D5), and GTR+I +I model of sequence evolution (Final Ln = -25426.19528). Terminal branches are colored by occurrence of a taxon in a given biogeographic region, which corresponds to the map legend. Numbers in gray circles refer to nodes discussed in the text. Bootstrap (BS) values ≥ 50 are reported below branches, with the exception that BS values are reported next to the node of *Microstemmaatoides atrocyanea* and *Centraspis ducalis*, as well as the node of the clade containing *Microstemmaatoides atrocyanea* and *Ectrichodia crux*.

Figure 3

Fig. 3. Strict consensus tree from 938 parsimonious trees from a TNT analysis of 110 taxa (outgroups removed from figure) and a combined morphological and molecular dataset (length = 7250 steps; RI = 0.587; CI = 0.356). Terminal branches are colored by taxon distribution, which corresponds to the map legend. Numbers in gray circles refer to nodes discussed in the text and SM10. Bootstrap values ≥ 50 are reported below branches. Examples of morphological characters that are discussed in the text are given: **A.–D.** Dorsal (**A.**, **B.**) and lateral (**C.**, **D.**) head morphology: **A.** *Glymmatophora crassipes*, **B.** *Toxopus tibialis*, **C.** *Maraenaspis bidens*, **D.** *Tanindrazanus kathrynae*; **E.** Lateral antennal morphology of *Toxopus insignis*; Dorsal (**F.–H.**), lateral (**I.**), and ventral (**J.**): **F.** *Maraenaspis bidens*, **G.** *Toxopus italaviana*, **H.** *Marojejycoris notadichroa*, **I.** *Gibbosella conisimilis*, **J.** *Tanindrazanus irwini*; **K.** Metathoracic gland evaporatorium of *Tanindrazanus varicolor* in lateral view; **L.** Ventral mid leg morphology of *Glymmatophora crassipes*. **M.** Hemelytral morphology of *Gibbosella planiscutum*; **N.** Ventral abdominal morphology of *Distirogaster tarsalis*. **A.**, **C.–G.**, and **I.–M.** modified from Forthman et al. (in review). Character numbers and character state codings are listed in Table 2. Abbreviations: 1A, first anal vein; aa, antennal articulation; ain, antennal insertion site; ap, antennal pseudoarticulation; apl, anterior pronotal lobe; app, anterolateral pronotal projection; asp, anterior femoral subapical protuberance; bf, basiflagellomere; cl, clypeus; co, corium; cp, corial pterostigma; Cu, cubitus; df, distiflagellomere; e, compound eye; exM, extension of M beyond M+Cu distal junction; fmp, femoral medial protuberance; g, gula; is, interocular sulcus; L2, labial segment II (first visible segment); L3, labial segment III (second visible segment); M, media; mc, metacoxa; mf, mid femur; mge, metathoracic gland evaporatorium; mms, transverse suture between meso- and metasterna; mss, mesosternum; mt, mid trochanter; mts, metasternum; p, pedicel; pa, papillae; pc, sternal paramedian carinae; pcd, postclypeal depression; ppl, posterior pronotal lobe; psp, posterior femoral subapical protuberance; pts, pronotal transverse suture; R, radius; s, scape; s2, sternite II (fused sternites I and II); s3, sternite III; sc, scutellum; sis, sternal intersegmental suture; sl, synthlipsis; sld, sternal medial longitudinal depression.

Figure 4

Fig. 4. Ancestral range reconstructions from DEC (left) and DEC+j (right) models for select nodes. Pie charts indicate the relative likelihoods of each reconstructed distribution, with geographical distributions color coded according to the legend at the top of the figure. The DEC+j model performed significantly better than the DEC: DEC+j LnL = -27.10; DEC LnL = -39.09; LRT p-value = 9.80e-7. Select dispersal events contributing to Madagascar's Ectrichodiinae diversity is shown for each model. Dispersal events unique to each model are indicated by dashed arrows.

Table 1

Table 1

Taxon sampling, USI codes, ethanol molecular specimen collection numbers (RCW), depositories, and GenBank accession numbers for sequenced specimens. Asterisks indicate sequences retrieved from GenBank.

Subfamily	Taxon	USI	RCW	Depository	COI	GenBank Accession No.		
						18S	28S D2	28S D3-D5
Ectrichodiinae	<i>Bannania</i> n. sp.	UCR_ENT 00119027	3064	UCR	KT221890	KT221910	KT221939	KT221968
	nr <i>Bannania</i> sp.	UCR_ENT 00004465	609	UCR	KT221908	KT221937	KT221966	KT221996
	<i>Brontostoma colossus</i>	UCR_ENT 00119017	3049	UCR	KT221891	KT221911	KT221940	KT221969
	<i>Brontostoma sanguinosum</i>	UCR_ENT 00002617	1243	UCR	KT221892	KT221912		KT221970
	<i>Caecina</i> sp.	UCR_ENT 00004440	2636	UCR	KT221893	KT221913	KT221941	KT221971
	<i>Centraspis ducalis</i>	UCR_ENT 00119021	3014	UCR	KT221894	KT221914	KT221942	KT221972
	<i>Cleptria corallina</i>	AMNH_PBI 00218770	14	UCR		FJ230462*	FJ230543*	FJ230621*, FJ230700*
	<i>Cryptonannus</i> n. sp.	UCR_ENT 00002757	1433	UCR	KT221895	KT221915		KT221973
	nr <i>Daraxa</i> sp.	UCR_ENT 00119016	3076	UCR	KT221909	KT221938	KT221967	KT221997
	<i>Distirogaster tarsalis</i>	UCR_ENT 00006366	2898	CAS	KR606396	KT221918	KT221945	KT221976
	<i>Distirogaster</i> n. sp.1	UCR_ENT 00007158	2881	UCR	KR606393	KT221916	KT221943	KT221974
	<i>Distirogaster</i> n. sp.2	UCR_ENT 00088090	3018	UCR	KR606395	KT221917	KT221944	KT221975
	<i>Ectrichodia crux</i>	UCR_ENT 00119022	3026	UCR	KT221896	KT221919	KT221946	KT221977
	<i>Ectrichodia lucida</i>	AMNH_PBI 00218769	13	UCR		FJ230461*	FJ230542*	FJ230620*, FJ230699*
	<i>Ectrychotes serdangensis</i>	AMNH_PBI 00218830	76	UCR		FJ230479*	FJ230560*	FJ230638*, FJ230717*
	<i>Ectrychotes</i> sp.1	UCR_ENT 00119028	571	UCR	KT221897	KT221920	KT221947	KT221978
	<i>Ectrychotes</i> sp.2	AMNH_PBI 00218932	188	UCR	JQ942322*	FJ230503*	FJ230584*	FJ230661*, FJ230740*
	<i>Gibbosella quadocris</i>	UCR_ENT 00044860	2938	UCR	KR606406			
	<i>Glymmatophora crassipes</i>	UCR_ENT 00088087	3020	UCR	KR606415	KT221921	KT221948	KT221979
	<i>Maraenaspis coccinea</i>	AMNH_PBI 00218772	16	UCR		FJ230463*	FJ230544*	
	<i>Marojejycoris brevifrons</i>	UCR_ENT 00006480	2923	CAS	KR606417	KT221922		KT221980
	<i>Mendis apicimaculata</i>	UCR_ENT 00119026	2647	UCR	KT221898	KT221923	KT221949	KT221981
	<i>Microstemmatoides atrocyanea</i>	UCR_ENT 00119029	3029	UCR	KT221899	KT221924	KT221950	KT221982
	<i>Neolibavius</i> n. sp.	UCR_ENT 00003965	1830	UCR	KT221900	KT221925	KT221951	KT221983
	<i>Pothea lugens</i>	UCR_ENT 00119018	3304	UCR	KT221902	KT221927	KT221953	

Table 1 (Continued)

Subfamily	Taxon	USI	RCW	Depository	COI	GenBank Accession No.		
						18S	28S D2	28S D3-D5
Ectrichodiinae	<i>Racelda</i> sp.	AMNH_PBI 00218801	41	UCR		FJ230472*	FJ230553*	FJ230631*, FJ230710*
	<i>Rhiginia aimara</i>	UCR_ENT 00119019	3077	UCR	KT221903	KT221928	KT221954	
	<i>Rhiginia cinctiventris</i>	AMNH_PBI 00218891	139	UCR	JQ942333*	FJ230490*	FJ230571*	FJ230648*, FJ230727*
	<i>Rhiginia</i> sp.	UCR_ENT 00119020	3047	UCR	KT221904	KT221929	KT221955	KT221985
	<i>Santosia</i> sp.	UCR_ENT 00004138	2046	UCR	KT221905	KT221930	KT221956	KT221986
	<i>Tanindrazanus marginatus</i>	UCR_ENT 00006887	2902	CAS	KR606423	KT221931	KT221957	KT221987
	<i>Tanindrazanus tenebricus</i>	UCR_ENT 00006723	2931	CAS	KR606430	KT221932	KT221958	KT221988
	<i>Tanindrazanus varicolor</i>	UCR_ENT 00006482	2949	UCR	KR606433	KT221933	KT221959	KT221989
	<i>Toxopus brucei</i>	UCR_ENT 00045338	2901	UCR	KR606438		KT221960	KT221990
	<i>Toxopus fisheri</i>	UCR_ENT 00045431	2910	CAS	KR606446		KT221961	KT221991
	<i>Toxopus griswoldi</i>	UCR_ENT 00006435	2916	CAS	KR606448	KT221934	KT221962	KT221992
	<i>Toxopus toamasina</i>	UCR_ENT 00007056	2951	SU	KR606457		KT221963	
	<i>Toxopus vazimba</i>	UCR_ENT 00006472	2953	CAS	KR606462	KT221935	KT221964	KT221993
	<i>Vilius macrops</i>	UCR_ENT 00119025	3068	UCR	KT221907	KT221936		KT221995
Emesinae	<i>Emesaya incisa</i>	AMNH_PBI 00219017	282	UCR	JQ942323*	FJ230515*	FJ230598*	FJ230672*, FJ230751*
Harpactorinae	<i>Apiomerus lanipes</i>	AMNH_PBI 00219016	281	UCR	JQ942298*	FJ230514*	FJ230597*	FJ230671*, FJ230750*
	<i>Apiomerus ochropterus</i>	AMNH_PBI 00218777	22	UCR		FJ230466*	FJ230548*	FJ230625*, FJ230704*
	<i>Micrauchenus lineola</i>	AMNH_PBI 00218790	35	UCR	JQ942329*	FJ230471*	FJ230552*	FJ230630*, FJ230709*
Peiratinae	<i>Peirates punctorius</i>	AMNH_PBI 00218960	216	UCR		FJ230508*	FJ230590*	FJ230666*, FJ230745*
Saicinae	<i>Kiskeyana palassaina</i>	AMNH_PBI 00190561	10	USNM		FJ230460*	FJ230541*	FJ230619*, FJ230698*
	<i>Saica</i> sp.	AMNH_PBI 00218796	42	UCR		FJ230473*	FJ230554*	FJ230632*, FJ230711*
Stenopodainae	<i>Oncocephalus</i> sp.	UCR_ENT 00000182	79	UCR		FJ230481*	FJ230562*	FJ230640*, FJ230719*

Table 1 (Continued)

Subfamily	Taxon	USI (UCR_ENT)	RCW	Depository	COI	GenBank Accession No.		
						18S	28S D2	28S D3-D5
Triatominae	<i>Stenopoda</i> sp.	AMNH_PBI 00218904	154	UCR		FJ230493*	FJ230574*	FJ230651*, FJ230730*
	<i>Stenopodessa</i> sp.	UCR_ENT 00000078	398	FCAP		FJ230532*	FJ230611*	FJ230688*, FJ230767*
	<i>Eratyrus mucronatus</i>	UCR_ENT 00052168	1811	UCR		JQ897555*	JQ897635*	JQ897711*
	<i>Panstrongylus lignarius</i>	UCR_ENT 00052166	1813	UCR		JQ897584*	JQ897656*	JQ897741*
	<i>Triatoma recurva</i>	AMNH_PBI 00218913	170	UCR		FJ230496*	FJ230577*	FJ230654*, FJ230733*
Tribelocephalinae	<i>Opistoplatys</i> sp.1	UCR_ENT 00119024	3056	UCR	KT221901	KT221926	KT221952	KT221984
	<i>Opistoplatys</i> sp.2	UCR_ENT 00052187	1592	UCR		JQ897612*	JQ897682*	JQ897767*
	<i>Tribelocephala peyrrierasi</i>	AMNH_PBI 00219033	287	CAS		FJ230521*	FJ230601*	
	<i>Tribelocephala</i> sp.	UCR_ENT 00119023	3048	UCR	KT221906		KT221965	KT221994

Table 2

Morphological characters and character state codings.

Character no.	Character	Character states
GENERAL		
0	Wing form	(0) Macropterous; (1) Brachypterous; (2) Apterous; (3) Micropterous
1	Body length	(0) Small; (1) Medium; (2) Large
2	General coloration	(0) Dull, ranging from pale to dark brown/black; (1) Metallic; (2) Red and black, sometimes with yellow or white markings; (3) Red and dull; (4) Orange and black; (5) Red and metallic
3	Bulbous hairs on integument	(0) Present; (1) Absent
4	Vestiture	(0) Very dense; (1) Moderately dense; (2) Sparse; (3) Glabrous
HEAD		
5	Relative head length in dorsal view	(0) As long as wide; (1) Longer than wide; (2) Wider than long
6	Ventral surface of head	(0) Flat; (1) Anteromedially depressed; (2) Medially depressed; (3) Paramedially depressed
7	Depth of ventral head depression	(0) Shallow; (1) Deep
8	Ventral tubercles on head	(0) Absent; (1) Present
9	Ventrolateral tubercle on gula	(0) Absent; (1) Present
10	Anteocular length	(0) Shorter than postocular; (1) As long as postocular; (2) Longer than postocular
11	Postocular-neck constriction	(0) Absent, gradually transition; (1) Present, distinctly constricted
12	Gula shape in lateral view	(0) Flat, conforming to rounded head shape; (1) Moderately swollen ventrolaterally, not produced beyond ventral head margin; (2) Greatly swollen ventrolaterally, produced beyond ventral head margin
13	Postclypeal depression	(0) Flat; (1) Depressed
14	Width of postclypeal depression	(0) Narrow; (1) Broad
15	Depth of postclypeal depression	(0) Shallow; (1) Deep
16	Extension of postclypeal depression	(0) Posterior margin of clypeus to middle of synthlipsis; (1) Posterior margin of clypeus to interocular sulcus
17	Maxillary plate in lateral view	(0) Reaching or nearly reaching dorsal clypeal margin; (1) Not reaching dorsal clypeal margin
18	Clypeal process	(0) Absent; (1) Present

Table 2 (*Continued*)

Character no.	Character	Character states
19	Shape of clypeus	(0) Without dorsal projection; (1) With dorsal projection
20	Clypeal apex relative to labral base	(0) Not elevated; (1) Elevated
21	Antennifer armature	(0) Unarmed; (1) With dorsolateral tubercle
22	Antennal shield	(0) Absent, not expanded; (1) Present, expanded
23	Antennal insertion site in lateral view	(0) Not concealed by antennal shield; (1) Concealed by antennal shield
24	Point of antennal insertion	(0) Sublateral or lateral; (1) Dorsal
25	Presence of ocellar lens	(0) Present; (1) Absent
26	Orientation of ocellar lens	(0) Medially; (1) Laterally
27	Ocellar lens size	(0) Large; (1) Small
28	Eye height in lateral view	(0) Less than or half of head height; (1) More than half of head height
29	Eye shape	(0) Adpressed; (1) Globbose
30	Anterior eye margin	(0) Convex; (1) Concave
31	Posterior eye margin	(0) Concave; (1) Straight
32	Dorsal eye margin	(0) Not attaining dorsal head surface; (1) Attaining dorsal head surface
33	Ventral eye margin	(0) Not attaining ventral head surface; (1) Attaining ventral head surface
34	Synthlipsis width	(0) About width of eye; (1) About 1.5 times width of eye; (2) About 2 times width of eye; (3) About 2.5 times width of eye; (4) About 3 times or more width of eye
35	Scapus length	(0) Not surpassing clypeal apex; (1) Surpassing clypeal apex
36	Pedicel curvature	(0) Straight; (1) Curved
37	Scapo-pedicellar articulation	(0) Slightly bent; (1) Strongly bent, pedicel and flagellomeres point posteriad
38	Antennal segmentation	(0) 4-segmented; (1) 6-segmented; (2) 7-segmented; (3) 8-segmented; (4) >8 segments
39	Labrum structure	(0) Subdivided by transverse membrane; (1) Completely sclerotized
40	Ventral margin of labial segment III	(0) Straight; (1) Convex; (2) Concave
41	Labial segment III length	(0) Longer than labial segment II; (1) Shorter than labial segment II; (2) Subequal to labial segment II
THORAX		
42	Pronotal length	(0) Wider than long; (1) As long as wide; (2) Longer than wide

Table 2 (*Continued*)

Character no.	Character	Character states
43	Pronotal collar	(0) Dinstinct in dorsal view; (1) Indistinct in dorsal view
44	Anterior pronotal margin shape	(0) Medially concaved; (1) Straight or nearly straight
45	Anterior pronotal lobe shape	(0) Flat; (1) Dorsally elevated, conical
46	Anterior pronotal lobe structure	(0) Rugose; (1) Smooth; (2) Tuberculate
47	Anterolateral armature of pronotum	(0) Absent; (1) Small tubercle; (2) Strongly projecting process
48	Lateral carinae on anterior pronotal lobe	(0) Absent; (1) Present
49	Anterior pronotal lobe length	(0) Longer than posterior lobe; (1) Shorter than posterior lobe; (2) As long as posterior lobe
50	Anterior pronotal lobe width	(0) More than half as wide but not as wide as posterior lobe; (1) As wide as posterior lobe
51	Posterior pronotal lobe structure	(0) Striated; (1) Smooth; (2) Punctate; (3) Tuberculate; (4) Longitudinal ridges
52	Posterior pronotal lobe lateral depressions	(0) Present; (1) Absent
53	Posterior pronotal lobe lateral depression structure	(0) Not foveate; (1) Foveate
54	Extension of pronotal longitudinal depression on anterior lobe	(0) Reaching or nearly reaching anterior margin; (1) Not reaching anterior margin; (2) Absent
55	Presence of pronotal longitudinal depression on posterior lobe	(0) Absent; (1) Present
56	Structure of pronotal longitudinal depression on posterior lobe	(0) Not foveate; (1) Foveate
57	Pronotal transverse suture	(0) Incomplete, divided by paramedian ridges; (1) Complete, not divided by paramedian ridges
58	Pronotal transverse suture structure	(0) Not foveate; (1) Foveate
59	Scutellar processes	(0) None, scutellum triangular; (1) Two paramedian apical processes; (2) Two paramedian and one medial apical processes; (3) Two paramedian and two lateral processes
60	Distance between paramedian scutellar processes	(0) Narrow; (1) Broad
61	Length of paramedian scutellar processes	(0) Short; (1) Long
62	Orientation of paramedian scutellar processes	(0) Horizontally directed; (1) Dorsally directed
63	Scutellar disc surface	(0) Medially depressed; (1) Flat

Table 2 (*Continued*)

Character no.	Character	Character states
64	Prosternal stridulatory process length	(0) Short; (1) Long, surpassing posterior margin of fore coxal cavity
65	Shape of apex of prosternal stridulatory process	(0) Acute; (1) Rounded
66	Transverse suture between meso- and metasterna	(0) Complete; (1) Incomplete, at least partially
67	Mesosternal surface	(0) Medially and laterally depressed; (1) Medially depressed; (2) Flat
68	Metasternal surface	(0) Medially depressed; (1) Flat or convex
69	Presence of metathoracic gland	(0) Absent; (1) Present
70	Presence of metathoracic gland evaporatorium	(0) Present; (1) Absent
71	Metathoracic gland evaporatorium size	(0) Small, not visible in lateral view; (1) Large, visible in lateral view
LEG		
72	Shape of fore coxa	(0) Globular; (1) Elongate
73	Shape of fore femur	(0) Slender; (1) Incrassate
74	Presence of papillae on fore trochanter	(0) Absent; (1) Present
75	Presence of papillae on mid trochanter	(0) Absent; (1) Present
76	Presence of papillae on hind trochanter	(0) Absent; (1) Present
77	Presence of anteroventral row of spines/tubercles on fore femur	(0) Absent; (1) Present
78	Presence of posteroventral row of spines/tubercles on fore femur	(0) Absent; (1) Present
79	Presence of anterior subapical tubercle on fore femur	(0) Absent; (1) Present
80	Presence of posterior subapical tubercle on fore femur	(0) Absent; (1) Present
81	Presence of medial tubercle on fore femur	(0) Absent; (1) Present
82	Presence of carina on fore femur	(0) Absent; (1) Ventrally entirely carinate; (2) Ventrally carinate basally
83	Presence of papillae on fore femur	(0) Absent; (1) Present
84	Presence of anteroventral row of spines/tubercles on mid femur	(0) Absent; (1) Present
85	Presence of posteroventral row of spines/tubercles on mid femur	(0) Absent; (1) Present
86	Presence of anterior subapical tubercle on mid femur	(0) Absent; (1) Present
87	Presence of posterior subapical tubercle on mid femur	(0) Absent; (1) Present
88	Presence of medial tubercle on mid femur	(0) Absent; (1) Present

Table 2 (*Continued*)

Character no.	Character	Character states
89	Presence of papillae on mid femur	(0) Absent; (1) Present
90	Presence of anteroventral row of spines/tubercles on hind femur	(0) Absent; (1) Present
91	Presence of anterior subapical tubercle on hind femur	(0) Absent; (1) Present
92	Presence of posterior subapical tubercle on hind femur	(0) Absent; (1) Present
93	Presence of subapical medial tubercle on hind femur	(0) Absent; (1) Present
94	Presence of medial tubercle on hind femur	(0) Absent; (1) Present
95	Presence of fossula spongiosa on fore tibia	(0) Present; (1) Absent
96	Presence of fossula spongiosa on mid tibia	(0) Present; (1) Absent
97	Protibial groove	(0) Absent; (1) Present
FORE WING		
98	Extension of corium on fore wing	(0) Restricted to areas adjacent to basal wing veins and with pterostigma-like appearance; (1) Well-developed corium
99	Vestiture of corium on fore wing	(0) Long, simple setae; (1) Glabrous
100	Basal part of M and Cu on fore wing	(0) Separate veins; (1) Forming one vein, at least partially
101	Presence of cubital cell on fore wing	(0) Absent; (1) Present
102	Shape of cubital cell on fore wing	(0) Quadrate; (1) Hexagonal
103	Distal part of M and Cu on fore wing	(0) Separate veins; (1) Fused basally
104	Extension of distal part of M on fore wing	(0) Extends beyond distal M+Cu junction; (1) No extension beyond distal M+Cu junction
ABDOMEN		
105	Color pattern of dorsal laterotergites	(0) Concolor; (1) Transversely bicolor; (2) Longitudinally bicolor
106	Expansion of dorsal laterotergite II	(0) Not expanded; (1) Laterally expanded; (2) Posterior tubercle
107	Dorsal laterotergite III armature	(0) Unarmed; (1) Posterior tubercle
108	Dorsal laterotergite IV armature	(0) Unarmed; (1) Posterior tubercle
109	Dorsal laterotergite V armature	(0) Unarmed; (1) Posterior tubercle
110	Dorsal laterotergite VI armature	(0) Unarmed; (1) Posterior tubercle
111	Connection of dorsal laterotergite and mediotergite	(0) Separated by membrane; (1) Sclerotized
112	Connection of ventral laterotergites and mediotergites III-VII	(0) Separated by membrane; (1) Sclerotized
113	Ventral connexival suture on sternite II	(0) Absent; (1) Present

Table 2 (*Continued*)

Character no.	Character	Character states
114	Carinulation of intersegmental suture between abdominal sternites II and III	(0) Not carinulate; (1) Entirely carinulate
115	Carinulation of intersegmental suture between abdominal sternites III and IV	(0) Not carinulate; (1) Entirely carinulate; (2) Laterally carinulate
116	Carinulation of intersegmental suture between abdominal sternites IV and V	(0) Not carinulate; (1) Entirely carinulate; (2) Laterally carinulate
117	Carinulation of intersegmental suture between abdominal sternites V and VI	(0) Not carinulate; (1) Entirely carinulate; (2) Laterally carinulate
118	Carinulation of intersegmental suture between abdominal sternites VI and VII	(0) Not carinulate; (1) Entirely carinulate; (2) Laterally carinulate
119	Presence of paramedian carinae on abdominal sternites	(0) Absent; (1) Present
120	Abdominal sternite surface	(0) Flat; (1) Medially depressed; (2) Convex or keeled
121	Spiracle shape	(0) Ovoid or elliptical; (1) Circular
MALE GENITALIA		
122	Anteroposterior thickness of median pygophore process	(0) Flat; (1) Thickened, at least partially
123	Lateral thickness of median pygophore process	(0) Flat; (1) Thickened, at least partially
124	Lateral shape of median pygophore process	(0) Subtriangular; (1) Blade-like; (2) Hook-like; (3) Knob-like; (4) Rod-like
125	Shape of median pygophore process in caudal view	(0) Subtriangular; (1) Subquadrate; (2) Rod-like; (3) Spade-like
126	Apex of median pygophore process	(0) Entire; (1) Medially notched, bifid
127	Process on ponticulus basilaris	(0) Absent; (1) Present
128	Median process on ductifer	(0) Absent; (1) Present
129	Length of basal plate extension	(0) Shorter than basal plate; (1) As long as basal plate; (2) Longer than basal plate
130	Posterior margin of dorsal phallothecal sclerite	(0) Rounded; (1) Medially notched
131	Armature of dorsal phallothecal sclerite	(0) Unarmed; (1) Armed with denticle-like processes
132	Presence of vesica on endosoma	(0) Absent; (1) Present
133	Sclerotization of posterior endosomal margin	(0) Membranous; (1) Entirely sclerotized; (2) Paramedian sclerites; (3) Slender sclerotized processes; (4) Sclerotized lateral margin; (5) Sclerotized lateral hooks adjacent to fan-like lobes and with a medial dorsally projecting spine
134	Medial sclerotization of posterior endosomal region	(0) Membranous; (1) Slightly sclerotized; (2) Distinctly sclerite

Table 2 (*Continued*)

Character no.	Character	Character states
135	Lateral sclerotization of posterior endosomal region	(0) Membranous; (1) Sclerotized
136	Medial sclerotization of medial endosomal region	(0) Membranous; (1) Slightly sclerotized; (2) Paramedial sclerites
137	Lateral sclerotization of medial endosomal region	(0) Membranous; (1) Sclerotized
138	Medial sclerotization of anterior endosomal region	(0) Membranous; (1) Sclerotized
139	Lateral sclerotization of anterior endosomal region	(0) Membranous; (1) Sclerotized
GLANDS		
140	Presence of Brindley's gland	(0) Present; (1) Absent
141	Dorsal abdominal scent gland (DAG) I in adults	(0) Absent; (1) Present
142	DAG II in adults	(0) Absent; (1) Present
143	DAG III in adults	(0) Absent; (1) Present
144	Size of DAG III relative to DAGs I and II, external view	(0) Similar size; (1) Larger

*Graphical Abstract

Neotropical Afrotropical Madagascar Oriental Neotropical+Afrotropical Neotropical+Oriental Neotropical+Madagascar
Madagascar+Oriental Afrotropical+Oriental Afrotropical+Madagascar