**Development of an *Agrobacterium*-mediated transformation system
for the cold-adapted fungi *Pseudogymnoascus destructans* and *P.
pannorum***

Tao Zhang¹, Ping Ren¹, Vishnu Chaturvedi^{1, 2} and Sudha Chaturvedi^{1, 2*}

¹Mycology Laboratory, Wadsworth Center, New York State Department of
Health, Albany, New York, USA

²Department of Biomedical Sciences, School of Public Health, University at
Albany, Albany, New York, USA

*Corresponding authors: Sudha Chaturvedi
<sudha.chaturvedi@health.ny.gov>

Key words

Agrobacterium tumefaciens-mediated transformation (ATMT); white-nose
syndrome (WNS); homologous recombination; urease

Running title

Targeted gene deletion in cold-adapted fungi

ABSTRACT

The mechanisms of cold adaptation by fungi remain unknown. This topic is of high interest due to the emergence of white-nose syndrome (WNS), a skin infection of hibernating bats caused by *Pseudogymnoascus destructans* (*Pd*). Recent studies indicated that apart from *Pd*, there is an abundance of other *Pseudogymnoascus* species in the hibernacula soil. We developed an *Agrobacterium tumefaciens*-mediated transformation (ATMT) system for *Pd* and a related fungus *Pseudogymnoascus pannorum* (*Pp*) to advance experimental studies. *URE1* gene encoding the enzyme urease was used as an easy to screen marker to facilitate molecular genetic analyses. A Uracil-Specific Excision Reagent (USER) Friendly pRF-HU2 vector containing *Pd* or *Pp ure1::hygromycin (HYG)* disruption cassette was introduced into *A. tumefaciens* AGL-1 cells by electroporation and the resulting strains were co-cultivated with conidia of *Pd* or *Pp* for various durations and temperatures to optimize the ATMT system. Overall, 680 *Pd* (0.006%) and 1800 *Pp* (0.018%) transformants were obtained from plating of 10^7 conidia; their recoveries were strongly correlated with the length of the incubation period (96 h for *Pd*; 72h for *Pp*) and with temperature (15-18°C for *Pd*; 25°C for *Pp*). The homologous recombination in transformants was 3.1% for *Pd* and 16.7% for *Pp*. The availability of a standardized ATMT system would allow future molecular genetic analyses of *Pd* and related cold-adapted fungi.

INTRODUCTION

Pseudogymnoascus destructans (*Pd*) is the etiologic agent of white-nose syndrome (WNS), a skin infection of hibernating bats. This pathogen has caused unprecedented reductions in the abundance of the bat populations in the northeastern United States since 2006, with up to 95% mortality in some hibernacula (Blehert *et al.*, 2009, Chaturvedi *et al.*, 2010, Frick *et al.*, 2010, Blehert, 2012). *Pd* is a newly recognized fungus that is well adapted to growing under cold conditions found in caves and mines. It secretes proteolytic enzymes documented in other pathogenic fungi to cause vertebrate skin infections. The pathogen appears to have a clonal population in the US (Rajkumar *et al.*, 2011, Ren *et al.*, 2012). However, the precise mechanisms by which *Pd* survives and invades the skin of the hibernating bats are not clear. Phylogenetic analyses of hibernacula soil revealed large number of *Pseudogymnoascus* spp., which have not been implicated in WNS (Minnis & Lindner, 2013). These results indicated that *Pd* possess certain unique virulence repertoire to cause disease in bats, which seems to be missing in other *Pseudogymnoascus* spp. It is also conceivable that mammalian or non-mammalian hosts are yet to be identified for other *Pseudogymnoascus* spp.

Recently, integrated genome profile analyses of *Pd* and *Geomyces pannorum* (re-named as *Pseudogymnoascus pannorum*; accession no.

AYKR000000000) yielded preliminary information on the fundamental biological capabilities including pathogenicity, cold adaptation, and regulation mechanisms (Chibucos *et al.*, 2013). However, follow-up experimental studies are not yet possible due to unavailability of an efficient transformation system. The lack of molecular tools also extends to other fungi inhabiting the cold environment.

The *Agrobacterium tumefaciens*-mediated transformation system (ATMT) has been successfully used for molecular genetic studies of yeasts and filamentous fungi, including *Saccharomyces* (Piers *et al.*, 1996), *Cryptococcus* (McClelland *et al.*, 2005), *Coccidioides* (Abuodeh *et al.*, 2000), *Aspergillus* (Gouka *et al.*, 1999, Michielse *et al.*, 2005), *Trichophyton* (Yamada *et al.*, 2009), *Fusarium* (Mullins *et al.*, 2001), *Glomus* (Helber & Requena, 2008), and *Oculimacula* (Eckert *et al.*, 2005) and *Verticillium* (Dobinson *et al.*, 2004). ATMT also has been shown to be an important alternative to other fungal transformation systems, as it yields a high number of transformants, is easier to perform, does not require special equipment, and is characterized by a high frequency of homologous recombination (Frandsen, 2011).

The enzyme urease encoded by *URE1* gene is an important virulence factor for the fungal pathogens *Cryptococcus neoformans* and *Coccidioides*

posadasii (Olszewski *et al.*, 2004, Mirbod-Donovan *et al.*, 2006) and the bacterial pathogens *Helicobacter pylori* and *Proteus mirabilis* (Jones *et al.*, 1990, Eaton *et al.*, 1991, Tsuda *et al.*, 1994). It has been reported that *Pd* also secretes significant amounts of urease, but its precise function in either the saprophytic or parasitic life of this fungus is not clear (Chaturvedi *et al.*, 2010, Reynolds & Barton, 2014). In this report, we have used the urease-encoding gene *URE1* as an easily screenable marker to develop an ATMT system for cold-adapted fungi, which would prove valuable for future molecular genetics and pathogenic studies of *Pd* and other related species in this genus.

MATERIALS AND METHODS

Strains, plasmids and media

All strains and plasmids used in this study are listed in Table 1. *Pseudogymnoascus destructans* (*Pd*) and *P. pannorum* (*Pp*) strains were routinely maintained on yeast extract peptone dextrose (YPD) agar and stored in 15% glycerol at -70°C. Potato dextrose agar (PDA; Difco), Christensen's urea agar (Sigma-Aldrich), YNB medium [2% glucose, 0.17% yeast nitrogen base without amino acids and ammonium sulfate (Cat. No.

233520) and 2% agar], YNB-U (0.227% urea as sole source of nitrogen) were used to assess phenotypes of mutants and wild type (WT) strains of *Pd* and *Pp*. The induction medium (IM) used for co-cultivation experiments was composed of K₂HPO₄·3H₂O 1.9 g, K₂HPO₄ 2.05 g, MgSO₄·7H₂O 0.5 g, (NH₄)₂SO₄ 0.5 g, glucose 1.8 g, 4% (w/v) glycerol and a final concentration of 40 mM MES hydrate (Sigma-Aldrich) per liter (Zhang *et al.*, 2013). Yeast extract broth (YEB) used for culture of *Agrobacterium tumefaciens* (AGL-1) cells contained 10 g peptone, 1 g yeast extract, 5 g sucrose and 0.5 g MgSO₄·7H₂O per liter and adjusted to pH 7.0.

URE1* gene characterization from *Pd* and *Pp

DNA sequences of *PdURE1* and *PpURE1* were first obtained from *Pd* (http://broadinstitute.org/annotation/genome/Geomyces_destructans/MultiHome.html) and *Pp* (<http://fungalgenomics.concordia.ca/fungi/Gpan.php>) genome databases and then blasted against *Pd* (<http://www.ncbi.nlm.nih.gov/nuccore/AYKP000000000>) and *Pp* (<http://www.ncbi.nlm.nih.gov/nuccore/AYKR000000000>) databases to get full-length sequences (Chibucos *et al.*, 2013). Open reading frame (ORF) analysis was carried out using the programs AUGUSTUS (Stanke *et al.*, 2004, Stanke & Morgenstern, 2005) and FGENESH (www.softberry.com).

Gene models were manually curated using exon/intron boundary predictions from SPLICEPORT (Dogan *et al.*, 2007). Multiple sequence alignments were created with reference to selected GenBank sequences using the MAFFT program (Kato *et al.*, 2009). Phylogenetic analysis using the neighbor-joining (NJ) method with 1,000 bootstrap replicates was done using MEGA 4.1 (Tamura *et al.*, 2007).

Vector construction

Genomic DNA from *Pd* and *Pp* cultures was extracted using a modified protocol (Möller *et al.*, 1992). In brief, approximately two loopfuls of the fungal colony was removed from the agar plate and placed in 300 µl of genomic DNA extraction buffer (100 mM Tris [pH 8.0], 10 mM EDTA, 2% SDS, 1.4 M NaCl, 1% CTAB, 0.4 µg/ml proteinase K). The mixture was incubated at 65°C for 1 h followed by chloroform: isoamyl alcohol (24:1) extraction, and precipitation with isopropanol and washing with 70% ethanol. The precipitated DNA was centrifuged at 12,000 rpm and the resulting pellet was dried under air and dissolved in 50 µl of Tris-EDTA (TE) buffer. The 5'- and 3'- prime arms of the *URE1* gene (approximately 1.5 kb) from *Pd* and *Pp* were PCR amplified using primers that contained 5' - deoxyuridine extensions (Table 1). These PCR amplicons were cloned into the Uracil-Specific Excision Reagent (USER)-Friendly cloning vector pRF-HU2

flanking the hygromycin B resistance (*HYG*) gene under the control of the *Aspergillus nidulans trpC* promoter and terminator. The vector contained the necessary elements (*oriV* and *trfA*) for replication in *Escherichia coli* and *Agrobacterium tumefaciens* cells (Frandsen *et al.*, 2008, Zhang *et al.*, 2013). The resulting plasmids were designated as *pPdure1::HYG* and *pPpure1::HYG* gene disruption cassettes.

ATMT transformation procedure

The ATMT transformation of *Pd* and *Pp* was carried out as per published protocols (Gouka *et al.*, 1999, Mullins *et al.*, 2001). In brief, *A. tumefaciens* strain AGL-1 was transformed with *pPdure1::HYG* and *Ppure1::HYG* disruption cassettes by electroporation (Mozo & Hooykaas, 1991). The transformed AGL-1 cells were subsequently grown in YEB broth containing kanamycin (50 µg/ml) for 48 h at 28°C. One milliliter of the cell culture was spun down and washed with induction medium (IM). Bacterial cells were diluted in IM medium (10 ml) with or without 200 µM acetosyringone (AS; Sigma-Aldrich) to an OD₆₀₀ = 0.30 and incubated (200 RPM) at 28°C until an OD₆₀₀ = of approximately 0.80 (1 x 10⁸ cells/ml) reached.

Conidia from *Pd* and *Pp* cultures grown in PDA agar for four weeks were harvested by gently scraping fungal growth from the agar surface and

176 passing it through 27G needle to get a suspension of single conidia (Figs.
177 S1A & 1B). Conidia were counted microscopically and adjusted to
178 concentrations as needed for various experiments. To determine optimum
179 concentration of hygromycin B required for complete inhibition of *Pd* or *Pp*
180 growth, approximately 100 μ l of conidial suspension (10^8) were spread on
181 PDA plates containing various concentration of hygromycin B (50-200 μ g/ml)
182 and plates were incubated at 15° C for 1-4 weeks and presence or absence
183 of growth of WT strains of *Pd* or *Pp* was assessed.

184 For transformation studies, conidia were adjusted to 1×10^7 cells/ml in
185 IM broth with or without AS. Aliquots of 100 μ l of *Pd* or *Pp* conidial
186 suspension (1×10^6 conidia) were mixed with an equal volume of AGL-1 cells
187 (1×10^7 cells) carrying either *pPdure1::HYG* or *pPpure1::HYG*, and were
188 evenly spread onto sterilized nitrocellulose membranes (Cat. no. D9527,
189 Sigma-Aldrich) placed on IM agar with or without AS. A total of 3
190 membranes per condition were used. The co-cultivation experiments were
191 performed at various temperatures (10°C-30°C) and time intervals (24 h to
192 96 h). The membranes were then transferred to PDA containing hygromycin
193 (100 μ g/ml) as a selection agent for the recovery of putative hygromycin
194 positive (Hyg⁺) transformants and cefotaxime (200 μ g/ml) to kill the
195 *Agrobacterium tumefaciens* cells. Representative plates containing putative
196 Hyg⁺ *Pd* and *Pp* transformants are shown (Supplementary Figs. S1C & S1D).
197 The putative Hyg⁺ transformants were then streaked on selection medium

and plates were incubated at 15°C for 2 weeks. Representative plates with putative Hyg⁺ colonies are shown (Supplementary Figs. S1E & S1F).

DNA extraction and analysis

Genomic DNA from the putative Hyg⁺ transformants was extracted as described above for the *Pd* and *Pp* WT strains. The transformants were first screened for the presence of the *HYG* gene and then for *ure1::HYG* disruption cassette integration at the homologous site by PCR (Table 1). The PCR conditions were: initial denaturation at 95°C for 3 min followed by 40 cycles at 94°C for 30 sec, annealing at 55°C to 63°C for 30 sec (based on primer pair) and extension at 72°C for 40 sec to 3 min (based on amplicon size). The transformants with single *ure1::HYG* integration at the homologous site were further confirmed by Southern analysis. In brief, genomic DNA (~5 µg) was digested with restriction enzymes, fractionated by electrophoresis (1% agarose gel), transferred to a HybondTM-XL nylon membrane (Cat. no. RPN303 S, GE Healthcare), and fixed by UV cross-linking. The membranes were hybridized overnight at 55°C with *HYG* probe [PCR amplified fragment of *HYG* labeled with ³²P-dCTP by random priming (Perkin Elmer)]. Following hybridization, these membranes were washed, dried and visualized using Image Quant Software (Molecular Dynamics Inc.).

Urease enzyme activity assay

The urease enzyme activities of mutants and WT strains were assessed by inoculating 10 μ l of conidial suspensions (10^7 conidia/ml) on Christensen's urea agar tubes and these tubes were incubated at 15°C for 2 weeks. A color change to bright magenta was indicative of urease production.

For assay of urease enzyme activity, the *ure1* mutant and WT strains were first inoculated in YNB-U broth and incubated at 15°C for 10 days. Cells were harvested by centrifugation at 10,000 g for 10 min. The cell pellets were washed twice in 50 mM sodium phosphate buffer (pH 7.6) and re-suspended in the same buffer. Cells were disrupted at a setting of 6.5 for 45 s with a FastPrep FP120 apparatus (Bio101 Savant Instruments, Inc., Holbrook, NY). Cellular debris was removed by centrifugation at 12,000 g at 4°C for 15 min, and the supernatant assayed for urease enzyme activity according to the protocol described in the QuantiChrom™ Urease Assay Kit (BioAssay Systems, Hayward, CA, USA). One unit of urease enzyme activity is defined as the amount of enzyme required to hydrolyze one μ mol of urea per min at 25°C. Hydrolysis was measured by monitoring the rate of release of ammonium ions (NH_4^+) plus ammonia (NH_3) from the urea substrate as determined by the Bertholet reaction. All analyses were conducted in triplicates.

242

243 Next, we tested if *URE1* gene is involved in urea metabolism.
244 Approximately 10 µl aliquots of serial dilutions of conidial suspension of *Pd*
245 and *Pp* WT and their respective *ure1* mutant strains were spotted on YNB-U
246 medium and plates were incubated at 15°C for 7 days.

247

248 **Growth and sporulation studies**

249

250 The *ure1* mutant and WT strains were assessed for growth at various
251 temperatures. Conidial suspensions were prepared as described above, and
252 adjusted to 10⁷ cells/ml. Serial 10-fold dilutions were prepared, and 10 µl of
253 each dilution was spotted onto YPD and PDA agar plates. All agar plates
254 were incubated at 6°C, 15°C and 25°C for 7-14 days. For sporulation,
255 mutant and WT strains were grown on PDA agar at 15°C for 25 days, scotch
256 tape mounts were prepared and assessed microscopically.

257

258 **Statistical analysis**

259 The student t-test was used for the analysis of efficiency of ATMT
260 transformation including temperature and duration of incubation. A *p*-value
261 of <0.05 was considered significant.

262

263 **Nucleotide Sequence Accession Numbers**

The nucleotide sequence of *PdURE1* and *PpURE1* were deposited in GenBank under accession numbers KP195072 and KP195073, respectively.

RESULTS

Identification of the *Pd/Pp URE1* gene and bioinformatic analyses

Based on *Pd/Pp* genome databases and sequence alignments, putative urease orthologues, designated *PdURE1* and *PpURE1* were obtained. The coding sequences of *PdURE1* spans 3,127 bp, corresponding to a predicted open reading frame of 837 amino acids, consisting of six exons interrupted by five introns (KP195072). In contrast, the sequence of *PpURE1* spans 3,904 bp coding for a predicted open reading frame of 882 amino acids with six exons and five introns (KP195073). The putative amino acid sequence of *PdURE1* shared 94% identity with putative *PpURE1*, and close homology to the metallo-dependent urease of *Glarea lozoyensis* ATCC 20868 (EPE30823, 77% identity). A similar level of identity was also noted with the putative urease of *Trichoderma reesei* QM6a (GenBank accession No. EGR47585, 75% identity) and *Coccidioides posadasii* 735 (GenBank accession No. AAC49868, 71% identity). The sequences also showed conservation of all of the histidine residues thought to be involved in binding to the nickel metallocenter, binding to the substrate, and catalysis (Cox *et al.*, 2000) (Fig.

1A). These results strengthened the observation that *URE1* gene function in urea metabolism is conserved across fungal species, including cold-adapted fungi. The phylogenetic analysis using neighbor-joining method (Saitou & Nei, 1987) revealed that *PdURE1* clustered closely together with the ortholog of *PpURE1*, consistent with their common origin within the *Pseudeurotiaceae* (Fig. 1B).

Optimization of ATMT for *Pd* and *Pp*

Prior to experimental procedures with ATMT, the hygromycin B (HygB) minimum inhibitory concentration was determined for *Pd* and *Pp*. Hygromycin concentration of 100 µg/ml showed complete inhibition of the growth of both *Pd* and *Pp*, whereas a hygromycin concentration of 75 µg/ml showed restricted growth of both these fungi (data not shown). Other antibiotics tested, including cefotaxime (200 µg/ml) and kanamycin (50 µg/ml), did not have any adverse effect on the growth of either *Pd* or *Pp* (data not shown).

To determine whether acetosyringone (AS), a natural wound response molecule, is essential for the ATMT of *Pd* or *Pp*, this chemical was removed from the IM medium in which the *Agrobacterium* cells were grown prior to co-cultivation or was removed from the co-cultivation medium. *Pd* or *Pp*

Hyg⁺ transformants were not recovered in the absence of AS, suggesting that this inducer molecule is critical for ATMT of both *Pd* and *Pp*. In the presence of AS and at a ratio of 1:10 *Pd* or *Pp* conidia and AGL-1 cells carrying either *Pdure1::HYG* or *Ppure1::HYG* plasmid, a minimum of 24 h to 36 h of co-cultivation was required to obtain *Pd* and *Pp* Hyg⁺ transformants (1~4 per 3 nitrocellulose membranes), respectively. However, the yields of Hyg⁺ transformants were substantially increased when the co-cultivation period was extended to 72 h for *Pp* ($p < 0.01$) and 96 h for *Pd* ($p < 0.01$) (Figs. 2A & 2B).

Next, we determined the influence of co-cultivation temperature on ATMT efficiency. The 1:10 ratio of fungal conidia and AGL-1 cells spread on membranes were incubated at various temperatures for 72 h, followed by transfer of these membranes to selection medium (Figs. 2C & 2D). The optimal temperature for the recovery of *Pd* Hyg⁺ transformants ranged from 15 to 18°C. Temperatures below 15°C or above 18°C were detrimental to the recovery of *Pd* transformants (Fig. 2C). In contrast, the optimal temperature for recovery of *Pp* Hyg⁺ transformants was 25°C, followed by 20°C. The recovery of *Pp* transformants at 15°C was comparable to that of *Pd* transformants, but temperatures below 15°C or above 25°C were detrimental to the recovery of *Pp* transformants (Fig. 2D).

Finally, based on ATMT optimizations, the co-cultivation duration of 72 h was used for both *Pd* and *Pp* and co-cultivation temperatures of 15°C for *Pd* and 25°C for *Pp* were used. A total of 620 *Pd* Hyg⁺ transformants and 1,800 *Pp* Hyg⁺ transformants were obtained from cultivation of approximately 1×10^7 cells in each conidial suspension, providing transformant yield of 0.006% and 0.018% for *Pd* and *Pp*, respectively.

Disruption of *PdURE1* and *PpURE1* genes

Of 128 putative Hyg⁺ *Pd* transformants analyzed by PCR with primer set (V2205-V2206) targeting *URE1* flanking *HYG* gene, four displayed a single amplicon of 3,000 bp in size, whereas the WT strain displayed a single amplicon of 800 bp in size, indicative of a homologous transformation of 3.1%. Results of the single integration event are shown for two of these mutants (Fig. 3A, 3Bi). All other transformants had two amplicons of 860 bp and 3,000 bp in size, indicative of ectopic integration (data not shown). The transformants that had single amplicon were further confirmed for the integration of *pPdure1::HYG* at the homologous site by two additional primer sets. The primer set (V2281-V2282) located within the *HYG* gene and on the right border outside of the disruption cassette, yielded a 1.8 kb PCR amplicon for mutants, but not for the WT strain (Fig. 3A, 3Bii). The primer set (V2203-V2204) located within the deleted region of *URE1* yielded a 318-

bp PCR amplicon for WT strain, but not for mutants (Fig. 3A, 3Biii). These mutants were passed through several generations of growth on YPD agar, which is a rich non-selective medium and thus, likely to facilitate the loss of any unstable genetic constructs. Southern hybridization of *Pst*I digested gDNA yielded single band of 3.9 kb for both the mutants, further confirming gene replacement at the homologous site without ectopic integration (Fig. 3A, 3C). These mutants were labeled as *Pdure1-10* and *Pdure1-21*. The rationale for selection of two *Pdure1* mutants was to test if both mutants had consistent phenotypes, as these mutants did not contain a functional *URE1*.

Of 36 *Pp* Hyg⁺ transformants analyzed by PCR with primer set (V2201-V2202) targeting *URE1* flanking *HYG* gene, six displayed a single amplicon of 2,694 bp in size, whereas the WT strain had a single amplicon of 957 bp in size, indicative of a homologous integration of 16.7%. Results of the single integration event are shown for two of these mutants (Fig. 4A, 4Bi). All other Hyg⁺ *Pp* transformants had two amplicons of 957 bp and 2,694 bp in size, indicative of ectopic integration (data not shown). Two of the six mutants were further confirmed for *URE1* deletion at the homologous site by additional primer sets as described for *Pd* (Fig. 4A, 4Bii-iii). The Southern analysis of gDNA digested with *Sac*II revealed a single band of expected size of 6.5. kb in both the mutant strains, further confirming gene replacement at the homologous site without ectopic integration (Fig. 4A, 4C). These mutants

were labeled as *Ppure1-4* and *Ppure1-11*, and the rationale for selection of two *Ppure1* mutants was same as described above for *Pdure1*.

The *Pdure1* and *Ppure1* mutants were assessed for urease enzyme activity along with their respective WT strains. It was interesting to note that the *Pd* WT and *Pdure1* mutant strains grew relatively better than the *Pp* WT and *Ppure1* mutant strains on Christensen's urea agar. Despite growth differences, no color development after 2 weeks of incubation, confirmed that the *URE1* gene was functionally deleted in both *Pdure1* and *Ppure1* mutants (Fig. 5A & Fig. 6A). These results were further corroborated by measurements of urease enzyme activity in cell lysates prepared from *Pdure1* or *Ppure1* mutants. Urease activity of the WT strains of *Pd* and *Pp* was 974 ± 113 U/g and $1,205 \pm 121$ U/g, respectively. In contrast, urease activity was undetectable in *Pdure1* and *Ppure1* mutants (Fig. 5B & Fig. 6B). Finally, we tested whether the *URE1* gene is involved in urea metabolism in these two organisms. *Ppure1* mutants failed to grow on YNB-U as compared to the *Pp* WT strain (Fig. 6C). In contrast, poor but detectable growth of *Pdure1* mutants was observed on YNB-U (Fig. 5C). To further analyze this discrepant result, we tested *Pd* WT and *Pdure1* mutant strains on YNB agar devoid of urea. Similar poor growth of WT and mutant strains observed indicating that the glucose in the YNB medium possibly supports minimal growth of *Pd* but not *Pp* strains (Supplementary Fig. S2). These results

further corroborated our findings regarding *Pdure1* mutants, as *Ppure1* mutants were unable to utilize urea, confirming that the *URE1* gene is involved in urea metabolism in both *Pd* and *Pp*.

Growth and sporulation

The *URE1* gene was found to be dispensable for the growth of *Pd* or *Pp*, as *Pdure1* mutants grew efficiently on YPD and PDA media between 6-15°C while *Ppure1* mutants grew as well between 6-25°C (Fig. 7). Similarly, *URE1* gene was not required for sporulation, as both *Pdure1* and *Ppure1* mutants produced spores as efficiently as their respective WT strains on PDA agar (Supplementary Fig. S3). These results indicated that *URE1* is dispensable for both cold adaptation and sporulation in *Pd* and *Pp*.

DISCUSSION

The highlight of this study was the successful use of the ATMT system for targeted gene disruption in cold-adapted fungi *Pseudogymnoascus destructans* (*Pd*) and *Pseudogymnoascus pannorum* (*Pp*). Among the parameters studied for ATMT, the incubation temperature seems to be the critical factor in the transformation of *Pd* and *Pp*. These results are not surprising as *Pd* is a psychrophilic fungus, which grows best between 15°C to 18°C (Chaturvedi *et al.*, 2010, Verant *et al.*, 2012). The temperature below

20°C has been shown to be detrimental to the expression of some *vir* (tumor forming) genes of *Agrobacterium* necessary for pilus production and conjugation for T-DNA transfer from donor to recipient cells (Fullner & Nester, 1996, Baron *et al.*, 2001 Sugui, 2005 #871). Apart from temperature, *Agrobacterium*-mediated transformation showed strong correlation with the duration of co-cultivation and the presence of AS. These results are in agreement with other fungal pathogens indicating that the pathways leading to T-DNA insertion are conserved across fungal species (Piers *et al.*, 1996, Gouka *et al.*, 1999, Abuodeh *et al.*, 2000, Yamada *et al.*, 2009).

We also found that homologous gene integration was lower in *Pd* (3.1%) than in *Pp* (16.7%). It is important to point out that the homologous arm of the *URE1* gene was chosen from the same strain used for transformation, as strict sequence identity is required for high efficiency homologous recombination in other fungal pathogens (Sudarshan *et al.*, 1999, Narasipura *et al.*, 2003, Narasipura *et al.*, 2005). The G+C content appears to be identical between *Pd* and *Pp* (Chibucos *et al.*, 2013), such that this does not appear to be a likely factor affecting low homologous gene integration observed in *Pd*.

The availability of a simple phenotypic test (color change in

Christensen's urea agar) was a consideration for choosing the *URE1* gene encoding urease for gene disruption studies in the present investigation. Others and we have demonstrated that *Pd* secretes a large amount of urease into the growth medium (Chaturvedi *et al.*, 2010, Raudabaugh & Miller, 2013). The role of urease in the fungal and bacterial virulence has been investigated using different vertebrate models (Jones *et al.*, 1990, Eaton *et al.*, 1991, Cox *et al.*, 2000, Mirbod-Donovan *et al.*, 2006). Currently, there are no laboratory models available for WNS or for testing the pathogenic potential of *Pdure1* and *Ppure1* mutant strains. Although, a recent study demonstrated that experimentally infected bats with *Pd* develop WNS (Lorch *et al.*, 2011), the set-up described is not amenable to fungal mutant comparisons in the laboratory. Therefore, the evaluation of the role of urease in the virulence of *Pd* and *Pp* awaits the availability of *in vivo* or *ex vivo* model.

ACKNOWLEDGEMENTS

This study was partly supported with funds from the National Science Foundation (16-0039-01). We are grateful to Dr. Seogchan Kang (University Park, PA, USA) for providing *Agrobacterium tumefaciens* AGL-1 strain. Dr. Rasmus JN Frandsen (Frederiksberg, Denmark) is thanked for the generous gift of plasmid pRF-HU2. We also thank Dr. Harry Taber for the editing of the manuscript.

REFERENCES

- Abuodeh RO, Orbach MJ, Mandel MA, Das A & Galgiani JN (2000) Genetic transformation of *Coccidioides immitis* facilitated by *Agrobacterium tumefaciens*. *The J Infect Dis* **181**: 2106-2110.
- Baron C, Domke N, Beinhofer M & Hapfelmeier S (2001) Elevated temperature differentially affects virulence, VirB protein accumulation, and T-pilus formation in different *Agrobacterium tumefaciens* and *Agrobacterium vitis* strains. *J Bacteriol* **183**: 6852-6861.
- Blehert DS (2012) Fungal disease and the developing story of bat white-nose syndrome. *PLoS Pathogen* **8**: e1002779.
- Blehert DS, Hicks AC, Behr M, *et al.* (2009) Bat white-nose syndrome: an emerging fungal pathogen? *Science* **323**: 227.
- Chaturvedi V, Springer DJ, Behr MJ, *et al.* (2010) Morphological and molecular characterizations of psychrophilic fungus *Geomyces destructans* from New York bats with White Nose Syndrome (WNS). *PLoS One* **5**: e10783.
- Chibucos MC, Crabtree J, Nagaraj S, Chaturvedi S & Chaturvedi V (2013) Draft Genome Sequences of Human Pathogenic Fungus *Geomyces pannorum* *Sensu Lato* and Bat White Nose Syndrome Pathogen *Geomyces (Pseudogymnoascus) destructans*. *Genome Announcement* **1**: e01045-01013.
- Cox GM, Mukherjee J, Cole GT, Casadevall A & Perfect JR (2000) Urease as a virulence factor in experimental cryptococcosis. *Infect Immun* **68**: 443-448.
- Dobinson KF, Grant SJ & Kang S (2004) Cloning and targeted disruption via *Agrobacterium tumefaciens*-mediated transformation of a trypsin protease gene from the vascular wilt fungus *Verticillium dothiae*. *Current genetics* **45**: 104-110.
- Dogan RI, Getoor L, Wilbur WJ & Mount SM (2007) SplicePort--an interactive splice-site analysis tool. *Nucleic Acids Res* **35**: W285-291.
- Eaton KA, Brooks CL, Morgan DR & Krakowka S (1991) Essential role of urease in pathogenesis of gastritis induced by *Helicobacter pylori* in gnotobiotic piglets. *Infect Immun* **59**: 2470-2475.
- Eckert M, Maguire K, Urban M, Foster S, Fitt B, Lucas J & Hammond-Kosack K (2005) *Agrobacterium tumefaciens*-mediated transformation of *Leptosphaeria* spp. and *Oculimacula* spp. with the reef coral gene DsRed and the jellyfish gene gfp. *FEMS Microbiol Lett* **253**: 67-74.
- Frandsen RJ, Andersson JA, Kristensen MB & Giese H (2008) Efficient four fragment cloning for the construction of vectors for targeted gene replacement in filamentous fungi. *BMC Mol Biol* **9**: 70.
- Frandsen RJN (2011) A guide to binary vectors and strategies for targeted genome modification in fungi using *Agrobacterium tumefaciens*-mediated transformation. *J Microbiol Methods* **87**: 247-262.

- Frick WF, Pollock JF, Hicks AC, Langwig KE, Reynolds DS, Turner GG, Butchkoski CM & Kunz TH (2010) An emerging disease causes regional population collapse of a common North American bat species. *Science* **329**: 679-682.
- Fullner KJ & Nester EW (1996) Temperature affects the T-DNA transfer machinery of *Agrobacterium tumefaciens*. *J Bacteriol* **178**: 1498-1504.
- Gouka RJ, Gerk C, Hooykaas PJ, Bundock P, Musters W, Verrips CT & de Groot MJ (1999) Transformation of *Aspergillus awamori* by *Agrobacterium tumefaciens*-mediated homologous recombination. *Nat Biotechnol* **17**: 598-601.
- Helber N & Requena N (2008) Expression of the fluorescence markers DsRed and GFP fused to a nuclear localization signal in the arbuscular mycorrhizal fungus *Glomus intraradices*. *The New phytologist* **177**: 537-548.
- Jones BD, Lockatell CV, Johnson DE, Warren JW & Mobley HL (1990) Construction of a urease-negative mutant of *Proteus mirabilis*: analysis of virulence in a mouse model of ascending urinary tract infection. *Infect Immun* **58**: 1120-1123.
- Katoh K, Asimenos G & Toh H (2009) Multiple alignment of DNA sequences with MAFFT. *Methods Mol Biol* **537**: 39-64.
- Lorch JM, Meteyer CU, Behr MJ, et al. (2011) Experimental infection of bats with *Geomyces destructans* causes white-nose syndrome. *Nature* **480**: 376-378.
- McClelland CM, Chang YC & Kwon-Chung KJ (2005) High frequency transformation of *Cryptococcus neoformans* and *Cryptococcus gattii* by *Agrobacterium tumefaciens*. *Fungal Genet Biol* **42**: 904-913.
- Michielse CB, Hooykaas PJ, van den Hondel CA & Ram AF (2005) *Agrobacterium*-mediated transformation as a tool for functional genomics in fungi. *Curr Genet* **48**: 1-17.
- Minnis AM & Lindner DL (2013) Phylogenetic evaluation of *Geomyces* and allies reveals no close relatives of *Pseudogymnoascus destructans*, comb. nov., in bat hibernacula of eastern North America. *Fungal Biol* **117**: 638-649.
- Mirbod-Donovan F, Schaller R, Hung CY, Xue J, Reichard U & Cole GT (2006) Urease produced by *Coccidioides posadasii* contributes to the virulence of this respiratory pathogen. *Infect Immun* **74**: 504-515.
- Möller EM, Bahnweg G, Sandermann H & Geiger HH (1992) A simple and efficient protocol for isolation of high molecular weight DNA from filamentous fungi, fruit bodies, and infected plant tissues. *Nucleic Acids Res* **20**: 6115-6116.
- Mozo T & Hooykaas PJ (1991) Electroporation of mega plasmids into *Agrobacterium*. *Plant molecular biology* **16**: 917-918.
- Mullins ED, Chen X, Romaine CP, Raina R, Geiser DM & Kang S (2001) *Agrobacterium*-mediated transformation of *Fusarium oxysporum*: an

efficient tool for insertional mutagenesis and gene transfer. *Phytopathology* **91**: 173-180.

Narasipura SD, Chaturvedi V & Chaturvedi S (2005) Characterization of *Cryptococcus neoformans* var. *gattii* *SOD2* reveals distinct roles of the two superoxide dismutases in fungal biology and virulence. *Molecular microbiology* **55**: 1782-1800.

Narasipura SD, Ault JG, Behr MJ, Chaturvedi V & Chaturvedi S (2003) Characterization of Cu,Zn superoxide dismutase (*SOD1*) gene knockout mutant of *Cryptococcus neoformans* var. *gattii*:role in biology and virulence. *Molecular microbiology* **47**: 1681-1694.

Olszewski MA, Noverr MC, Chen GH, Toews GE, Cox GM, Perfect J & Huffnagle GB (2004) Urease expression by *Cryptococcus neoformans* promotes microvascular sequestration, thereby enhancing central nervous system invasion. *Am J Pathol* **164**: 1761-1771.

Piers KL, Heath JD, Liang X, Stephens KM & Nester EW (1996) *Agrobacterium tumefaciens*-mediated transformation of yeast. *Proc Natl Acad Sci U S A* **93**: 1613-1618.

Rajkumar SS, Rudd RJ, Okoniewski JC, Xu J, Chaturvedi S & Chaturvedi V (2011) Dispersal of a single, clonal genotype of *Geomyces destructans* among New York bats with geomycosis (White Nose Syndrome). *Emerg Infect Dis* **17**: 1273-1276.

Raudabaugh DB & Miller AN (2013) Nutritional Capability of and Substrate Suitability for *Pseudogymnoascus destructans*, the Causal Agent of Bat White-Nose Syndrome *Plos One* **8**: e78300.

Ren P, Haman KH, Last LA, Rajkumar SS, Keel MK & Chaturvedi V (2012) Clonal Spread of *Geomyces destructans* among Bats, Midwestern and Southern United States. *Emerging infectious diseases* **18**: 883-885.

Reynolds HT & Barton HA (2014) Comparison of the white-nose syndrome agent *Pseudogymnoascus destructans* to cave-dwelling relatives suggests reduced saprotrophic enzyme activity. *PLoS One* **9**: e86437.

Saitou N & Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**: 406-425.

Stanke M & Morgenstern B (2005) AUGUSTUS: a web server for gene prediction in eukaryotes that allows user-defined constraints. *Nucleic Acids Res* **33**: W465-467.

Stanke M, Steinkamp R, Waack S & Morgenstern B (2004) AUGUSTUS: a web server for gene finding in eukaryotes. *Nucleic Acids Res* **32**: W309-312.

Sudarshan S, Davidson RC, Heitman J & Alspaugh JA (1999) Molecular analysis of the *Cryptococcus neoformans* *ADE2* gene, a selectable marker for transformation and gene disruption. *Fungal Genet Biol* **27**: 36-48.

Tamura K, Dudley J, Nei M & Kumar S (2007) MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Mol Biol Evol* **24**: 1596-1599.

- 597 Tsuda M, Karita M, Morshed MG, Okita K & Nakazawa T (1994) A urease-
 598 negative mutant of *Helicobacter pylori* constructed by allelic exchange
 599 mutagenesis lacks the ability to colonize the nude mouse stomach. *Infect*
 600 *Immun* **62**: 3586-3589.
- 601 Verant ML, Boyles JG, Waldrep WJ, Wibbelt G & Blehert DS (2012)
 602 Temperature-dependent growth of *Geomyces destructans*, the fungus
 603 that causes bat white-nose syndrome. *PLoS One* **7**: e46280.
- 604 Yamada T, Makimura K, Satoh K, Umeda Y, Ishihara Y & Abe S (2009)
 605 *Agrobacterium tumefaciens*-mediated transformation of the
 606 dermatophyte, *Trichophyton mentagrophytes*: an efficient tool for gene
 607 transfer. *Med Mycol* **47**: 485-494.
- 608 Zhang T, Zhuo Y, Jia X, Liu J, Gao H, Song F, Liu M & Zhang L (2013)
 609 Cloning and characterization of the gene cluster required for beauvericin
 610 biosynthesis in *Fusarium proliferatum*. *Sci China Life Sci* **56**: 628-637.

Table 1 Strains, plasmids and oligonucleotides used in this study.

Strains	Description	Source
<i>E. coli</i> Top10	<i>lacx74 recA1 deoR F – mcrA Δ (mrr-hsdRMS-mcrBC) φ80 lacZΔM15Δ araD139Δ (ara-leu)7697 galU galK</i>	Invitrogen
<i>Agrobacterium tumefaciens</i> (AGL-1)	Genotype - AGL0 <i>recA::bla</i> pTiBo542deltaT Mop+ CbR	Dr. Seogchan Kang, University Park, PA, USA
<i>Pseudogymnoascus destructans</i> (Pd) M1379	Wild type strain	Bat skin (Chibucos <i>et al.</i> , 2013)
<i>Pseudogymnoascus pannorum</i> (Pp) M1372	Wild type strain	Soil (UAMH1062) (Chibucos <i>et al.</i> , 2013)
<i>Pdure1-10</i>	<i>ure1</i> mutant of <i>Pd</i>	This study
<i>Pdure1-25</i>	<i>ure1</i> mutant of <i>Pd</i>	This study
<i>Ppure1-4</i>	<i>ure1</i> mutant of <i>Pp</i>	This study
<i>Ppure1-11</i>	<i>ure1</i> mutant of <i>Pp</i>	This study
Plasmid	Description	Source
pRF-HU2	Vector for targeted gene replacement containing <i>HYG</i> gene as selection marker	(Frandsen <i>et al.</i> , 2008)
<i>pPdure1::HYG</i>	5' & 3' prime arms of <i>PdURE1</i> gene cloned into pRF-HU2	This study
<i>pPpure1::HYG</i>	5' & 3' prime arms of <i>PpURE1</i> gene cloned into pRF-HU2	This study
Oligonucleotides	Sequence (5'-3')	Purpose
V2149a	<u>GGTCTTAA</u> UTTTAGGCGGAGCTATGAC	<i>pPdure1::HYG</i> vector construction
V2150a	<u>GGCATTAAUA</u> AAGGGCGTTGATGATGAC	<i>pPdure1::HYG</i> vector construction
V2151a	<u>GGACTTAAU</u> TATGTTGCAGGCTTTGGA	<i>pPdure1::HYG</i> vector construction
V2152a	<u>GGGTTTAAU</u> GATCTGGCATCATATCGTC	<i>pPdure1::HYG</i> vector construction
V2171	AGCTGCGCCGATGGTTTCTACAA	Hyg ⁺ transformant screening and <i>HYG</i> probe for Southern analysis of <i>Ppure1</i> mutants
V2172	GCGCGTCTGCTGCTCCATACAA	Hyg ⁺ transformant screening and <i>HYG</i> probe

		for Southern analysis of <i>Ppure1</i> mutants
V2173	<u>GGGTTTAA</u> UTGCCACCGCCAATGTATC	<i>pPpure1::HYG</i> vector construction
V2174	<u>GGACTTAA</u> UTCATCATCAACGCCCTTA	<i>pPpure1::HYG</i> vector construction
V2175	<u>GGCATTAA</u> UACGGTGACGAGGTAGGTT	<i>pPpure1::HYG</i> vector construction
V2176	<u>GGTCTTAA</u> UGACGAGGCGGTTGAGGAA	<i>pPpure1::HYG</i> vector construction
V2199	CCATACCGTCACGCAGAG	<i>Ppure1</i> mutant screening
V2200	AATAAAGGGCGGAAGAGG	<i>Ppure1</i> mutant screening
V2201	TAAATCCGTTCTTGATGCCGATATC	<i>Ppure1</i> mutant screening
V2202	AATGTTGCGGATCTCATG	<i>Ppure1</i> mutant screening
V2203	ATTGTTGATTGGAGTGGTATT	<i>Ppure1</i> mutant screening
V2204	CCGCATGTTGTTCTTGCC	<i>Ppure1</i> mutant screening
V2205	CATCATCAACGCCCTTAT	<i>Ppure1</i> mutant screening
V2206	AGCTCCCATGTCGTGTAA	<i>Ppure1</i> mutant screening
V2281	GACCAGTTGCCTAAATGAA	<i>Ppure1</i> mutant screening
V2282	GTAAGTATGGGAAGCAGAAA	<i>Ppure1</i> mutant screening
V2285	TTTCGGCGTGCGGTATGGT	<i>Ppure1</i> mutant screening
V2286	TCGGATTCAAGCGACTGG	<i>Ppure1</i> mutant screening
V2370	TTCAGTAACGTTAAGTGGATCC	<i>HYG</i> probe for Southern analysis of <i>Ppure1</i> mutants
V2372	CGCAAGGAATCGGTCAATAC	<i>HYG</i> probe for Southern analysis of <i>Ppure1</i> mutants

613 The underlined sequences denote 9-bp long 2-deoxyuridine containing
614 overhangs for directional cloning into pRF-HU2 vector (Frandsen et al.,
615 2008).

616

FIGURE LEGENDS

Fig. 1 Multiple alignment and phylogenetic relationship of deduced amino acid sequences of *URE1* of *Pd* and *Pp* with other fungal *URE1* genes

A) Portions of the amino acid sequences of URE1p were used for multiple alignments, and the sequences corresponding to amino acids 399 to 412 and 513 to 525 are numbered based on *Pp*URE1p. The conserved histidine residues important in binding to the nickel metallocenter (amino acid 406), binding to the substrate (amino acid 408), and catalysis (amino acid 518) are in boldface and marked with a star.

B) The full-length amino acid sequences of URE1p were used to construct a phylogenetic tree by the neighbor-joining method. The bootstrap scores are based on 1,000 reiterations. The URE1p from *Penicillium chrysogenum* is used as an outgroup.

Fig. 2 Co-cultivation parameters for transformation efficiency of *Pd* and *Pp*

A-B) Effect of co-cultivation duration. *A. tumefaciens* AGL-1 cells grown in IM medium containing AS were mixed with *Pd* or *Pp* conidial suspension at

10:1 ratio and of these suspension, 200 μ l was inoculated per plate followed by incubation at 15°C for *Pd* (A) and at 25°C for *Pp* (B) for various durations prior to hygromycin selection. Data presented as the average of three plates per treatment.

C-D) Effect of co-cultivation temperatures. *A. tumefaciens* AGL-1 cells grown in IM medium containing AS were mixed with *Pd* or *Pp* conidial suspension at 10:1 ratio and of these 200 μ l was inoculated per plate followed by incubation at various temperatures for 72 h prior to hygromycin selection.

Fig. 3 Disruption of *Pd URE1* by ATMT

A) Restriction map of the *Pdure1*-disruption cassette. 5' *URE1* (1.4 kb) and 3' *URE1* (1.5 kb) were cloned into pRF-HU2 flanking hygromycin B resistance (*HYG*) gene under the control of *Aspergillus nidulans* promoter (PTrpC) and terminator (TTrpC). The arrows represent primers used for PCR and solid grey line represent *HYG* probe used for Southern analysis. The dashed line (thick) represents genomic sequences outside of the disruption cassette. The location of restriction enzyme *Pst*I within the *HYG* gene of the disruption cassette and at the 5' end of the *URE1* locus, outside of the disruption cassette is shown. The dashed line (thin) represents expected size product for mutant strains following enzyme digestion and Southern analysis.

B) Analysis of *ure1* transformants by PCR. Primer pair (V2205-V2206) targeting *URE1* flanking *HYG* gene (Bi-3.0 kb in size for mutants and 860 bp in size for WT), primer pair (V2281-V2282) located in the *HYG* gene and right border of the *URE1* gene (Bii-1.8 kb in size for mutants and no PCR amplicon for WT), and primer pair (V2203-V2204) located within the deletion region of *URE1* gene (Biii-318 bp in size for WT and no PCR amplicon for mutants).

C) Southern hybridization analysis. Genomic DNA from the WT and the mutant strains was digested with *Pst*I and hybridized with *HYG* probe (3.9 kb band for both the mutants and no band for the WT).

Fig. 4 Disruption of *PpURE1* by ATMT

A) Restriction map of the *Ppure1*-disruption cassette. 5' *Ure1* (1.5 kb) and 3' *Ure1* (1.5 kb) were cloned into pRF-HU2 flanking hygromycin B resistance (*HYG*) gene under the control of *Aspergillus nidulans* promoter (PTrpC) and terminator (TTrpC). The arrows represent primers used for PCR and solid grey line represents *HYG* probe used for Southern analysis. The dashed line represents genomic sequences outside of the disruption cassette. The location of restriction enzyme *Sac*II within the *HYG* gene and at the 5' end of the *URE1* locus, outside of the disruption cassette is shown.

681 B) Analysis of *ure1* transformants by PCR. Primer pair (V2201-V2202)
682 targeting *URE1* flanking *HYG* gene (Bi-957 bp in size for WT and 2,694 bp in
683 size for mutant strains), primer pairs (V2285-V2286) located in the *HYG*
684 gene and in the right border of *URE1* gene (Bii-1.6 kb in size for mutants
685 and no PCR amplicon for WT), and primer pair (V2199-V220) within the
686 deletion region of *URE1* gene (Biii-466 bp in size for WT and no PCR
687 amplicon for mutants).

688 C) Southern hybridization analysis. Genomic DNA from the WT and the
689 mutant strains was digested with *SacII* and hybridized with *HYG* probe (6.4
690 kb band for both the mutants and no band for the WT).

691

692 **Fig. 5 Characterization of *Pdure1* mutants**

693

694 A) Approximately 10 μ l aliquot of *Pd* conidial suspensions (10^7 /ml) from WT
695 and mutant strains were placed on Christensen's urea agar and incubated for
696 2 weeks at 15°C. Absence of the magenta color with mutant strains indicated
697 loss of urease enzyme activity. Tube without culture served as quality
698 control for medium.

699 B) The urease enzyme activity from cell lysates was determined according to
700 the Urease Assay Kit (BioAssay Systems). The mutant strains did not
701 produce urease enzyme while WT did.

C) Approximately 10 µl aliquot of the dilution series of conidial suspensions of the WT and *Pdure1* mutant strains were spotted on two separate YNB-U agar plates to avoid any residual effect of the growth of WT strain on the mutant strains. Plates were incubated at 15°C for 7-days. A significant difference of growth between mutant and WT strains confirmed that *URE1* is required when nitrogen is the sole source of nutrient.

Fig. 6 Characterization of *Ppure1* mutants

A) Approximately 10 µl aliquots of *Pp* conidial suspensions from the WT and mutant strains were placed on Christensen's urea agar and incubated for 2 weeks at 15°C. Absence of the magenta color with mutant strains indicated loss of urease enzyme activity. Tube without culture served as quality control for medium

B) The urease enzyme activity from cell lysates was determined according to Urease Assay Kit (BioAssay Systems). The *Ppure1* mutants did not produce urease enzyme while WT did.

C) Approximately 10 µl aliquots of dilution series of the conidial suspensions of the WT and mutant strains were spotted on two separate YNB-U agar plates to avoid any residual effect of the growth of WT strain on the mutant strains. Plates were incubated at 15°C for a 7-day period. No growth of

mutants indicated that the *URE1* is required when nitrogen is the sole source of nutrient.

Fig. 7 Growth of *Pdure1* and *Ppure1* mutants at different temperatures

A. Approximately 10 µl aliquots of the dilution series of the conidial suspension of WT and *Pdure1* mutant strains were spotted on YPD and PDA agar. Plates were incubated at 6°C for 14-day, and at 15°C and 25°C for 7-day periods. No growth of the WT and mutant strains observed at 25°C.

B. Approximately 10 µl aliquots of the dilution series of the conidial suspension of *Pp* WT and *Ppure1* mutant strains were spotted on YPD and PDA agar. Plates were incubated at 6°C for 14-day, and at 15°C and 25°C for 7-day periods. Excellent growth of both *Pp* WT and *Ppure1* mutant strains was observed at all temperatures tested

Table 1 Strains, plasmids and oligonucleotides used in this study.

Strains	Description	Source
<i>E. coli</i> Top10	<i>lacx74 recA1 deoR F – mcrA Δ (mrr-hsdRMS-mcrBC) ϕ80 lacZΔM15Δ araD139Δ (ara-leu)7697 galU galK</i>	Invitrogen
<i>Agrobacterium tumefaciens</i> (AGL-1)	Genotype - AGL0 <i>recA::bla</i> pTiBo542deltaT Mop+ CbR	Dr. Seogchan Kang, University Park, PA, USA
<i>Pseudogymnoascus destructans</i> (Pd) M1379	Wild type strain	Bat skin (Chibucos <i>et al.</i> , 2013)
<i>Pseudogymnoascus pannorum</i> (Pp) M1372	Wild type strain	Soil (UAMH1062) (Chibucos <i>et al.</i> , 2013)
<i>Pdure1-10</i>	<i>ure1</i> mutant of <i>Pd</i>	This study
<i>Pdure1-25</i>	<i>ure1</i> mutant of <i>Pd</i>	This study
<i>Ppure1-4</i>	<i>ure1</i> mutant of <i>Pp</i>	This study
<i>Ppure1-11</i>	<i>ure1</i> mutant of <i>Pp</i>	This study
Plasmid	Description	Source
pRF-HU2	Vector for targeted gene replacement containing <i>HYG</i> gene as selection marker	(Frandsen <i>et al.</i> , 2008)
<i>pPdure1::HYG</i>	5' & 3' prime arms of <i>PdURE1</i> gene cloned into pRF-HU2	This study
<i>pPpure1::HYG</i>	5' & 3' prime arms of <i>PpURE1</i> gene cloned into pRF-HU2	This study
Oligonucleotides	Sequence (5'-3')	Purpose
V2149a	<u>GGTCTTAAU</u> TTTAGGCGGAGCTATGAC	<i>pPdure1::HYG</i> vector construction
V2150a	<u>GGCATTAAU</u> AAGGGCGTTGATGATGAC	<i>pPpure1::HYG</i> vector

		<i>construction</i>
V2151a	<u>GGACTTAA</u> TATGTTGCAGGCTTTGGA	<i>pPdure1::HYG</i> vector construction
V2152a	<u>GGGTTTAA</u> UGATCTGGCATCATATCGTC	<i>pPdure1::HYG</i> vector construction
V2171	AGCTGCGCCGATGGTTTCTACAA	Hyg ⁺ transformant screening and <i>HYG</i> probe for Southern analysis of <i>Ppure1</i> mutants
V2172	GCGCGTCTGCTGCTCCATACAA	Hyg ⁺ transformant screening and <i>HYG</i> probe for Southern analysis of <i>Ppure1</i> mutants
V2173	<u>GGGTTTAA</u> UTGCCACCGCCAATGTATC	<i>pPpure1::HYG</i> vector construction
V2174	<u>GGACTTAA</u> UTCATCATCAACGCCCTTA	<i>pPpure1::HYG</i> vector construction
V2175	<u>GGCATTAA</u> UACGGTGACGAGGTAGGTT	<i>pPpure1::HYG</i> vector construction
V2176	<u>GGTCTTAA</u> UGACGAGGCGGTTGAGGAA	<i>pPpure1::HYG</i> vector construction
V2199	CCATACCGTCACGCAGAG	<i>PpURE1</i> mutant screening
V2200	AATAAAGGGCGGAAGAGG	<i>PpURE1</i> mutant screening
V2201	TAAATCCGTTCTTGATGCCGATATC	<i>Ppure1</i> mutant screening
V2202	AATGTTGCGGATCTCATG	<i>Ppure1</i> mutant screening
V2203	ATTGTTGATTGGAGTGGTATT	<i>Pdure1</i> mutant screening
V2204	CCGCATGTTGTTCTTGCC	<i>Pdure1</i> mutant screening
V2205	CATCATCAACGCCCTTAT	<i>Pdure1</i> mutant screening
V2206	AGCTCCCATGTCGTGTAA	<i>Pdure1</i> mutant screening

V2281	GACCAGTTGCCTAAATGAA	<i>Pdure1</i> mutant screening
V2282	GTAAGTATGGGAAGCAGAAA	<i>Pdure1</i> mutant screening
V2285	TTTCGGCGTGGGTATGGT	<i>Ppure1</i> mutant screening
V2286	TCGGATTCAAGCGACTGG	<i>Ppure1</i> mutant screening
V2370	TTCAGTAACGTTAAGTGGATCC	<i>HYG</i> probe for Southern analysis of <i>Pdure1</i> mutants
V2372	CGCAAGGAATCGGTCAATAC	<i>HYG</i> probe for Southern analysis of <i>Pdure1</i> mutants

The underlined sequences denote 9-bp long 2-deoxyuridine containing overhangs for directional cloning into pRF-HU2 vector (Frandsen et al.,

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7