RNA-Sequencing of *Cercospora beticola* DMI-sensitive and -resistant isolates after treatment with tetraconazole identifies common and contrasting pathway induction

Melvin D. Bolton ^{a,b}, Malaika K. Ebert ^{a,b,c,1}, Luigi Faino ^{c,1}, Viviana Rivera-Varas ^b, Ronnie de Jonge ^{d,e}, Yves Van de Peer ^{d,e}, Bart P. H. J. Thomma ^c, and Gary A. Secor ^b

^a *USDA – ARS, Northern Crop Science Laboratory, Fargo, ND, USA*

^b *North Dakota State University, Department of Plant Pathology, Fargo, ND, USA*

^c *Wageningen University, Laboratory of Phytopathology, Wageningen, The Netherlands*

^d *Department of Plant Systems Biology, VIB, Ghent, Belgium*

^e *Department of Plant Biotechnology and Bioinformatics, Ghent University, Ghent Belgium*

Correspondence: Email: Melvin.Bolton@ars.usda.gov

¹ The contribution of these authors should be considered equal.

ABSTRACT

Cercospora beticola causes Cercospora leaf spot of sugar beet. Cercospora leaf spot management measures often include application of the sterol demethylation inhibitor (DMI) class of fungicides. The reliance on DMIs and the consequent selection pressures imposed by their widespread use has led to the emergence of resistance in *C. beticola* populations. Insight into the molecular basis of tetraconazole resistance may lead to molecular tools to identify DMI-resistant strains for fungicide resistance management programs. Previous work has shown that expression of the gene encoding the DMI target enzyme (CYP51) is generally higher and inducible in DMI-resistant *C. beticola* field strains. In this study, we extended the molecular basis of DMI resistance in this pathosystem by profiling the transcriptional response of two *C. beticola* strains contrasting for resistance to tetraconazole. A majority of the genes in the ergosterol biosynthesis pathway were induced to similar levels in both strains with the exception of *CbCyp51*, which was induced several-fold higher in the DMI-resistant strain. In contrast, a secondary metabolite gene cluster was induced in the resistance strain, but repressed in the sensitive strain. Genes encoding proteins with various cell membrane fortification processes were induced in the resistance strain. Site-directed and ectopic mutants of candidate DMI-resistance genes all resulted in significantly higher EC₅₀ values than the wild-type strain, suggesting that the cell wall and/or membrane modified as a result of the transformation process increased resistance to tetraconazole. Taken together, this study identifies important cell membrane components and provides insight into the molecular events underlying DMI resistance in *C. beticola*.

Keywords: ergosterol, sterol demethylation inhibitor, RTA1, cell-membrane, transformation, EC₅₀, tetraconazole

1. INTRODUCTION

Cercospora beticola causes Cercospora leaf spot (CLS) of sugar beet (*Beta vulgaris*). The fungus is considered to be the most destructive foliar pathogen of sugar beet worldwide. CLS is of significant concern to the \$3 billion United States sugar beet industry, which is the primary provider of refined sucrose used for natural sweetening domestically and has a growing interest to use sugar beet for biofuel production (Panella, 2010). Yield and quality losses from CLS together with costs associated with fungicide application result in annual losses in excess of several million dollars in the United States alone. The disease causes a reduction in root harvest weight and extractable sucrose due to loss of photosynthetic capacity and diversion of photosynthate from roots for foliage regeneration (Franc, 2010). Individual leaf spots are most commonly found on older leaves initially, but all leaves in advanced stages of disease may have numerous 3-5 mm circular leaf spot lesions harboring fungal stroma (Franc, 2010). Infection is initiated when sporulation from over-wintered stomata results in conidia being delivered to sugar beet leaf surfaces primarily by wind and water splash (Khan et al., 2008). After spore germination, the non-symptomatic biotrophic stage of disease development occurs when the fungus penetrates stomata to reach the apoplast where effectors released by invading hyphae facilitate infection (Daub and Ehrenshaft, 2000; Steinkamp et al., 1979). As early as five days after initial infection, the necrotrophic stage ensues and conidia produced in the resulting necrotic tissue can initiate another disease cycle provided environmental conditions are conducive (Franc, 2010). Although no sexual stage of *C. beticola* has been identified to our knowledge, population studies have shown that *C. beticola* populations carry the hallmarks of sexual reproduction (Bolton et al., 2014; Bolton et al., 2012c). Besides sugar beet, several hosts for *C. beticola* have been described (Lartey et al., 2005). Although the biological importance of these hosts is not clear, we cannot exclude the possibility that a sexual cycle occurs on hosts besides sugarbeet. Disease management measures for CLS include the use

of tolerant sugar beet varieties and crop rotation, but the disease is managed effectively only when combined with timely fungicide applications (Khan et al., 2007; Secor et al., 2010).

A CLS outbreak occurred in 1998 that led to estimated losses of \$45 million in north central U.S. due to reduced yield and fungicide application costs (Secor et al., 2010). The subsequent year, the U.S. Environmental Protection Agency granted an emergency exemption for growers to use the non-registered triazole fungicide tetraconazole, the first use of an azole to manage CLS, to help contain another CLS outbreak. Since then, tetraconazole and other triazoles have been popular because of protective and curative properties, and low levels of phytotoxicity. Reliance on triazole chemistries for CLS management has become even more pronounced with recent development of resistance to other chemistries including benzimidazole (Weiland and Halloin, 2001) and quinone outside inhibitors (Birla et al., 2012; Bolton et al., 2013a; Bolton et al., 2013b). The reliance on triazoles and the consequent selection pressures imposed by their widespread use has led to a shift in sensitivity in *C. beticola* populations (Bolton et al., 2012b; Secor et al., 2010).

Triazoles are the largest group of fungicides of the sterol demethylation inhibitors (DMIs) and act through inhibition of cytochrome P450-dependent sterol 14 α -demethylase (CYP51), an enzyme required for biosynthesis of the fungal cell membrane component ergosterol (Cools et al., 2013). Resistance to DMIs is known to occur through a variety of mechanisms including amino acid exchanges in CYP51 (Carter et al., 2014; Cools et al., 2010; Leroux et al., 2007), elevated expression of *Cyp51* (Cools et al., 2012; Ma and Tredway, 2013), and/or active efflux of fungicide from the cell due to increased expression of genes encoding membrane ATP-binding cassette (ABC) and/or major facilitator superfamily transporters (Kretschmer et al., 2009; Zwiers et al., 2002). Recently, melanization has also been shown to play a role in DMI resistance (Lendenmann et al., 2015). DMI resistance in *C. beticola* is associated with over-expression of *CbCyp51* (Bolton et al., 2012a). Although mutations in *Cyp51*

regulatory elements have been shown to drive *Cyp51* over-expression in other pathosystems (Carter et al., 2014; Ma et al., 2006), no similar promoter mutation could be identified in *C. beticola* (Bolton et al., 2012a). Since a mutation in *Cyp51* regulatory elements is generally associated with constitutive over-expression (Cools et al., 2012; Hamamoto et al., 2000; Ma et al., 2006; Sun et al., 2013), the lack of a similar mutation in *CbCyp51* may explain why *CbCyp51* is inducible in DMI-resistant strains with expression levels that correlate with tetraconazole rates used in *in vitro* assays (Bolton et al., 2012a). The mechanism responsible for *CbCyp51* inducible expression is not known.

In this study, we used RNA-seq technology to profile the transcriptional response of a *C. beticola* DMI-resistant and -sensitive strains to tetraconazole and acetone (control) treatments. Pathway analysis showed that flux through the ergosterol biosynthesis pathway is a response shared by both strains after exposure to tetraconazole. A majority of the genes in the ergosterol biosynthesis pathway were induced to similar levels in both strains with the exception of *CbCyp51*, which was induced several-fold higher in the DMI-resistant strain. Surprisingly, site-directed and ectopic mutants of candidate DMI-resistance genes all resulted in significantly higher EC₅₀ values than the wild-type strain, suggesting that the transformation process increased resistance to tetraconazole. The implications of these genes for DMI resistance in the *C. beticola* pathosystem are discussed.

2. Materials and methods

2.1. Fungal strains and culture conditions

Sugar beet leaves with *Cercospora* leaf spot were collected from commercial fields of the Red River Valley region of MN and ND. Conidia were harvested by adding 50 μ L of T-water (0.06% (v/v) Tween 20 (Sigma-Aldrich, St. Louis, USA), 0.02% (w/v) filter-sterilized ampicillin added after the solution had been autoclaved) to the surface of a *C. beticola* leaf spot lesion. The lesion was gently scraped with a pipette tip to liberate conidia into the T-water. The conidia-laden T-water was collected and transferred to water agar plates (1.5% (w/v) agar (BD, Franklin Lakes, USA), 0.02% (w/v) filter-sterilized ampicillin added after the solution had been autoclaved) for 24 h at 22 °C ($\pm$ 1 °C) to initiate conidia germination. One germinated conidium was transferred to a clarified V8-medium plate (10% (v/v) clarified V8 juice (Campbell's Soup Co., Camden, USA), 0.5% (w/v) CaCO₃, and 1.5% (w/v) agar (BD)) and was incubated at 25 °C for two weeks. This single spore-derived colony was the source inoculum plate for subsequent fungicide sensitivity assays.

EC₅₀ calculations were carried out as described by Secor and Rivera (2012). Briefly, agar plugs (5 mm each) were taken from the leading edge of colony growth of the original non-amended V8-agar plates described above and placed on V8-agar plates amended with serial ten-fold dilutions of active ingredient from 0.001 to 1.0 μ g ml⁻¹. After 15 days, inhibition of radial growth was measured and compared to growth on non-amended medium and was used to calculate an EC₅₀ value for each isolate. *Cercospora beticola* strains 09-32 (DMI-sensitive) and 10-73-4 (DMI-resistant) were used in this study based on EC₅₀ value determination.

Liquid cultures of 09-32 and 10-73-4 were initiated by placing an agar plug taken from the leading edge of the clarified V8-medium plates (described above) into a 250 mL flask containing 50 mL potato dextrose broth (BD). Each isolate was allowed to grow for 96 h after which 5 μ L of tetraconazole stock

solution ($10 \mu\text{g mL}^{-1}$ in acetone) was added to each flask for a final concentration of $1.0 \mu\text{g mL}^{-1}$ as described by Bolton et al. (2012a). As a control, each isolate was also grown as described above except $5 \mu\text{L}$ acetone was added in place of tetraconazole. Flasks were shaken (125 rpm) for 48 h at 22°C ($\pm 1^\circ\text{C}$) after which mycelium was harvested using a Büchner funnel and immediately frozen in liquid nitrogen. Total RNA was isolated from mycelium using the RNeasy Plant Mini kit (Qiagen; Venlo; Netherlands) following the manufacturer's instructions that included an on-column DNase treatment (Qiagen). There was two flasks of each fungal strain/treatment for each of two biological replicates.

2.2. Genome assembly and gene prediction

Two sequence libraries (500-bp and 5-kb insert size) of *Cercospora beticola* strain 10-73-4 were prepared and sequenced at the BGI Americas (Cambridge, USA) using the Illumina high throughput-sequencing platform. Genome assembly was performed using A5 pipeline (default settings) (Tritt et al., 2012). Gene prediction was performed by aligning sequences from the predicted transcriptome of the DMI-sensitive *C. beticola* strain 09-40 (de Jonge and Bolton, *unpublished data*) to the genome of *C. beticola* 10-73-4 by GMAP software (settings: -fulllength ; -format=gff3_gene) (Wu and Nacu, 2010).

2.3. Expression analysis

Two biological replicates for each treatment were used for RNA-seq analysis. RNA-seq libraries (180 bp insert size) were prepared and sequenced (100- bp reads) by BGI Americas. RNA-seq data were aligned to the genome of *C. beticola* strain 10-73-4 using STAR (v2.5) software (Dobin et al., 2013) using the following options: --alignIntronMax 1000 --alignMatesGapMax 1000 --outFilterMismatchNmax 15 --outSAMstrandField intronMotif --outFilterIntronMotifs RemoveNoncanonical --outSAMattrIHstart 1. Cuffdiff, within the Cufflinks (v2.2.1) software package (Roberts et al., 2011), was used to estimate transcript level changes of the replicate experiments. The dataset for this project can be obtained at <http://www.ncbi.nlm.nih.gov/bioproject/294383>.

2.4. Transformation and disruption of target genes

Split-marker PCR constructs for targeted gene replacement were prepared as described by Catlett et al. (2003) using genomic DNA of 10-73-4 wild type *C. beticola* and pDAN (Friesen et al., 2006) as PCR templates. Primers for split-marker development are listed in Table 4.

A 5 mm plug was taken from the actively growing zone of *C. beticola* wild type strain 10-73-4 on potato dextrose agar (PDA; BD). Liquid cultures were initiated by grinding the plug with 500 μ L Fries media (Friesen and Faris, 2012), which was subsequently transferred to a 125 mL flask containing 50 mL Fries media. Flasks were wrapped in aluminum foil and shaken at 150 rpm at 21 °C for four days. Cultures were then transferred to a sterile blender cup, ground for 10 s, and transferred to a 500 mL flask containing 200 mL Fries media. Cultures were grown as described above for an additional 24 hours after which mycelium was harvested with two layers of Miracloth (Calbiochem, Billerica, USA) using a Büchner funnel. The mycelium was rinsed with mycelium wash solution (0.7M KCl, 10mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$), broken into small pieces with a sterile spatula, transferred to deep petri dish containing 40 mL of osmoticum (20 g L⁻¹ lysing enzymes from *T. harzianum* (Sigma-Aldrich) and 12.5 g L⁻¹ Driselase (Sigma-Aldrich), and incubated at 30 °C for 6 to 8 h while shaking at 50 rpm. Protoplasts were harvested by filtering the osmoticum solution through two layers of Miracloth and collected by centrifugation (2000 x g for 5 min). After a washing step with 15 mL STC (1 M Sorbitol, 10 mM Tris-HCl pH 7.5, 10 mM CaCl_2), protoplasts were adjusted to 10⁸ mL⁻¹ with STC:PEG (4:1) and divided into 200- μ L aliquots. Subsequently, PEG-based transformation was performed essentially as described by Liu and Friesen (2012) using 70 μ g of each construct per 200 μ L protoplast aliquot. As a regeneration step, protoplasts were mixed with 10 mL of regeneration medium agar, poured into a petri dish, and kept in the dark at 21 °C. After 14 h, the plate was overlaid with 10 mL regeneration medium agar containing 150 μ g mL⁻¹ Hygromycin B (Roche Diagnostics, Indianapolis, USA) for transformant selection. After 10 days, putative transformants were transferred to PDA plates amended with 150 μ g mL⁻¹ hygromycin.

DNA was extracted from PDA cultures using a modified version of the microprep protocol published by Fulton et al. (1995), replacing chloroform:isoamyl (24:1) with phenol:chloroform:isopropanol (25:24:1). PCR amplifications were carried out using the GoTaq Flexi DNA Polymerase kit (Promega, Madison, WI) following the manufacturer's suggested protocol. Site-directed transformation was confirmed by the presence of PCR products generated with a forward primer designed to anneal upstream of the 5' flanking region of each individual target gene (5F) and a reverse primer annealing to the 5' end of the hygromycin resistance encoding gene (HYG R2) (Table 4). Additionally, knock-out mutants were verified by the absence of a PCR product when using forward primer (1F) of the target gene 5' flank and a reverse primer designed on the target gene coding sequence (6R) (Table 4). The presence of a 1F/6R PCR product and absence of a 5F/HYG R2 PCR product indicated a mutant with an ectopic integration of the construct. Verified isolates were stored at 22°C in the dark until used for fungicide sensitivity assays, which were carried out within three weeks of verification.

3. RESULTS

3.1. *Cercospora beticola* transcript level changes after exposure to tetraconazole

Two isolates harvested from the north central U.S. sugar beet growing area contrasting for DMI sensitivity were used in this study. *Cercospora beticola* strain 09-32 is sensitive to all DMIs used for CLS management exhibiting an EC_{50} of $0.006 \mu\text{g ml}^{-1}$ to tetraconazole while DMI-resistant strain 10-73-4 has an EC_{50} to tetraconazole of $35.9 \mu\text{g ml}^{-1}$. The genome sequence of *C. beticola* strain 10-73-4 was determined using sequencing-by-synthesis on the Illumina HiSeq-2000, which served as the reference sequence in this study. To this end, ~ 1.5 Gb of paired-end (PE) library-derived reads (500-bp insert size; 100-bp read length) was produced in combination with ~ 1.5 Gb of mate-pair (MP) library-derived reads (5-kb insert size; 100-bp read length). The PE and MP sequences were assembled using the A5 pipeline (Tritt et al., 2012) and 105 scaffolds were generated with an N50 of about 405 kb and the longest contig of $\sim 2,475$ kb. The assembled genome was used for gene annotation. In total, 12,440 genes were predicted on the *C. beticola* strain 10-73-4 strain using the coding sequences from the manual gene annotation of *C. beticola* strain 09-40 (de Jonge and Bolton, *unpublished data*). Both 09-32 and 09-40 are 98.3% identical on the nucleotide level to 10-73-4. An average of 98.0% and 99.4% of reads were mapped to the reference sequence across replicates and treatments for 09-32 and 10-73-4, respectively (Supplemental Table 1). Out of 12,440 genes, about 11,587 have RNA-seq support for more than 95% of the gene length.

Both isolates were grown *in vitro* followed by a tetraconazole or acetone treatment as described by Bolton et al. (2012a). Although any pair-wise comparison could be made between isolates and treatments, we sought to identify transcript level changes solely associated with the response to tetraconazole in each strain. Since *C. beticola* isolates are typically genetically diverse (Bolton et al., 2012c), comparisons within a single genetic background negate any between-genotype background

variation that may inadvertently identify genes not associated with tetraconazole resistance. Therefore, we employed a univariate analysis (t tests, Q -value ≤ 0.001 , and $\log_2$ fold regulation differences ≥ 1.0) between mock- and tetraconazole-treatments for each *C. beticola* strain.

We identified 414 genes that were differentially-expressed between mock- and-tetraconazole-treatments in the DMI-sensitive isolate 09-32 (Fig. 1). Of the 286 genes uniquely identified in 09-32, 115 were induced and 171 were repressed (Supplemental Table 2). To characterize the biological processes underlying these 286 genes, Of these genes, 213 were induced and 216 were repressed (Supplemental Table 2). To characterize the biological processes that occurred in the DMI-sensitive strain in response to tetraconazole, we used gene ontology (GO) term enrichment analysis which revealed 11 and 21 GO terms significantly over and under-represented, respectively (Supplemental Table 3). However, since previous work in our laboratory revealed that *CbCyp51* was induced upon exposure to tetraconazole in DMI-resistant isolates (Bolton et al., 2012a), we were primarily interested in identifying genes with expression profiles similar to *CbCyp51* that were differentially-expressed after exposure to tetraconazole in the DMI-resistant strain 10-73-4 when compared to the control treatment. To that end, we identified 244 genes differentially-expressed between mock- and tetraconazole-treatments in 10-73-4 (Fig. 1) of which 172 were induced and 72 repressed. In total, 128 genes were differentially-expressed in both isolates as a consequence of tetraconazole exposure (Fig. 1).

3.2. Classification of genes differentially-expressed in both strains in response to tetraconazole

Although we were primarily interested in identifying genes induced in response to tetraconazole in the DMI-resistant isolate, identifying pathways induced in both strains as a consequence of tetraconazole exposure may characterize the background and perhaps a more universal response to tetraconazole in this pathosystem. We identified 128 genes differentially-expressed in both *C. beticola* strains (Fig. 1). These genes were analyzed with InterPro (Mitchell et al., 2014) and PFAM (Finn et al., 2014) to aid in protein functional analysis by classifying into protein families by predicting domains and

other important sites. GO terms (Ashburner et al., 2000) were further applied to classify genes into specific processes to aid in pathway reconstruction (Table 1). GO term enrichment analysis revealed four GO terms significantly over-represented in both strains, which included lipid metabolic processes (GO:0006629), oxidoreductase activity (GO:0016491), catalytic activity (GO:0003824), and the cellular component endoplasmic reticulum (GO:0005783) (Supplemental Table 4). In contrast, cellular processes (GO:0009987), cellular metabolic processes (GO:0044237), and protein metabolic processes (GO:0019538) were repressed in both strains (Supplemental Table 4). In total, InterPro and PFAM classifications were available for 94 and 80 genes, respectively, which enabled the classification of nearly half of the identified genes into 27 GO term categories (Table 1).

Four genes involved with the isoprenoid biosynthetic process (Table 1) were induced in both strains. Isoprenoids, also known as terpenoids, are a large class of ubiquitous metabolites synthesized through the condensation of C₅ units that are precursors for ergosterol biosynthesis in fungi. Further analysis of the gene list (Table 1) identified an additional gene (*Erg10*) in the terpenoid backbone biosynthesis pathway (KEGG Pathway map00900) involved in ergosterol production (Fig. 2A). The last gene in the terpenoid backbone biosynthesis pathway leading to steroid biosynthesis (KEGG Pathway map00100) is farnesyl diphosphate synthase (FDS), giving rise to farnesyl diphosphate, the initial substrate for the steroid biosynthesis pathway (Fig. 2A). In total, we identified 11 genes in the steroid biosynthesis pathway leading to the production of ergosterol that were induced in both strains (Fig. 2A). Additionally, the *Erg26* gene that is involved with ergosterol production was only differentially-expressed in 10-73-4 (see below) but was also included in Figure 2. To compare the expression for each gene in both treatments for each isolate, we plotted FPKM reads for each gene identified as differentially-expressed (Fig. 2B). Although *Cyp51* was shown to have a higher relative expression in the DMI-sensitive strain 09-32 (5.10 log₂ fold change) compared to the DMI-resistant strain (3.57 log₂ fold change) (Table 1), analysis of the FPKM values illustrate that *Cyp51* was induced to a much higher level

in DMI-resistant strain 10-73-4 after tetraconazole exposure (Fig. 2B). To confirm that differential *Cyp51* expression was not due sequence polymorphism between strains, we confirmed that the coverage of RNA-seq reads was uniform for 09-32 and 10-73-4 along the *Cyp51* coding region (Supplementary Fig. 1). Similarly, expression of *Erg3* was much higher in 10-73-4 than 09-32 after tetraconazole exposure (Fig. 2B). However, most genes had similar FPKM values between the two strains for each treatment (Fig. 2B). In total, 17 of the 23 genes in the ergosterol biosynthesis pathway were induced in both strains upon tetraconazole exposure (Fig. 2), suggesting ergosterol biosynthesis is a universal response to tetraconazole in *C. beticola*. In support of this hypothesis, two key regulators of sterol homeostasis, sterol regulatory element-binding protein ortholog Sre1 and sterol regulatory element-binding protein cleavage-activating protein, were induced in both strains after tetraconazole treatment (Table 1). In addition, a gene with a CRAL/TRIO and Sec14p-like domain was induced in both strains (Table 1). Proteins with these domains are known to be involved with lipid binding (Panagabko et al., 2003) and have been associated with resistance to DMIs in yeast (Šimová et al., 2013).

Polyketide synthases (PKSs) catalyze the elongation of polyketides and are among the most prominent and well-studied fungal secondary metabolite biosynthetic genes. Genes involved with the modification of intermediate products formed from such enzymes are typically found in a contiguous gene cluster that may also include a transport-related gene involved with SM efflux and/or a pathway-specific transcription factor that control expression of the cluster (Thomma et al., 2015). A polyketide biosynthesis gene cluster (CBET3_11350-RA to CBET3_11356; Table 1) was differentially-expressed in response to tetraconazole between the two isolates. CBET3_11350-RA is a predicted non-reducing polyketide synthase (de Jonge and Bolton, *unpublished*) and together with five other genes in the cluster was induced in the tetraconazole-resistant strain 10-73-4 but were all repressed in the sensitive strain 09-32 (Table 1). [Additionally, genes encoding a PKS \(CBET3_09638-RA\) and a tetrahydroxynaphthalene reductase \(CBET3_09636-RA\) that belong to the same PKS cluster \(de Jonge and Bolton, *unpublished*\),](#)

were induced in the resistant strain but repressed in the sensitive strain (Table 1). *CBET3_009636* is an ortholog to *PKS1* from *Zymoseptoria tritici*, which was recently shown to be associated with increased melanization and DMI-resistance (Lendenmann et al., 2015).

3.3. Classification of genes differentially-expressed in DMI-resistant strain in response to tetraconazole

InterPro and PFAM classification was available for 86 and 77 genes, respectively, permitting the classification of nearly half of the differentially-expressed genes identified in the DMI-resistant strain into 21 GO term categories (Table 2). No obvious pathways unique to 10-73-4 were up-regulated in response to tetraconazole. GO term enrichment analysis revealed a single GO term (secondary metabolic process; GO:0019748) over-represented (Supplemental Table 5). Several genes involved directly or indirectly with the fungal cell membrane or cell wall were induced in response to tetraconazole. For example, a gene containing an RTA1 domain known to encode a plasma membrane protein (Manente and Ghislain, 2009) was the most highly differentially-expressed gene in the study (Table 2). RTA1 encodes a protein with seven potential transmembrane domains and has been shown to mediate resistance to 7-amincholesterol, a class of sterol biosynthesis inhibitors (Soustre et al., 1996). Another gene belonging to the Gpr1/Fun34/YaaH family encoding a plasma membrane protein was induced in 10-73-4 (Table 2). Although there is no sequence conservation with RTA1, this protein is also an integral protein of the plasma membrane with six transmembrane domains as has been shown in the yeast *Yarrowia lipolytica* (Augstein et al., 2003). Additionally and as mentioned above, *Erg26*, which is in the ergosterol pathway, was specifically induced in 10-73-4 (Table 2).

Several genes encoding proteins involved with transmembrane transport were differentially-expressed in 10-73-4. Of particular interest are major facilitator superfamily (MFS) and ATP-binding cassette (ABC) transporters, which have been shown to be associated with fungicide resistance by active efflux of xenobiotics (Cools et al., 2013; Roohparvar et al., 2008; Zwiers et al., 2002). We identified five

MFS and two ABC transporters induced in 10-73-4 in response to tetraconazole (Table 2). Several genes with a more indirect putative role in the fungal cell wall were also implicated in fungicide resistance in *C. beticola* strain 10-73-4. For example, numerous genes encoding glycoside hydrolase family proteins were differentially-expressed in 10-73-4 (Table 2). Glycoside hydrolases are a widespread group of enzymes that hydrolyze the glycosidic bond between two or more carbohydrates or between a carbohydrate and a non-carbohydrate moiety, and are classified largely on substrate specificity but occasionally by mode of action. Seven genes encoding four glycoside hydrolase families were induced in response to tetraconazole in 10-73-4 (Table 2). Three glycoside hydrolase family 16 genes were induced, which are known to encode enzymes that break down β -1,3-glucans, a major component of fungal cell walls (Bowman and Free, 2006).

Since there is no *CbCyp1* promoter regulatory polymorphism associated with enhanced DMI resistance (Bolton et al., 2012a) as has been shown in other fungal systems (Hamamoto et al., 2000; Ma et al., 2006; Schnabel and Jones, 2001), we are interested in identifying transcription factors responsible for *CbCyp51* and other DMI-associated gene up-regulation. In this study, we identified one transcription factor uniquely induced in 10-73-4 (Table 2). This transcription factor is of the $\text{Zn(II)}_2\text{Cys}_6$ class and shares homology to *Aspergillus* spp. AflR (Table 2). AflR transcription factor orthologs are known to regulate biosynthesis of several secondary metabolites including aflatoxin, sterigmatocystin, and dothistromin (Bradshaw et al., 2013; Brown et al., 1996).

Five genes encoding cytochrome P450s (CYPs) were induced in 10-73-4 after tetraconazole treatment (Table 2). Fungal CYPs carry out a wide range of functions dependent on their enzymatic properties and substrate specificity in both primary and secondary metabolism (Lah et al., 2008). Similarly, other genes with a diversity of functions were induced including α/β hydrolases, carbohydrate esterases, and genes with FAD/NAD(P)-binding domains (Table 2).

3.4. Generation of mutants in candidate genes associated with DMI-resistance

To aid in establishing a role in DMI-resistance, several genes were chosen for further characterization based on expression values and/or confirmed roles in DMI resistance in other pathosystems. Genes were characterized by replacing the gene of interest from the genome of DMI-resistant strain 10-73-4 with a gene encoding hygromycin B resistance, followed by analysis of EC₅₀ values compared to the ectopic mutants of the same gene and the progenitor wild-type strain (Table 3). Average EC₅₀ values were calculated based on individual EC₅₀ values measured from three independent ectopic and site-directed transformants. The average EC₅₀ of three independent measurements of the wild-type progenitor strain 10-73-4 was 35.9 µg ml⁻¹. Regardless of the gene chosen for replacement or whether the transformation event resulted in a site-directed or ectopic mutant, average EC₅₀ values of all mutants was significantly higher than the progenitor wild-type strain (Table 3), suggesting that the transformation process alone increased resistance to tetraconazole. Transformations and subsequent EC₅₀ evaluations were repeated, which provided similar results (not shown).

To investigate this phenomenon further, we produced protoplasts from two additional *C. beticola* strains and compared their EC₅₀ values to their progenitor strains that did not go through the protoplast process. These experiments were identical to previous transformation procedures except no gene encoding a selection marker was included and therefore protoplasts were recovered from plates lacking hygromycin selection. Protoplasting caused a 25.7% and 6.4% increase in EC₅₀ value, respectively, compared to the non-protoplasted progenitor strain. To investigate if the elevated EC₅₀ values were transient in nature, we cycled mutant strains through two cycles of PDA growth, but observed similar levels of fungicide resistance (not shown).

4. Discussion

The fungal pathogen *C. beticola* is the most important foliar pathogen of sugar beet and requires well-timed fungicide applications for disease management. Due to widespread use of tetraconazole, resistance has been increasing in *C. beticola* populations in recent years (Bolton et al., 2012b; Secor et al., 2010). Insight into the molecular basis of tetraconazole resistance may lead to molecular tools to identify DMI-resistant strains for fungicide resistance management programs. In this study, we profiled the response of two *C. beticola* strains contrasting for resistance to the important DMI fungicide tetraconazole. Although our initial interest was to identify induced genes in the DMI-resistant strain, we observed a large overlap of genes differentially-expressed in both strains (Fig. 1) suggesting a common background response when exposed to tetraconazole. Notably, 17 of the 23 genes in the ergosterol biosynthesis pathway were induced in both strains upon tetraconazole exposure (Fig. 2). The clear induction of ergosterol pathway genes suggests maintenance or reinforcement of cell membrane integrity is a universal response when exposed to tetraconazole. Similarly, a strain of *Aspergillus niger* responded to the ergosterol-targeting fungicide fenpropimorph with enhanced expression of several ergosterol pathway genes identified using an Affymetrix microarray (Meyer et al., 2007). Cools et al. (2007) used a cDNA microarray approach to identify three induced genes in an epoxiconazole-sensitive strain of *Zymoseptoria Z. tritici*, all of which were ergosterol biosynthesis pathway genes (*Erg25*, *Erg24*, and *Erg5*). These same genes were induced in the *Z. tritici* epoxiconazole-resistant strain, but with lower transcript levels (Cools et al., 2007). In the human pathogen *Candida albicans*, induced expression of *Cyp51* and other genes involved with ergosterol biosynthesis has been shown to be associated with DMI exposure (De Backer et al., 2001; Dunkel et al., 2008; Liu et al., 2005). Although most ergosterol biosynthesis genes were induced to similar levels in our study, *CbCyp51* was induced to much higher levels in the DMI-resistant strain compared to the sensitive strain (Fig. 2), underlying a key difference between sensitive and resistant *C. beticola* strains and supporting our previous results that showed native *CbCyp51* expression is generally higher and expression is inducible in DMI-resistant field strains

(Bolton et al., 2012a). Notably, the *C. beticola* *HMGCR* gene, which encodes the rate-controlling enzyme of the mevalonate pathway in terpenoid backbone biosynthesis leading to ergosterol production, was induced in both strains in response to tetraconazole (Table 1). The mechanism for the up-regulation of *CbCyp51* or other ergosterol biosynthesis genes is currently not known in *C. beticola*. In the human pathogen *Candida albicans*, the zinc cluster transcription factor Upc2p has been shown to regulate the expression of *Cyp51* and other genes involved in ergosterol biosynthesis upon exposure to DMIs (MacPherson et al., 2005). Likewise, the *Saccharomyces cerevisiae* transcriptional activators Upc2p and Ecm22p positively regulate the expression of ergosterol biosynthesis genes (Vik and Rine, 2001). In our study, we identified several transcription factors induced in response to tetraconazole (Tables 1 and 2). Although it is currently not possible to fully characterize these genes with gene replacement studies (discussed below), it will be interesting to investigate whether these transcription factors regulate expression of ergosterol biosynthesis genes in *C. beticola*.

Similar to DMI exposure, cellular oxygen deficiency has been shown to induce ergosterol biosynthesis genes. For example, several ergosterol biosynthesis genes were up-regulated in the rice blast fungus *Magnaporthe oryzae* in response to hypoxia (Choi et al., 2015). Another example of fungal response to hypoxia is the induction of the sterol regulatory element-binding protein (SREBP) pathway, which induces ergosterol biosynthesis and hyphal growth in order to scavenge more oxygen in human fungal pathogens (Hughes et al., 2005). SREBPs are transcription factors that are activated when sterols are depleted due to triggers such as hypoxia and iron limitation (Blatzer et al., 2011). Active SREBP turns on the expression of sterol synthesis enzymes and other oxygen-dependent proteins (Blatzer et al., 2011; Choi et al., 2015). Recently, Liu et al. (2015) identified a novel SREBP gene (*sreA*) that was required for DMI-resistance and *Cyp51* expression in *Penicillium digitatum*. In our study, we also identified a gene encoding SREBP induced along with ergosterol biosynthesis genes in both strains (Table 1), perhaps suggesting that oxygen is limiting and/or ergosterol deficiency is sensed in *C. beticola* cells upon DMI

exposure. Ergosterol biosynthesis is thought to be impaired under hypoxic conditions because of decreased synthesis of heme and heme-containing proteins (Manente and Ghislain, 2009). Hypoxia and heme-deficiency are also known to induce *RTA1* expression (Protchenko et al., 2008), which was the most highly-induced gene in the DMI-resistant strain in our study (Table 2). The *RTA1* gene was originally identified during a screen for genes that confer resistance to the sterol biosynthesis inhibitor 7-amincholesterol in *S. cerevisiae* (Soustre et al., 1996). RTA1 proteins encode a protein with six or seven transmembrane spans (Manente and Ghislain, 2009). Although the function of RTA1 is not fully understood, RTA1 is not thought to be involved with efflux of 7-amincholesterol and *S. cerevisiae* null mutants are still viable under normal growth conditions (Soustre et al., 1996). Recent evidence suggests several gene networks involved with diverse cellular responses including hypoxia, responses to some cytotoxic drugs, and ergosterol biosynthesis converge at the promoter of the *RTA1* gene (Kořaczowska et al., 2012), suggesting that RTA1 is involved with the response to various stresses perhaps by fortifying the cell membrane. Similar to *S. cerevisiae* (Manente and Ghislain, 2009), the *C. beticola* genome encodes nine RTA1-like proteins that are similar only on the amino acid level (de Jonge and Bolton, *unpublished*). Taken together, this may suggest that functional redundancy of RTA1 is necessary to ready the cell for a variety of stresses.

The basic protocol for transformation of fungal protoplasts by the addition of a high concentration of polyethylene glycol (PEG) is well-established and remains the most common method to conduct transformation in filamentous fungi (Liu and Friesen, 2012). In our study, we used PEG-mediated transformation of protoplasts to introduce constructs to remove genes putatively associated with DMI-resistance. Surprisingly, DMI-resistance as measured by EC_{50} value was significantly increased in all mutants generated in this study regardless of the target gene or whether the integration was site-directed or ectopic (Table 3). To further characterize this phenomenon, we generated mutants in two other strains and found that EC_{50} values increased in both. This may suggest that *C. beticola* responds to

protoplast formation in the transformation process with a new cell wall that is particularly fortified, which results in enhanced DMI-resistance. Likewise, the cell membrane may be modified as a result of PEG exposure which results in increased DMI resistance. To our knowledge, this phenomenon has not been described in filamentous fungi. Since there are no other reported transformation alternatives for *C. beticola*, future research directed towards developing an *Agrobacterium tumefaciens*-based transformation protocol will be necessary to characterize genes associated with DMI resistance in this pathogen.

In conclusion, we used next generation sequencing to compare the response of two strains contrasting with DMI-resistance in the sugar beet pathogen *C. beticola*. The genes induced in response to tetraconazole highlight the important role that the cell membrane plays in DMI-resistance. This is further supported by our finding that mutants generated from *C. beticola* protoplasts exhibited higher DMI-resistance than progenitor strains. Future studies directed at dissecting particular cell-wall and -membrane components specifically involved with tetraconazole resistance will be of particular interest to understand DMI resistance in this pathosystem.

Acknowledgements

We thank Keshav Birla and Jon Neubauer (USDA-ARS) for excellent technical assistance. M. D. Bolton is supported by USDA-ARS CRIS project 3060-22000-047-00 and grants from the Beet Sugar Development Foundation and the Sugar beet Research and Education Board of MN and ND.

REFERENCES

- Ashburner, M., Ball, C. A., Blake, J. A., Botstein, D., Butler, H., Cherry, J. M., Davis, A. P., Dolinski, K., Dwight, S. S., Eppig, J. T., 2000. Gene Ontology: tool for the unification of biology. *Nat. Genet.* 25, 25-29.
- Augstein, A., Barth, K., Gentsch, M., Kohlwein, S. D., Barth, G., 2003. Characterization, localization and functional analysis of Gpr1p, a protein affecting sensitivity to acetic acid in the yeast *Yarrowia lipolytica*. *Microbiology*. 149, 589-600.
- Birla, K., Rivera-Varas, V., Secor, G. A., Khan, M. F. R., Bolton, M. D., 2012. Characterization of *cytochrome b* from European field isolates of *Cercospora beticola* with quinone outside inhibitor resistance. *Eur. J. Plant Pathol.* 134, 475-488.
- Blatzer, M., Barker, B. M., Willger, S. D., Beckmann, N., Blosser, S. J., Cornish, E. J., Mazurie, A., Grahl, N., Haas, H., Cramer, R. A., 2011. SREBP coordinates iron and ergosterol homeostasis to mediate triazole drug and hypoxia responses in the human fungal pathogen *Aspergillus fumigatus*. *PLoS Genet.* 7, e1002374.
- Bolton, M. D., Birla, K., Rivera-Varas, V., Rudolph, K., Secor, G. A., 2012a. Characterization of *CbCyp51* from field isolates of *Cercospora beticola*. *Phytopathol.* 102, 298-305.
- Bolton, M. D., de Jonge, R., Inderbitzin, P., Liu, Z., Birla, K., Van de Peer, Y., Subbarao, K. V., Thomma, B. P. H. J., Secor, G. A., 2014. The heterothallic sugarbeet pathogen *Cercospora beticola* contains exon fragments of both *MAT* genes that are homogenized by concerted evolution. *Fungal Genet. Biol.* 62, 43-54.
- Bolton, M. D., Rivera-Varas, V., del Río Mendoza, L. E., Khan, M. F. R., Secor, G. A., 2012b. Efficacy of variable tetraconazole rates against *Cercospora beticola* isolates with differing in vitro sensitivities to DMI fungicides. *Plant Dis.* 96, 1749-1756.
- Bolton, M. D., Rivera-Varas, V., Secor, G. A., Cattanaach, A. W., Metzger, M. S., 2013a. Identification of the G143A mutation in cytochrome b associated with Q₀I resistance in *Cercospora beticola* isolates from the Red River Valley. *Plant Health Progress*.
- Bolton, M. D., Rivera, V., Secor, G., 2013b. Identification of the G143A mutation associated with Q₀I resistance in *Cercospora beticola* field isolates from Michigan, United States. *Pest Manag. Sci.* 69, 35-39.
- Bolton, M. D., Secor, G. A., Rivera, V., Weiland, J. J., Rudolph, K., Birla, K., Rengifo, J., Campbell, L. G., 2012c. Evaluation of the potential for sexual reproduction in field populations of *Cercospora beticola* from USA. *Fungal Biol.* 116, 511-521.
- Bowman, S. M., Free, S. J., 2006. The structure and synthesis of the fungal cell wall. *BioEssays*. 28, 799-808.
- Bradshaw, R. E., Slot, J. C., Moore, G. G., Chettri, P., de Wit, P. J. G. M., Ehrlich, K. C., Ganley, A. R. D., Olson, M. A., Rokas, A., Carbone, I., Cox, M. P., 2013. Fragmentation of an aflatoxin-like gene cluster in a forest pathogen. *New Phytol.* 198, 525-535.
- Brown, D. W., Yu, J. H., Kelkar, H. S., Fernandes, M., Nesbitt, T. C., Keller, N. P., Adams, T. H., Leonard, T. J., 1996. Twenty-five coregulated transcripts define a sterigmatocystin gene cluster in *Aspergillus nidulans*. *Proc. Natl. Acad. Sci. U. S. A.* 93, 1418-1422.
- Carter, H. E., Fraaije, B. A., West, J. S., Kelly, S. L., Mehl, A., Shaw, M. W., Cools, H. J., 2014. Alterations in the predicted regulatory and coding regions of the sterol 14 α -demethylase gene (*CYP51*) confer decreased azole sensitivity in the oilseed rape pathogen *Pyrenopeziza brassicae*. *Mol. Plant Pathol.* 15, 513-522.
- Catlett, N. L., Lee, B. N., Yoder, O. C., Turgeon, B. G., 2003. Split-marker recombination for efficient targeted deletion of fungal genes. *Fungal Genetics Reports*. 50, 9-11.

- Choi, J., Chung, H., Lee, G.-W., Koh, S.-K., Chae, S.-K., Lee, Y.-H., 2015. Genome-wide analysis of hypoxia-responsive genes in the rice blast fungus, *Magnaporthe oryzae*. PLoS ONE. 10, e0134939.
- Cools, H. J., Bayon, C., Atkins, S., Lucas, J. A., Fraaije, B. A., 2012. Overexpression of the sterol 14 α -demethylase gene (*MgCYP51*) in *Mycosphaerella graminicola* isolates confers a novel azole fungicide sensitivity phenotype. Pest Manag. Sci. 68, 1034-1040.
- Cools, H. J., Fraaije, B. A., Bean, T. P., Antoniow, J., Lucas, J. A., 2007. Transcriptome profiling of the response of *Mycosphaerella graminicola* isolates to an azole fungicide using cDNA microarrays. Mol. Plant Pathol. 8, 639-651.
- Cools, H. J., Hawkins, N. J., Fraaije, B. A., 2013. Constraints on the evolution of azole resistance in plant pathogenic fungi. Plant Pathol. 62, 1-7.
- Cools, H. J., Parker, J. E., Kelly, D. E., Lucas, J. A., Fraaije, B. A., Kelly, S. L., 2010. Heterologous expression of mutated eburiol 14 α -demethylase (CYP51) proteins of *Mycosphaerella graminicola* to assess effects on azole fungicide sensitivity and intrinsic protein function. Appl. Environ. Microbiol. 76, 2866-2872.
- Daub, M. E., Ehrenschaft, M., 2000. The photoactivated *Cercospora* toxin cercosporin: contributions to plant disease and fundamental biology. Annu. Rev. Phytopathol. 38, 461-490.
- De Backer, M. D., Ilyina, T., Ma, X.-J., Vandoninck, S., Luyten, W. H. M. L., Vanden Bossche, H., 2001. Genomic profiling of the response of *Candida albicans* to itraconazole treatment using a DNA microarray. Antimicrob. Agents Chemother. 45, 1660-1670.
- Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., Gingeras, T. R., 2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics. 29, 15-21.
- Dunkel, N., Liu, T. T., Barker, K. S., Homayouni, R., Morschhäuser, J., Rogers, P. D., 2008. A gain-of-function mutation in the transcription factor Upc2p causes upregulation of ergosterol biosynthesis genes and increased fluconazole resistance in a clinical *Candida albicans* isolate. Eukaryot. Cell. 7, 1180-1190.
- Finn, R. D., Bateman, A., Clements, J., Coggill, P., Eberhardt, R. Y., Eddy, S. R., Heger, A., Hetherington, K., Holm, L., Mistry, J., Sonnhammer, E. L. L., Tate, J., Punta, M., 2014. Pfam: the protein families database. Nucleic Acids Res. 42, D222-D230.
- Franc, G. D., Ecology and epidemiology of *Cercospora beticola*. In: R. T. Lartey, et al., (Eds.), *Cercospora leaf spot of sugar beet and related species*. The American Phytopathological Society, St. Paul, Minnesota, 2010, pp. 7-19.
- Friesen, T. L., Faris, J. D., Characterization of plant-fungal interactions involving necrotrophic effector-producing plant pathogens. In: M. D. Bolton, B. P. H. J. Thomma, (Eds.), *Plant Fungal Pathogens: Methods and Protocols*. Springer, 2012, pp. 191-207.
- Friesen, T. L., Stukenbrock, E. H., Liu, Z., Meinhardt, S., Ling, H., Faris, J. D., Rasmussen, J. B., Solomon, P. S., McDonald, B. A., Oliver, R. P., 2006. Emergence of a new disease as a result of interspecific virulence gene transfer. Nat. Genet. 38, 953-956.
- Fulton, T. M., Chunwongse, J., Tanksley, S. D., 1995. Microprep protocol for extraction of DNA from tomato and other herbaceous plants. Plant Mol. Biol. Report. 13, 207-207.
- Goswami, R., Targeted Gene Replacement in Fungi Using a Split-Marker Approach. In: M. D. Bolton, B. P. H. J. Thomma, (Eds.), *Plant Fungal Pathogens*. Humana Press, 2012, pp. 255-269.
- Hamamoto, H., Hasegawa, K., Nakaune, R., Lee, Y. J., Makizumi, Y., Akutsu, K., Hibi, T., 2000. Tandem repeat of a transcriptional enhancer upstream of the sterol 14 α -demethylase gene (*CYP51*) in *Penicillium digitatum*. Appl. Environ. Microbiol. 66, 3421.
- Hughes, A. L., Todd, B. L., Espenshade, P. J., 2005. SREBP pathway responds to sterols and functions as an oxygen sensor in fission yeast. Cell. 120, 831-842.
- Khan, J., del Río, L. E., Nelson, R., Khan, M. F. R., 2007. Improving the *Cercospora* leaf spot management model for sugar beet in Minnesota and North Dakota. Plant Dis. 91, 1105-1108.

- Khan, J., Rio, L. E. d., Nelson, R., Rivera-Varas, V., Secor, G. A., Khan, M. F. R., 2008. Survival, dispersal, and primary infection site for *Cercospora beticola* in sugar beet. *Plant Dis.* 92, 741-745.
- Kończakowska, A., Manente, M., Kończakowski, M., Laba, J., Ghislain, M., Wawrzycka, D., 2012. The regulatory inputs controlling pleiotropic drug resistance and hypoxic response in yeast converge at the promoter of the aminosterol resistance gene *RTA1*. *FEMS Yeast Res.* 12, 279-292.
- Kretschmer, M., Leroux, P., Mosbach, A., Walker, A.-S., Fillinger, S., Mernke, D., Schoonbeek, H.-J., Pradier, J.-M., Leroux, P., De Waard, M. A., Hahn, M., 2009. Fungicide-driven evolution and molecular basis of multidrug resistance in field populations of the grey mould fungus *Botrytis cinerea*. *PLoS Pathog.* 5, e1000696.
- Lah, L., Kraševac, N., Trontelj, P., Komel, R., 2008. High diversity and complex evolution of fungal cytochrome P450 reductase: Cytochrome P450 systems. *Fungal Genet. Biol.* 45, 446-458.
- Lartey, R. T., Caesar-TonThat, T. C., Caesar, A. J., Shelver, W. L., Sol, N. I., Bergman, J. W., 2005. Safflower: A New Host of *Cercospora beticola*. *Plant Dis.* 89, 797-801.
- Lendenmann, M. H., Croll, D., McDonald, B. A., 2015. QTL mapping of fungicide sensitivity reveals novel genes and pleiotropy with melanization in the pathogen *Zymoseptoria tritici*. *Fungal Genet. Biol.* 80, 53-67.
- Leroux, P., Albertini, C., Gautier, A., Gredt, M., Walker, A.-S., 2007. Mutations in the *CYP51* gene correlated with changes in sensitivity to sterol 14 α -demethylation inhibitors in field isolates of *Mycosphaerella graminicola*. *Pest Manag. Sci.* 63, 688-698.
- Liu, J., Yuan, Y., Wu, Z., Li, N., Chen, Y., Qin, T., Geng, H., Xiong, L., Liu, D., 2015. A novel sterol regulatory element-binding protein gene *sreA* identified in *Penicillium digitatum* is required for prochloraz resistance, full virulence and *erg11* (*cyp51*) regulation. *PLoS ONE.* 10, e0117115.
- Liu, T. T., Lee, R. E. B., Barker, K. S., Lee, R. E., Wei, L., Homayouni, R., Rogers, P. D., 2005. Genome-wide expression profiling of the response to azole, polyene, echinocandin, and pyrimidine antifungal agents in *Candida albicans*. *Antimicrob. Agents Chemother.* 49, 2226-2236.
- Liu, Z., Friesen, T. L., Polyethylene glycol (PEG)-mediated transformation in filamentous fungal pathogens. In: M. D. Bolton, B. P. H. J. Thomma, Eds.), *Plant Fungal Pathogens: Methods and Protocols*. Springer, 2012, pp. 365-375.
- Ma, B., Tredway, L. P., 2013. Induced overexpression of cytochrome P450 sterol 14 α -demethylase gene (*CYP51*) correlates with sensitivity to demethylation inhibitors (DMIs) in *Sclerotinia homoeocarpa*. *Pest Manag. Sci.* 69, 1369-1378.
- Ma, Z., Proffer, T. J., Jacobs, J. L., Sundin, G. W., 2006. Overexpression of the 14 α -demethylase target gene (*CYP51*) mediates fungicide resistance in *Blumeriella jaapii*. *Appl. Environ. Microbiol.* 72, 2581-2585.
- MacPherson, S., Akache, B., Weber, S., De Deken, X., Raymond, M., Turcotte, B., 2005. *Candida albicans* zinc cluster protein Upc2p confers resistance to antifungal drugs and is an activator of ergosterol biosynthetic genes. *Antimicrob. Agents Chemother.* 49, 1745-1752.
- Manente, M., Ghislain, M., 2009. The lipid-translocating exporter family and membrane phospholipid homeostasis in yeast. *FEMS Yeast Res.* 9, 673-687.
- Meyer, V., Damveld, R. A., Arentshorst, M., Stahl, U., van den Hondel, C. A. M. J. J., Ram, A. F. J., 2007. Survival in the presence of antifungals: genome-wide expression profiling of *Aspergillus niger* in response to sublethal concentrations of caspofungin and fenpropimorph. *J. Biol. Chem.* 282, 32935-32948.
- Mitchell, A., Chang, H.-Y., Daugherty, L., Fraser, M., Hunter, S., Lopez, R., McAnulla, C., McMenamin, C., Nuka, G., Pesseat, S., Sangrador-Vegas, A., Scheremetjew, M., Rato, C., Yong, S.-Y., Bateman, A., Punta, M., Attwood, T. K., Sigrist, C. J. A., Redaschi, N., Rivoire, C., Xenarios, I., Kahn, D., Guyot, D., Bork, P., Letunic, I., Gough, J., Oates, M., Haft, D., Huang, H., Natale, D. A., Wu, C. H., Orengo,

- C., Sillitoe, I., Mi, H., Thomas, P. D., Finn, R. D., 2014. The InterPro protein families database: the classification resource after 15 years. *Nucleic Acids Res.*
- Panagabko, C., Morley, S., Hernandez, M., Cassolato, P., Gordon, H., Parsons, R., Manor, D., Atkinson, J., 2003. Ligand Specificity in the CRAL-TRIO Protein Family. *Biochemistry.* 42, 6467-6474.
- Panella, L., 2010. Sugar beet as an energy crop. *Sugar Tech.* 12, 288-293.
- Protchenko, O., Shakoury-Elizeh, M., Keane, P., Storey, J., Androphy, R., Philpott, C. C., 2008. Role of PUG1 in inducible porphyrin and heme transport in *Saccharomyces cerevisiae*. *Eukaryot. Cell.* 7, 859-871.
- Roberts, A., Pimentel, H., Trapnell, C., Pachter, L., 2011. Identification of novel transcripts in annotated genomes using RNA-Seq. *Bioinformatics.* 27, 2325-2329.
- Roohparvar, R., Mehrabi, R., Van Nistelrooy, J. G., Zwiers, L. H., De Waard, M. A., 2008. The drug transporter MgMfs1 can modulate sensitivity of field strains of the fungal wheat pathogen *Mycosphaerella graminicola* to the strobilurin fungicide trifloxystrobin. *Pest Manag. Sci.*
- Schnabel, G., Jones, A. L., 2001. The 14 α -demethylase (*CYP51A1*) gene is overexpressed in *Venturia inaequalis* strains resistant to myclobutanil. *Phytopathol.* 91, 102-110.
- Secor, G. A., Rivera, V. V., Fungicide resistance assays for fungal plant pathogens. In: M. D. Bolton, B. P. H. J. Thomma, Eds.), *Plant Fungal Pathogens: Methods and Protocols*. Humana Press, New York, 2012, pp. 385-392.
- Secor, G. A., Rivera, V. V., Khan, M. F. R., Gudmestad, N. C., 2010. Monitoring fungicide sensitivity of *Cercospora beticola* of sugar beet for disease management decisions. *Plant Dis.* 94, 1272-1282.
- Šimová, Z., Poloncová, K., Tahotná, D., Holíč, R., Hapala, I., Smith, A. R., White, T. C., Griač, P., 2013. The yeast *Saccharomyces cerevisiae* Pdr16p restricts changes in ergosterol biosynthesis caused by the presence of azole antifungals. *Yeast.* 30, 229-241.
- Soustre, I., Letourneux, Y., Karst, F., 1996. Characterization of the *Saccharomyces cerevisiae* *RTA1* gene involved in 7-amincholesterol resistance. *Curr. Genet.* 30, 121-125.
- Steinkamp, M. P., Martin, S. S., Hoefert, L. L., Ruppel, E. G., 1979. Ultrastructure of lesions produced by *Cercospora beticola* in leaves of *Beta vulgaris*. *Physiol. Plant Pathol.* 15, 13-26.
- Sun, X., Xu, Q., Ruan, R., Zhang, T., Zhu, C., Li, H., 2013. PdMLE1, a specific and active transposon acts as a promoter and confers *Penicillium digitatum* with DMI resistance. *Environmental Microbiology Reports.* 5, 135-142.
- Thomma, B. P. H. J., Seidl, M. F., Shi-Kunne, X., Cook, D., Bolton, M. D., Van Kan, J. A. L., Faino, L., 2015. Mind the gap; seven reasons to close fragmented genome assemblies. *Fungal Genet. Biol.*
- Tritt, A., Eisen, J. A., Facciotti, M. T., Darling, A. E., 2012. An integrated pipeline for de novo assembly of microbial genomes. *PLoS One.* 7, e42304.
- Vik, Å., Rine, J., 2001. Upc2p and Ecm22p, dual regulators of sterol biosynthesis in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 21, 6395-6405.
- Weiland, J. J., Halloin, J. M., 2001. Benzimidazole resistance in *Cercospora beticola* sampled from sugarbeet fields in Michigan, USA. *Can. J. Plant Pathol.* 23, 78-82.
- Wu, T. D., Nacu, S., 2010. Fast and SNP-tolerant detection of complex variants and splicing in short reads. *Bioinformatics.* 26, 873-881.
- Zwiers, L.-H., Stergiopoulos, I., Van Nistelrooy, J. G. M., De Waard, M. A., 2002. ABC transporters and azole susceptibility in laboratory strains of the wheat pathogen *Mycosphaerella graminicola*. *Antimicrob. Agents Chemother.* 46, 3900-3906.

Table 1. Differentially-expressed genes¹ identified in both DMI-sensitive (09-32) and -resistant (10-73-4) *Cercospora beticola* strains after exposure to tetraconazole.

Gene	Predicted function	Log ₂ FC		E-value	Accession	InterPro	PFAM
		09-32	10-73-4				
ATP binding (GO:0005524)							
CBET3_05020-RA	AAA family ATPase	-2.39	-2.19	0.0	XP_007600097	IPR003959	PF00004
CBET3_11559-RA	AAA family ATPase	2.29	1.83	0.0	EMF10161	IPR003959	PF00004
CBET3_09831-RA	ABC transporter, ABC-B family, MDR type	2.29	2.21	0.0	XP_007927232	IPR003439	PF00005
CBET3_05101-RA	P-loop containing nucleoside triphosphate hydrolase	-1.82	-1.88	0.0	EMF08438	IPR003959	PF00004
Biological process (GO:0008150)							
CBET3_00433-RA	FAD/NAD-P-binding domain-containing protein	1.55	1.37	8.35E-64	KJX96668	IPR002938	
Biosynthetic process (GO:0009058)							
CBET3_01606-RA	Squalene synthase (ERG9)	2.63	2.08	0.0	EMF16463	IPR002060	PF00494
CBET3_09638-RA	Polyketide synthase	-2.54	1.99	0.0	XP_003848644	IPR001031	PF00975
Carbohydrate metabolic process (GO:0005975)							
CBET3_07796-RA	Glycoside hydrolase family 16 protein	-1.47	-1.57	9E-75	XP_007930006	IPR000757	PF00722
CBET3_08404-RA	Glycoside hydrolase family 29 protein	3.98	3.13	0.0	KEQ97150	IPR000933	PF01120
Catalytic activity (GO:0003824)							
CBET3_09907-RA	Lanosterol synthase (ERG7)	3.65	3.41	0.0	XP_003850188	IPR001330	PF00432
Coenzyme A biosynthetic process (GO:0015937)							
CBET3_06021-RA	Pantothenate kinase	1.69	2.29	0.0	KEQ64276	IPR004567	PF03630
Fatty acid biosynthetic process (GO:0006633)							
CBET3_06384-RA	Methylsterol monooxygenase (ERG25)	2.74	3.08	0.0	XP_003854964	IPR006694	PF04116
CBET3_09933-RA	C-5 sterol desaturase (ERG3)	3.31	2.82	0.0	KEQ58244	IPR006694	PF04116
Hydrolase activity, acting on carbon-nitrogen (but not peptide) bonds (GO:0016810)							
CBET3_03701-RA	Cyanide hydratase	2.75	1.98	0.0	CDM33757	IPR000132	PF00795
Integral component of membrane (GO:0016021)							
CBET3_07811-RA	DUF887-domain-containing protein	1.61	1.42	0.0	EMF14457	IPR006634	PF03798
CBET3_10695-RA	Potassium/sodium efflux P-type ATPase like protein	-2.09	-1.52	0.0	KJX92451	IPR001757	
CBET3_00214-RA	Sterol regulatory element-binding protein cleavage-activating	2.25	2.42	0.0	KJX95407	IPR000731	
Isoprenoid biosynthetic process (GO:0008299)							
CBET3_09160-RA	Farnesyl diphosphate synthase (FDS)	2.23	2.38	0.0	XP_003850094	IPR000092	PF00348
CBET3_06225-RA	Hydroxymethylglutaryl-CoA synthase (ERG13)	3.45	2.85	0.0	EMF11041	IPR010122	PF01154

CBET3_06742-RA	Isopentenyl-diphosphate Delta-isomerase (IDI1)	1.73	2.09	0.0	EMF09571	IPR011876	
CBET3_11582-RA	Mevalonate kinase (ERG12)	2.58	1.92	0.0	EMF10221	IPR006205	
Membrane (GO:0016020)							
CBET3_01327-RA	Delta(14)-sterol reductase (ERG24)	4.28	4.10	0.0	EMF17378	IPR001171	PF01222
Metabolic process (GO:0008152)							
CBET3_04233-RA	Acetyl-CoA C-acetyltransferase (ERG10)	3.28	3.52	0.0	XP_003852634	IPR002155	
CBET3_11684-RA	α/β hydrolase	5.21	5.88	2E-164	KJX97729	IPR013094	PF07859
CBET3_02550-RA	α/β hydrolase	4.55	1.54	0.0	EMF15641	IPR013094	PF07859
CBET3_05716-RA	Arylamine N-acetyltransferase 2	-1.98	-1.61	25E-174	CBL43313	IPR00144	PF00797
CBET3_03452-RA	Glutathione-dependent formaldehyde-activating enzyme	4.24	3.25	1E-18	EQB44398	IPR006913	PF04828
CBET3_01835-RA	NAD(P)-binding protein	4.85	5.15	24E-151	EMF16324	IPR002198	PF00106
CBET3_02822-RA	NAD(P)-binding protein	-4.23	1.52	7E-174	KEQ67599	IPR002198	PF00106
CBET3_11388-RA	NAD(P)-binding protein	-2.10	2.25	3E-103	KEQ60389	IPR002198	PF00106
CBET3_03710-RA	Pyridoxal phosphate-dependent transferase	-1.58	-1.84	0.0	EMF15374	IPR000192	PF00266
CBET3_02385-RA	Short-chain dehydrogenase/reductase	6.29	6.26	1E-75	WP_026299182	IPR002198	PF00106
CBET3_09491-RA	Short-chain dehydrogenase/reductase	1.93	2.03	4E-170	EKG11072	IPR002198	PF00106
Organic acid metabolic process (GO:0006082)							
CBET3_08424-RA	Acetate kinase	1.98	2.17	0.0	EMF13292	IPR004372	
Oxidation-reduction process (GO:0055114)							
CBET3_07492-RA	3-hydroxy-3-methylglutaryl-coenzyme A reductase (HMGCR)	1.52	1.68	0.0	EMF14786	IPR002202	PF00368
CBET3_11380-RA	4-carboxymuconolactone decarboxylase	3.20	4.56	39E-98	KJX97620	IPR003779	PF02627
CBET3_11379-RA	14- α -demethylase (CYP51/ERG11)	5.10	3.57	0.0	AEU04152	IPR001128	PF00067
CBET3_00597-RA	Alcohol dehydrogenase II	3.93	2.58	0.0	EMF17075	IPR002085	PF08240
CBET3_03149-RA	C-22 sterol desaturase (ERG5)	3.30	2.65	0.0	XP_003855007	IPR001128	PF00067
CBET3_10532-RA	Copper amine oxidase	-1.75	-1.77	0.0	EKG14169	IPR000269	PF01179
CBET3_02500-RA	Cytochrome P450	2.06	2.21	1E-168	XP_008596070	IPR001128	PF00067
CBET3_09194-RA	Cytochrome P450	4.41	5.35	0.0	EMF12198	IPR001128	PF00067
CBET3_10815-RA	Cytochrome P450	2.49	3.13	0.0	EMF12954	IPR001128	PF00067
CBET3_04462-RA	Cytochrome P450	6.15	3.05	2E-163	XP_007581051	IPR001128	PF00067
CBET3_10351-RA	D-3-phosphoglycerate dehydrogenase	1.96	1.43	1E-116	GAD98572	IPR006140	PF02826
CBET3_11680-RA	Flavin-containing amine oxidase like	3.16	4.03	0.0	KJX96742	IPR001613	PF01593
CBET3_10667-RA	Hypothetical protein with FAD-binding domain	2.69	6.73	0.0	KJX98909	IPR016166	

CBET3_11975-RA	Hypothetical protein with oxidoreductase domain	-1.87	-2.52	1E-147	KJX96592	IPR005123	PF13640
CBET3_11351-RA	O-methylsterigmatocystin oxidoreductase	-3.14	3.37	0.0	GAM33870	IPR001128	PF00067
CBET3_08405-RA	Squalene monooxygenase (ERG1)	2.00	2.20	0.0	XP_003854443	IPR013698	PF08491
CBET3_11355-RA	Zn-dependent oxidoreductase	-2.99	2.59	4E-72	WP_042369849	IPR013149	PF00107
Oxidoreductase (GO:0016491)							
CBET3_01462-RA	L-lactate dehydrogenase	1.92	1.70	0.0	EMF17491	IPR000262	PF01070
CBET3_09636-RA	Tetrahydroxynaphthalene reductase	-4.25	1.32	1E-136	XP_007931484	IPR002198	PF00106
Phosphopantetheine binding (GO:0031177)							
CBET3_11350-RA	Polyketide synthase	-3.74	2.79	0.0	AAR92210	IPR020806	
Protein binding (GO:0005515)							
CBET3_04298-RA	Dihydroxyacid dehydratase	-1.91	-1.81	0.0	EQB43606	IPR001841	
CBET3_08724-RA	F-box domain protein	-1.43	-1.98	0.0	KJX95423	IPR001810	
CBET3_10269-RA	Galactose oxidase	1.90	2.34	0.0	EMF13012	IPR006652	
CBET3_11629-RA	Glutathione s-transferase	3.94	3.76	1E-51	XP_007581437	IPR004045	PF13417
CBET3_11956-RA	Glutathione s-transferase	3.24	3.27	2E-107	KJX95768	IPR004045	PF13417
CBET3_04747-RA	Thioredoxin reductase	2.22	1.37	0.0	EMF13895	IPR001327	PF07992
Protein dimerization activity (GO:0046983)							
CBET3_00556-RA	Sterol regulatory element binding protein sre1	3.26	2.63	0.0	XP_007579378	IPR011598	
Proteolysis (GO:0006508)							
CBET3_04384-RA	Metallopeptidase MepB like protein	2.24	1.95	0.0	KJX93553	IPR001567	PF01432
CBET3_05080-RA	Serine carboxypeptidase	-1.89	-1.45	0.0	KJX94244	IPR001563	
CBET3_02384-RA	zinc carboxypeptidase like	4.05	3.56	8E-142	KJX93403		
Pyridoxal phosphate binding (GO:0030170)							
CBET3_04693-RA	2,2-dialkylglycine decarboxylase	-4.33	-1.79	0.0	AAM18795	IPR005814	
CBET3_09485-RA	Hypothetical protein with MOSC N-terminal beta barrel domain	1.76	1.53	0.0	EMF11755	IPR005302	PF03473
Regulation of transcription, DNA-templated (GO:0006355)							
CBET3_05836-RA	bZIP transcription factor	1.66	1.80	9E-138	XP_007922535	IPR004827	PF00170
CBET3_09356-RA	Zn(II) ₂ Cys ₆ fungal-type transcription factor	2.04	2.01	1E-125	EMF12171	IPR001138	PF00172
Self proteolysis (GO:0097264)							
CBET3_00643-RA	RHS Repeat protein	2.93	3.57	0.0	GAD94323	IPR022385	
Steroid biosynthetic process (GO:0006694)							
CBET3_06157-RA	Sterol 24-C-methyltransferase (ERG6)	4.84	4.00	0.0	EMF11257.1	IPR013705	PF08498

Transcription, DNA-templated (GO:0006351)

CBET3_08827-RA	Transcription elongation factor S-II	4.86	3.34	1E-29	EMR80493	IPR003618	PF07500
----------------	--------------------------------------	------	------	-------	----------	-----------	---------

Transmembrane transport (GO:0055085)

CBET3_03438-RA	Cation diffusion facilitator 1	2.28	2.58	0.0	KEQ77619	IPR002524	PF01545
CBET3_02677-RA	Hexose transporter-like protein	1.94	3.02	0.0	ENH76856	IPR003663	PF00083
CBET3_03495-RA	Lactose permease	1.69	1.54	0.0	XP_001933299	IPR003663	PF00083
CBET3_11354-RA	Major facilitator superfamily multidrug resistance transporter	-2.71	2.57	0.0	XP_960834	IPR011701	PF07690

Transport (GO:0006810)

CBET3_08499-RA	Major intrinsic protein	2.29	1.57	0.0	XP_001258974	IPR000425	PF00230
----------------	-------------------------	------	------	-----	--------------	-----------	---------

Ubiquitin-dependent protein catabolic process (GO:0006511)

CBET3_08917-RA	Cullin repeat-containing protein	-1.70	-1.75	6E-23	KEQ88354	IPR001373	PF00888
----------------	----------------------------------	-------	-------	-------	----------	-----------	---------

Unclassified

CBET3_11353-RA	2-polyprenyl-6-methoxyphenol hydroxylase	-3.40	3.20	0.0	AJW29601	IPR002938	PF01494
CBET3_08812-RA	3-keto-steroid reductase (ERG27)	2.43	2.13	0.0	EMF13199		
CBET3_11356-RA	AflY	-3.15	2.93	4E-143	ACH72894		
CBET3_00169-RA	α/β -hydrolase	3.62	3.13	8E-103	XP_007309066	IPR000073	PF12697
CBET3_09693-RA	α/β -hydrolase	3.14	2.60	7E-151	EMF12025	IPR000073	PF12697
CBET3_11622-RA	α/β -hydrolase	1.80	1.57	1E-50	KJX96893	IPR000073	PF12697
CBET3_00444-RA	Amino acid transporter like protein	2.70	2.86	0.0	KJX92975	IPR013057	PF01490
CBET3_06377-RA	C-8 sterol isomerase (ERG2)	4.58	4.74	2E-127	EMF11494.1	IPR006716	PF04622
CBET3_03624-RA	Cell wall serine-threonine-rich galactomannoprotein Mp1	3.34	2.01	3E-13	XP_746510	IPR021054	PF12296
CBET3_02310-RA	CRAL/TRIO domain-containing protein	1.78	2.36	0.0	EMF15692	IPR001251	PF00650
CBET3_10982-RA	Cyclohexanone monooxygenase	4.14	5.85	3E-174	KKA19221		
CBET3_02376-RA	DUF1768-domain-containing protein	-2.08	-3.17	6E-54	KEQ83024	IPR012816	PF08719
CBET3_08738-RA	Enoyl-CoA hydratase	3.88	5.76	1E-22	XP_001273472		
CBET3_11689-RA	Flavin-binding monooxygenase-like	4.27	4.19	0.0	EMF14477		PF13738
CBET3_01022-RA	Glyoxalase	-2.02	-1.49	8E-102	KJX97387	IPR025870	PF13468
CBET3_05419-RA	HAD-superfamily hydrolase like	1.59	1.44	6E-156	KJX99196	IPR006353	PF13344
CBET3_02577-RA	Hypothetical protein	-2.17	-1.88	1E-141	EMF15696		
CBET3_02847-RA	Hypothetical protein	-1.61	-2.01	3E-40	XP_007923422		
CBET3_02907-RA	Hypothetical protein	4.24	4.20	2E-109	EMF15962		
CBET3_03327-RA	Hypothetical protein	1.87	1.37	0.0	EMF15447		

CBET3_03903-RA	Hypothetical protein	4.41	4.49	2E-38	XP_007922746		
CBET3_05291-RA	Hypothetical protein	-1.86	-1.98	0.0	XP_007672588		
CBET3_05777-RA	Hypothetical protein	4.36	4.09	4E-94	EMF08815		
CBET3_06684-RA	Hypothetical protein	1.68	2.41	0.0	KIW92900		
CBET3_06743-RA	Hypothetical protein	2.38	2.35	3E-49	EME40860		
CBET3_07300-RA	Hypothetical protein	3.97	5.25	2E-87	EMF09684		
CBET3_08175-RA	Hypothetical protein	3.06	2.92	7E-76	EMF10381		
CBET3_08918-RA	Hypothetical protein	2.59	2.61	3E-84	XP_007793395		
CBET3_09355-RA	Hypothetical protein	2.17	2.10	2E-23	EMF12169		
CBET3_09832-RA	Hypothetical protein	3.77	3.19	0.0	EMF11737		
CBET3_10368-RA	Hypothetical protein	3.14	2.29	2E-89	EME42939		
CBET3_10369-RA	Hypothetical protein	2.87	2.19	2E-81	EME42938		
CBET3_10412-RA	Hypothetical protein	2.20	1.65	0.0	EMF12762		
CBET3_11736-RA	Hypothetical protein	-1.33	-1.77	0.0	NP_001135720		
CBET3_11756-RA	Hypothetical protein	-1.87	-1.83	1E-52	EME46083		
CBET3_12127-RA	Hypothetical protein	-5.65	-2.51	3E-25	KJX96086	IPR029226	PF14856
CBET3_09910-RA	Hypothetical protein with Rad9-like Rad53-binding domain	2.07	1.61	0.0	XP_007927419	IPR013914	PF08605
CBET3_11389-RA	Hypothetical protein with SnoL-like domain	-2.89	2.80	6E-47	EFQ36840		PF13577
CBET3_03989-RA	Integral membrane protein	1.81	1.89	1E-90	EMF10635	IPR016964	PF10914
CBET3_03405-RA	Lysine decarboxylase-like	-1.63	-1.41	5E-110	KEQ65240	IPR005269	PF03641
CBET3_07048-RA	Mitochondrial carrier protein	2.02	1.61	1E-137	EMF09651		
CBET3_10011-RA	Phosphoglycerate mutase	-2.62	-1.90	1E-134	EMF08719	IPR013078	PF00300
CBET3_03623-RA	ribose-phosphate pyrophosphokinase	3.38	1.95	4E-145	XP_008602384	IPR000073	PF12697
CBET3_01326-RA	S-adenosyl-L-methionine-dependent methyltransferase	2.48	2.40	40E-145	EMF17377	IPR029063	PF13489
No annotation available							
CBET3_01151-RA	No significant similarity	2.82	2.01				
CBET3_02311-RA	No significant similarity	2.01	2.57				
CBET3_03047-RA	No significant similarity	3.81	3.59				
CBET3_06020-RA	No significant similarity	2.97	2.86				
CBET3_06156-RA	No significant similarity	5.53	4.47				
CBET3_07294-RA	No significant similarity	2.09	2.31				
CBET3_10038-RA	No significant similarity	-2.38	-2.94				

*Biological process GO term ontology was given preference when grouping genes. If no Biological Process GO term was identified for a gene, Molecular function and Cellular component GO terms were used for classification.

¹Genes were considered differentially expressed if Q-values ≤ 0.001 and $\log_2$ fold regulation differences ≥ 1.0 using *t*-tests of expression values between mock- and tetraconazole-treatments for each *C. beticola* strain.

Table 2. Differentially-expressed genes¹ unique to *Cercospora beticola* DMI-resistant strain 10-73-4 after exposure to tetraconazole.

Gene	Predicted function	Log ₂ FC	E-value	Accession	InterPro	PFAM
ATP binding (GO:0005524)						
CBET3_00527-RA	AAA family ATPase	-1.55	4E-83	XP_007921050	IPR003959	PF00004
Biosynthetic process (GO:0009058)						
CBET3_06542-RA	Phospho-2-dehydro-3-deoxyheptonate aldolase	-1.80	0.0	EMF11415	IPR006218	PF00793
CBET3_11935-RA	Serine palmitoyl CoA transferase subunit LcbA	-2.41	0.0	XP_003849027	IPR004839	PF00155
Carbohydrate metabolic process (GO:0005975)						
CBET3_06057-RA	Beta glucan-binding protein	1.73	7E-155	EMR86206	IPR000757	PF00722
CBET3_00682-RA	Glycoside hydrolase family 16	3.79	0.0	EMF17292	IPR000757	PF00722
CBET3_05663-RB	Glycoside hydrolase family 16	3.48	1E-134	EMF17702	IPR000757	PF00722
CBET3_01742-RA	Glycoside hydrolase family 16	1.85	6E-161	EMF16313	IPR000757	PF00722
CBET3_07968-RA	Glycoside hydrolase family 17	1.95	1E-177	EMF13167	IPR000490	PF00332
CBET3_09944-RA	Glycoside hydrolase family 17	1.59	6E-156	EMF08721	IPR000490	
CBET3_09115-RA	Glycoside hydrolase family 18	1.68	1E-158	EMF11844	IPR001223	PF00704
CBET3_10389-RA	Glycoside hydrolase family 72	1.95	0.0	XP_007927960	IPR004886	PF03198
CBET3_00012-RA	Transaldoase	4.38	1E-157	KJZ75321.1	IPR001585	PF00923
Catalytic activity (GO:0003824)						
CBET3_06881-RA	Glycoside hydrolase family 92	-2.40	0.0	EMF09713	IPR012939	PF07971
Heme binding (GO:0020037)						
CBET3_11439-RA	Indoleamine 2,3-dioxygenase	1.88	0.0	EMF10116	IPR000898	PF01231
Hydrolase activity (GO:0016787)						
CBET3_10362-RA	5'/3'-nucleotidase SurE family	2.12	2E-107	XP_001267521	IPR002828	PF01975
CBET3_09066-RA	Carbon-nitrogen hydrolase	1.59	1E-135	EMF12014	IPR003010	PF00795
CBET3_11455-RA	Metallo-dependent hydrolase	-1.41	1E-100	EHA27811	IPR006680	PF01979
Integral component of membrane (GO:0016021)						
CBET3_04709-RA	GPR1/FUN34/YaaH-class plasma membrane protein	1.73	1E-157	KJX99469	IPR000791	PF01184
Lipid metabolic process (GO:0006629)						
CBET3_07897-RA	α/β hydrolase	3.68	2E-148	EMF14093	IPR002921	PF01764
Metabolic process (GO:0008152)						
CBET3_02504-RA	17- β -hydroxysteroid dehydrogenase	4.51	1E-166	3IS3_A	IPR002198	PF00106

CBET3_12252-RA	Carboxypeptidase S1	1.93	0.0	EGE05173	IPR001563	PF00450
CBET3_08150-RA	Trypsin-like serine protease	1.64	1E-57	XP_006961656	IPR001254	PF00089
Regulation of transcription, DNA-templated (GO:0006355)						
CBET3_02344-RA	Zn(II) ₂ Cys ₆ transcription factor (AflR)	3.28	1E-24	KKA18106	IPR002409	PF08493
Response to stress (GO:0006950)						
CBET3_11653-RA	RTA1 domain protein	8.00	2E-117	KJX94617	IPR007568	PF04479
Steroid biosynthetic process (GO:0006694)						
CBET3_00902-RA	C-3 sterol dehydrogenase (Erg26)	1.47	0.0	EMF17180	IPR002225	PF01073
Transmembrane transport (GO:0055085)						
CBET3_02206-RA	Hexose transporter	-2.20	0.0	ACR78273	IPR003663	PF00083
CBET3_08810-RA	Histidine permease	-1.77	0.0	EMF13360	IPR004841	PF00324
CBET3_06018-RA	Major facilitator superfamily transporter	2.44	6E-138	XP_007274312	IPR011701	PF07690
CBET3_01824-RA	Major facilitator superfamily transporter	2.26	0.0	XP_003065775	IPR011701	PF07690
CBET3_03348-RA	Major facilitator superfamily transporter	1.34	1E-167	EME48032	IPR011701	PF07690
CBET3_10856-RA	Major facilitator superfamily general substrate transporter	-3.13	0.0	EMF09454	IPR011701	PF07690
CBET3_11268-RA	Major facilitator superfamily general substrate transporter	1.54	0.0	EMF09835	IPR011701	PF07690
CBET3_04140-RA	Peptide transporter	-1.72	0.0	ENH84505	IPR004813	PF03169
Transport (GO:0006810)						
CBET3_05001-RA	ABC transporter	5.52	0.0	EMF08365	IPR003439	PF00005
CBET3_03349-RA	ABC transporter, ABC-A family	1.48	0.0	XP_007924170	IPR026082	PF00005
Ubiquitin-dependent protein catabolic process (GO:0006511)						
CBET3_10720-RA	Cullin repeat-containing protein	-1.87	2E-32	KEQ88354	IPR001373	PF00888
Unclassified						
CBET3_05042-RA	Acetyltransferase-like protein	-2.32	4E-90	KJX94255	IPR016181	
CBET3_06118-RA	Autophagy protein	-1.40	0.0	EMF11171	IPR024671	PF11700
CBET3_07833-RA	Beta lactamase-like protein	-3.16	0E+00	EME48565	IPR001466	PF00144
CBET3_11652-RA	C6 zinc finger domain-containing protein	3.33	3E-28	KJX94621		
CBET3_11585-RA	Carbohydrate esterase family 3 protein	4.93	2E-99	EMF10206	IPR013830	PF13472
CBET3_11060-RA	Fasciclin domain-containing protein	1.89	3E-48	XP_007295171	IPR000782	PF02469
CBET3_00365-RA	Glycosyl transferase	-2.53	9E-83	EKV16648	IPR029044	
CBET3_05558-RA	Hypothetical protein	4.83	8E-159	EMF08800		
CBET3_00511-RA	hypothetical protein	2.97	9E-93	EME49690		

CBET3_10668-RA	Hypothetical protein	2.73	1E-98	XP_007928244		
CBET3_05254-RA	Hypothetical protein	2.27	0.0	EMF08527		
CBET3_03147-RA	Hypothetical protein	2.15	8E-18	EMR84685		
CBET3_02508-RA	Hypothetical protein	2.10	1E-166	XP_007696463	IPR025337	PF14027
CBET3_06857-RA	Hypothetical protein	1.91	5E-115	XP_007930785		
CBET3_01561-RA	Hypothetical protein	1.79	3E-22	XP_001932896	IPR011050	
CBET3_08337-RA	Hypothetical protein	1.77	1E-80	EMF13537		
CBET3_09241-RA	Hypothetical protein	1.72	3E-90	KJX99495	IPR010730	PF06985
CBET3_01612-RA	Hypothetical protein	1.79	5E-11	EMF16235		
CBET3_06982-RA	Hypothetical protein	2.60	1E-94	EME39020		
CBET3_02518-RA	Hypothetical protein	-1.42	3E-138	EMF16076	IPR025187	PF13430
CBET3_02546-RA	Hypothetical protein	-1.53	2E-15	XP_007923133		
CBET3_09459-RA	Hypothetical protein	-1.58	0.0	KJX99536	IPR021840	PF11915
CBET3_11415-RA	Hypothetical protein	-1.60	0.0	BAN91486		
CBET3_11454-RA	Hypothetical protein	-1.61	0.0	EMF09894	IPR009571	PF06687
CBET3_11257-RA	Hypothetical protein	-1.65	5E-75	EMF10176	IPR018392	PF01476
CBET3_12197-RA	Hypothetical protein	-1.76	3E-104	EQL33191		
CBET3_01494-RA	Hypothetical protein	-1.82	1E-33	EWG50720		
CBET3_08277-RA	Hypothetical protein	-1.80	4E-75	EMF10352		
CBET3_02870-RA	Hypothetical protein	-1.84	3E-66	EMF15383		
CBET3_05929-RA	Hypothetical protein	-2.14	6E-13	EME46074		
CBET3_10955-RA	Hypothetical protein	-2.13	4E-179	EMF09808		
CBET3_05246-RA	Hypothetical protein	-2.23	2E-21	XP_007777459		
CBET3_05560-RA	Hypothetical protein	-2.22	2E-130	EMF08810		
CBET3_10855-RA	Hypothetical protein	-1.76	1E-14	EMF09452		
CBET3_05304-RA	Hypothetical protein	-2.45	6E-151	EME38618		
CBET3_03830-RA	Hypothetical protein	-3.29	1E-104	XP_003857461	IPR007310	PF04183
CBET3_00215-RA	Hypothetical protein with HET domain	1.29	9E-29	EWZ48880	IPR010730	PF06985
CBET3_03842-RA	Hypothetical protein with tetratricopeptide repeat domain	2.99	1E-104	XP_002118036	IPR001440	PF00515
CBET3_00212-RA	Major facilitator superfamily, general substrate transporter	4.31	2E-117	KGO53278	IPR020846	
CBET3_05545-RA	NPP1 domain-containing protein	1.64	8E-120	XP_007588068	IPR008701	PF05630
CBET3_04143-RA	P-loop containing nucleoside triphosphate hydrolase	1.58	4E-145	EMF10878	IPR027417	PF09439

CBET3_00211-RA	Siderophore esterase IroE-like	4.76	2E-125	XP_001276792	IPR000801	PF00756
CBET3_03652-RA	ThiJ/Pfpl family protein	-1.85	5E-37	KFG86071	IPR025628	PF13278
CBET3_02345-RA	Toxin biosynthesis regulatory protein	2.75	2E-65	CBF90109	IPR011991	
CBET3_11414-RA	Zn-dependent hydrolase	-2.17	4E-155	ENH83839	IPR001279	PF00753
No annotation available						
CBET3_05717-RA	No significant similarity	1.34				
CBET3_02219-RA	No significant similarity	2.53				
CBET3_02217-RA	No significant similarity	2.40				
CBET3_09434-RA	No significant similarity	-1.82				
CBET3_03832-RA	No significant similarity	-2.26				
CBET3_10751-RA	No significant similarity	-2.27				
CBET3_02070-RA	No significant similarity	-2.27				
CBET3_10954-RA	No significant similarity	-3.19				

*Biological process GO term ontology was given preference when grouping genes. If no Biological Process GO term was identified for a gene,

Molecular function and Cellular component GO terms were used for classification.

¹Genes were considered differentially expressed if Q-values ≤ 0.001 and $\log_2$ fold regulation differences ≥ 1.0 using *t*-tests of expression values between mock- and tetraconazole-treatments.

Table 3. EC₅₀ analyses of *Cercospora beticola* strains containing ectopic or site-directed mutations in candidate DMI-resistance genes.

Gene	Predicted function	Average EC ₅₀ value (µg ml ⁻¹)*	
		Ectopic	Site-directed
CBET3_11380-RA	4-Carboxymuconolactone decarboxylase	52.2 (± 13.7)	57.8 (±9.1)
CBET3_11653-RA	RTA1 domain protein	48.7 (±7.1)	52.5 (±10.5)
CBET3_11585-RA	Carbohydrate esterase family 3 protein	56.1 (±3.1)	60.2 (±16.4)
CBET3_11652-RA	C6 zinc finger domain-containing protein	53.4 (±5.7)	51.0 (±6.1)
CBET3_02344-RA	C6 transcription factor (AflR)	51.9 (±8.4)	55.9 (±10.5)

*The EC₅₀ value of the progenitor strain 10-73-4 was 35.9 (±2.6) µg ml⁻¹. Values in parenthesis indicate standard deviation of the mean of three individual mutants.

Table 4. Primers used in this study.

Primer name	Specificity/role*	Sequence (5' to 3')
MDB-277	Split-marker HYG-F	GACGTTGTAACGACGGCCAGTG
MDB-258	Split-marker HY-R	GGATGCCCTCCGCTCGAAGTA
MDB-259	Split-marker YG-F	CGTTGCAAGACCTGCCTGAA
MDB-278	Split-marker HYG-R	CACAGGAAACAGCTATGACCATGA
MDB-1145	HYG R2	GGCAGGTAGATGACGACCAT
MDB-1317	CBET3_11653-RA 1F	TCTGAACCCTCAACGAGTCC
MDB-1318	CBET3_11653-RA 2R	CACTGGCCGTCGTTTTACAACGTCGTTGCCGACACTTCTGTACA
MDB-1319	CBET3_11653-RA 3F	TCATGGTCATAGCTGTTTCCTGTGTTCCCGTCTCCAATGCTTCT
MDB-1320	CBET3_11653-RA 4R	GTCTCTCCGCTTGCCAATC
MDB-1429	CBET3_11653-RA 5F	AGTGCCTGGTCCTTGATCAA
MDB-1396	CBET3_11653-RA 6R	AACGGAAGCATGCAAGGT
MDB-1325	CBET3_11585-RA 1F	GGCGTTTTCTGGCATAACGAA
MDB-1326	CBET3_11585-RA 2R	CACTGGCCGTCGTTTTACAACGTCGTCGCTGACTTGATTGCTT
MDB-1327	CBET3_11585-RA 3F	TCATGGTCATAGCTGTTTCCTGTGAGGCTACCGAGGAGATGGTA
MDB-1328	CBET3_11585-RA 4R	CCTGGCAGATGTCTCCTCAA

MDB-1431	CBET3_11585-RA 5F	CAATCAACAGATCAGCGCGA
MDB-1398	CBET3_11585-RA 6R	GTCTCCGTGTTTCAGCGATC
MDB-1333	CBET3_11652-RA 1F	CTGTGTAGGCCTCGACTTTC
MDB-1334	CBET3_11652-RA 2R	CACTGGCCGTCGTTTTACAACGTCCCGGGCTGATAAGGTGGAAT
MDB-1335	CBET3_11652-RA 3F	TCATGGTCATAGCTGTTTCCTGTGAGAAGTTGTACACCTGGGCA
MDB-1336	CBET3_11652-RA 4R	CAGCACCTCGCATTCTTC
MDB-1433	CBET3_11652-RA 5F	ATCACAGGAAGAACGGCTCA
MDB-1400	CBET3_11652-RA 6R	TTCACCGTAGCCCAAGTGAT
MDB-1337	CBET3_02344-RA 1F	CTTCTCGAACATTGGGCCAC
MDB-1338	CBET3_02344-RA 2R	CACTGGCCGTCGTTTTACAACGTCTCTTGGTAATTGGAGGCGGT
MDB-1339	CBET3_02344-RA 3F	TCATGGTCATAGCTGTTTCCTGTGCACGCGCTATCATTCAAGTCG
MDB-1340	CBET3_02344-RA 4R	TACTTCCGACACGAAGCCAT
MDB-1434	CBET3_02344-RA 5F	AACCTCTTCTCGGGACACTG
MDB-1401	CBET3_02344-RA 6R	CCGCTAAGAATGCTGTCACC
MDB-1297	CBET3_11380-RA 1F	ACAGGAGGAAGAAGGTGGTG
MDB-1298	CBET3_11380-RA 2R	CACTGGCCGTCGTTTTACAACGTCTTACCATCATCGGCTGTGC
MDB-1299	CBET3_11380-RA 3F	TCATGGTCATAGCTGTTTCCTGTGCAGCTTCCAGGACCACAAC
MDB-1300	CBET3_11380-RA 4R	GCAAAGAGAGTGATAGCCGC
MDB-1422	CBET3_11380-RA 5F	GAACTTGACGGGTCGATGC
MDB-1391	CBET3_11380-RA 6R	CAGTCCACTCATACGTTGCG

*Primer nomenclature and design essentially as described by Goswami (2012).

FIGURE LEGENDS

Figure 1. Overlap of genes identified as differentially-expressed (Q-value ≤ 0.001 , and $\log_2$ fold regulation differences ≥ 1.0) between mock- and tetraconazole-treatments for each *Cercospora beticola* strain. *C. beticola* strain 09-32 is DMI-sensitive and 10-73-4 is DMI-resistant.

Figure 2. Ergosterol pathway in *Cercospora beticola*. **A.** DMI-exposure in both strains causes an induction (red) of ergosterol pathway genes. **B.** Fragments per kilobase of transcript per million (FPKM) reads for each gene identified as differentially-expressed in the ergosterol pathway in both *C. beticola* strains. *C. beticola* strain 09-32 is DMI-sensitive and 10-73-4 is DMI-resistant. Values represent the average FPKM values of two biological replicates. Error bars are standard deviations obtained via CummeRbund.

Figure 1.

Figure 2.