REVIEW

Integrating physiology, behavior, and energetics: biologging in a free-living arctic hibernator

Cory T. Williams^a, Brian M. Barnes^b, C. Loren Buck^{a,*}

^aCenter for Bioengineering Innovation and Department of Biological Sciences, Northern Arizona University, Flagstaff AZ, 86011, USA

^bInstitute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, AK 99775, USA,

Running title: Biologging in an arctic hibernator

Ms. Has 23 pages and 5 figures

*Author for Correspondence:

C. Loren Buck

Center for Bioengineering Innovation

Northern Arizona University

Box: 4185

Flagstaff AZ 86011

Phone: 928-523-5976

ABSTRACT

The use of animal-borne instruments (ABIs), including biologgers and biotransmitters, has played an integral role in advancing our understanding of adjustments made by animals in their physiology and behavior across their annual and daily cycles and in response to weather and environmental change. Here, we review our research employing body temperature (T_b), light, and acceleration biologgers to measure patterns of physiology and behavior of a free-living, semi-fossorial hibernator, the arctic ground squirrel (*Urocitellus parryii*). We have used these devices to address a variety of physiological, ecological, and evolutionary questions within the fields of hibernation physiology, phenology, behavioral ecology, and chronobiology. We have also combined biologging with other approaches, such as endocrinology and tracking the thermal environment, to provide insights into the physiological mechanisms that underlie fundamental questions in biology including physiological performance trade-offs, timing and functional energetics. Finally, we explore the practical and methodological considerations that need to be addressed in biologging studies of free-living vertebrates and discuss future technological advancements that will increase the power and potential of biologging as a tool for assessing physiological function in dynamic and changing environments.

Key Words

Accelerometry; Biologger; Body temperature logger; Circadian rhythms; Ground squirrel; Light logger; Overall Dynamic Body Acceleration; Phenology; Thyroid Hormone; Torpor

Abbreviations: ABI, Animal-Borne Instruments; AGS, Arctic Ground Squirrel; ECG, Electrocardiogram; T_b , Body Temperature; SCN, Suprachiasmatic Nucleus; T3, Triiodothyronine; T4, Thyroxine

1. Introduction

Determining how individuals function within and interact with their environment is critical to understand behavioral and physiological adaptations and plasticity and by extension, limits to organismal responses to environmental change. The importance of understanding animal-environment interactions and physiological function is evidenced by the emergence of a new field in biology, known as “conservation physiology”, which seeks to incorporate this knowledge into ecological models designed to predict population, community and ecosystem-level responses to environmental and land use change (Cooke et al. 2013). The physiological mechanisms that underlie phenotypic plasticity remain elusive, however, and therefore represent a significant impediment to predicting the ecological effects of a changing climate (Denny & Helmuth 2009).

Controlled laboratory experiments will always play an instrumental role in environmental physiology, but only by investigating physiological function and behavior in free-living individuals will we be able to fully understand how individuals regulate various aspects of their annual cycle including, as examples, reproduction, migration, molt, and hibernation (Bartholomew 1986). It is well known that captivity can result in psychological and physiological stresses in animals as evidenced by impaired function of the hypothalamus-pituitary-adrenal axis, depressed immune function, reduced growth rates, and disturbed reproductive cycling (e.g., Romero and Wingfield 1999, Morgan and Tromborg 2007, Buehler et al. 2008). Limitations in our ability to replicate natural habitat and natural diets in captive situations constrains the usefulness of data collected in a laboratory setting to develop a thorough understanding of physiological responses to environmental challenges under natural conditions. For these reasons, studies of free-living animals are central to understanding how individuals alter their physiology and behavior across their annual cycle and in response to unpredictable environmental change.

Biologging and biotelemetry, which involve the collection of data from animal-borne instruments (hereafter: ABIs), are advancing our fundamental understanding of physiological adaptation and responsiveness to environmental change. Specifically, this technology allows us to answer questions regarding physiology, behavior, and ecology of free-living animals under natural conditions that would have previously been limited to tests on model organisms under controlled conditions. To date, the greatest impact of ABIs in ecology and environmental physiology has been through the use of devices that enable quantitative measurement of animal movement through space and time, particularly via global positioning systems (GPS), satellite telemetry, and geolocators (Rutz and Hays 2009, Cagnacci et al. 2010). However, a wide variety of ABIs have been developed that measure a range of physiological, behavioral, and environmental parameters including, but not limited to, body temperature (T_b), heart rate, acceleration, pressure (depth), salinity, light, heat flux, EEG, and PO_2 (Butler et al. 2004, Block 2005, Vyssotski et al. 2006, McDonald & Ponganis 2013). Some of these parameters, such as heart rate and

acceleration, correlate strongly with metabolic rate which provides insight into how metabolism and daily energy expenditure are influenced by both intrinsic and extrinsic factors, including weather (Green et al. 2009, Halsey et al. 2011). Combining biologging with other techniques, such as endocrinology and immunology, allows for an integrative approach to understanding how physiology mitigates the effects of environmental change and influences behavior, performance, and ultimately lifetime fitness (Wingfield et al. 1997, Semeniuk et al. 2009). Thus, the use and further development of ABIs is critical to understanding the complex interactions between physiology, behavior, climate, and the environment (Evans et al. 2016).

Here, we review our use of ABIs that measure temperature, light, and acceleration in a free-living semi-fossorial hibernator, the arctic ground squirrel (hereafter: AGS; *Urocitellus parryii*). First we provide a brief overview of the practical and methodological issues associated with the use of these particular ABIs. Then, we use our work on AGSs as an example to illustrate the diversity of physiological and ecological questions that can be addressed using these relatively simple devices. We also demonstrate how combining biologging with endocrinology can provide insight into the functional mechanisms that underlie individual differences in behavior and physiology. The promise and power of ABIs is enormous and continued technological advancements of ABIs will undoubtedly provide for a more integrative understanding of the physiological and behavioral mechanisms that underlie vertebrate responses to environmental change.

2. ABIs – Practical and Methodological Considerations

According to “Moore’s Law”, which is actually an empirical observation, the number of transistors on an integrated circuit doubles approximately every 18 months and this has important implications for processing speeds and memory (Schaller 1997). Battery technology is also improving, though at a much slower rate, such that the battery has become the greatest limiting technical factor in terms of continuing to reduce the size, mass and lifespan of ABIs. Nevertheless, the combined evolution of processing speed, storage capacity and battery technology has resulted in modern devices that are much smaller and better performing than their predecessors of just a few years ago. As a result, technologies that were once only suitable for deployment on relatively large animals, such as accelerometry, are now being used on increasingly smaller organisms (Rutz and Hayes 2009, Brown et al. 2013). This has led to widespread use of ABIs, but there are a variety of practical and methodological considerations that need to be addressed when initiating an ABI-study. These considerations include cost and robustness of the device, sampling resolution, method of attachment, capture/handling effects, effects of the device itself, and methods of analysis. Despite the continued progress in miniaturizing and improving ABI technology, the commercial demand is relatively small and thus further development/refinement of this technology requires continued

support from funding agencies. The high cost of cutting-edge ABIs also means that many researchers continue to use older, more affordable technologies. Selection of an appropriate sampling interval, resolution and method of analyses is also very important and dependent on the question being addressed; this subject is too lengthy to discuss here, but see Ropert-Coudert & Wilson (2004).

Whether an ABI is worn externally or implanted is an important consideration. In general, we suggest following the taxon-specific guidelines provided by various societies (birds: Fair et al. 2010; fish: Nickum et al. 2004; mammals: Sikes et al. 2011; reptiles and amphibians: Beaupre et al. 2004). Physical characteristics of the device such as its size, shape, mass, and buoyancy require important consideration and are dependent upon the size and ecology of the species; drag and buoyancy, for example, are much more important than mass for aquatic animals (Ponganis 2007). Mass is much more important for flying birds and bats than it is for animals that use terrestrial locomotion (Barron et al. 2010, Rojas et al. 2010, O'Mara et al. 2014). It should also be noted that even within a particular taxonomic group, some species may be more sensitive to capture/handling stress and/or carrying the ABIs, and therefore attempts should be made to validate a lack of deleterious effects on natural physiology and behavior of a given species (e.g., Whidden et al. 2007, Jepson et al. 2015). In some cases the decision regarding implantation or external-mounting will depend on the ABI itself. ABIs that measure environmental parameters (e.g., light) will obviously need to be mounted externally whereas some ABIs that measure physiological variables (e.g., PO₂) must be implanted; however, many ABIs can be either mounted externally or implanted. One should not assume that external devices are necessarily preferred simply because they avoid invasive surgeries as some externally-affixed ABIs may be more likely to cause negative behavioral effects (Saraux et al. 2011). In a recent meta-analysis of biologging effects in birds, White et al. (2013) found that externally attached, but not implanted ABIs, were consistently detrimental to the birds. They concluded that implantation of ABI's is preferable to external attachment, provided the risk of mortality associated with the anaesthesia and surgery required for implantation can be mitigated. We are not aware of a similar analysis for mammals, but when deploying collar-mounted devices it is crucial to be aware that episodes of growth or fattening must be accounted for in the design of the attachment to minimize irritation, abrasion, and infection (Sikes et al. 2011). A collar that is affixed too loosely can also cause harm if it is prone to becoming caught on sticks, branches, or other foreign objects. The choice of whether an ABI is implanted or affixed-externally will also depend on how the device downloads and its battery life and memory capacity which influences the length of deployment (see "The Future of Biologging" below).

2.1 Measuring Body Temperature (T_b)

On February 16, 1959, Dr. Robert Rausch crawled into the winter den of a black bear in Anchorage, Alaska and obtained a measurement of its rectal T_b during hibernation (Hock 1960). The single data point he obtained from this presumably nerve-racking event confirmed earlier studies of captive bears indicating these large carnivores do not exhibit the substantial reductions in T_b observed in small mammals (Hock 1951, Lyman & Chatfield 1956, Bartholomew & Cade 1957, Hock 1960). Needless to say, technological advances in biotelemetry and biologging have greatly increased our ability to obtain relatively continuous measurements of T_b in free-living animals throughout their annual cycles (e.g., Boyer and Barnes, 1999, Körtner et al. 2000, Dausman 2005, McCafferty et al. 2015). ABIs that measure T_b have played a particularly important role in studies designed to determine if, when and which species exhibit heterothermy, including hyperthermia (Fuller et al., 1999) and daily torpor and hibernation, and determining the conditions under which heterothermy is employed (reviewed in Ruf and Geiser 2015). Recent studies have also begun to focus on the use of a controlled moderate decrease in T_b as an energy-saving mechanism in animals typically regarded as strict endotherms (Arnold et al. 2004, Hetem et al. 2014). In addition, because daily cycles of T_b are generated by the master circadian clock in the suprachiasmatic nucleus (SCN; Buhr and Takahashi 2010), T_b is a potentially useful proxy for assessing the persistence of rhythms in free-living individuals (Williams et al 2012a).

Early laboratory studies of torpor and hibernation were reliant on captive animals because T_b was measured using a thermocouple inserted into the rectum (e.g., Benedict and Lee 1938) or implanted subcutaneously (Hock 1951) and connected to a large external chart recorder that was not attached to the animal. While these studies provided important insight into the physiology of heterothermy in species that exhibit daily torpor or prolonged hibernation, they were limited to the laboratory because animals were tethered to the external data recorder. Further, the disruptive nature of the process meant that measured patterns of heterothermy were not absolutely indicative of natural patterns. Captivity itself can also influence thermoregulation in unexpected ways (Geiser et al. 2000). For example, captive hibernators will often go through a step-wise series of “test-drops” - short (<24 h) bouts of torpor with consecutively lower T_b interrupted by arousals to euthermic T_b between each short torpor bout - prior to entering their first multiday torpor; these test-drops were postulated to be necessary for resetting or reprogramming the hypothalamic regions controlling thermoregulation (Strumwasser 1958, Hammel 1967, Luecke and South 1972; Pivorum 1976). However, these “test-drops” rarely occur in free-living animals and appear to be a side-effect of captivity itself (Sheriff et al. 2011), at least in some species.

In the late 80’s and early 90’s, field researchers began to deploy newly developed and smaller T_b loggers and transmitters that could be either surgically implanted into the abdominal cavity or affixed to a collar (e.g., Young 1990, Brigham 1992, Michener 1992). Widespread use of these devices has led to a variety of discoveries, from supercooling of body fluids in hibernating AGS (Barnes 1989) to hibernation

in tropical primates (Dausmann et al. 2004) to a species of dormouse that can hibernate for 11 months of the year when there is insufficient food for reproduction (Hoezl et al. 2015). Biologging of T_b has also had important conservation implications; mortality associated with white nose syndrome in bats, for example, has been shown to be associated with the increased energetic costs of frequent arousals from torpor, which, in turn, is driven by hypotonic dehydration associated with wing damage (Warnecke et al. 2013).

Technological advances are leading to the development of even smaller devices with a range of capabilities (see Future of Biologging below). Nevertheless, as previously mentioned, a key consideration for many field biologists is cost of device. For example, despite their fairly poor resolution and relatively small memory capacity, Thermochron iButtons (Maxim Integrated, Dallas, TX, USA), which first became available almost 2 decades ago, remain the most commonly deployed device for measuring T_b in small mammals. The largest capacity iButton that measures across the range of T_b characteristic of heterotherms (DS1922L) can collect only 4096 datapoints at its highest resolution setting (0.125°C) and 8192 datapoints at the low resolution setting (0.5°C). Preference for iButtons stems from their low cost ($\sim \$55/\text{device}$) and relatively long battery life ($\sim 5\text{yr}$); cost is low because these devices have commercial applications and are manufactured in bulk. To accommodate the use of these devices on smaller animals, particularly bats, researchers have developed methods of modifying the devices to decrease their size and mass (Robert & Thompson 2003, Lovegrove 2009) and to insulate them because they emit ultrasonic noise (Willis et al. 2009). Much smaller temperature sensitive transponders have also been developed, but these devices must be in close proximity to an external reader ($\sim 1\text{m}$) and the error in measured values increases at low T_b (Wacker et al. 2012, Langer and Fietz 2014).

In our research, we have used modified TidBits (12–13g, Stowaway model TBICU32-05+44, coated weight 14–15 g, Onset Computer Corp, Pocasset, MA) which have greater memory and resolution (32520 datapoints with 0.2°C resolution), as well as the smaller iButtons (3g; 4–5g coated), with our selection of devices being dependent on our research question; the modified TidBits are no longer being produced, however. To assess whether very low amplitude circadian rhythms persist during deep torpor we have also recently deployed very high capacity, high resolution dataloggers (TidBit v2, Onset Computer Corp) that can measure 42000 datapoints at 0.02°C resolution (Fig. 1). These devices are more expensive ($\$132$) and quite large (23g) such that they are not generally suitable for small mammals. Again, the selection of an appropriate ABI will depend on the question being addressed.

In addition to selecting the most appropriate device, one must consider whether to a) surgically implant the device in the abdominal cavity, b) implant the T_b logger/transmitter subcutaneously, or c) affix the device to adjacent to the skin. Several studies have demonstrated that skin temperature closely tracks core T_b during heterothermy although ambient temperature can have slight, but significant effects

on skin temperature (Barclay et al. 1996, Körtner and Geiser 2000, Dausmann 2005). Differences between core T_b and skin temperature are also expected to be evident during interbout arousals during hibernation with the periphery warming later and cooling faster, although this difference is likely not detectable at the sampling resolution used in most field studies. Skin temperature is also likely to be much less reflective of core T_b in animals exposed to highly fluctuating environmental conditions. Even for core T_b , no gold standard exist for measuring deep-body temperature though measurements should be from sites that track temperature changes in the central volume (Taylor et al. 2014). Variation in T_b can be substantial both within the body and over time, particularly in species for which the heterothermic response involves peripheral cooling, reductions in thermal core volume, and/or temporary abandonment of high core T_b (Eichhorn et al. 2011). Overall, the determination of whether to use an implanted or externally mounted T_b logger will depend on the question(s) being addressed, the potential for bias due to methods of attachment or surgery, and which approach does the least harm to the animal.

2.2 Light Loggers and Accelerometers – Quantifying Activity and Energy Expenditure

Light sensitive radio-transmitters were first utilized to measure patterns of above-ground activity by Hut et al. (1999) who showed that European ground squirrels (*Spermophilus citellus*) are below ground in early morning and evening and thus do not perceive the rapid natural light-dark transitions that occur at civil twilight. Long et al. (2005) subsequently used these devices, in combination with T_b loggers, to understand how environmental conditions and activity patterns influenced core T_b in free-living AGS. Despite the utility of light-sensitive radio-transmitters (reviewed in Long et al. 2007), use of these devices never became widespread, likely because of their large size (~10 g) and expense, as well as the significant challenges associated with the continuous collection of telemetry data in heavily vegetated or topographically complex environments. However, these issues have been overcome with the development of lightweight and rugged light loggers that were originally designed for the geolocation of migratory birds based on the timing of dawn and dusk (Phillips et al. 2004). We have used these devices affixed to collars to measure patterns of above-ground activity in AGS (Williams et al. 2012b) as well as patterns of nest use in an arboreal rodent, the red squirrel (*Sciurus vulgaris*; Williams et al. 2014b). Our experience to date suggests that most of these devices are not sensitive enough for measuring detailed activity patterns in nocturnal rodents.

Light loggers are useful in assessing timing and duration of above-ground activity (or outside of a nest), but they are not necessarily informative as to the intensity of activity of an animal while it is aboveground. To estimate intensity of activity, accelerometers, which typically measure acceleration along 3 orthogonal axes, are potentially useful (Wilson et al. 2006, Halsey et al. 2009, Brown et al. 2013). Use of accelerometers generally falls into two categories. The first approach involves quantifying fine-

scale movements and body postures using high resolution sampling (>10hz) to deduce specific behaviors. This approach requires validation data (direct observations of behavior) combined with an automated system to categorize waveform patterns and assign them to different behaviors; although manual assignment is sometimes used, a variety of machine-learning algorithms have been devised to convert waveforms to behavioral categories (reviewed in Brown et al. 2013). The second approach is to utilize accelerometry data as an index of energy expenditure. The relationship between body acceleration and oxygen consumption (as a proxy for metabolic rate) has been examined in a wide variety of taxa and, although the strength of the relationship varies somewhat, it has held across all species examined to date (reviewed in Brown et al. 2013). The most commonly used acceleration-based index of metabolic rate is Overall Dynamic Body Acceleration (ODBA), which involves removing the static effect of gravity for each axis and then summing the absolute values of dynamic acceleration for each axis (Wilson 2006). Although ODBA is typically calculated using measurements of acceleration at a frequency of 10hz or higher, sampling frequencies as low as 1 Hz provide reliable estimates of energy expenditure, even in small animals (Halsey et al. 2009). While accelerometers have been long used in the study of medium to large-sized animals, including humans, reductions in the size of these devices have recently enabled deployments on small mammals. We are currently using axy-3 accelerometers (2g; Technosmart, Italy) affixed to collars on free-living AGS; at a 10 Hz setting, we obtain up to 40 days of data on a fully charged battery.

2.3 The Future of Biologging

In many cases, researchers have switched from biotelemetry to biologging to limit practical difficulties related to interference in the reception of transmitted data and to reduce the mass or size of devices deployed by eliminating the battery demands and antenna associated with data transmission. The move to biologgers, in combination with technical advances that have greatly miniaturized circuit boards, memory, and batteries, have enabled the deployment of increasingly sophisticated devices on smaller organisms (Rutz & Hays 2009). Further, the development of individual devices that can measure many physiological and behavioral parameters simultaneously is providing greater insight into how animals take advantage of predictable change, as well as cope with environmental stochasticity (Wilson et al. 2008). However, a key limitation of biologgers is that the animal must be recaptured to obtain the data. This limitation is particularly severe for implanted devices, because a second surgery must be performed to recover the device.

In addition to biologgers and biotelemetry devices, a hybrid class of ABIs exist which archive the data onboard the device and later transmit data for retrieval. These devices, known as “pop-up tags”, are typically used on fish in marine systems where salt water prevents the transmission of data; the tags are

buoyant and transmit data to satellites only after they are released from the animal and are floating on the water's surface (Musyl et al. 2011). Because the tags are not continuously transmitting data, they have reduced battery consumption relative to a typical transmitter. Similar temperature logging devices have now been developed that can be programmed and downloaded through organic tissue using RFID (radio-frequency identification) technology allowing data recovery without a secondary surgery (Weepit loggers, Alpha Mach, Sainte-Julie, Canada). Additionally, devices that transmit data via radio-waves rather than via satellite, are now being utilized in laboratory experiments with rodents; these implanted devices are designed to measure a variety of physiological parameters including T_b , heart rate, ECG, and acceleration (e.g., MouseMonitor, Indus Industries, Texas, USA). These devices can be programmed and downloaded remotely and, importantly, the batteries can be charged inductively extending the life of the devices. Although the limited memory and, in particular, battery power, of these new inductively chargeable ABIs make them unsuitable for use on free-living animals, the development of devices with greater memory and battery life in the future will provide a powerful tool for measuring physiological function under field conditions.

3. Biologging in Arctic Ground Squirrels

3.1 Life history and hibernation physiology

AGSs are obligate hibernators that spend ~7-8 months of the year sequestered below ground in a hibernacula and have a relatively short ~4-5 month active season during which they reproduce, molt, and grow and fatten again for the subsequent hibernation bout (Buck and Barnes 1999a). The presence of continuous permafrost in the Arctic means that, unlike for most other hibernators, the hibernacula of AGS lies within the active layer of soil which will freeze mid-way through their hibernation season reaching average temperatures of -10 C that require AGS to defend a large thermal gradient between ambient temperature and T_b (Buck and Barnes 1999b). This is in contrast with most small mammalian hibernators that thermoconform, with T_b approximating ambient temperature during torpor. Lab studies indicate AGS are capable of maintaining steady state torpor at ambient temperatures as low as -26°C via a 36-fold increase in metabolic rate during torpor associated with heat generation in brown adipose tissue (Richter et al. 2014). Abdominal T_b while animals are torpid, yet thermogenic, is consistently below 0°C (Buck et al. 2008) and some tissues and their fluids will even decrease to below their equilibrium freezing point and thus are supercooled (Barnes 1989). T_b is not uniform during deep torpor, however, as animals will exhibit regional heterothermy with the head and thoracic region being warmer than the abdominal cavity and periphery (Barnes 1989). This is evident in our data from free-living animals as collar temperatures (skin) exceed abdominal T_b during torpor (Fig. 1). This likely occurs because heat produced by brown

adipose tissue, which is concentrated in periarterial regions in the thoracic cavity, is transferred by blood directly to the brain resulting in a relatively higher neck and brain temperature. Perhaps more surprisingly, we found that collar temperature appears to warm slightly across each torpor bout during the portion of the year that animals are continually thermogenic (Fig. 1b). This suggests that, unlike abdominal T_b , heat production by brown fat may not be constant during deep torpor. Whether this indicates a slight change in brain temperature across the torpor bout requires further investigation. Similar to most small mammalian hibernators, torpor is interrupted by regular arousal intervals during which animals become euthermic for ~14h (Buck et al. 2008; Fig 1). Light logging at a frequency of once per minute indicates that animals do not broach the surface during these intervals of euthermia but rather remain sequestered within their below ground hibernacula throughout their 6-9 month hibernation season. However, female AGS will sometimes re-enter torpor and extend hibernation in spring following particularly long euthermic intervals (>24h) which can include episodes of above-ground activity (CTW, BMB, CLB unpub. data).

3.2 Phenology

Phenology is the study of the timing of annually recurring life-cycle events and how this timing is influenced by climate conditions. Most long-term phenological studies of hibernators rely on trapping data to infer when animals have completed hibernation. However, data may be inaccurate if animals have terminated hibernation but inclement weather reduces their activity and likelihood of being captured (Williams et al. 2014b). Additionally, some species of ground squirrel exhibit prolonged below-ground intervals of euthermia immediately prior to and following the heterothermic interval of hibernation (reviewed in Williams et al. 2014a). This is particularly important in male AGS, which average 7-9 and 15-25 days of below-ground euthermia in fall and spring, respectively (Sheriff et al. 2011). In response to environmental variation or differences in gender (Barnes 1996), hibernators may alter the amount of time spent torpid or the duration of these below-ground euthermic intervals; which approach is used has important implications for energetics as metabolic rate during torpor can be reduced by as much as 98% relative to basal euthermic levels (Ruf and Geiser 2015). Conventional trapping approaches, however, cannot determine whether changes in the timing of fall immergence or spring emergence are associated with a reduction in the duration of the heterothermic interval *versus* the episodes of below-ground euthermia. For these reason, we have been using T_b as a means of assessing the timing of seasonally recurring life-cycle events and determining how this timing is influenced by spatial and temporal heterogeneity in environmental conditions.

In fall, the amplitude and mean of the daily rhythm of T_b decreases when animals first enter their hibernacula (Fig. 1a; Williams et al. 2011); in males, but not females, this occurs well in advance of

initiating torpor (~7-9 days ; Sheriff et al. 2011). In spring, T_b of euthermic AGS are arrhythmic until they emerge to the surface (Fig 1c; Williams et al. 2012a, 2012b). Measuring T_b using implanted devices also provides a means of assessing if and when females give birth because core T_b decreases during gestation and increases by 1-1.5°C on the date of parturition (Williams et al. 2011). Although collecting collar temperature reflects core T_b data during heterothermy (Barclay et al. 1996, Dausmann 2005; Fig. 1), we have found that collar T_b deviates substantially from abdominal T_b when animals are active in their burrows or on the surface (Fig. 1), and thus it is not a suitable means of estimating core T_b during the active season. This also means that collar T_b is not useful for determining the duration of the below-ground intervals of euthermia that precede and follow heterothermy. However, ABIs that measure both light and T_b , such as the Intigeo geolocators (1g; Migrate Technology Ltd., Cambridge, UK) that we utilize, provide a means of capturing all of this information using a collar mounted device (Fig. 1). Furthermore, the light logging approach provides much more detailed data on time active on the surface. For example, males may emerge from hibernation but then reenter and remain in their burrow for several days if weather conditions are poor such that rates of thermal exchange (i.e., heat loss) on the surface are high (Williams et al. 2012b).

Hibernation is obligate in AGS, with the timing of initiation and termination of torpor governed by an endogenous circannual clock (reviewed in Williams et al. 2014a). Nevertheless, hibernators have been shown to exhibit phenotypic plasticity in the timing of annually recurring life-cycle events (Williams et al. 2014a). We have been examining how AGS respond to spatial and annual variation in environmental conditions, particularly with respect to variation in the timing of snow cover patterns which influence access to forage in the spring and fall (Sheriff et al 2011, 2013, 2015). Our research has focused on two populations near Toolik Field Station (68° 38' N, 149° 38' W) that are only ~20 km apart but differ substantially in their snow cover regime. We have found that females at Toolik Lake terminate heterothermy an average of 11 days later than squirrels at Atigun River, where snow melt occurs an average of 27 days earlier (Sheriff et al. 2011). Delaying the termination of hibernation results in delayed parturition (13 days) such that juveniles have less time to grow and fatten prior to hibernation (Fig 2a-b; Williams et al. 2012c); this situation is exacerbated by conditions in the fall as snow cover also accumulates 14 days earlier at Toolik Lake (Sheriff et al. 2015). While females appear to be plastic in their spring phenology, there are no site-differences in when heterothermy is terminated in males; however, the below-ground euthermic interval that follows heterothermy is, on average, 10 days longer in males at the high snow cover site (Sheriff et al. 2011, 2013). In addition to variation in timing between sites and between years, individuals from the same location can differ substantially in their timing within a year (Fig. 2); females from the same site can differ in both the timing and duration of their active season (Fig. 3). For females, individuals that terminate hibernation and

give birth earlier tend to enter hibernation earlier the subsequent fall (Fig. 2c). Additionally, evidence is now emerging that females will enter hibernation earlier if they lose their litter during early lactation (Fig. 3). Thus, the onset of hibernation in females is driven by an endogenous clock but is also influenced by carry-over effects from hibernation and reproduction. In contrast, the initiation of hibernation in males is not affected by conditions during the prior spring. Altogether, it is clear that annual timing in the AGS is sensitive to environmental conditions but sensitivity is a sex-dependent trait. More work is needed however, to elucidate the mechanisms that underlie this sensitivity as well as to define the limits of phenotypic plasticity. The contribution of genetic variation to individual differences in timing is also unknown. Ultimately, climate-driven changes in phenology are likely to occur through microevolution, in addition to phenotypic plasticity. Phenotypic plasticity itself has an underlying genetic basis and thus plasticity may also evolve under climate change (Chown et al. 2010).

3.2.1. Predicting Phenology under Climate Change

Climate change is predicted to be more rapid and more severe in the Arctic compared to most regions of the world. Specifically, the Arctic is predicted to warm more rapidly (Ding et al. 2014), have reduced sea ice (Notze and Marotzke 2012), increased atmospheric humidity and precipitation (Kopec et al. 2016), and expansion of deciduous woody shrubs (Sturm et al. 2001). Physiological plasticity and the potential for microevolution are important components of population resilience and can potentially be useful in predicting the ‘winners’ and ‘losers’ under climate change (Somero 2010). However, like many species, AGS are likely to be impacted by climate change in a number of ways making clear predictions difficult (Wheeler and Hik 2013).

Snow cover directly affects access to food for AGS and is therefore an important driver of spring phenology (Sheriff et al. 2011). Warming temperatures could lead to earlier snow melt resulting in advances in the timing of reproduction. This is likely to be beneficial to hibernators as it provides increased time for juveniles to grow and fatten prior to their first hibernation (Morton and Sherman 1978, Williams et al. 2014a). In the Rocky Mountains of Colorado, for example, yellow-bellied marmots (*Marmota flaviventris*) have advanced their phenology by almost 30 days over the past three decades and this has led to higher population densities (Ozgul et al. 2010). However, the Arctic is also predicted to experience increases in precipitation (Kopec et al. 2016), and if this precipitation comes in the form of late spring snowstorms or results in higher winter snowpack it could result in delayed snow melt. Either of these scenarios could lead to delays in the termination of hibernation and onset of reproduction, which could potentially be manifest in lower juvenile survival and recruitment. Lane et al. (2012), for example, found that later snowmelt in the Rocky Mountains of Alberta led to delayed phenology and lower fitness for Columbian ground squirrels (*Urocitellus columbianus*).

Given the widespread range distribution of AGS (patchily distributed throughout NE Siberia in Eurasia and from Alaska to Hudson Bay in North America) and their broad ecological niche (from arctic tundra to boreal forest), it seems likely that effects of climate change on phenology will be highly variable and tied to impacts on localized climate/weather and other stressors. Nevertheless, in regions subject to later snow melt or increases in spring snowstorm events, our data suggests females are more phenologically plastic and are therefore likely to be buffered from late springs to a greater degree. Because the endocrine initiation of gonadal maturation triggers the earlier termination of torpor in sexually maturing males (Barnes 1996), only females and non-reproductive males appear to be capable of re-entering torpor if spring foraging conditions deteriorate. Such a sex-specific response could have implications for demography, including population sex-ratios, if male survival decreases as a consequence of increased snow cover and lower resource availability following the termination of hibernation. However, male survival is also likely to be influenced by the size and quality of their food caches, conditions during fall fattening, and male density during the mating season. Overall, phenology of AGS will probably be strongly influenced by climate change, but effects are likely to vary widely across their range and predicting the consequences of these phenological responses is not yet possible.

3.3 Circadian Rhythms

The question as to whether circadian rhythms persist throughout the year in polar environments, which seasonally lack a light:dark cycle, has long been of interest to chronobiologists (reviewed in Williams et al. 2015). Evidence to date suggests that polar vertebrates, including year-round residents and seasonal migrants, exhibit diverse behavioral and physiological responses to the relatively continuous lighting conditions that occur during summers and winters at high latitudes. Differences among species appear to reflect variance in the ecological niches they occupy, as well as differences in their evolutionary histories (Williams et al. 2015). ABIs have played an important role in assessing the persistence and period of rhythms in free living vertebrates. For example, accelerometers have shown that circadian rhythms of activity are lost in free-living Svalbard reindeer during the polar day and polar night (van Oort et al. 2005). Radio-transmitters have demonstrated that substantial diversity exists in the persistence of daily activity rhythms in shorebirds depending on species, sex and breeding stage; remarkably, in some species a free-running behavioral rhythm that is synchronized between pair mates has been detected (Steiger et al. 2013). However, measuring behavioral data alone is potentially problematic because rhythms may be produced as animals respond directly to external cues rather than being generated by an endogenous clock, a phenomenon known as “masking”. The continued development of ABIs that can measure a variety of physiological and behavioral variables, in combination with experimental field

manipulations, should shed light on whether observed rhythms are driven by endogenous clocks or are the result of masking.

In addition to assessing the timing of seasonally recurring life-cycle events, T_b data are also very useful in examining questions associated with circadian timing. In fact, while most studies investigating circadian rhythmicity in free-living polar organisms rely on activity data or hormone data, we argue that T_b is a better metric. The advantage of T_b over activity data stems from the fact that, in mammals, the master circadian clock in the SCN drives circadian rhythms of T_b which subsequently entrain peripheral oscillators found throughout the body (Buhr and Takahashi 2010). Because the SCN controls these T_b rhythms, they are likely to be less affected by masking. Although melatonin rhythms are also generated by the SCN, obtaining continuous measures of hormones in free-living animals is not currently possible.

We have shown that during the active season, daily rhythms of activity and T_b in AGS persist throughout the arctic summer, which includes 6 weeks of continuous daylight while the sun remains above the horizon at our study site (Long et al. 2005, Williams et al. 2012a, Williams et al. 2012b). How AGS maintain rhythmicity in the absence of a light-dark cycle remains unclear. Although AGS might be sensitive to daily rhythms in the intensity of light, we suspect that the spectral quality of light may be more important, as rhythms in the color temperature of light are relatively robust in the Arctic and are less sensitive to disruption by changing cloud cover compared with total light intensity (Ashley et al. 2013). A recent study demonstrating that the mammalian circadian clock can be entrained by changes in color also strengthens this hypothesis (Walmsley et al. 2015).

We have also shown that abdominal T_b is constant during deep torpor, remaining within the 0.2°C resolution of the loggers we were then using (Williams et al. 2012a). In fact, T_b measured during torpor using much higher resolution loggers (0.02°C) also lacks circadian organization (Fig. 1). In free-living males, T_b also remains arrhythmic during the 21-28 day intervals of below-ground euthermia that follow heterothermy; daily rhythms do not resume until exposure to light following emergence (Williams et al. 2012a, 2012b). Collectively, our results suggest that circadian clocks in AGS remain functional and entrained under the midnight sun, but are either arrested or disrupted during the constant darkness of hibernation. These results are consistent with the finding that clock gene expression is arrested during deep torpor in the European hamster, *Cricetus cricetus* (Revel et al. 2007).

3.3 Activity & Energetics

Data generated from ABIs (T_b and light loggers) we have deployed on free-living AGS indicate that the timing of key seasonal events differs by gender and age and is plastic and adjusted depending on environmental conditions. Further, these data clearly demonstrate that AGS maintain entrained daily rhythms of physiology and activity throughout the arctic summer. For the past several years we have also

507 been using light loggers to examine how the timing and duration of daily above-ground activity are
508 modified in response to changing weather conditions across the active season. We are also interested in
509 understanding whether differences in reproductive requirements between males and females results in
510 sex-differences in either the amount of time spent above ground each day or the plastic response to
511 changing environmental conditions. Finally, we have begun examining whether plasma level of thyroid
512 hormone, a potent mediator of metabolism, is correlated with daily and seasonal variation in activity.

513 Perhaps not surprisingly, we have found that the amount of time spent above ground each day is
514 strongly influenced by environmental parameters that affect thermal exchange, including ambient
515 temperature, precipitation, and solar radiation (Fig. 4; Williams et al. 2014). While we were not surprised
516 to find an effect of weather conditions on the amount of time animals spent above ground, the size of the
517 effect was surprisingly large; the duration of time above ground each day has a coefficient of variation
518 between 26-34%, depending on site, and weather conditions explain more than 50% of the day-to-day
519 variation (Williams et al. 2014; in review). Our finding that >50% of the day to day variation in above
520 ground activity is affected by weather that influences thermal exchange conditions suggest that effects of
521 climate change on weather conditions during the active season could have important population-level
522 implications. However, we currently lack data linking patterns of activity and energy expenditure to
523 measures of fitness, such as fecundity and survival.

524 Our studies have also revealed that differences between males and females in how and when
525 energy is allocated towards reproduction produces sex-differences in time spent above-ground. Light
526 logger data revealed that females spend less time above ground during early lactation but slightly more
527 time aboveground during late lactation (Williams et al. in review). This result appears to be at odds with
528 earlier studies of ground squirrels which indicate daily energy expenditure, as measured using the doubly-
529 labelled water approach, is much higher in lactating females compared to males (Kenagy et al. 1989).
530 However, time spent above ground does not necessarily reflect activity levels or efficiency in foraging.
531 To address this issue, we estimated daily movement-based energy expenditure by calculating ODBA from
532 long-term (up to 60day) deployments of accelerometers. We found that mean daily ODBA was positively
533 correlated with time spent above ground, but females were more active per unit time spent above ground
534 and had higher movement-based energy expenditure than males throughout gestation and late lactation
535 (Williams et al. in review). Similar to time spent above ground, ODBA varies substantially from day-to-
536 day with more than 50% of this variation being attributable to variation in thermal exchange conditions.
537 Interestingly, while Fletcher et al. (2012) recently showed that seasonal stage differences overwhelm
538 environmental factors as determinants of energy expenditure in red squirrels, the opposite appears to be
539 true in our system, at least in terms of movement-based energy expenditure.

Total energy expenditure includes basal metabolism, movement-based energy expenditure, as well as energy allocated to other processes (e.g., thermoregulation, digestion, lactation, growth, etc.). Maintaining a high core T_b makes up a substantial portion of RMR, and in AGS, levels of core T_b varies substantially across the season (Fig. 3; Williams et al. 2011). Recently, we showed that circulating total triiodothyronine (T3; the active form of thyroid hormone), a potent modulator of basal metabolism, varies substantially across the active season in free-living AGS (Wilsterman et al. 2015). When we compare T3 levels to abdominal T_b in free-living AGS, we see they are positively correlated (Fig. 5) which suggests T3 may contribute to increases in RMR through modulation of core T_b . Perhaps more surprising than the associations between T3 and T_b was our finding that both T3 and thyroxine (T4; a precursor of T3) were positively correlated with time spent above ground each day, as measured using light loggers (Wilsterman et al. 2015). These data support the common assumption that basal metabolism, total energy expenditure, and activity levels of individuals are physiologically linked (Biro and Stamps 2010). However, the relationship between thyroid hormones and time spent above ground was context dependent and only occurred during particular life-history stages (Wilsterman et al. 2015). Additionally, we now know that time spent above ground explains only ~40% of the variation in movement-based energy expenditure in female AGS (as measured by ODBA; Williams et al. in review). Nevertheless, these findings provide a framework for understanding thyroid hormone function in free-living animals and provide the first link between thyroid hormones and behavior in free-living mammals. The continued use of field endocrinology, in combination with ABIs, particularly those that can measure multiple physiological and behavioral parameters simultaneously (Wilson et al. 2008), will help shed light on the physiological links between endocrine drivers and their behavioral and physiological outputs.

4. Conclusions

The world is changing on global, regional, and local scales and understanding how individuals, populations, species, and communities will be affected requires integrative studies designed to measure behavior and physiology in free-living animals. Our studies on AGS illustrate that ABIs provide a powerful tool to address a wide variety of behavioral, physiological, and ecological questions, from understanding the role of circadian clocks in polar environments to assessing how heterotherms might cope with climate change. The development of small ABIs that can measure many parameters, store data on an onboard high capacity chip, and later be downloaded, reprogrammed, and charged remotely, will greatly enhance our ability to monitor behavior and physiology in free-ranging, undisturbed subjects, thereby advancing basic and applied biological research.

Acknowledgments

This research was supported by grants from the National Science Foundation to CTW (IOS-1147187), CLB (EF- 0732755 and IOS-1147232), and BMB (EF-0732763 and IOS-1147187). Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation.

References

- Arnold, W., Ruf, T., Reimoser, S., Tataruch, F., Onderscheka, K., Schober, F., 2004. Nocturnal hypometabolism as an overwintering strategy of red deer (*Cervus elaphus*). *Am. J. Physiol.-Reg. Integr. Comp. Physiol.* 286, R174-R181.
- Ashley, N.T., Schwabl, I., Goymann, W., Buck, C.L., 2013. Keeping time under the midnight sun: Behavioral and plasma melatonin profiles of free-living Lapland longspurs (*Calcarius lapponicus*) during the arctic summer. *J. Exp. Zool. A* 319, 10-22.
- Barclay, R.M., Kalcounis, M.C., Crampton, L.H., Stefan, C., Vonhof, M.J., Wilkinson, L., Brigham, R.M., 1996. Can external radiotransmitters be used to assess body temperature and torpor in bats? *J. Mamm.* 77, 1102-1106.
- Barnes, B.M., 1989. Freeze avoidance in a mammal: body temperatures below 0 degree C in an Arctic hibernator. *Science* 244, 1593-1595.
- Barnes, B.M., 1996. Relationships between hibernation and reproduction in male ground squirrels. In: *Adaptations to the Cold: Tenth International Hibernation Symposium*. Geiser F et al, eds. University of New England Press, Armidale, Australia. 71-80.
- Barron, D.G., Brawn, J.D., Weatherhead, P.J., 2010. Meta-analysis of transmitter effects on avian behaviour and ecology. *Methods Ecol. Evol.* 1, 180-187.
- Bartholomew, G.A., 1986. The role of natural history in contemporary biology. *BioScience* 36, 324-329.
- Bartholomew, G.A., Cade, T.J., 1957. Temperature regulation, hibernation, and aestivation in the little pocket mouse, *Perognathus longimembris*. *J. Mamm.* 38, 60-72.
- Beaupre, S.J., Jacobson, E.R., Lillywhite, H.B., Zamudio, K., 2004. Guidelines for the use of live amphibians and reptiles in field and laboratory research. American Society of Ichthyologists and Herpetologists. Miama, FL.
- Benedict, F.G., Lee, R.C., 1938. Hibernation and marmot physiology. Carnegie Inst. of Washington, Washington, D. C.
- Biro, P.A., Stamps, J.A., 2010. Do consistent individual differences in metabolic rate promote consistent individual differences in behavior? *Trends Ecol. Evol.* 25, 653-659.
- Block, B.A., 2005. Physiological ecology in the 21st century: advancements in biologging science. *Integr. Comp. Biol.* 45, 305-320.

608 Boyer, B.B., Barnes, B.M., 1999. Molecular and metabolic aspects of hibernation. *Bioscience* 49, 713-
 609 724.
 610 Brigham, R.M., 1992. Daily torpor in a free-ranging goatsucker, the common poorwill (*Phalaenoptilus*
 611 *nuttallii*). *Physiol. Zool.* 65, 457-472.
 612 Brown, D.D., Kays, R., Wikelski, M., Wilson, R., Klimley, A.P., 2013. Observing the unwatchable
 613 through acceleration logging of animal behavior. *Anim. Biotelem.* 1, 1-16.
 614 Buck, C.L., Barnes, B.M., 1999a. Annual cycle of body composition and hibernation in free-living arctic
 615 ground squirrels. *J. Mamm.* 80, 430-442.
 616 Buck, C.L., Barnes, B.M., 1999b. Temperatures of hibernacula and changes in body composition of arctic
 617 ground squirrels over winter. *J. Mamm.* 80, 1264-1276.
 618 Buck, C.L., Breton, A., Kohl, F., Toien, O., Barnes, B.M., 2008. Overwinter body temperature patterns in
 619 free-living arctic squirrels (*Spermophilus parryii*). *Hypometabolism in animals: torpor*
 620 *hibernation and cryobiology*. University of KwaZulu-Natal, Pietermaritzburg, 317-326.
 621 Buehler, D.M., Piersma, T., Tieleman, B.E., 2008 Captive and free-living red knots *Calidris canutus*
 622 exhibit differences in non-induced immunity that suggest different immune strategies in different
 623 environments. *J. Avian Biol.* 39, 560-566.
 624 Buhr, E.D., Yoo, S.H., Takahashi, J.S., 2010. Temperature as a universal resetting cue for mammalian
 625 circadian oscillators. *Science* 330, 379-385.
 626 Butler, P.J., Green, J.A., Boyd, I.L., Speakman, J.R., 2004. Measuring metabolic rate in the field: the pros
 627 and cons of the doubly labelled water and heart rate methods. *Funct. Ecol.* 18, 168-183.
 628 Cagnacci, F., Boitani, L., Powell, R.A., Boyce, M. S., 2010. Animal ecology meets GPS-based
 629 radiotelemetry: a perfect storm of opportunities and challenges. *Phil. Trans. Roy. Soc. B* 365,
 630 2157-2162.
 631 Chown, S.L., Hoffmann, A.A., Kristensen, T.N., Angilletta Jr, M.J., Stenseth, N.C., Pertoldi, C., 2010.
 632 Adapting to climate change: a perspective from evolutionary physiology. *Climate Res.* 43, 3.
 633 Cooke, S.J., Sack, L., Franklin, C.E., Farrell, A.P., Beardall, J., Wikelski, M., Chown, S.L., 2013. What is
 634 conservation physiology? Perspectives on an increasingly integrated and essential science.
 635 *Conserv. Physiol.* 1, cot001.
 636 Dausmann, K.H., Glos, J., Ganzhorn, J.U., Heldmaier, G., 2004. Physiology: hibernation in a tropical
 637 primate. *Nature* 429, 825-826.
 638 Dausmann, K.H., 2005. Measuring body temperature in the field—evaluation of external vs. implanted
 639 transmitters in a small mammal. *J. Thermal Biol.* 30, 195-202.
 640 Denny, M., Helmuth, B., 2009. Confronting the physiological bottleneck: a challenge from ecomechanics.
 641 *Integr. Comp. Biol.* 49, 197-201.

642 Ding, Q., Wallace, J.M., Battisti, D.S., Steig, E.J., Gallant, A.J., Kim, H.J., Geng, L., 2014. Tropical
 643 forcing of the recent rapid Arctic warming in northeastern Canada and Greenland. *Nature* 509,
 644 209-212.

645 Eichhorn, G., Groscolas, R., Le Glaunec, G., Parisel, C., Arnold, L., Medina, P., Handrich, Y., 2011.
 646 Heterothermy in growing king penguins. *Nature Comm.* 2, 435.

647 Evans, A.L., Singh, N.J., Friebe, A., Arnemo, J.M., Laske, T.J., Fröbert, O., Swenson, J.E., Blanc, S.,
 648 2016. Drivers of hibernation in the brown bear. *Frontiers Zool.* 13, 1.

649 Fair, J.M., Jones, J., 2010. Guidelines to the use of wild birds in research. Ornithological Council.

650 Fletcher, Q.E., Speakman, J.R., Boutin, S., McAdam, A.G., Woods, S.B., Humphries, M.M., 2012.
 651 Seasonal stage differences overwhelm environmental and individual factors as determinants of
 652 energy expenditure in free-ranging red squirrels. *Funct Ecol.* 26, 677-687.

653 Fuller, A., Moss, D.G., Skinner, J.D., Jessen, P.T., Mitchell, G., Mitchell, D., 1999. Brain, abdominal and
 654 arterial blood temperatures of free-ranging eland in their natural habitat. *Pflügers Archiv.* 438,
 655 671-680.

656 Geiser, F., Holloway, J.C., Körtner, G., Maddocks, T.A., Turbill, C., Brigham, R.M., 2000. Do patterns of
 657 torpor differ between free-ranging and captive mammals and birds?. In Heldmaier, G.,
 658 Klingenspor, M. (Eds), *Life in the Cold*. Springer Berlin Heidelberg, pp. 95-102.

659 Green, J.A., Halsey, L.G., Wilson, R.P., Frappell, P.B., 2009. Estimating energy expenditure of animals
 660 using the accelerometry technique: activity, inactivity and comparison with the heart-rate
 661 technique. *J. Exp. Biol.* 212, 471-482.

662 Halsey, L.G., Shepard, E.L.C., Quintana, F., Laich, A.G., Green, J.A., Wilson, R.P., 2009. The
 663 relationship between oxygen consumption and body acceleration in a range of species. *Comp.*
 664 *Biochem. Physiol. A* 152, 197-202.

665 Halsey, L.G., Shepard, E.L., Wilson, R.P., 2011. Assessing the development and application of the
 666 accelerometry technique for estimating energy expenditure. *Comp. Biochem. Physiol. A* 158,
 667 305-314.

668 Hammel, H.T., 1967. Temperature regulation and hibernation. In: Fisher, K.C., Dawe, A.R., Lyman, C.P.,
 669 Schonbaum, E., South, F.E. (eds) *Mammalian hibernation III*. Elsevier, Amsterdam, pp 86–96.

670 Hetem, R.S., Maloney, S.K., Fuller, A., Mitchell, D., 2014. Heterothermy in large mammals: inevitable or
 671 implemented? *Biological Reviews* DOI: 10.1111/brv.12166

672 Hock, R.J., 1951. The metabolic rates and body temperatures of bats. *Biol. Bull.* 101, 289-299.

673 Hock, R.J., 1960. Seasonal variations in physiologic functions of arctic ground squirrels and black bears.
 674 *Bull. Mus. Comp. Zool.* 124, 155-171.

675 Hoelzl, F., Bieber, C., Cornils, J.S., Gerritsmann, H., Stalder, G. L., Walzer, C., Ruf, T., 2015. How to
 676 spend the summer? Free-living dormice (*Glis glis*) can hibernate for 11 months in non-
 677 reproductive years. *J. Comp. Physiol. B* 185, 931-939.

678 Hut, R.A., van Oort, B.E., Daan, S., 1999. Natural entrainment without dawn and dusk: the case of the
 679 European ground squirrel (*Spermophilus citellus*). *J. Biol. Rhythms* 14, 290-299.

680 Jepsen, N., Thorstad, E.B., Havn, T., Lucas, M.C., 2015. The use of external electronic tags on fish: an
 681 evaluation of tag retention and tagging effects. *Anim. Biotelem.* 3, 1.

682 Kenagy, G.J., Sharbaugh, S.M., Nagy, K.A., 1989. Annual cycle of energy and time expenditure in a
 683 golden-mantled ground squirrel population. *Oecologia* 78, 269-282.

684 Kopec, B.G., Feng, X., Michel, F.A., Posmentier, E.S., 2016. Influence of sea ice on Arctic precipitation.
 685 *Proc. Nat. Acad. Sci.* 113, 46-51.

686 Körtner, G., Geiser, F., 2000. Torpor and activity patterns in free-ranging sugar gliders *Petaurus*
 687 *breviceps* (Marsupialia). *Oecologia* 123, 350-357.

688 Körtner, G., Brigham, R.M., Geiser, F., 2000. Metabolism: Winter torpor in a large bird. *Nature* 407, 318-
 689 318.

690 Lane, J.E., Kruuk, L.E., Charmantier, A., Murie, J.O., Dobson, F.S., 2012. Delayed phenology and
 691 reduced fitness associated with climate change in a wild hibernator. *Nature*, 489, 554-557.

692 Langer, F., Fietz, J., 2014. Ways to measure body temperature in the field. *J. Thermal Biol.* 42, 46-51.

693 Long, R.A., Martin, T.J., Barnes, B.M., 2005. Body temperature and activity patterns in free-living arctic
 694 ground squirrels. *J. Mamm.* 86, 314-322.

695 Long, R.A., Hut, R.A., Barnes, B.M., 2007. Simultaneous collection of body temperature and activity
 696 data in burrowing mammals: a new technique. *J. wildl. Manage.* 71, 1375-1379.

697 Lovegrove, B.G., 2009. Modification and miniaturization of Thermochron iButtons for surgical
 698 implantation into small animals. *J. Comp. Physiol. B* 179, 451-458.

699 Luecke, R.H., South, F.E., 1972. A possible model for thermoregulation during deep hibernation. In:
 700 South, F.E., Hannon, J.P., Willis, J.R., Pengelley, E.T., Alpert, N.R. (eds) *Hibernation and*
 701 *hypothermia, perspectives and challenges*. Elsevier, New York, pp 577–605.

702 Lyman, C.P., Chatfield, P.O., 1956. Physiology of hibernation in mammals. *Physiol. Induced*
 703 *Hypothermia* 1, 80-122.

704 McCafferty, D.J., Gallon, S., Nord, A., 2015. Challenges of measuring body temperatures of free-ranging
 705 birds and mammals. *Anim. Biotelem.* 3, 1.

706 McDonald, B.I., Ponganis, P.J., 2013. Insights from venous oxygen profiles: oxygen utilization and
 707 management in diving California sea lions. *J. Exp. Biol.* 216, 3332-3341.

708 Michener, G.R., 1992. Sexual differences in over-winter torpor patterns of Richardson's ground squirrels
709 in natural hibernacula. *Oecologia* 89, 397-406.

710 Morgan, K.N., Tromborg, C.T., 2007. Sources of stress in captivity. *Applied Anim. Behav. Sci.* 102, 262-
711 302.

712 Morton, M.L., Sherman, P.W., 1978. Effects of a spring snowstorm on behavior, reproduction, and
713 survival of Belding's ground squirrels. *Can. J. Zool.* 56, 2578-2590.

714 Musyl, M.K., Domeier, M.L., Nasby-Lucas, N., Brill, R.W., McNaughton, L.M., Swimmer, J.Y.,
715 Lutcavage, M.S., Wilson, S.G., Galuardi, B., Liddle, J.B., 2011. Performance of pop-up satellite
716 archival tags. *Mar. Ecol. Prog. Ser.* 433, 1-28.

717 Nickum, J.G., 2004. Guidelines for the use of fishes in research. American Fisheries Society

718 Notz, D., Marotzke, J., 2012. Observations reveal external driver for Arctic sea-ice retreat. *Geophys. Res.*
719 *Lett.* 39, L08502.

720 O'Mara, M.T., Wikelski, M., Dechmann, D.K., 2014. 50 years of bat tracking: device attachment and
721 future directions. *Methods Ecol. Evol.* 5, 311-319.

722 Ozgul, A., Childs, D.Z., Oli, M.K., Armitage, K.B., Blumstein, D.T., Olson, L.E., Tuljapurkar, S.,
723 Coulson, T., 2010. Coupled dynamics of body mass and population growth in response to
724 environmental change. *Nature* 466, 482-485.

725 Phillips, R.A., Silk, J.R.D., Croxall, J.P., Afanasyev, V., Briggs, D.R., 2004. Accuracy of geolocation
726 estimates for flying seabirds. *Mar. Ecol. Prog. Ser.* 266, 265-272.

727 Pivorun, E.B., 1976. A biotelemetry study of the thermoregulatory patterns of *Tamias striatus* and
728 *Eutamias minimus* during hibernation. *Comp. Biochem. Physiol.* 53, 265-271.

729 Ponganis, P.J., 2007. Bio-logging of physiological parameters in higher marine vertebrates. *Deep Sea*
730 *Res. Part II* 54, 183-192.

731 Revel, F.G., Herwig, A., Garidou, M.L., Dardente, H., Menet, J.S., Masson-Pévet, M., Simonneaux, V.,
732 Saboureau, M., Pévet, P., 2007. The circadian clock stops ticking during deep hibernation in the
733 European hamster. *Proc. Nat. Acad. Sci.* 104, 13816-13820.

734 Richter, M.M., Williams, C.T., Lee, T.N., Tøien, Ø., Florant, G.L., Barnes, B.M., Buck, C.L., 2015.
735 Thermogenic capacity at subzero temperatures: how low can a hibernator go? *Physiol. Biochem.*
736 *Zool.* 88, 81-89.

737 Robert, K.A., Thompson, M.B., 2003. Reconstructing Thermochron iButtons to reduce size and weight as
738 a new technique in the study of small animal thermal biology. *Herpetological Rev.* 34, 130-173.

739 Rojas, A.D., Körtner, G., Geiser, F., 2010. Do implanted transmitters affect maximum running speed of
740 two small marsupials? *J. Mammal.* 91, 1360-1364.

741 Romero, L.M., Wingfield, J.C., 1999. Alterations in hypothalamic–pituitary–adrenal function associated
 742 with captivity in Gambel’s white-crowned sparrows (*Zonotrichia leucophrys gambelii*). *Comp.*
 743 *Biochem. Physiol. B* 122, 13–20.
 744 Ropert-Coudert, Y., Wilson, R.P., 2004. Subjectivity in bio-logging science: do logged data mislead.
 745 *Mem. Natl. Inst. Polar Res.* 58, 23–33.
 746 Ruf, T., Geiser, F., 2015. Daily torpor and hibernation in birds and mammals. *Biol. Rev.* 90, 891–926.
 747 Rutz, C., Hays, G.C., 2009. New frontiers in biologging science. *Biol. Lett.* 5, 289–292.
 748 Saraux, C., Le Bohec, C., Durant, J.M., Viblanc, V.A., Gauthier-Clerc, M., Beaune, D., Park, Y.H.,
 749 Yoccoz, N.G., Stenseth, N.C., Le Maho, Y., 2011. Reliability of flipper-banded penguins as
 750 indicators of climate change. *Nature* 469, 203–206.
 751 Schaller, R.R., 1997. Moore’s law: past, present and future. *Spectrum IEEE* 34, 52–59.
 752 Semeniuk, C.A., Bourgeon, S., Smith, S.L., Rothley, K.D., 2009. Hematological differences between
 753 stingrays at tourist and non-visited sites suggest physiological costs of wildlife tourism. *Biol.*
 754 *Conserv.* 142, 1818–1829.
 755 Sheriff, M.J., Kenagy, G.J., Richter, M., Lee, T., Tøien, Ø., Kohl, F., Buck, C.L., Barnes, B.M., 2011.
 756 Phenological variation in annual timing of hibernation and breeding in nearby populations of
 757 arctic ground squirrels. *Proc. Roy. Soc. Lond. B* 278, 2369–2375.
 758 Sheriff, M.J., Williams, C.T., Kenagy, G.J., Buck, C.L., Barnes, B.M., 2012. Thermoregulatory changes
 759 anticipate hibernation onset by 45 days: data from free-living arctic ground squirrels. *J. Comp.*
 760 *Physiol. B* 182, 841–847.
 761 Sheriff, M.J., Richter, M.M., Buck, C.L., Barnes, B.M., 2013. Changing seasonality and phenological
 762 responses of free-living male arctic ground squirrels: the importance of sex. *Phil. Trans. Roy.*
 763 *Soc. Lond. B* 368, 20120480.
 764 Sheriff, M.J., Buck, C.L., Barnes, B.M., 2015. Autumn conditions as a driver of spring phenology in a
 765 free-living arctic mammal. *Clim. Change Resp.* 2, 4.
 766 Sikes, R.S., Gannon, W.L., 2011. Guidelines of the American Society of Mammalogists for the use of
 767 wild mammals in research. *J. Mamm.* 92, 235–253.
 768 Somero, G.N., 2010. The physiology of climate change: how potentials for acclimatization and genetic
 769 adaptation will determine ‘winners’ and ‘losers’. *J. Exp. Biol.* 213, 912–920.
 770 Steiger, S.S., Valcu, M., Spoelstra, K., Helm, B., Wikelski, M., Kempenaers, B., 2013. When the sun
 771 never sets: diverse activity rhythms under continuous daylight in free-living arctic-breeding birds.
 772 *Proc. Roy. Soc. Lond. B* 280(1764), 20131016.
 773 Strumwasser, F., 1958. Factors in the pattern, timing and predictability of hibernation in the squirrel,
 774 *Citellus beecheyi*. *Am. J. Physiol.* 196, 8–14.

775 Sturm, M., Racine, C., Tape, K., 2001. Climate change: increasing shrub abundance in the Arctic. *Nature*
776 411, 546-547.

777 Taylor, N.A., Tipton, M.J., Kenny, G.P., 2014. Considerations for the measurement of core, skin and
778 mean body temperatures. *J. Thermal Biol.* 46, 72-101.

779 van Oort, B.E., Tyler, N.J., Gerkema, M.P., Folkow, L., Blix, A.S., Stokkan, K.A., 2005. Circadian
780 organization in reindeer. *Nature* 438, 1095-1096.

781 Vyssotski, A.L., Serkov, A.N., Itskov, P.M., Dell'Omo, G., Latanov, A.V., Wolfer, D.P., Lipp, H.P.,
782 2006. Miniature neurologgers for flying pigeons: multichannel EEG and action and field
783 potentials in combination with GPS recording. *J. Neurophys.* 95, 1263-1273.

784 Wacker, C.B., Rojas, A.D., Geiser, F., 2012. The use of small subcutaneous transponders for quantifying
785 thermal biology and torpor in small mammals. *J. Thermal Biol.* 37, 250-254.

786 Walmsley, L., Hanna, L., Moulard, J., Martial, F., West, A., Smedley, A.R., Bechtold, D.A., Webb, A.R.,
787 Lucas, R.J., Brown, T.M., 2015. Colour as a signal for entraining the mammalian circadian clock.
788 *PLoS Biol.* 13, e1002127.

789 Warnecke, L., Turner, J.M., Bollinger, T.K., Misra, V., Cryan, P.M., Blehert, D.S., Wibbelt, G., Willis, C.
790 K., 2013. Pathophysiology of white-nose syndrome in bats: a mechanistic model linking wing
791 damage to mortality. *Biol. Lett.* 9, 20130177.

792 Wheeler, H.C., Hik, D.S., 2013. Arctic ground squirrels *Urocitellus parryi* as drivers and indicators of
793 change in northern ecosystems. *Mamm. Rev.* 43, 238-255.

794 Whidden, S.E., Williams, C.T., Breton, A.R., Buck, C.L., 2007. Effects of transmitters on the
795 reproductive success of tufted puffins. *J. Field Ornithol.* 78, 206-212.

796 White, C.R., Cassey, P., Schimpf, N.G., Halsey, L.G., Green, J.A., Portugal, S. J., 2013. Implantation
797 reduces the negative effects of bio-logging devices on birds. *J. Exp. Biol.* 216, 537-542.

798 Williams, C.T., Sheriff, M.J., Schmutz, J.A., Kohl, F., Tøien, Ø., Buck, C.L., Barnes, B.M., 2011. Data
799 logging of body temperatures provides precise information on phenology of reproductive events
800 in a free-living arctic hibernator. *J. Comp. Physiol. B* 181, 1101-1109.

801 Williams, C.T., Barnes, B.M., Buck, C.L., 2012a. Daily body temperature rhythms persist under the
802 midnight sun but are absent during hibernation in free-living arctic ground squirrels. *Biol. Lett.* 8,
803 31-34.

804 Williams, C.T., Barnes, B.M., Richter, M., Buck, C.L., 2012b. Hibernation and circadian rhythms of body
805 temperature in free-living Arctic ground squirrels. *Physiol. Biochem. Zool.* 85, 397-404.

806 Williams, C.T., Sheriff, M.J., Kohl, F., Barnes, B.M., Buck, C.L., 2012c. Interrelationships among timing
807 of hibernation, reproduction, and warming soil in free-living female arctic ground squirrels. In
808 *Living in a Seasonal World* (pp. 63-72). Springer Berlin Heidelberg.

- Williams, C.T., Barnes, B.M., Kenagy, G.J., Buck, C.L., 2014a. Phenology of hibernation and reproduction in ground squirrels: integration of environmental cues with endogenous programming. *J. Zool.* 292, 112-124.
- Williams, C.T., Wilsterman, K., Kelley, A.D., Breton, A.R., Stark, H., Humphries, M.M., Humphries, A.G., Barnes, B.M., Boutin, S., Buck, C.L., 2014b. Light loggers reveal weather-driven changes in the daily activity patterns of arboreal and semifossorial rodents. *J. Mamm.* 95, 1230-1239.
- Williams, C.T., Barnes, B.M., Buck, C.L., 2015. Persistence, entrainment, and function of circadian rhythms in polar vertebrates. *Physiology* 30, 86-96.
- Williams, C.T., Wilsterman, K., Zhang, V., Moore, J., Barnes, B.M., Buck, C.L. In Review. Sex-differences in above-ground activity and movement-based energy expenditure in a semi-fossorial mammal.
- Willis, C.K., Jameson, J.W., Faure, P.A., Boyles, J.G., Brack Jr, V., Cervone, T.H., 2009. Thermocron iButton and iBBat temperature dataloggers emit ultrasound. *J. Comp. Physiol. B* 179, 867-874.
- Wilson, R.P., White, C.R., Quintana, F., Halsey, L.G., Liebsch, N., Martin, G.R., Butler, P.J., 2006. Moving towards acceleration for estimates of activity-specific metabolic rate in free-living animals: the case of the cormorant. *J. Anim. Ecol.* 75, 1081-1090.
- Wilson, R.P., Shepard, E.L.C., Liebsch, N., 2008. Prying into the intimate details of animal lives: use of a daily diary on animals. *Endangered. Spec. Res.* 4, 123-137.
- Wilsterman, K., Buck, C.L., Barnes, B.M., Williams, C.T., 2015. Energy regulation in context: Free-living female arctic ground squirrels modulate the relationship between thyroid hormones and activity among life history stages. *Horm. Behav.* 75, 111-119.
- Wingfield, J.C., Hunt, K., Breuner, C., Dunlap, K., Fowler, G.S., Freed, L., Lepson, J., 1997. Environmental stress, field endocrinology, and conservation biology. *Behavioral approaches to conservation in the wild.* Cambridge University Press, Cambridge, 95-131
- Young, P.J., 1990. Hibernating patterns of free-ranging Columbian ground squirrels. *Oecologia* 83, 504-511.

837

838 **Fig 1.** Minimum (blue line) and maximum (red line) collar temperature both closely track core T_b (black
 839 line), as assessed using an abdominally-implanted temperature logger, throughout the heterothermic
 840 interval of hibernation. The light measurements (green line) indicate when the AGS is above vs. below
 841 ground. The more variable collar temperatures reflect the lower resolution of the collar mounted logger
 842 relative to the implanted T_b logger ($\pm 0.5^\circ\text{C}$ vs. $\pm 0.02^\circ\text{C}$ resolution). Abdominal T_b can be used to assess
 843 the timing of immergence in fall, because the amplitude of the daily T_b rhythm decreases after the final
 844 day spent above ground (arrow in Panel a). Abdominal T_b can also be used to determine the timing of
 845 emergence because a daily T_b rhythm is initiated with exposure to light (arrow in Panel c). Variation in
 846 collar temperature is much higher than in abdominal T_b when animals are euthermic and therefore, unlike
 847 core T_b , is not useful in assessing the intervals of below-ground euthermia that precede and follow
 848 hibernation; however, this can be assessed with a collar-mounted logger than measures both light and
 849 temperature. Abdominal T_b remains stable throughout torpor bouts when the animal is thermogenic
 850 whereas collar temperature is slightly warmer and exhibits a slight rise across the torpor bout (Panel b).

Fig 2. The relationship between the timing of parturition and (a) last day of heterothermy, (b) date of emergence, and (c) date of subsequent immersion into the hibernacula for a single year in female arctic ground squirrels. Phenology data was acquired using abdominally-implanted T_b loggers. Snow melt occurs one month later at Toolik Lake (open circles) compared to Atigun River (closed circles). Figure from Williams et al. (2012c).

Fig 3. Core body temperature (T_b) of 5 females throughout their active season in 2014. Although females remained euthermic throughout the active season, T_b is dynamic and exhibits 24 h daily rhythms with an amplitude of 2-3°C. The abrupt increase in T_b indicated by the arrows delineate when females gave birth (Williams et al. 2011). T_b begins to decline well before the onset of hibernation; the decline begins ~45 days before first entering torpor in late summer (Sheriff et al. 2012). The length of the active season is affected by the reproductive status of the females; female D, which had a much shorter active season, lost her litter in early lactation. On average, females that give birth earlier will subsequently enter hibernation significantly earlier (Williams et al. 2012c).

Fig. 4. The relationship between total time spent aboveground by AGS during the day (calculated based on light logging) and ambient temperature conditions in 2014 on days without precipitation (black circles) and days with >2mm of precipitation (open circles).

874

875

876

877 **Fig 5.** Total triiodothyronine (tT₃: mean ± SE) vs. mean abdominal body temperature (T_b: mean ± SE) of
 878 female arctic ground squirrels during four stages of the active season. Different lowercase letters indicate
 879 significant ($p < 0.05$) differences in tT₃ between groups whereas different uppercase letters indicate
 880 significant differences in T_b between groups. Samples sizes for tT₃ data shown in brackets. Core T_b was
 881 measured in 6 females. Thyroid hormone data from Wilsterman et al. (2015).