Claudin gene expression patterns do not associate with interspecific differences in paracellular nutrient absorption

Edwin R. Price^a, Katherine H. Rott^a, Enrique Caviedes-Vidal^{b,c}, and William H. Karasov^a

^a*Department of Forest and Wildlife Ecology, University of Wisconsin-Madison, Madison WI 53706, USA*

^b*Departamento de Bioquímica y Ciencias Biológicas y Laboratorio de Biología “Professor E. Caviedes Codelia”, Universidad Nacional de San Luis, 5700 San Luis, Argentina.*

^c*Laboratorio de Biología Integrativa, Instituto Multidisciplinario de Investigaciones Biológicas de San Luis, Consejo Nacional de Investigaciones Científicas y Técnicas, 5700 San Luis, Argentina.*

*Correspondence and current address:
Edwin R. Price
Department of Biological Sciences
1511 W. Sycamore
Denton TX 76203, USA
Edwin.price@unt.edu
telephone: 608-234-2665
fax 940-565-3821

Abstract

Bats exhibit higher paracellular absorption of glucose-sized molecules than non-flying mammals, a phenomenon that may be driven by higher permeability of the intestinal tight junctions. The various claudins, occludin, and other proteins making up the tight junctions are thought to determine their permeability properties. Here we show that absorption of the paracellular probe L-arabinose is higher in a bat (*Eptesicus fuscus*) than a vole (*Microtus pennsylvanicus*) or a hedgehog (*Atelerix albiventris*). Furthermore, histological measurements demonstrated that hedgehogs have many more enterocytes in their intestines, suggesting that bats cannot have higher absorption of arabinose simply by having more tight junctions. We therefore investigated the mRNA levels of several claudins and occludin, because these proteins may affect permeability of tight junctions. To assess the levels of expression of claudins per tight junction, we normalized the mRNA levels of the claudins to the constitutively expressed tight junction protein ZO-1, and combined these with measurements previously made in a bat and a rodent to determine if there were among-species differences in the mRNA levels of these genes associated with the tight junction. Although expression ratios of several genes varied among species, there was not a consistent difference between bats and non-flyers in the expression ratio of any particular gene. Protein expression patterns may differ from mRNA expression patterns, and might better explain differences among species in arabinose absorption.

Key Words: bat, claudin, gene expression, intestine, nutrients, occludin, paracellular absorption, tight junction, comparative transcriptomics

48 **Introduction**

49 In the intestine, adjacent enterocytes are linked by tight junctions, which impede the
50 movement of molecules across the epithelium and thereby form a barrier to solute flux. Tight
51 junctions are complex structures composed of proteins such as claudins and occludin (OCLN)
52 that span the cell membrane and interact in the extracellular space between cells, and also
53 connect with intracellular scaffolding proteins via interactions with zonula occludens 1 (ZO-1)
54 (Shen et al., 2011). These several proteins are thought to mediate the permeability
55 characteristics of tight junctions, determining for example, the size and charge of solutes that can
56 pass through the tight junction (Günzel and Yu, 2013). Many studies of tight junctions have
57 focused on the movement of ions across the epithelium, although a few have investigated the
58 permeability to larger, macronutrient-sized molecules. For example, overexpression and deletion
59 studies of OCLN and claudin-1 (CLDN1) have shown that these genes are associated with
60 increased permeability to mannitol (Amasheh et al., 2002; McCarthy et al., 2000; Tamura et al.,
61 2011; Van Itallie et al., 2001), although this is not a consistent finding across all studies (see for
62 example (Schulzke et al., 2005)). The effects of altered expression of any tight junction protein
63 may be context-specific (depending on starting expression level, expression of other proteins,
64 tissue type, etc.), and it is still not certain whether any claudins affect the permeability to
65 macronutrients in a specific way (Günzel and Yu, 2013).

66 Variation in tight junction permeability might be able to provide a mechanistic
67 explanation for the variation among species in their reliance on paracellular nutrient absorption.
68 Whereas non-flying mammals such as rodents rely heavily on the transcellular, transporter-
69 mediated pathway of glucose absorption, small birds and bats use the paracellular pathway (i.e.,
70 movement of glucose through tight junctions) for a majority of glucose absorption (Brun et al.,

2014; Caviedes-Vidal et al., 2007; Karasov et al., 2012; Price et al., 2014). This has been hypothesized to help birds and bats compensate for their smaller intestines (Caviedes-Vidal et al., 2007; Price et al., 2015).

Although high paracellular glucose absorption has been documented in all bat species studied (Brun et al., 2014; Caviedes-Vidal et al., 2008; Fasulo et al., 2013a; Keegan, 1984, 1980; Price et al., 2014; Tracy et al., 2007), the mechanism by which this occurs is still unclear. Bats might simply have more tight junctions in their intestines, achieved either by greater villous amplification or smaller enterocytes. Alternatively, bats might have tight junctions that are more permeable to macronutrient-sized molecules. We recently suggested that both may be occurring (Price et al., 2014). Little brown bats (*Myotis lucifugus*) have a higher density of cells (and presumably tight junctions) than white-footed mice (*Peromyscus leucopus*), but the total number of enterocytes in the small intestine was lower. We suggested that the higher paracellular absorption of nutrients by little brown bats must instead be driven by characteristics of the tight junction, and we demonstrated differences in tight junction gene expression between the species. In particular, expression of CLDN1 and CLDN15 were higher, and CLDN2 lower in little brown bat intestine compared to the white-footed mouse (Price et al., 2014).

This single species-pair comparison was intriguing, but begs further testing. For example, some of those expression differences might be related more to diet, phylogeny, or mere chance, than to the intestinal permeability characteristics associated with those species. In the present study, we therefore made integrated measurements on 3 more species: the insectivorous big brown bat (*Eptesicus fuscus*), the herbivorous meadow vole (*Microtus pennsylvanicus*), and the insectivorous hedgehog (*Atelerix albiventris*). These species provide a new comparison of a bat and a rodent, and the hedgehog provides a non-flying species that is not only shares an

insectivorous diet with the bat, but as part of the Laurasiatheria, is more closely related to the bat than to the vole (Nery et al., 2012). First, we demonstrate that the big brown bat has higher paracellular absorption of glucose-sized molecules than the non-flying species, thus confirming the pattern of high paracellular permeability in bats. Next we examine the anatomy and histology of the intestine, and show that it is unlikely that the high paracellular permeability in the bat can be explained by simply having more tight junctions than non-flyers. Finally, we measured tight junction gene expression in an attempt to understand how claudins and OCLN control paracellular permeability to glucose. For these gene expression measurements, we make comparisons among these three species and also with two previously measured species, the little brown bat (*Myotis lucifugus*) and the white-footed mouse (*Peromyscus leucopus*) (Price et al., 2014).

Methods:

Animals

Big brown bats (*Eptesicus fuscus*) are common North American insectivorous bats (Kurta and Baker, 1990) (Table 1). We captured them in Dane County, Wisconsin, using mistnets placed near streams or over the exit of bats' day roosts in human habitations. Bats were used in experiments immediately following capture. We obtained domesticated hedgehogs (*Atelerix albiventris*, all over 6 months age) from breeders in Wisconsin. Hedgehogs (order Erinaceomorpha) are primarily insectivorous (Santana et al., 2010), and were maintained on a diet of Purina Cat Chow Complete supplemented occasionally with mealworms (larvae of *Tenebrio molitor*). They were kept under 12h:12h L:D lighting conditions with food and water

provided *ad libitum* between experiments. Meadow voles (*Microtus pennsylvanicus*) are common grassland rodents of the Midwest and northeastern United States and have a primarily herbaceous diet (Lindroth and Batzli, 1984). We captured meadow voles in the Biocore Prairie and community gardens of the Lakeshore Nature Preserve, University of Wisconsin-Madison. Voles were maintained under similar conditions as the hedgehogs except that their diet consisted of a commercial rodent chow (Purina 5010 Rodent Diet) supplemented daily with fresh produce (kale, carrots, and apples). No animals were obviously pregnant at the time of capture or testing. The sample sizes in Table 1 are for gene expression measurements. Smaller subsets of animals (noted in other tables and figures) were used for whole-animal and histology measurements so as to conserve resources. Experiments were approved by the University of Wisconsin-Madison Animal Care and Use Committee (#A1441). Bats and voles were caught with permission from the Wisconsin Department of Natural Resources (permits E/T 704 and SCP-SOD-2011).

Measurement of paracellular nutrient absorption

We used two radiolabeled probes to assess nutrient absorption. L-arabinose (M_r 150) is a carbohydrate that is somewhat smaller than D-glucose (M_r 180) but its absorption is not transporter-mediated in bats or rodents (Lavin et al., 2007; Price et al., 2014). Its absorption was therefore used as an estimate of the paracellular component of D-glucose absorption. The difference between arabinose and glucose in molecular size likely causes this to be an overestimation, a point we consider further in our discussion. To estimate total glucose absorption, we used 3-O-methyl-D-glucose (3OMD-glucose), a molecule that is absorbed by both mediated and non-mediated mechanisms like D-glucose, but unlike glucose, is not substantially metabolized, and thus can be readily recovered. In the non-flying species, we also measured absorption of lactulose (M_r 342) and creatinine (M_r 113), and we present those data in

supplementary table S1. Creatinine and lactulose absorptions were not assessed in the big brown bats due to a scarcity of experimental animals of that species.

Animals were dosed orally with [^{14}C]-L-arabinose and [^3H]-3OMD-glucose at the same time. The gavage vehicle was 50 mM glucose in water. Although Na^+ is required for Na^+ -coupled glucose transport, Na^+ is secreted into the gut with bicarbonate and can readily diffuse from the blood (Brody, 1999). Thus, the lack of sodium in the gavage solution should not have affected our results, and indeed, glucose absorption was complete (see Results). After dosing, animals were placed in a metabolic chamber for collection of urine over the following several hours, where they had access to 50 mM glucose in water but no food. Voles and hedgehogs were placed in standard rat-sized wire-bottomed metabolic chambers, but for hedgehogs, we modified the chamber to have a smaller (5.7 mm) mesh size which seemed to make walking around the cage easier and more comfortable. Previously, we have favored a serial blood sampling technique in bats because of the difficulty of collecting urine (and separating it from feces) in bats (Brun et al., 2014; Caviedes-Vidal et al., 2008; Tracy et al., 2007). However, we tested and adopted a urine collection technique for bats in this study. We designed a small (15×15×5 cm) plastic metabolic chamber with metal screening glued to the top from which the bats could easily hang, and a sealable door near the bottom through which we could collect urine. The bats rarely produced feces during the experiments, and it was generally easy to separate from urine. Bats were occasionally offered water with or without glucose using a ball-tipped syringe, although they rarely drank. There were few urination events, and this likely led to some experimental variation, but this technique had some advantages over blood collection: 1) we were able to use the same calculation of fractional absorption for bats and the non-flying species, 2) we could

avoid taking multiple blood samples from a small animal, and therefore, 3) we were able to use fewer bats because we could conduct two separate trials on individual bats.

In separate trials, we injected the same probes intraperitoneally (saline vehicle) and collected urine in a similar manner. Urine subsamples were counted with a Wallac 1414 liquid scintillation counter (PerkinElmer, Waltham, MA, USA). Fractional absorption (f) of probes was determined by dividing the recovery (percentage of dose recovered in urine) in gavage trials by the recovery in injection trials. We found that we achieved asymptotic recovery of probes within about 3 hours in bats. We therefore waited at least 6 hours between trials in bats to provide an extra buffer period, and we also alternated the order of injection and gavage for the various bats to control for any potential carryover effects. In the non-flying species, trials were separated by at least 3 days.

Tissue Histology

Following the absorption trials, we euthanized the animals with isoflurane anesthesia and pneumothorax. The small intestine was removed, measured for length, divided into thirds, cut longitudinally, and laid flat to measure circumference using calipers. From each third, we preserved a central 2-cm segment in 10% formalin. At this time, we also placed in RNAlater a small segment from the distal end of the proximal third (see Gene Expression). Later, 5- μ m sections were cut and stained with haematoxylin and eosin. We measured villus length, villus width, and crypt width using 200x magnification and counting 25 villi per section using NIS-Elements D software (Nikon Instruments, Melville, NY). We then calculated amplification ratio according to Kisielinki et al. (Kisielinski et al., 2002).

We measured enterocyte size by counting the number of enterocytes per length along a villus (10 villi per section) at 400x magnification. The inverse of this measurement is the average width of an enterocyte. Enterocyte width was squared to determine the luminal surface area of an average enterocyte, and we multiplied this by the villous amplification ratio to determine the number of enterocytes per nominal surface area. For each section, this was multiplied by nominal surface area to determine the number of enterocytes per section, and sections were summed to calculate the total number of enterocytes in the intestine.

Gene Expression

We determined mRNA expression profiles for several tight junction-associated genes that are expressed in the intestine (ZO-1, CLDN1, CLDN2, CLDN4, CLDN7, and CLDN15) in 11 *Eptesicus fuscus*, 6 *Atelerix albiventris*, and 9 *Microtus pennsylvanicus*. We also measured expression of two common housekeeper genes (RPLP0 and EEF1A1). Tissue that had been stored in RNAlater was extracted using TRIzol (Life Technologies, Carlsbad, CA). RNA concentration was determined from absorbance at 260 nm using a NanoDrop spectrophotometer (Thermo Scientific, Wilmington, DE), and purity was checked by the A_{260}/A_{280} ratio (≈ 2 for all samples). 10 μ g RNA was digested using DNase I (New England Biolabs, Ipswich, MA) to remove genomic DNA, and RNA was then reverse transcribed to cDNA (iScript cDNA synthesis kit, Bio-Rad, Hercules, CA).

Degenerate primers were designed using known or predicted sequences for mammals in GenBank (Benson et al., 2012) and were used to determine partial coding sequences for our target genes. Specific primer sets were then designed with the aid of computer programs (Larkin et al., 2007; Rozen and Skaletsky, 2000)(Table 2). We verified that specific primers amplified a

single target by examination of melt curves, the presence of a single band when the product is electrophoresed, and subsequent sequencing of that band.

For the best comparisons across species, we sought to design specific primers that amplified the target genes in all 3 species. This should make it likely that the amplification efficiency is similar across species. Further, it should make the fluorescence per amplicon (which varies with amplicon length) similar across species. However, for most genes, the sequences varied too much among species to design a single set of primers that would amplify in all 3 species. Additionally, we wished to make comparisons with data from 2 previously reported species (the little brown bat, *Myotis lucifugus*, n=9; and the white-footed mouse, *Peromyscus leucopus*, n=8)(Price et al., 2014), in which amplicon length sometimes differed substantially from the present study. We therefore accounted for differences among species in the following additional two ways.

First, we used LinRegPCR (Ruijter et al., 2009) for calculation of starting concentrations. The LinRegPCR algorithm uses the log-linear amplification phase to calculate efficiency for each sample. We could therefore obtain many semi-independent estimates of amplification efficiency for each gene in each species. As recommended by the program designers (Ruijter et al., 2009), for each gene (i.e., for each amplicon), we averaged the efficiency for each species and applied that average to the calculation for that gene for every sample within the species. Reaction efficiency exceeded 1.75 for all genes in all species. For each gene, we chose a fluorescence threshold that was common to all 5 species (including the 2 previously reported (Price et al., 2014)) before calculating the starting concentration.

Secondly, we corrected for differences in fluorescence per amplicon by dividing the starting concentration (calculated above) by the amplicon length (see mathematical derivation in Appendix). Because we amplified similar sections of the genes in the various species, this correction was generally small (coefficients of variation ranging 0-1%, depending on the gene). However, comparing to our previous study where amplicons were often of different lengths, the correction could be substantially greater (CVs: 0-37%).

In order to examine the composition of the tight junctions, we calculated expression ratios of 2 tight junction genes, thus controlling for the number of tight junctions. The expression ratios we report here are calculated as: (corrected starting concentration of the target gene)/(corrected starting concentration of the denominator gene). Although 2 standard housekeeper genes (EEF1A1 and RPLP0) were available to us, we were concerned that these do not provide appropriate denominators for understanding the composition of tight junctions. This is because whole tissue was used for extractions, and the species may differ in the ratio of enterocytes to other cells (e.g., muscle and connective tissue) in the intestinal samples. A ratio of CLDN2 to EEF1A1 expression, for example, might be informative about interspecific differences in the amount of CLDN2 or the number of tight junctions per mass of intestine, but will not necessarily be informative regarding the compositional makeup of each tight junction, the variable of interest in this study. Similarly, the ratio of a claudin gene to standard housekeepers might be driven by the total number of tight junctions per enterocyte, and not by the composition of the tight junction per se. Ultimately, we wanted to assess the compositional makeup of claudins per tight junction, and therefore we needed to use a gene that is specific to tight junctions in the denominator of the expression ratio. This has the effect of controlling for the total number of tight junctions. Because ZO-1 is thought to be constitutively expressed in

tight junctions (Holmes et al., 2006), we preferred ZO-1 as this denominator gene. Obviously, this choice could affect our results if the species vary in the amount of ZO-1 expressed per tight junction. Therefore we also investigated the use of other tight junction genes as well as more standard housekeeper genes (RPLP0 and EEF1A1) in the denominator of the expression ratios. The consequences of this choice are addressed further in the discussion.

We used Platinum Taq polymerase (Life Technologies; conditions: 1x buffer, 2.5 mM MgCl₂, 0.2 mM dNTPs, 0.25 µM each primer, 1 unit Taq, 0.7x SYBR Green 1, 1 µl template in 20 µl rxn volume). We ran samples in duplicate with a cycling program of 94°C for 10 min, then 40 cycles of 94°C for 30 sec, 55°C for 30 sec, and 72°C for 30 seconds, with fluorescence acquisition at the end of the 72°C step.

Statistics

Differences among species were detected using ANOVA and Tukey's post-hoc tests. Significance was set at $P < 0.05$. Tests were performed with the aid of R software (R Development Core Team, 2010). Principal components analysis was performed in R using expression ratios. When data were missing for the principal components analysis, we imputed data based on species averages, or variable averages when that was not available. Data are presented as means + SE.

Results

Absorption of paracellular probes in intact animals

Absorption of 3OMD-glucose was essentially complete (f not significantly different from unity) in the bat and the vole ($P > 0.6$ for both)(Figure 1). There was no statistical difference in

fractional absorption of 3OMD-glucose among species, but in the hedgehog, 3OMD-glucose absorption was somewhat less than unity ($f=0.84 \pm 0.05$, $P=0.021$). In the bat, arabinose absorption was $85 \pm 14.2\%$, which was significantly higher ($F_{2,17}=8.9$, $P=0.0025$) than the hedgehog ($45 \pm 4.7\%$; $P=0.037$) or the vole ($22 \pm 2.3\%$; $P=0.0025$) (Figure 1).

Morphometry and Histology

The bats had shorter intestines than both the voles and hedgehogs ($F_{2,23}=160$, $P<0.0001$, $P<0.0001$)(Table 1). Hedgehogs also had wider intestines than the voles and bats ($F_{2,24}=94$, $P<0.0001$; $P<0.0001$), although intestinal circumference was not significantly different between the voles and bats ($P=0.17$)(Table 1).

Due to their longer villi, hedgehogs had the highest amplification ratio of the 3 species (Table 3)(Figure 2). Enterocyte width was somewhat lower in the bats compared to the two non-flying species, and the bat had more enterocytes per nominal surface area than the vole. However, due largely to their longer intestines, voles and hedgehogs had a much greater number of cells (and thus tight junctions) summed over the small intestine (Table 3). The nominal surface area of vertebrates scales with $(\text{body mass})^{0.76}$ (Karasov, 2012). If we correct for this scaling relationship, the bats have summed nominal surface areas ($0.92 \pm 0.4 \text{ cm}^2 \text{ g}^{-.76}$) ($F_{2,23}=10.4$, $P=0.0005$) that are lower than voles ($1.36 \pm 0.17 \text{ cm}^2 \text{ g}^{-.76}$)($P=0.012$) and not significantly different from hedgehogs ($0.59 \pm 0.02 \text{ cm}^2 \text{ g}^{-.76}$)($P=0.14$). Similarly we can calculate the number of cells per normalized nominal surface area by dividing the number of cells in the intestine by $(\text{body mass})^{0.76}$. The bats had significantly higher ($F_{2,8}=5.6$, $P=0.03$)

total cells per intestine normalized for body mass (25 ± 3 million cells $\text{g}^{-.76}$) than the voles (17 ± 1 million cells $\text{g}^{-.76}$; $P=0.03$) but not hedgehogs (22 ± 2 million cells $\text{g}^{-.76}$; $P=0.55$).

RNA expression

We tried several ways of calculating the expression ratio, i.e., the ratio of the target gene divided by a denominator gene. As we note in the methods, we prefer using ZO-1 as the denominator gene, because it: 1) is thought to be expressed constitutively in the tight junctions; 2) could be measured in all species and therefore can be used to construct an expression ratio in all species; and 3) is a gene associated with the tight junction, and therefore, using it in the denominator provides a measure of the relative expression of the target gene while controlling for the total number of tight junctions, which may vary among species. The following results are based on using ZO-1 as the denominator gene, but we also show the results expressed using other denominator genes (Supplementary Figures S1-S7).

The CLDN1 expression ratio varied among species ($F_{3,33}=2.9$, $P=0.0474$)(Figure 3). The expression ratio in *P. leucopus* was significantly lower than in *E. fuscus* ($P=0.041$). The CLDN2 expression ratio also varied significantly among species ($F_{3,32}=5.1$, $P=0.0053$), being significantly lower in *M. lucifugus* ($P=0.003$) compared to *E. fuscus*. The CLDN4 expression ratio did not vary significantly among species ($F_{2,26}=2.1$, $P=0.14$). For CLDNs 1, 2, and 4, we tried two primer sets for the hedgehog (*A. albiventris*) but could not produce suitable melt curves.

The CLDN7 expression ratio varied significantly among species ($F_{4,38}=12.6$, $P<0.0001$), with the bats and hedgehog tending to display higher expression ratios than the rodents. The

CLDN15 expression ratio varied significantly among species ($F_{4,38}=4.6$, $p=0.0039$), with *M. lucifugus* displaying higher expression than *A. albiventris* and *P. leucopus* ($P<0.024$ for both). The OCLN expression ratio varied among species ($F_{4,38}=11.6$, $P=0.0013$), with expression in *A. albiventris* and *M. lucifugus* tending to be higher than the rodents (Figure 3).

Given that we used new correction factors, we were interested in whether the expression patterns we reported previously (Price et al., 2014) changed. Although statistical significance was sometimes lost in the current study, the patterns observed were the same (*M. lucifugus* had lower CLDN2 and higher CLDN1 and CLDN15 expression ratios compared to *P. leucopus*). The loss of significance can likely be attributed to lower statistical power due to the addition of the new species. Regardless of statistical significance, however, the expression differences we previously observed between *M. lucifugus* and *P. leucopus* were not replicated in the 3 new species. The CLDN1 expression ratio, for example, was just as high in *M. pennsylvanicus* as the two bats (Figure 3). The expression ratio of CLDN2 was relatively high in *E. fuscus*, whereas it was low in *M. lucifugus*. The pattern of differences between bats and rodents for the CLDN15 expression ratio held up in the new analysis, but the hedgehog had a CLDN15 expression ratio that was similar to that of the bats. We also examined our results using different genes in the denominator of the expression ratios (for example EEF1A1, RPLP0, and CLDN genes), but in no case was there a clear difference between bats and non-flyers (Supplementary Figures S1-S7). Using an average of the expression of EEF1A1 and RPLP0 as the denominator resulted in expression ratios that are qualitatively similar to those using ZO-1 as the denominator (compare Figure 3 with Figure S7).

Our principal components analysis found 2 components that together account for 63% of the variance (Table 4). The first PCA axis had high negative loadings for CLDN2 and CLDN4

and a high positive loading for OCLN. The second axis had a strong negative loading for CLDN15 (Table 4). Plotting the individual scores for the first and second axes, the species did not segregate into distinct groups (Figure 4). The bats tended to have strongly negative PC2 scores, while the vole had positive PC2 scores, although there was substantial overlap among all species (Figure 4).

Discussion

Paracellular nutrient absorption in intact animals

As we have observed in previous comparisons of bats and rodents (Brun et al., 2014; Price et al., 2014), the big brown bat absorbed the majority of the L-arabinose dose, whereas the vole absorbed only a minority. Although we hasten to point out that we have not performed a fully phylogenetically controlled analysis, the hedgehog, which is more closely related to bats than rodents (Nery et al., 2012), provides extra confidence that, among mammals, the phenomenon of high paracellular absorption of nutrient-sized probes is unique to bats. L-arabinose absorption in the hedgehog (45%) and the vole (22%) were near or within in the range observed previously in rodents (15-40%)(Brun et al., 2014; Fasulo et al., 2013b; Price et al., 2014, 2013). Thus, our results lend further support to the hypothesis that flying vertebrates have evolved high intestinal permeability to nutrients to compensate for their smaller intestines (Caviedes-Vidal et al., 2007; Price et al., 2015). The species in this study varied in their source (wild vs. domestic) and duration in captivity (bats were used immediately after capture). Additionally, the natural and captive diets varied among species. These differences may have

affected the animals' stress levels and intestinal function during our experiments. However, the interspecific differences in paracellular nutrient absorption we observed are consistent with those previously observed in other species representing a range of diets and sources (Price et al., 2015).

Due to the size-sieving effect of tight junctions, absorption of larger molecules is inhibited (Anderson and Van Itallie, 2009; Bijlsma et al., 1995; Chediack et al., 2003; Delahunty and Hollander, 1987; Fasulo et al., 2013b). Because of this, and the size differential between L-arabinose (M_r 150) and D-glucose (M_r 180), our measurement of L-arabinose absorption overestimates the paracellular portion of glucose absorption. Nonetheless, our measurements show distinct differences among species in the paracellular absorption of nutrient-sized molecules, and this very likely extends to glucose as well.

Histological Measurements

The high level of L-arabinose absorption in bats could potentially derive from having more tight junctions across which L-arabinose can absorb. Compared to the voles and hedgehogs, big brown bats have shorter and narrower intestines, i.e., the bats have a smaller nominal surface area (the surface area considering the intestine a smooth-bore tube). This could potentially be offset by greater villous amplification or enterocyte density in the bats. However, while the bats had somewhat smaller (and therefore more densely packed) enterocytes than either of the non-flyers, villous amplification was similar in the bats and voles, and even higher in the hedgehogs. Thus, for a given quantity of nominal surface area, bats and hedgehogs had a similar number of enterocytes, and presumably, tight junctions. Although the number of cells per

nominal surface area was greater in the bats than the voles, when we take into account the greater surface area of the voles, the total number of enterocytes in the small intestine is greater in voles than bats. Moreover, the total number of enterocytes in hedgehog small intestine was approximately 10 fold greater than the bats. When controlling for the effect of body mass on nominal surface area, bats had more total enterocytes than the vole but not the hedgehog. However, the proximate factors controlling absorption of paracellular probes should be the permeability of individual tight junctions, the absolute number of tight junctions, and the transit time, the latter of which was not measured here but is generally much faster in bats (Buchler, 1975; Klite, 1965; Price et al., 2015) than non-flyers. Bats cannot, therefore, obtain their high rates of paracellular nutrient absorption simply by having more junctions between cells through which the nutrients can be absorbed.

mRNA expression of tight junction-associated genes

An alternative mechanism for explaining the high level of paracellular nutrient absorption in bats is that the tight junctions of bats are somehow more permeable to nutrient-sized molecules. While it is unclear what characteristics determine tight junction permeability to nutrients, it is thought that the relative expression of the proteins that form the tight junction – CLDNs and OCLN – could be responsible (Günzel and Yu, 2013). We therefore investigated the expression of several tight junction genes that are known to be highly expressed in the small intestine (Holmes et al., 2006). For this, we focused on the ratios of expression of the several CLDNs and OCLN to expression of ZO-1, so as to control for any differences among species in the number of tight junctions per sampled tissue. This should have the effect of comparing among species the relative amount of each CLDN or OCLN expressed per tight junction.

The patterns of gene expression we observed in these 3 species, in combination with those previously reported (Price et al., 2014), did not demonstrate a clear difference between bats and non-flying mammals. We previously reported that the bat *M. lucifugus* had higher CLDN1 expression than the cricetid mouse *P. leucopus* (Price et al., 2014). While CLDN1 could not be measured in hedgehogs, its expression ratio in the vole was similar to that in both bat species. We also reported a low expression ratio of CLDN2 in *M. lucifugus* relative to *P. leucopus* (Price et al., 2014), but the CLDN2 expression ratio was much higher in the bat *E. fuscus*, and was similar to that of the vole. We did observe a replication of the pattern in CLDN15, to the extent that both bats had (non-significantly) higher CLDN15 expression ratios than both rodents. However, the hedgehog had a CLDN15 expression ratio similar to the bats, and therefore there was not a general difference between the flyers and non-flyers. The principal components analysis confirmed these findings. The five species had substantial overlap on their PC axes, and the 2 bats did not form a coherent and distinct group apart from the non-flyers.

Although these data do not support the hypothesis that interspecific differences in paracellular nutrient absorption are determined by the composition of claudins in the tight junction, we caution several caveats. We only measured expression at a single point along the intestine. If expression patterns vary considerably along the length of the intestine (Holmes et al., 2006), or if the point we chose is not analogous among species, then our data may not be easily comparable across species. Additionally, the composition of tight junctions, in terms of proteins present, may differ from gene expression patterns due to differences in translation and protein catabolism, as well as shuttling of proteins between tight junctions and intracellular pools (Shen et al., 2011, 2008).

425 *Methodological considerations*

426 Gene expression is most commonly compared within a species across treatments. This
427 design allows certain assumptions in the methodology. These include assumptions that the
428 fluorescence per amplicon is constant and that the reaction efficiency is constant, at least within a
429 range of starting concentrations. Our cross-species design necessitated evaluation of these
430 assumptions and/or adjustments for deviations. When possible, we designed primers that could
431 be used to amplify genes in several species. This methodological step alone should largely
432 satisfy both assumptions, because reaction efficiency is primarily a function of primer binding,
433 and similar-length amplicons of the same gene should have similar fluorescence. Using the same
434 primers for multiple species also has the advantage of being able to use a single mastermix for all
435 samples.

436 When different primer sets are necessary, adjustments must be made for both the
437 fluorescence per amplicon and the reaction efficiency. Determining the fluorescence per
438 amplicon is perhaps the most difficult aspect. While amplicon length affects the fluorescence of
439 Sybr Green, it is not clear exactly why this is. Sybr Green exhibits both intercalation and surface
440 binding to DNA (Zipper et al., 2004). The increase in fluorescence with increasing amplicon
441 length has been found to be driven mostly by adenine-thymine (AT) content (Colborn et al.,
442 2008; Zipper et al., 2004), but contrasting studies attribute the most of the increase in
443 fluorescence to guanine-cytosine (GC) content (Giglio et al., 2003; Gudnason et al., 2007). The
444 fact that various studies have found opposing results suggests that SYBR Green
445 binding/fluorescence may vary with specific DNA sequences. In light of this possibility, we
446 believe our practice of using total amplicon length (which will also tend to correlate with other
447 variables such as AT content and GC content) is the best way to normalize among different

amplicons. As further confirmation of the robustness of our results, normalizing using AT content instead of total amplicon length would not have changed our conclusions (data not shown). We also note that our conclusions would be substantially similar even if we had restricted our analysis to only the 3 new species of this study, which all had quite similar amplicons.

An additional consideration when comparing across species is the choice of an appropriate denominator gene to form an expression ratio. We used ZO-1 as our primary denominator gene to assess the compositional makeup of the tight junctions. This choice was based primarily on the idea that using a ratio of two tight junction genes should control for any among-species differences in tight junctions, and thus, this ratio should reflect the proportion that any particular claudin or occludin makes up in the tight junction. Alternatively, we could have used standard reference genes, such as the *EEF1A1* and *RPLP0* that we measured, in the denominator. Using ratios based on standard reference genes should reflect the amount of each claudin per tissue that was sampled, which would be affected by both the abundance of the claudin in each tight junction and the total number of tight junctions. This might give results that are difficult to interpret if, for example, species varied in the amount of muscular or connective tissue per amount of epithelial layer. In our data, we would have come to similar conclusions using either denominator (compare Figure 3 with Figure S7), although the pattern of gene expression would be slightly different. The similarity of these two results may serve as confirmation that ZO-1 is constitutively expressed in tight junctions, or might indicate that our species have a fairly similar ratio of tight junctions to total intestine.

Conclusions

In agreement with previous studies, the big brown bat absorbed much more of an L-arabinose dose than either the vole or hedgehog. This difference cannot be explained by the bat having more enterocytes, and presumably more tight junctions, in their intestines; in fact, hedgehogs have 10-fold more enterocytes than the bat. The differences in paracellular permeability of the intestine to nutrient-sized molecules is therefore likely mediated by differences in the permeability of the tight junctions in the various species. Our gene expression profiles of tight junction-associated genes failed to detect a consistent pattern of expression differences between bats and non-flyers, although a larger dataset might yet shed more light on this problem.

Acknowledgements

We thank Jim Gammeter, Jerry Goth (Swamp Lover's Conservancy), Zach DeQuattro, the UW Lakeshore Nature Preserve, and the City of Madison Parks Division for providing access to lands and trapping assistance. Zach Peery graciously shared his lab equipment. Cherry Tsai provided laboratory assistance and we thank Becky Kirby for helpful discussions of reaction efficiency. We thank Chelsea Magruder for her mouse image. An anonymous reviewer provided helpful comments on the manuscript. Funding was provided by the National Science Foundation (IOS-1025886 to WHK and ECV) and the Department of Forest and Wildlife Ecology at the University of Wisconsin-Madison.

491 **Literature Cited:**

- 492 Amasheh, S., Meiri, N., Gitter, A.H., Schöneberg, T., Mankertz, J., Schulzke, J.D., Fromm, M., 2002.
 493 Claudin-2 expression induces cation-selective channels in tight junctions of epithelial cells. *J. Cell*
 494 *Sci.* 115, 4969–4976. doi:10.1242/jcs.00165
- 495 Anderson, J.M., Van Itallie, C.M., 2009. Physiology and function of the tight junction. *Cold Spring Harb.*
 496 *Perspect. Biol.* 1, a002584. doi:10.1101/cshperspect.a002584
- 497 Benson, D.A., Karsch-Mizrachi, I., Clark, K., Lipman, D.J., Ostell, J., Sayers, E.W., 2012. GenBank.
 498 *Nucleic Acids Res.* 40, D48–53. doi:10.1093/nar/gkr1202
- 499 Bijlsma, P.B., Peeters, R.A., Groot, J.A., Dekker, P.R., Taminiau, J.A., Van Der Meer, R., 1995.
 500 Differential in vivo and in vitro intestinal permeability to lactulose and mannitol in animals and
 501 humans: a hypothesis. *Gastroenterology* 108, 687–696.
- 502 Brody, T., 1999. *Nutritional Biochemistry*. Academic Press, San Diego.
- 503 Brun, A., Price, E.R., Gontero-Fourcade, M.N., Fernandez-Marinone, G., Cruz-Neto, A.P., Karasov,
 504 W.H., Caviades-Vidal, E., 2014. High paracellular nutrient absorption in intact bats is associated
 505 with high paracellular permeability in perfused intestinal segments. *J. Exp. Biol.* 217, 3311–3317.
- 506 Buchler, E.R., 1975. Food transit time in *Myotis lucifugus* Chiroptera: Vespertilionidae. *J. Mammal.* 56,
 507 252–255.
- 508 Caviades-Vidal, E., Karasov, W.H., Chediack, J.G., Fasulo, V., Cruz-Neto, A.P., Otani, L., 2008.
 509 Paracellular absorption: a bat breaks the mammal paradigm. *PLoS One* 3, e1425.
 510 doi:10.1371/journal.pone.0001425
- 511 Caviades-Vidal, E., McWhorter, T.J., Lavin, S.R., Chediack, J.G., Tracy, C.R., Karasov, W.H., 2007. The
 512 digestive adaptation of flying vertebrates: high intestinal paracellular absorption compensates for
 513 smaller guts. *Proc. Natl. Acad. Sci. U. S. A.* 104, 19132–19137.
- 514 Chediack, J.G., Caviades-Vidal, E., Fasulo, V., Yamin, L.J., Karasov, W.H., 2003. Intestinal passive
 515 absorption of water-soluble compounds by sparrows: effect of molecular size and luminal nutrients.
 516 *J. Comp. Physiol. B.* 173, 187–197. doi:10.1007/s00360-002-0314-8
- 517 Colborn, J.M., Byrd, B.D., Koita, O.A., Krogstad, D.J., 2008. Estimation of copy number using SYBR
 518 Green: confounding by AT-rich DNA and by variation in amplicon length. *Am. J. Trop. Med. Hyg.*
 519 79, 887–892.
- 520 Delahunty, T., Hollander, D., 1987. A comparison of intestinal permeability between humans and three
 521 common laboratory animals. *Comp. Biochem. Physiol. A. Comp. Physiol.* 86, 565–567.
- 522 Fasulo, V., Zhang, Z., Chediack, J.G., Cid, F.D., Karasov, W.H., Caviades-Vidal, E., 2013a. The capacity
 523 for paracellular absorption in the insectivorous bat *Tadarida brasiliensis*. *J. Comp. Physiol. B* 183,
 524 289–96. doi:10.1007/s00360-012-0696-1

525 Fasulo, V., Zhang, Z., Price, E.R., Chediack, J.G., Karasov, W.H., Caviedes-Vidal, E., 2013b.
 526 Paracellular absorption in laboratory mice: Molecule size-dependent but low capacity. *Comp.*
 527 *Biochem. Physiol. Part A.* 164, 71–76.

528 Giglio, S., Monis, P.T., Saint, C.P., 2003. Demonstration of preferential binding of SYBR Green I to
 529 specific DNA fragments in real-time multiplex PCR. *Nucleic Acids Res.* 31, e136.

530 Gudnason, H., Dufva, M., Bang, D.D., Wolff, A., 2007. Comparison of multiple DNA dyes for real-time
 531 PCR: effects of dye concentration and sequence composition on DNA amplification and melting
 532 temperature. *Nucleic Acids Res.* 35, e127.

533 Günzel, D., Yu, A.S.L., 2013. Claudins and the modulation of tight junction permeability. *Physiol. Rev.*
 534 93, 525–69. doi:10.1152/physrev.00019.2012

535 Holmes, J.L., Van Itallie, C.M., Rasmussen, J.E., Anderson, J.M., 2006. Claudin profiling in the mouse
 536 during postnatal intestinal development and along the gastrointestinal tract reveals complex
 537 expression patterns. *Gene Expr. Patterns* 6, 581–588. doi:10.1016/j.modgep.2005.12.001

538 Karasov, W.H., 2012. Terrestrial vertebrates, in: Sibly, R.M., Brown, J.H., Kodric-Brown, A. (Eds.),
 539 *Metabolic Ecology: A Scaling Approach*. Wiley-Blackwell, Chichester, UK, pp. 212–224.

540 Karasov, W.H., Caviedes-Vidal, E., Bakken, B.H., Izhaki, I., Samuni-Blank, M., Arad, Z., 2012. Capacity
 541 for Absorption of Water-Soluble Secondary Metabolites Greater in Birds than in Rodents. *PLoS*
 542 *One* 7, e32417. doi:10.1371/journal.pone.0032417

543 Keegan, D.J., 1984. Glucose absorption in the fruit bat studied using the intestinal ring method (abstract).
 544 *S. Afr. J. Sci.* 80, 132.

545 Keegan, D.J., 1980. The lack of an active glucose transport system in the bat intestine (abstract). *S. Afr. J.*
 546 *Sci.* 76, 570–571.

547 Kisielinski, K., Willis, S., Prescher, A., Klosterhalfen, B., Schumpelick, V., 2002. A simple new method
 548 to calculate small intestine absorptive surface in the rat. *Clin. Exp. Med.* 2, 131–135.

549 Klite, P.D., 1965. Intestinal bacterial flora and transit time of three neotropical bat species. *J. Bacteriol.*
 550 90, 375–379.

551 Kurta, A., Baker, R.H., 1990. *Eptesicus fuscus*. *Mamm. Species* 356, 1–10.

552 Larkin, M.A., Blackshields, G., Brown, N.P., Chenna, R., McGettigan, P.A., McWilliam, H., Valentin, F.,
 553 Wallace, I.M., Wilm, A., Lopez, R., Thompson, J.D., Gibson, T.J., Higgins, D.G., 2007. Clustal W
 554 and Clustal X version 2.0. *Bioinformatics* 23, 2947–2948.

555 Lavin, S.R., McWhorter, T.J., Karasov, W.H., 2007. Mechanistic bases for differences in passive
 556 absorption. *J. Exp. Biol.* 210, 2754–2764. doi:10.1242/jeb.006114

557 Lindroth, R.L., Batzli, G.O., 1984. Food habits of the meadow vole (*Microtus pennsylvanicus*) in
 558 bluegrass and prairie habitats. *J. Mammal.* 65, 600–606.

559 McCarthy, K.M., Francis, S.A., McCormack, J.M., Lai, J., Rogers, R.A., Skare, I.B., Lynch, R.D.,
 560 Schneeberger, E.E., 2000. Inducible expression of claudin-1-myc but not occludin-VSV-G results in
 561 aberrant tight junction strand formation in MDCK cells. *J. Cell Sci.* 113, 3387–3398.

562 Nery, M.F., González, D.J., Hoffmann, F.G., Opazo, J.C., 2012. Resolution of the laurasian phylogeny:
 563 Evidence from genomic data. *Mol. Phylogenet. Evol.* 64, 685–689.

564 Price, E.R., Brun, A., Caviedes-Vidal, E., Karasov, W.H., 2015. Digestive adaptations of aerial lifestyles.
 565 *Physiology* 30, 69–78.

566 Price, E.R., Rott, K.H., Caviedes-Vidal, E., Karasov, W.H., 2014. Paracellular nutrient absorption is
 567 higher in bats than rodents: integrating from intact animals to the molecular level. *J. Exp. Biol.* 217,
 568 3483–3492.

569 Price, E.R., Ruff, L.J., Guerra, A., Karasov, W.H., 2013. Cold exposure increases intestinal paracellular
 570 permeability to nutrients in the mouse. *J. Exp. Biol.* 216, 4065–70. doi:10.1242/jeb.088203

571 R Development Core Team, 2010. R: A language and environment for statistical computing.

572 Rozen, S., Skaletsky, H.J., 2000. Primer3 on the WWW for general users and for biologist programmers,
 573 in: *Bioinformatics Methods and Protocols: Methods in Molecular Biology*. Humana Press, Totowa,
 574 NJ, pp. 365–386.

575 Ruijter, J.M., Rarakers, C., Hoogaars, W.M.H., Karlen, Y., Bakker, O., van den Hoff, M.J.B., Moorman,
 576 A.F.M., 2009. Amplification efficiency: linking baseline and bias in the analysis of quantitative
 577 PCR data. *Nucleic Acids Res.* 37, e45.

578 Santana, E.M., Jantz, H.E., Best, T.L., 2010. *Atelerix albiventris* (Erinaceomorpha: Erinaceidae). *Mamm.*
 579 *Species* 42, 99–110. doi:10.1644/857.1.Key

580 Schulzke, J.D., Gitter, A.H., Mankertz, J., Spiegel, S., Seidler, U., Amasheh, S., Saitou, M., Tsukita, S.,
 581 Fromm, M., 2005. Epithelial transport and barrier function in occludin-deficient mice. *Biochim.*
 582 *Biophys. Acta* 1669, 34–42. doi:10.1016/j.bbamem.2005.01.008

583 Shen, L., Weber, C.R., Raleigh, D.R., Yu, D., Turner, J.R., 2011. Tight junction pore and leak pathways:
 584 a dynamic duo. *Annu. Rev. Physiol.* 73, 283–309. doi:10.1146/annurev-physiol-012110-142150

585 Shen, L., Weber, C.R., Turner, J.R., 2008. The tight junction protein complex undergoes rapid and
 586 continuous molecular remodeling at steady state. *J. Cell Biol.* 181, 683–695.

587 Tamura, A., Hayashi, H., Imasato, M., Yamazaki, Y., Hagiwara, A., Wada, M., Noda, T., Watanabe, M.,
 588 Suzuki, Y., Tsukita, S., 2011. Loss of claudin-15, but not claudin-2, causes Na⁺ deficiency and
 589 glucose malabsorption in mouse small intestine. *Gastroenterology* 140, 913–923.
 590 doi:10.1053/j.gastro.2010.08.006

591 Tracy, C.R., McWhorter, T.J., Korine, C., Wojciechowski, M.S., Pinshow, B., Karasov, W.H., 2007.
 592 Absorption of sugars in the Egyptian fruit bat (*Rousettus aegyptiacus*): a paradox explained. *J. Exp.*
 593 *Biol.* 210, 1726–1734. doi:10.1242/jeb.02766

594 Van Itallie, C., Rahner, C., Anderson, J.M., 2001. Regulated expression of claudin-4 decreases
595 paracellular conductance through a selective decrease in sodium permeability. J. Clin. Invest. 107,
596 1319–1327.

597 Zipper, H., Brunner, H., Bernhagen, J., Vitzthum, F., 2004. Investigations on DNA intercalation and
598 surface binding by SYBR Green I, its structure determination and methodological implications.
599 Nucleic Acids Res. 32, e103.

600

601

Figure Captions:

Figure 1: Fractional absorption of a glucose analog (3-O-methyl-D-glucose; 3-OMG) and a paracellularly absorbed probe (L-arabinose) in 8 big brown bats (*Eptesicus fuscus*), 6 hedgehogs (*Atelerix albiventris*), and 5 meadow voles (*Microtus pennsylvanicus*). Fractional absorption is calculated as oral bioavailability, and represents the proportion of an oral dose that is absorbed by the animal. Fractional absorption of L-arabinose was significantly higher in the bat than either of the non-flying mammals. Data are means + SE.

Figure 2: Representative views of intestinal villi in *Eptesicus fuscus* (top row), *Microtus pennsylvanicus* (middle row), and *Atelerix albiventris* (bottom row). Left to right, sections from the proximal, medial, and distal thirds of the intestine are shown. The bar shows 100 μ m.

Figure 3: Expression ratios of several claudins (CLDN) and occludin (OCLN) in the intestine of 2 bats, 2 rodents, and a hedgehog. Expression ratio is presented as the ratio of target gene expression to expression of ZO-1. Expression was not measured in the hedgehog (*A. albiventris*) for CLDN1, CLDN2, and CLDN4, nor in *P. leucopus* for CLDN4. For a given gene, bars sharing similar letters do not differ significantly ($P>0.05$). $N_{M. lucifugus}=9$, $N_{E. fuscus}=11$, $N_{P. leucopus}=8$, $N_{M. pennsylvanicus}=9$, $N_{A. albiventris}=6$. Data on *M. lucifugus* and *P. leucopus* are recalculated from Price et al. (2014). Data are means + SE.

622 Figure 4: Principal components analysis of gene expression ratios of several claudins and
623 occludin in the intestine of 2 bats, 2 rodents, and a hedgehog. The first principal component is
624 plotted on the horizontal axis and the second principal component is plotted on the vertical axis.
625 Ellipses are 95% confidence ellipses for each species.

626

627 Supplementary Figures S1-S7: qPCR data presented with various choices of denominators to
628 make the expression ratio. All data are ratios of the target gene (labeled on the horizontal axis)
629 divided by the denominator gene and are means + SE. Denominator genes are as follows: Fig
630 S1: CLDN1; Fig S2: CLDN2; Fig S3: CLDN4; Fig S4: CLDN7; Fig S5: CLDN15; Fig S6:
631 OCLN; Fig S7: a geometric average of EEF1A1 and RPLP0.

632

633 **Table 1: Animal Attributes**

	<i>Eptesicus fuscus</i>	<i>Atelerix albiventris</i>	<i>Microtus</i> <i>pennsylvanicus</i>
N (#♂ / #♀)	5 / 6	0 / 6	7 / 2
Body Mass (g)	17.9 ± 1.1	439 ± 33	35.4 ± 4.1
Body Length (cm; snout to base of tail)	7.1 ± 0.2	18.8 ± 0.8	10.86 ± 0.5
Small Intestine Length (cm)	12.4 ± 0.6 ¹	54.1 ± 3.6 ¹	26.3 ± 1.3
Small Intestine Circumference (mm)	6.58 ± 0.22	11.06 ± 0.19	7.59 ± 0.33
Cecum Mass (g wet; including contents)	Absent	Absent	1.75 ± 0.17
Large Intestine Length (cm)	n.m. ¹	n.m. ¹	11.6 ± 0.73

634 ¹n.m.=not measured. In *E. fuscus* and *A. albiventris*, the large intestine is short and difficult to
635 distinguish from the small intestine macroscopically. For these animals, the small intestine
636 length listed in this table is the length of the whole intestine.

Table 2: Primers used for qPCR and the GenBank accessions upon which they were based¹.

Gene	Forward Primer (5' to 3')	Reverse Primer (5' to 3')	GenBank Accessions
Zonula occludens 1 (ZO-1)	<i>M. pennsylvanicus</i> and <i>A. albiventris</i> : CGTAGCTCTGGCATTATTCG <i>E. fuscus</i> : CGTAGCTCCGGCATCATCCG	<i>M. pennsylvanicus</i> and <i>A. albiventris</i> : CCTGGCACTTTTCCGAGA <i>E. fuscus</i> : CCTGGCACTCTTCCGAGA	<u>KJ420409-</u> <u>KJ420412</u>
Occludin (OCLN)	<i>M. pennsylvanicus</i> : ATCCTGGGCATCCTGGTGT <i>A. albiventris</i> : GTCCTGGGCATTATGGTGT <i>E. fuscus</i> : ATCCTGGGCTTCATGGTGT	All 3 species: GGGATCCACCACACAGTAGT	<u>KJ451437-</u> <u>KJ451442</u>
Claudin-1 (CLDN1)	<i>E. fuscus</i> and <i>A. albiventris</i> : TGCGGATGGCTGTCATTG <i>M. pennsylvanicus</i> : TGTGGATGGCTGTCATTG	All 3 species: CAGCCCAGCCAGTGAAGAG	<u>KJ420416-</u> <u>KJ420420</u>
Claudin-2 (CLDN2)	<i>M. pennsylvanicus</i> and <i>A. albiventris</i> : CAAGTTCTTATGTTGGTGCCAGC <i>E. fuscus</i> : CAAGTTCTTATGTTGGTGCTAGC	<i>M. pennsylvanicus</i> and <i>A. albiventris</i> : CTGGGCAGCCTGGATGT <i>E. fuscus</i> : CTGAGCAGCCTGGATGT	<u>KJ451447-</u> <u>KJ451450</u>
Claudin-4 (CLDN4)	All 3 species: AACTGCGTGGTGCAAGAGC	<i>E. fuscus</i> and <i>M. pennsylvanicus</i> : GCAGTTGGTGCACTTGCC	<u>KJ405716-</u> <u>KJ406719</u>

	<i>A. albiventris</i> :		
	GCAGTTGGTACATTTGCC		
Claudin-7	All 3 species:	All 3 species:	<u>KJ451444-</u>
(CLDN7)	TACGACTCGGTGCTCGC	AGCAAGACCTGCCACGAT	<u>KJ451446</u>
Claudin-15	<i>M. pennsylvanicus</i> :		<u>KJ502667-</u>
(CLDN15)	GGAGTTCCCTTCCATGCT	<i>M. pennsylvanicus</i> and <i>E. fuscus</i> :	<u>KJ502669</u>
	<i>E. fuscus</i> : GGAGTTCCCGTCCATGCT	CCAGGAAGCCCAGGAGG	
	<i>A. albiventris</i> :	<i>A. albiventris</i> :	
	CGAGTTCCCGTCCCTGC	CCAGACCGCCCATCAG	
EEF1A1	All 3 species:	All 3 species:	<u>KJ396960-</u>
	CAGCACCTACATTAAGAAAATTGG	CCCTTGAACCAAGGCATATT	<u>KJ396962</u>
RPLP0	All 3 species:	All 3 species	<u>KJ396955-</u>
	GCGACCTGGAAGTCCAATA	ATCTGCTGCATCTGCTTGG	<u>KJ396959</u>

¹Primers and accession numbers for *M. lucifugus* and *P. leucopus* can be found in Price et al. (2014).

Table 3: Histological measurements from small intestine of big brown bats (*Eptesicus fuscus*), meadow voles (*Microtus pennsylvanicus*) and hedgehogs (*Atelerix albiventris*).

	<i>E. fuscus</i> (n=3)	<i>M. pennsylvanicus</i> (n=5)	<i>A. albiventris</i> (n=6)	F	P-value
Villus length (µm)					
Proximal	571 ± 45 ^a	602 ± 44 ^a	1220 ± 41 ^b	F _{2,11} =74	<0.0001
Medial	445 ± 40 ^a	583 ± 27 ^a	999 ± 37 ^b	F _{2,11} =66	<0.0001
Distal	313 ± 38 ^a	432 ± 35 ^a	731 ± 63 ^b	F _{2,8} =20	0.0008
Villus width (µm)					
Proximal	72 ± 8.0 ^a	94 ± 6.6 ^a	166 ± 4.9 ^b	F _{2,11} =64	<0.0001
Medial	74 ± 5.6 ^a	86 ± 2.6 ^a	159 ± 4.0 ^b	F _{2,11} =138	<0.0001
Distal	73 ± 7.3 ^a	77 ± 4.0 ^a	134 ± 5.7 ^b	F _{2,8} =35	0.0001
Crypt Width (µm)					
Proximal	73 ± 6.9 ^a	86 ± 3.3 ^{a,b}	155 ± 1.3 ^b	F _{2,11} =12	0.0017
Medial	37 ± 1.8 ^a	43 ± 1.7 ^a	49 ± 1.4 ^b	F _{2,11} =14.7	0.0008
Distal	39 ± 2.3	39 ± 1.3	48 ± 1.2	F _{2,8} =0.36	0.7079
Amplification Ratio					
Proximal	14.8 ± 2.0 ^{a,b}	12.6 ± 0.80 ^a	18.0 ± 0.77 ^b	F _{2,11} =8	0.0072
Medial	11.0 ± 1.3 ^a	13.3 ± 0.48 ^{a,b}	15.2 ± 0.52 ^b	F _{2,11} =8.7	0.0054
Distal	8.1 ± 1.1 ^a	10.7 ± 0.48 ^{a,b}	13.1 ± 0.90 ^b	F _{2,8} =9.35	0.0081
Enterocyte Width (µm)					
Proximal	6.39 ± 0.42 ^a	8.33 ± 0.15 ^c	7.47 ± 0.11 ^b	F _{2,11} =20.5	0.0001
Medial	6.34 ± 0.32 ^a	8.23 ± 0.18 ^c	7.49 ± 0.12 ^b	F _{2,11} =21.6	0.0002
Distal	6.53 ± 0.33 ^a	8.14 ± 0.18 ^b	7.64 ± 0.04 ^b	F _{2,8} =14.9	0.0020
Cells per nominal surface area (10 ⁶)					

cells per cm²)

Proximal	36.2 ± 3.8^a	18.2 ± 1.0^b	32.4 ± 1.7^a	$F_{2,11}=22.2$	0.0001
Medial	27.2 ± 0.6^a	19.7 ± 0.8^b	27.3 ± 1.6^a	$F_{2,11}=10.5$	0.0028
Distal	$18.8 \pm 1.3^{a,b}$	16.2 ± 0.8^a	22.4 ± 1.7^b	$F_{2,8}=6.6$	0.0199
Summed cells of					
the small intestine					
(10 ⁶ cells)	206 ± 42^a	308 ± 24^a	2308 ± 115^b	$F_{2,8}=355$	<0.0001

F and P values refer to an ANOVA for differences among species.

Within a row, values that share letters do not differ significantly ($P>0.05$) with Tukey's HSD test.

Table 4: Factor loadings for the first two axes of a principal components analysis of tight junction gene expression.

	PC1	PC2
Proportion Explained:	0.368	0.293
Factor loadings:		
Cldn1/ZO1	0.892	-0.575
Cldn2/ZO1	-1.2279	-0.717
Cldn4/ZO1	-1.1777	-0.984
Cldn7/ZO1	0.594	-0.612
Cldn15/ZO1	0.231	-1.375
Ocln/ZO1	1.118	-0.7573

Appendix:

After correction for background fluorescence (Ruijter et al., 2009), the equation relating the fluorescence at threshold (F_t) to the number of starting copies (N_0) is:

$$F_t = N_0 \cdot E^{C_t} \cdot S \quad (\text{eq. 1})$$

, where E = reaction efficiency, with range [1,2];

C_t = cycle count when the fluorescence threshold is reached; and

S = fluorescence per single copy of the amplicon

Rearranging,

$$N_0 = F_t / (E^{C_t} \cdot S) \quad (\text{eq. 2})$$

With typical qPCR, in which differences in expression are measured within a species, S is often dropped in practice. This is because it is an unknown quantity, and because it is a constant across treatments, and therefore is not necessary for calculating “fold difference”. For example, the fold difference of individual 1 versus individual 2 is:

$$N_{0(1)} / N_{0(2)} = \{F_t / (E^{C_{t(1)}} \cdot S)\} / \{F_t / (E^{C_{t(2)}} \cdot S)\} = E^{C_{t(2)}/E^{C_{t(1)}}} = E^{\{C_{t(2)}-C_{t(1)}\}} \quad (\text{eq. 3})$$

The LinRegPCR program (Ruijter et al., 2009), for example, presents starting concentrations as:

$$N_{0\text{LinRegPCR}} = F_t / E^{C_t} \quad (\text{eq. 4})$$

However, the primers and amplicons of different species (denoted here by species A, B, etc.) may differ in both E and S . We can chose to set F_t equal for both species, and we can calculate E for each species individually. For example, for species A, eq. 4 will now be calculated:

$$N_{0\text{LinRegPCR}} = F_t / E_A^{C_t} \quad (\text{eq. 5})$$

, where E_A is the reaction efficiency for species A

However, we must still account for interspecific differences in S . Here, we assume S for species A is a function of the length of the amplicon for species A and a fluorescence constant k_F that describes the fluorescence of SYBR Green when bound to double-stranded DNA. This k_F is invariant across amplicons. Thus,

$$S_A = \text{Length}_A \cdot k_F \quad (\text{eq. 6})$$

, where Length_A = the amplicon length for species A

Measuring expression for species A then derives from substituting eq. 5 into eq. 2 while making it specific to species A:

$$N_0 = N_{0\text{LinRegPCR}}/S_A \quad (\text{eq. 7})$$

Substituting using eq. 6,

$$N_0 = N_{0\text{LinRegPCR}}/\{\text{Length}_A \cdot k_F\} \quad (\text{eq. 8})$$

Because k_F is constant across amplicons, we can drop it for all subsequent calculations of fold difference or expression ratios. Thus we simplify to our final equation:

$$N_0 = N_{0\text{LinRegPCR}}/\text{Length}_A \quad (\text{eq. 9})$$

Figure 1:

Figure 2:

Figure 3:

Figure 4: