PALEOMAGNETISM OF UPPER EDIACARAN CLASTICS FROM THE SOUTH URALS: IMPLICATIONS TO PALEO GEOGRAPHY OF BALTICA AND THE OPENING OF THE IAPETUS OCEAN

Natalia M. Levashova^{1, #}, Mikhail L. Bazhenov¹, Joseph G. Meert², Konstantin N. Danukalov³,
Inessa V. Golovanova³, Nikolay B. Kuznetsov¹, Natalia M. Fedorova¹

¹ Geological Institute, Russian Academy of Sciences, Pyzhevsky Lane, 7, Moscow, 119017 Russia

² Department of Geological Sciences, 355 Williamson Hall, Gainesville, FL 32611, USA

³ Institute of Geology, Ufa Scientific Center, Russian Academy of Sciences, Karl Marx st., 16/2, Ufa, 450000 Russia

Abstract

The progress in understanding the evolution of the Earth during the Ediacaran-Cambrian is greatly hindered by the scarcity and inconsistency of paleomagnetic data for this time interval. In order to acquire new data and clarify the confusing situation, Upper Ediacaran clastic rocks of the Basu Formation were sampled at several localities in the westernmost parts of the South Urals that is the deformed margin of Baltica at least since the beginning of the Neoproterozoic. With the aid of stepwise thermal demagnetization, a dual-polarity high-temperature component (HTC) was reliably isolated from grey and maroon sandstones and siltstones at 34/49 sites. The HTC mean direction $D^\circ = 55$, inclination $I^\circ = -35$, ($k = 31$, $\alpha_{95}^\circ = 4.5$) corresponds to a paleolatitude of $19^\circ \pm 3^\circ$. The reversal and fold tests are positive for the HTC. The slump test on two meter-sized slumps shows that the HTC predates slumping in one case and is coeval with it, in the other, thus convincingly indicating the primary origin of the HTC. Also, we demonstrate that inclination shallowing is either absent altogether, or, at worst, less than 10° , in these rocks; hence the position of Baltica can be reliably reconstructed for time 560-575 Ma. We reviewed paleomagnetic data with ages from 615 to 530 Ma for Baltica and Laurentia and come to the conclusion that there is still no uncontested scenario for the opening of the Iapetus Ocean that is based on non-controversial geologic and paleomagnetic data.

Key words: Late Ediacaran, Baltica, West Urals, paleomagnetism, primary remanence, Iapetus Ocean

[#] Corresponding author

INTRODUCTION

The Ediacaran-Early Ordovician interval is one of the most intriguing epochs of geologic time. The onset of the Ediacaran corresponds to the approximate end of the so-called “Snowball glaciations” and the terminal Ediacaran (~542 Ma) is followed by a great radiation of life forms commonly referred to as the “Cambrian Explosion”. It has been argued that Ediacaran-Cambrian time period was also a time where continents underwent rapid changes in their latitudinal positions due to unusually rapid plate motions (Meert et al., 1993), inertial interchange true polar wander (Kirschvink et al., 1997), true polar wander (Evans, 1998) or interactions with superplumes (Meert and Tamrat, 2004). Abrajevitch and Van der Voo (2010) argue that the seeming complexities of continental motion in the Ediacaran-Cambrian are the result of the geomagnetic field switching between a dominant axial dipole field and an equatorial dipole field. These changes may be interrelated and connected, at least in part, to the distribution of continents across the globe.

The chief tool in deciphering paleogeography is paleomagnetism, and there are numerous attempts to provide a robust Ediacaran-Cambrian paleogeography using paleomagnetic data. Yet, it is precisely the Ediacaran-Cambrian time period (~635-488 Ma) where the paleomagnetic data are the most controversial (McCausland et al., 2007; Meert et al., 2007; Meert et al., 1998, Pisarevsky et al., 2008; Abrajevitch and Van der Voo, 2010; Meert, 2014). The problems are particularly acute for Laurentia and Baltica; most probably because they have the largest dataset.

There are very few, if any, Cambrian - Ediacaran targets on the platform of Baltica that are suitable for paleomagnetic studies and remain unstudied. However, there is a nearly thousand-kilometer-long band of Neoproterozoic and Ediacaran rocks in the western part of the South and Middle Urals along the Baltica margin (Fig. 1a). In particular, a nearly two kilometer thick Ediacaran terrigenous sequence is located in the South Urals (Stratotype..., 1983; Bekker, 1988). These para-autochthonous units offer an opportunity to obtain new paleomagnetic data for the Ediacaran of Baltica. Because the units are considered to be para-autochthonous, the rocks must

yield high-quality paleomagnetic results (including the age of the rocks) that can be evaluated for post-remanent rotations and translations with respect to the craton.

The western parts of the Uralian fold belt are similar to those of cratonic Baltica. In particular, a more than ten kilometers thick succession of Meso- and Neoproterozoic mostly sedimentary rocks is exposed in the Bashkir anticlinorium (Uplift) in the western South Urals (Fig. 1b). This succession is reliably correlated with sections in the Ural Foredeep and the eastern parts of Baltica *sensu stricto*. Correlation is further supported by a series of matched seismic profiles (Stratotype..., 1983; Puchkov et al., 2001). Although eastward-dipping seismic boundaries under the Bashkir Uplift are interpreted as the consequence of westward thrusting of the Uralian units over the craton in Permian time (Puchkov et al., 2001), the inferred transport along the thrusts is not considered to be significant. In light of this, these units are considered to be part of the Baltica deformed margin since the early Neoproterozoic (~ 1 Ga; Puchkov, 2003).

Exposed within the Bashkir Uplift is a nearly ten kilometer thick succession of Meso- and Neoproterozoic age clastic sediments and carbonates with subordinate Mesoproterozoic volcanics (Stratotype..., 1983). The uppermost member of this succession, the Asha Series, comprises up to 2 km thick terrigenous clastics, presumably molasse (Bekker, 1988). It is divided into five members, the Bakeev, Uryuk, Basu, Kuk-Karauk and Zigan Formations in ascending order (Fig. 1c). The lower two units are predominately arkose, and the upper three formations are polymictic. Sandstone and siltstone prevail through the Series, with subordinate mudstone and rare gritstone.

There are no angular unconformities between the base of the Asha Series and mid-Permian rocks to the west of the Zilmerdak Fault (thick red line in Fig. 1b). Therefore, the only folding in the westernmost Urals took place in mid-Permian time (Kungurian, 272-279 Ma; Puchkov, 2003). Although Ediacaran rocks outcrop over a rather large area (Fig. 1b), the younger horizons of the Asha series with eastern dips are rare, being truncated by the regional Zilmerdak Fault. Consequently, most exposures of the Asha rocks have similar shallow dips to the west.

Until recently, the ages of the Asha Series and its members were not well constrained. The Asha Series para-conformably (and locally unconformably) overlies older Neoproterozoic rocks and is in turn para-conformably overlain by Middle Ordovician sediments in the south and Emsian (Early Devonian) sandstone elsewhere (Ivanushkin et al., 2009). A study of detrital zircons from the Kuk-Karauk Conglomerate and Basu Fm. (Kuznetsov et al., 2012a; 2012b; 2013) reveal a wide age range from >600 Ma to >2500 Ma and hence do not impose tight constraints on the age of the Asha Series. So far, the only isotopic age directly from the Asha Group (U–Pb method on zircons from a tuff bed) is that of 548.2 ± 3.5 Ma on the Zigan Fm. (Grazhdankin et al., 2011; Levashova et al., 2013). Ediacaran fossils were reported from several formations of the Asha Series, including the lowermost Bakeev Fm. (*Nemania Bakeevi* sp. nov.; *Beltanella zilimica* sp. nov.; *Intrites punctatus* Fed.; *Incertae sedis Bunyerichnus* sp.; *Beltanelloides* (?) sp.) and the Basu Fm. (*Pseudorhizostomites howchini* Sprigg; *Protodipleurosoma paulus* Beck.; *Paliella patelliformis* Fed.; *Medusinites* sp.) (Bekker, 1992). Also found were some trace fossils: *Palaeopascichnus delicatus* Palij; *Neonereites uniserialis* Seilacher, *Catellichnus oktonarius* Becker (Bekker, 1992; 1996). Grazhdankin et al. (2011) and Ronkin et al. (2006) suggest a correlation between the Chernykamen Formation (Sylvitsa Group, Central Urals) and the Zigan Formation (South Urals) on the basis of Ediacaran fossils, sequence boundaries and geochronology. These correlations (and known U–Pb ages) would narrowly constrain the age of the Basu Formation to between 570–560 Ma (Ronkin et al., 2006; Grazhdankin et al., 2011; Maslov et al. 2013).

Several attempts to acquire paleomagnetic data on the terrigenous rocks of the Asha Series were undertaken and are summarized in Table 1 (the data based on incomplete demagnetization and not subjected to principal component analysis are excluded). The oldest Bakeev Fm. of presumed Late Cryogenian (Early Ediacaran) age is exposed in several locations (Stratotype..., 1983) although no useful results were obtained from a pilot collection of samples collected by our research team.

The Uryuk Fm. is mostly composed of arkose gritstone and coarse-grained sandstones. The upper part of this formation comprises intercalating whitish and brick red medium-grained arkoses. Several sites yielded a component, which is directionally close to the widespread Late Paleozoic overprint in the Urals and was treated as such (Uryuk-red in Table 1).

A 500 to 900 meter thick greenish-gray and brown-gray sandstone with subordinate siltstone of the Basu Fm. were the subjects of several studies (Komissarova, 1963; 1970; Golovanova et al., 2011; Iosifidi et al., 2012). A dual-polarity remanence with northeastern declination and negative inclination was reliably identified by all researchers (Basu-1 in Table 1) and was considered a mid-Paleozoic prefolding overprint by Iosifidi et al. (2012). These authors have also isolated another bi-polar component (Basu-2 in Table 1) in five samples out of 30 studied and claimed it to be primary.

The most complicated dataset comes from the Kuk-Karauk Fm. that comprises mostly reddish-brown coarse-grained sandstones, often with millimeter-sized floating pebbles, and a regionally traceable conglomerate unit (Kuk-Karauk Conglomerate). This unit is up to 50 meters thick in the south of the Bashkir Uplift, thins out to a few meters in its central part and wedges out altogether further to the north. Most pebbles are made of white quartz; with subordinate granites, black volcanic rocks, red cherts, and sometimes reddish sedimentary rocks. Pisarevsky et al. (1999) obtained scattered component directions from 33 samples of medium grained sandstones both below and above the conglomerate at two localities but claimed the presence of three dual-polarity components after sorting the data with the aid of cluster analysis (Kuk-Karauk-1 to Kuk-Karauk-3, Table 1). Iosifidi et al. (2012) studied 70 samples from three localities and recognized two dual-polarity components, both of which agree with their results on the Basu Fm. We studied 34 samples from five sites at two localities and found a dispersed cloud ($k \sim 2$) of high-temperature components (HTC). All in all, neither the presence of any consistent component appears to be established in this formation. Finally, Golovanova et al. (2011) sampled thirty cobbles of magmatic rocks and sediments from the Kuk-Karauk Conglomerate,

most of which yielded a well defined and rather tightly grouped single component aligned with the present-day field.

The youngest member of the Asha Series, the Zigan Fm., yielded a well-defined dual-polarity remanence in maroon sandstones at three localities (Levashova et al., 2013).

Here we present new paleomagnetic results from the Basu Fm and examine the implications of these results for Ediacaran paleogeography.

SAMPLING

The 500 to 900 meter thick Basu Fm. comprises about half of the Asha Series (Fig. 1c). Within a ~30-40 meter thick transitional zone, red arkoses of the underlying Uryuk Fm. are gradually replaced with darker colored polymictic sandstones of the Basu Fm. Its upper boundary is imprecisely defined by gradual transition to predominantly brown red and generally coarser sandstones of the Kuk-Karauk Fm (Fig. 1c). The sections of the Basu Fm. vary laterally, but have no regionally recognizable marker horizons; this and limited patchy exposures along river beds and roads lead to a situation that even sections a few kilometers apart cannot be unambiguously matched.

The Basu Fm. mainly consists of fine- to medium-grained polymictic sandstones, with subordinate siltstones and millimeter-thick argillite films on bedding surfaces of coarser-grained varieties. Most sandstones and siltstones are grey to greenish-grey, sometimes maroon, whereas argillites are usually chocolate-colored, thus often making the outcrops look like red beds. The rocks are non-metamorphosed; even calcite veins are rare. Cross-bedding, ripple marks, sometimes slumps, testify to accumulation of the Basu rocks at shallow water depths. The polymictic part of the Asha Series, including the Basu Fm., is regarded as molasse by Bekker (1988). The scarcity of coarse-grained rocks points to accumulation in distal parts of a molasse basin.

We sampled the Basu Fm. at several localities that are spread over more than one hundred kilometers from north to south (red stars in Fig. 1b). Locality GB is at the northern periclinal

section of the Avdyrdak Anticline where it is cut by the Inzer River. This locality provided us with six sites from adjoining sections along with a slump fold that was several decimeters in length. Localities BSW and BSE are sampled from road cuts in the Basu River valley where it crosses the western and eastern limbs, respectively, of the same fold. Twenty-three sites that are spread over a considerable part of the formation thickness were studied. Three sites were collected along the eastern bank of the Takata River (Locality TK). At locality KK (in the Kuk-Karauk valley a total of 17 sites and two slump folds were sampled (see more detail below). The remaining sites yielded numerous samples, but no interpretable data (white stars in Fig 1b).

PALEOMAGNETISM

Methods

Paleomagnetic samples were collected as hand-sized blocks oriented with a magnetic compass, one sample being taken at a stratigraphic level. Four to twelve or even more samples collected from a 3 to 10, sometimes up to 30 meters thick section parts, depending on the exposure quality and the availability of suitable rock varieties, is considered a site. Cubic specimens of 8-cm³ volume were cut from each hand sample. The collection was studied in the Paleomagnetic laboratories of Geological Institute of the Russian Academy of Sciences in Moscow and at the Institute of Geology (Ufa Scientific Center, Russian Academy of Sciences) in the city of Ufa. Individual specimens were stepwise heated in 12 to 20 increments up to 700°C in either homemade ovens (Moscow) or utilizing an Analytical Services TD-48 thermal demagnetizer with internal residual fields of < 10 nT (Ufa); measurements of the natural remanent magnetization, NRM, were done with a JR-4 or JR-6 spinner magnetometer with a noise level of 0.05 mA/m. Demagnetization results were plotted in orthogonal vector diagrams (Zijderveld, 1967). Visually identified linear trajectories were used to determine directions of magnetic components by Principal Component Analysis (PCA), employing a least-squares fit comprising three or more demagnetization steps (Kirschvink, 1980), anchoring the fitting lines to the origin where

appropriate. Paleomagnetic software written by Jean-Pascal Cogné (2003) was used in the analysis.

Results from the main collection

The NRM values in the Basu rocks range from less than 1 mA/m to nearly 100 mA/m. In contrast, susceptibility values are uniformly low, below 0.5 mSI units in most samples (Komisarova, 1970). Stepwise thermal demagnetization revealed that the NRM could be accounted for by up to three components in varying proportion. A low-temperature component (LTC) was removed between 200-300 °C and may dominate in samples with low NRM values. The LTC directions are scattered at some sites or gravitate toward the present-day field direction. Thus, this component is likely a recent viscous overprint and is excluded from further analysis.

After removal of the LTC, an intermediate-temperature component, ITC, is present (Fig. 2a-b) in many cases. The ITC directions are scattered, with the exception of site BSW10 (Fig. 2b-c), where the ITC is adequately clustered with a mean direction in situ ($D^\circ = 243.8$, $I^\circ = -39.3$, $\alpha_{95}^\circ = 13.2^\circ$, $n = 8$ samples). This direction is similar to the regional mean of the Permian overprint ($D^\circ = 230.0$, $I^\circ = -40.7$, $\alpha_{95}^\circ = 8.7$; Levashova et al., 2013).

In approximately half of the collection, no ITC is observed and a single well-defined high temperature (HTC) shows a rectilinear decay to the origin (Figs. 2d-f). In samples with an ITC component, the HTC component can also be reliably isolated (Fig. 2a-b). It seems that the same HTC is present in the samples of Basu rocks, irrespective of the differences in demagnetization characteristics. It is worth noting that there are no samples with the HTC bypassing the origin and revealing a remanence of opposite polarity. The HTC is mostly unblocked between 580° and 700° (Fig. 2) and resides in hematite. Optical studies indicate that hematite grains are widespread in the Basu rocks (Kozlov, 1982). Rock magnetic studies of the Basu rocks confirm that hematite, sometimes with traces of magnetite, is the main remanence carrier (Komissarova, 1963; 1970; Iosifidi et al., 2012). This conclusion is valid for all rock varieties, and the samples

of very similar color and grain size may have the single HTC, both HTC and ITC, or no stable component at all. Dark maroon and purely gray samples may yield similar HTC directions.

HTC directions are acceptably grouped at 34 sites of 49 studied (Fig. 3a-b; Table 2). The fold test (McElhinny, 1964) is positive ($k_{TC}/k_{IS} = 5.15$; 95% critical value is 1.51), maximum data grouping is at 100% unfolding. Some site-means, however, are rather poorly defined, and the same test was repeated for better defined 24 sites with $\alpha_{95} < 15^\circ$. Using the more restrictive dataset, the fold test was also positive ($k_{TC}/k_{IS} = 6.45$; 95% critical value is 1.63) (Fig 3d). The positive fold test only constrains the age of magnetization to older than the mid-Permian (Puchkov, 2003). We note that the mean from sites with $\alpha_{95} > 15^\circ$ and the mean from sites with $\alpha_{95} < 15^\circ$ are statistically similar (called *imprecise* and *precise*, respectively; Table 2). We conclude that the lower precision data stem from poor averaging of internal scatter rather than from contamination by another remanence. Note also that the HTC directions are statistically similar over more than 100 km (Fig. 1b, Table 2) that negates local rotations between sampling sites.

The HTC reveals a dual-polarity magnetization (Figs. 2 and 3). At four sites, where relatively thick intervals were sampled, both polarities are found in several adjacent samples; such one-polarity intervals are treated as separate sites (Table 2). The HTC directions form two distinct polarity clouds at both site-mean and sample levels after tilt correction. The reversal test is “double” positive: the scatter of normal and reverse sites are statistically identical ($k_N=50.3$, $k_R=47.8$), and the polarity means are antipodal: the observed angle between them (inverted to one polarity) is much less than the critical one ($\gamma_{obs} = 3.8^\circ$, $\gamma_{crit} = 8.8^\circ$; class B; McFadden and McElhinny, 1990). The test is also positive at the sample level ($k_N=15.9$, $k_R=14.9$; $\gamma_{obs} = 2.8^\circ$, $\gamma_{crit} = 5.4^\circ$). Note that there is no trace of the Basu 2 component (Table 1) that was claimed to be of primary origin by Iosifidi et al. (2012), among our nearly two hundred HTC sample directions (Fig. 3c).

Results on slump folds and the origin of the high-temperature component

We sampled three slumps at different localities. In the lower part of the KK section, slump 1 is exposed in a ~ 2.5 m high wall of an abandoned small quarry that was used for road construction. In some places (gray spots in Fig. 4a), the rocks are strongly fractured, most probably during the quarry being in use. Slump 2 is found in the natural outcrop in the upper part of the KK section (Fig. 5a). Some fifteen samples were taken from each slump, mostly from fine-grained platy sandstone, and a standard site of 7-8 samples was taken from non-deformed beds close to each slump. Bedding was separately measured for each sample from the slumps; this was facilitated by platy character of sandstones that show the strongest degree of slumping. Finally, both slump 3 and host rocks at locality GB did not yield stable components.

Demagnetization characteristics are very similar for deformed and non-deformed rocks (Fig. 4b-d, 5b-d) and also similar to those for the Basu rocks in general (Fig. 2). The grouping of HTC data at two non-deformed sites is reasonably good, and the corresponding site-means agree with the other HTC directions after tilt correction (Tables 2 and 3). Thus we conclude that the remanences in non-deformed rocks near the slumps and in the main body of the Basu Fm. are of the same origin.

In slump 1, from the lower part of the KK section, the HTC directions are significantly better grouped after tilt correction, and the best data grouping is achieved after 100% unfolding although the overall scatter remains rather high (Fig. 4e-g). In slump 2, the best data grouping is achieved at 55% of unfolding (Fig. 5e-h). Moreover, the slump mean is displaced both from the mean for the adjacent site (Fig. 5e-f) and main dataset (Fig. 3b).

The fold test in paleomagnetism is based on an assumption that each fold limb rotates as a rigid body about a horizontal axis. When slumping in a pliable bed occurs very soon after deposition, this assumption is invalid. Hence the perfect alignment of paleomagnetic vectors is unlikely even if the remanence predates slumping. Moreover, the slumped block may be displaced as the whole, which is likely what happened with slump 2. Thus, the HTC is older than

slumping in the first case, while it appears to be coeval with deformation in the second. These slump-test results provide strong evidence for a primary remanence in the Basu Fm.

In sum, the positive fold and reversal tests and two slump tests indicate that the remanence was acquired very close to the age of deposition. In spite of this evidence for a primary remanence, it should be noted that the Basu pole falls close to the Late Ordovician - Silurian segment of the apparent polar wander path (APWP) for Baltica. This similarity was noted decades ago (Komissarova, 1970) and confirmed by recent studies (Golovanova et al., 2011). In the latter paper, the remagnetization hypothesis was analyzed, but could not be rejected as a possible explanation for the similarity in direction. Given the positive slump tests in our study, we conclude that the agreement between the APWP for Baltica and the Basu pole is fortuitous, and the dual-polarity HTC in the Basu rocks is primary.

INCLINATION SHALLOWING

Inclination shallowing was discovered many decades ago (King, 1955) but was thought to be relatively uncommon. In the 1990's evidence began to mount that inclination shallowing is more widespread in fine-grained redbeds and sparked a renewed interest in analytical and experimental techniques aimed at identifying the problem. Detection of inclination shallowing remains cumbersome and labor-intensive, while correcting for inclination shallowing was, and still is, a tricky business (Sun and Kodama, 1992; Tan and Kodama, 2002). In an effort to simplify both detection and correction of inclination shallowing, the elongation-inclination test was regarded as a major break-through and is widely utilized (Tauxe and Kent, 2004; Tauxe, 2005). There are, however, several points that may affect the validity of this approach:

1. Tauxe and Kent (2004) used a model of the geomagnetic field based on observations over the past 5 Ma. While the 0-5 Ma field data are fairly robust, the model has not been validated for earlier periods in Earth history and even during that interval, the fit of model predictions to observations often shows large discrepancies (Johnson et al., 2008).

2. The elongation of a paleomagnetic distribution is a statistical parameter that becomes acceptably “steady” for large datasets only. Tauxe and Kent (2004) recommend a minimum of 100 site means are required for proper analysis. This level of sampling density is seldom achieved with the exception of a series of thick lava flows.

3. The scatter of observed directions, especially in sediments, is due not to secular variation alone but to its sum with paleomagnetic noise from other sources. Hence, the distributions of paleomagnetic data may be more circular than predicted by the TK-model.

4. The correction for inclination shallowing is performed with the bootstrap that gives biased, often strongly biased (see Figure 5 in Tauxe et al., 1991), estimates for all parameters that are not normally distributed. In particular, elongation parameter has a clearly non-Gaussian distribution, and thus its estimates can be strongly biased.

Thus while the E-I method may prove useful in some younger studies, the limitations noted above preclude its blanket use as a corrective measure. An alternative may be the use of qualitative tests to determine if shallowing is present. In a purely geometric sense, inclination shallowing will affect the distribution of paleomagnetic data by transforming a circular or near circular distributions into a banana-like distribution with elongation in the east-west direction (Bazhenov, 1981; Tauxe, 2005). Hence the presence of strong inclination shallowing can be established by observation if the banana-like elongation is present and statistically significant.

In our case, the number of acceptable Basu sites (Table 2) is too limited for the E-I method. We compiled sample directions from the sites, where consistent HTC data are recovered from all or most samples (Fig. 3c, Table 4). As the samples are stratigraphically ordered at each site, each unit vector represents a spot reading of the field, albeit an imprecise one. The pre-shallowing form of the distribution can be reconstructed by stepwise varying the shallowing parameter f in the formula: $\tan I_m = f * \tan I_o$, where I is inclination ($^\circ$), subscripts m and o stand for measured and original values, respectively, f may vary from 1 to 0. After each step, we calculated an elongation parameter E value after Tauxe (2005) and a best-fitting great circle that

denotes the long axis of distribution. The latter must pass close to the projection pole (deviation of the great circle from the pole, ΔGC , is close to zero) if a distribution is elongated in vertical plane as predicted by the model of Tauxe and Kent (2004). In other words, the declination of the point where this great circle crosses the stereonet equator must coincide with the mean declination for the dataset in this model. In contrast, this great circle must deviate from the projection pole with $\Delta GC^\circ \sim (90 - I_m)$ for shallowed data. Note also that, to be statistically significant, the observed elongation must exceed the 95% critical value for this parameter; for 195 vectors used in analysis, the latter value is ~ 1.30 .

For the original dataset, i.e., for $f = 1.0$ (Fig. 6a, Table 4), the unit vectors form a well-defined cloud of nearly circular outline, E is statistically insignificant and much less than the value predicted by the TK-model. ΔGC of 40.5° is large but significantly less than the predicted value of $\sim 54^\circ$ for shallowed data. Hence this distribution is axially symmetric with statistically insignificant and spuriously oriented elongation that, by itself, is evidence against strong shallowing. The features predicted by the TK model are absent as well. For $f = 0.9$, the elongation slightly increases but remains insignificant; in contrast, ΔGC decreases sharply to $\sim 10^\circ$ and becomes nearly vertical. The deviation from the circularity is barely discernible (Fig. 6b, Table 4). After the next step ($f = 0.8$; Fig. 6c), the elongation becomes both visible and statistically significant, while ΔGC becomes negligible; elongation becomes similar to the TK model predictions. The situation becomes even clearer for $f = 0.7$ (Fig. 6d, Table 4), with E exceeding the prediction of the TK model. Hence we found no indication of strong inclination shallowing in the Basu data although it remains possible that something on the order of 10% or less shallowing may have occurred.

COHERENCE OF THE SOUTH URALS AND CRATONIC BALTICA

Can we extrapolate the paleomagnetic data from the western Urals to Baltica? The answer to this question requires evidence that the sampled region was not a far-traveled terrane that had docked to Baltica during the Paleozoic. We noted earlier that the Neoproterozoic sequences

(~900-542 Ma) are reliably traced from the fold belt to the Ural Foredeep and further to the eastern parts of Baltica; this correlation is supported by a series of seismic profiles (Stratotype..., 1983; Puchkov et al., 2001; Maslov, 2004; Kuznetsov et al., 2010). Despite the gaps in geologic record, no angular unconformities are found between Ediacaran through Paleozoic sedimentary rocks until the mid-Permian in the westernmost units of the South Urals (Puchkov, 2003). Thus, there are no traces of any collisional events from the late Neoproterozoic until the Permian. Taking these two lines of evidence together we are confident that the western tectonic units of the Urals are a part of the cratonic Baltica margin, deformed in Permian time (Puchkov, 2003; Maslov, 2004; Bogdanova et al., 2008; Levashova et al., 2013).

While the study area can be reliably tied to cratonic Baltica, it does not preclude rotation of these units with respect to the craton during the Permian orogenesis. In an effort to evaluate this possibility, Levashova et al. (2013) studied Paleozoic and Neoproterozoic rocks at several localities and have isolated a synfolding component with SW and up directions. The mean direction of this overprint was found to agree within few degrees with the reference directions recalculated from the apparent polar wander path of Baltica for the interval from 300 to 270 Ma. As major folding in the studied westernmost part of the South Urals took place in Kungurian time (Puchkov, 2003), the observed agreement speaks against relative rotation between this part of the Urals and cratonic Baltica (Levashova et al., 2013). Thus, both geologic and paleomagnetic data do not reveal significant relative motions between the study area and craton, and paleomagnetic data from these western Uralian units can be used to constrain the paleogeography of Baltica during the Ediacaran interval.

INTERPRETATION AND DISCUSSION

The aforementioned paleomagnetic and geological data allow us to conclude that: 1) the rocks of the Basu Formation were deposited between 570 and 560 Ma; 2) the high temperature component of magnetization in these rocks is of primary origin; 3) inclination shallowing is

negligible; 4) the Basu Formation pole can be extrapolated to Baltica. Thus, we can include this result in the list of Ediacaran paleomagnetic poles for Baltica (Table 5).

Cambrian - Late Precambrian –APWP’s for Laurentia and Baltica are a matter of considerable debate because the existing set of paleomagnetic data for this time interval is highly controversial (Meert, 2014). Thus, before speculating on Laurentia and Baltica relationships and the possible timing of the Iapetus opening, one must review the Late Precambrian – Cambrian paleomagnetic data for these two continents.

Early Ordovician – Ediacaran paleomagnetic data from Baltica

The well-defined Early Ordovician (480 Ma) mean pole (Torsvik et al., 2012) forms a useful ‘anchor-point’ for evaluating plate motion implied by the Ediacaran-Cambrian paleomagnetic data (EO in Table 5). The poles for Baltica for the interval from 615 to >480 Ma are listed in Table 5 and compared with the Baltic APWP in Figure 7.

A pole on Upper Cambrian (ca. 500 Ma) Andarum shale (AS, Table 5; Torsvik and Rehnstrom, 2001) is based on eleven samples only. The Narva sediments pole (NS; Table 5) is based on eleven samples of Lower Cambrian fine-grained sediments but is confirmed by the reversal test (Khramov and Iosifidi, 2009). The Early Cambrian (about 535 Ma) Tornetrask/Dividal results are derived from sediments along the Caledonian front (T; Table 5; Torsvik and Rehnstrom, 2001; Rehnstrom and Torsvik, 2003). The Nekso sandstone pole (NE; Table 5; Lewandowski and Abrahamsen, 2003) is derived from a horst block within the Caledonian deformational region.

Upper Ediacaran sediments outcrop along the southeastern coast of the White Sea; their stratigraphic age is confirmed by the ~ 555 Ma U-Pb age on zircons from ash beds (Martin et al., 2000; Grazhdankin, 2003). Four consistent poles (Z, VH, AR, WC; Table 5) from a narrow stratigraphic interval (~550-555 Ma) are available from this area (Popov et al., 2002; 2005; Iglesia Llanos et al., 2005).

In Volhynia (southwestern Baltica), the Ediacaran section comprises a several hundred meters thick succession of lava flows and tuffs (known mostly from boreholes) that is overlain by a 200 meter thick sedimentary cover (Velikanov, 1985). Few U-Pb ages on zircons from the volcanic rocks yield a late Ediacaran-Fortunian age of ca. 540-550 Ma (Compston et al., 1995), although older ages of ~ 580 Ma were inferred from $^{40}\text{Ar}/^{39}\text{Ar}$ data (Elming et al., 2007). Three teams independently studied paleomagnetism of these volcanics (Iosifidi et al., 2001; Nawrocki et al., 2004; Elming et al., 2007). Sampling was limited to two small quarries with no more than four cooling units in the northern part of Volhynia, and one or two cooling units only were sampled in two small quarries in the south. Elming et al. (2007) note that the “northern” sites (VN; Table 5) indicate a high-paleolatitude position for Baltica, whereas the “southern” sites (VS; Table 5) place central Baltica at a latitude of ~20°, N or S. Iosifidi et al. (2005) studied overlying Ediacaran sediments in the same area and recognized six different remanent directions, some of them bi-polar, in these rocks. Our re-evaluation of these data shows that the distributions of component directions strongly overlap and just two very diffuse clouds can be recognized (not illustrated). Therefore, there is a reason to question the validity of each mean direction reported by Iosifidi et al. (2005); we note, however, that none of their directions infers a high-latitude position for Baltica during the Late Ediacaran-Early Cambrian.

The Zigan pole (550-542 Ma; Z in Table 5; Levashova et al., 2013) was derived from the Zigan Fm. redbeds from the westernmost tectonic units of the Ural fold belt located at deformed eastern margin of Baltica. This remanence is likely primary as indicated by a positive reversal test and regional consistency test. Similar to the Basu data, this result can be extrapolated to cratonic Baltica. The Zigan pole falls into the cluster formed by the White Sea poles and South Volhynia pole (Fig. 7). Good grouping of the data from three remote parts of the craton makes the overall Late Ediacaran mean pole (VWZ, Table 5) a useful “anchor-point” for constructing the pre-Ordovician part of the Baltic APWP.

The Basu pole (575-560 Ma, BA in Table 5) is described in this study. Although the Asha Series is usually described as a conformable pile of five formations this is due primarily to a lack of precise ages and/or diagnostic fossils (Stratotype..., 1983). Paleomagnetic data are available only from the Basu and Zigan Fms., which are separated by the relatively thin Kuk-Karauk Fm with reportedly gradual upper and lower boundaries. The large difference between the corresponding poles (Table 5) must also be explained in terms of the available age constraints for the two units.

Paleomagnetic data are available on the Fen Complex in southern Norway (Poorter, 1972; Piper, 1988; Meert et al., 1998) and Alnö Complex in central Sweden (Piper, 1981; Meert et al., 2007). These carbonatite complexes are nearly coeval (~ 585 Ma), and their corresponding paleomagnetic data can place the study locations in Baltica at intermediate latitudes (Fen complex, F; Table 5), close to the geographic pole (Alnö steep component, A1; Table 5) or in the tropics (Alnö shallow component, A2; Table 5). Meert (2014) showed that the Fen paleomagnetic pole is most likely a remagnetization acquired during the opening of the Oslo Rift (Permo-Triassic) and that the Alno data are of too poor quality to merit further discussion.

The sole Early Ediacaran pole comes from the Egersund dykes in SW Norway (E; Table 5; Walderhaug et al., 2007). This pole is reliably dated at 616 ± 3 Ma (zircon U-Pb age) and is confirmed by a positive baked contact test.

Comparison of the Ediacaran-Cambrian and reference data for Baltica shows that the Andarum, Tornetrask, Alno1 and Fen poles fall directly on the Early Mesozoic-Permian segment of the APWP, while the Nekso pole falls within 10° of this overprint direction (Fig. 7). We feel that this similarity to a younger pole along with the lack of evidence for a primary remanence in these rocks, thus strongly undermine the validity of these data. The Egersund dykes pole also falls near the Early Ordovician mean pole for Baltica (within 11°) although the age of its remanence is supported by a baked contact test.

The Narva pole was considered by Khramov and Iosifidi (2009) as representative of a remagnetization at the Cambrian –Ordovician boundary, although it differs by more than 30° from the nearest reference pole. The Narva result is based on limited statistics and thus should not be used as a tie-point in any tectonic analysis. The same arguments pertain to the Volhynia North pole. The WZV appears to be the most trustworthy in the data list analyzed, while the primary origin and reliability of the Basu result have been already argued. Thus, for the Ediacaran - Early Ordovician interval, the paleoposition of Baltica can be ascertained from the four poles only, which are the 480 Ma reference, VWZ, Basu, and the Egersund poles. The resulting very schematic APWP (Fig. 7) infers the following kinematic scenario:

- 1) Egersund –Basu (616 - 565 Ma) interval. Baltica moves from high to low latitudes by 45° at velocity of $\sim 0.9^\circ/\text{Ma}$ (9 cm/year) and rotates by ca. 30° CW.
- 2) Basu- VWZ (565 - 550 Ma) interval. Baltica moves to a still lower latitude at velocity of $\sim 0.8^\circ/\text{Ma}$ (8 cm/year) and quickly rotates by ca. 50° CW
- 3) VWZ -Early Ordovician (550 – 480 Ma) interval. Baltica reverses its motion and drifts poleward at velocity of $\sim 0.7^\circ/\text{Ma}$ (7 cm/year). The sense of rotation is reversed and the craton rotates by ca. 100° CCW.

All inferred motions with poleward velocities between 7 and 9 cm/year are rather fast but similar and even larger velocities are not uncommon.

Early Cambrian - Ediacaran paleomagnetic data from Laurentia

The Early Ordovician and Middle to Late Cambrian parts of the Laurentia's APWP are based on a sufficient number of mutually consistent paleopoles, and we rely on the existing APWP (Torsvik et al., 2012). As for the Early Cambrian to Ediacaran poles, we mostly follow the choice of McCausland et al. (2007).

Two poles are listed in Table 6 for Early Cambrian time. McCausland et al. (2007) claim that the result on the 532 Ma Mont Rigaud stocks satisfies all quality criteria (Van der Voo,

1990), except for the clearly positive field test. Three poles are tabulated for Late Ediacaran time (Table 6), although none of them provides a well-resolved pole position for Laurentia. The 550±3 Ma Skinner Cove Fm. result (SC; McCausland and Hodych, 1998) is from western Newfoundland, which is allocthonous to Laurentia. This area probably never was far away from the continent margin (Cawood et al., 2001; Hodych et al., 2004) but the lack of relative rotations and, hence, full coherency with Laurentia was not demonstrated. Dual-polarity shallow directions are reported from the carbonate Johnnie Fm. (JR, Van Alstine and Gillet, 1979), but the rock age is poorly constrained. Basing on the assumed age of the overlying rocks, Meert et al. (1994) estimated these carbonates as ca. 555 Ma although more recent estimates constrain the age of the Johnnie to ‘younger than 580 Ma’ (Pettersen et al., 2011; Trower and Grotzinger, 2010; Kaufman et al., 2007). Finally, the 565 Ma Sept Iles intrusion has a shallow ‘A’ magnetization (SI-A) in the intrusion only and a steep dual-polarity ‘B’ magnetization (SI-B) present in four mafic dykes and several sites in the intrusion (Tanczyk et al., 1987). These researchers interpreted the ‘A’ remanence as primary because it has higher coercivity and unblocking temperature; this view is supported by a positive inverse contact test. Collectively, SC, JR and SI-A poles suggest that Laurentia was located at low latitudes in the Late Ediacaran.

For the Early Ediacaran epoch, the 572 ± 5 Ma Catoctin basalts carry three different components of magnetization (Meert et al., 1994). A steep Catoctin ‘A’ remanence (CT-A) passes the baked contact test and McCausland et al. (2007) suppose it to be of primary origin. However, the pre-folding dual-polarity shallow ‘B’ remanence (CT-B) is also found in Catoctin basalts and is interpreted as primary in other studies (Hodych et al., 2004; Murphy et al., 2004). The Catoctin ‘C’ component (CT-C) was isolated at intermediate temperatures, and Meert et al. (1994) argued that it was the result of Middle Paleozoic overprinting. The 577±1 Ma Callander complex carry a steeply directed dual-polarity remanence (CC) that passes the baked contact test (Symons et al., 1991; Kamo et al., 1995). The 590 Ma Grenville Dykes steep ‘B’ remanence (GD-B) was interpreted as primary (Murthy, 1971; Buchan et al., 2004), while the shallow ‘A’

directions in other Grenville Dykes (Murthy, 1971) do not have supporting field test and are supposed to be of secondary origin.

The first paleomagnetic study from the 615 Ma Long Range dykes (Murthy et al., 1992) was later recalculated by Hodych et al. (2004). This recalculated result (LR) for five out of the six dykes encompasses two polarities, and, despite poor resolution, suggests a low latitude location of Laurentia.

In full accord with earlier papers (e.g., McCausland et al., 2007), our analysis points to existence of two groups of poles: one of them suggests a low-latitude position for Laurentia, while the other group places it in high latitudes (Table 6; Fig. 8a). Four poles inferring a high-latitude position for Laurentia range in age from 590 Ma to 564 Ma and do not overlap its Phanerozoic APWP for any polarity option. In contrast, all low-latitude poles are in good to excellent agreement with the Early-Middle Paleozoic segment of this APWP (Fig. 8a). Moreover, the Laurentian low-latitude poles for the 615 – 530 Ma window do not outline any clear progression with age; instead, they reveal an erratic pattern with nearly identical start- and end-points (Fig. 8b). One way to explain the above features is to hypothesize several remagnetization events at different parts of east Laurentia, but testing this hypothesis against geological data is out of scope of this paper. An argument against this assumption is the positioning of the Johnnie pole from western Laurentia where no tectonic events are known in the Middle Paleozoic.

The subject of Laurentia's paleolatitude in the Ediacaran remains enigmatic. Until the early 1990's, Laurentia was placed at low latitudes until several studies pointed to a possible high-latitude position for Laurentia (Symons and Chiasson, 1991; Park, 1992, 1994; Meert et al., 1994). A high-latitude position was then generally accepted in the literature until questioned by Pisarevsky et al. (2000; see also comments by Meert et al., 2001 and reply by Pisarevsky et al., 2001). While not conclusive, the low-latitude position for Laurentia has become 'favored' in more recent literature and an explanation for the high latitude results remains elusive.

Tectonic implications

Paleogeographic and tectonic reconstructions require reasonably precise constraints on the timing of continental rifting and initiation of ocean spreading, timing of ocean closure and continental collision, and positions of the continents through time. To reconstruct the Laurentia and Baltica relationships and the process of the Iapetus Ocean opening and evolution in the Late Precambrian – Early Paleozoic one needs both paleomagnetic data from the two continents and geological evidence on the process of the Iapetus opening.

The Humber zone of the Appalachian orogen represents the eastern margin of Laurentia and preserves a record of continental margin initiation associated with opening of the Iapetus Ocean (Williams, 1995; Cawood et al., 1996; 2001). This zone consists of crystalline basement of the Grenville province unconformably covered by a sequence containing rift, continental margin, and foreland basin units. Rift sequences contain mafic to felsic volcanic flows and intrusions, while drift sequences are carbonate-dominated. Two pulses of rifting, from ca. 760 to 700 Ma and from ca. 620 to 550 Ma, are recognized along the eastern margin of Laurentia (Aleinikoff et al., 1995; Mitchell et al., 2011; O'Brien and Van der Pluijm, 2012). The first pulse is correlated to the opening of the proto-Pacific between the west Laurentia and East Australia-West Antarctica conjugate margins (Powell et al., 1993; Powell, 1995; Wingate and Giddings, 2000). Younger rift-related magmatic activity, widespread in the central to northeastern Appalachians (Aleinikoff et al., 1995), is associated with the opening of the Iapetus Ocean, first between Laurentia and Baltica, and then between Laurentia and elements of west Gondwana.

Geological data indicate that igneous activity at the Laurentian margin continued in a rift setting until ca. 550 Ma (cf. Kumarapeli et al., 1989; Williams, 1995; Cawood et al., 1996; Higgins and van Breemen, 1998). However, McCausland and Hodych (1998) suggested that post-570 Ma magmatism may reflect either igneous activity along the failed Ottawa-Bonnechere rift arm, or passive margin alkalic magmatism. Thus, the initiation of Iapetus spreading started either at about ca. 570 Ma, or at about ca. 550 Ma. A younger limit on the timing of the initiation

of seafloor spreading is provided by the age of the earliest drift-related sedimentation, which is probably not older than ca. 525 – 520 Ma (Cawood et al., 2001).

The Iapetus traces are more limited in Europe, being confined to some rift-related magmatic complexes in Norway (Dahlgren, 1994; Meert et al., 1998), Sweden (Meert et al., 2007) and the British Isles (van Staal et al., 1998) along with rift-related mafic dykes (Bingen et al., 1998).

The paleogeographic reconstructions for Early Ordovician interval that includes the Tremadocian and early Arenigian (beginning at 490 Ma, ending at 471 Ma) seem to be most reliable due to the large number of faunal and paleomagnetic studies that have been undertaken on the rocks of this age. In the Early Ordovician, both “Baltica” and “West Gondwana” arms of the Iapetus Ocean oceans were at their widest (Torsvik and Trench, 1991; Cocks and Torsvik, 2002). Hence the early Ordovician reconstructions (Fig. 9a) can be used as a starting point for evaluating the earlier paleogeography.

Gondwana was mostly assembled at around 550-540 Ma (Meert and Van der Voo, 1998; Meert, 2003; Meert and Lieberman, 2008), although the fusion of some cratonic elements (Amazonia, West Africa) with central Gondwana may have occurred during Middle to Late Cambrian times (Tohver et al., 2006). At the time of Iapetus Ocean opening, Gondwana did not exist as a single continent, and thus we will deal only with the Baltica – Laurentia relationships in our reconstructions.

The position of Baltica during the Cambrian is poorly known, as there are no unambiguous paleomagnetic data for this time interval. At the end of the Ediacaran (~550 Ma), Baltica’s position is defined by the group of paleomagnetic poles (VWZ, Table 5; Figure 9b). The position of Laurentia at that time is defined only by the 550±3 Ma Skinner Cove Formation (Table 6). According to the limited data, Laurentia is located at low latitudes at the end of the Ediacaran and the “Baltica” arm of the Iapetus Ocean is already rather wide (Fig. 9b).

For the 575-560 Ma interval, the position of Baltica is defined by the Basu pole (BA in Table 5), and there are three paleomagnetic results from Laurentia (Table 6). The 565 Ma Sept Iles intrusion has a shallow 'A' magnetization and a steep dual-polarity 'B' magnetization (Tanczyk et al., 1987). The 572 ± 5 Ma Catoctin basalts carry three different components of magnetization (Meert et al., 1994): a steep Catoctin 'A' remanence, a shallow Catoctin 'B' remanence and a Catoctin 'C' component that was isolated at intermediate temperatures and is supposed to be a Middle Paleozoic overprint. The 577 ± 1 Ma Callander complex carries a steeply directed dual-polarity remanence (Symons and Chiasson, 1991; Kamo et al., 1995).

The Sept Iles 'A' result and the Catoctin 'B' remanence suggest the low latitude paleoposition of Laurentia 575 – 565 Ma ago as does the imprecisely dated Johnnie Formation, while the Sept Iles 'B', Catoctin 'A' and Callander complex remanences imply a high latitude position for Laurentia at that time. If the low-latitude Sept Iles 'A' and Catoctin 'B' poles are used, Baltica and Laurentia can be positioned close to each other and suggest a relatively narrow Iapetus Ocean. If we suppose the Sept Iles 'B', Catoctin 'A' and Callander complex remanences to be of primary origin, Laurentia must have been far to the south of Baltica with a wide Iapetus Ocean (Fig. 9c).

There are no paleomagnetic data for Baltica between 575-560 Ma Basu and the 616 ± 3 Ma Egersund dykes poles (Table 5). The oldest Early Ediacaran result – the 615 Ma Long Range dykes pole suggests a low latitude position for Laurentia, but is based on only five dykes. According to geological data, the Iapetus Ocean had not yet opened at 615 Ma. Paleomagnetic data from the nearly coeval Long Range Dykes and Egersund Dykes poles are consistent with a relatively tight fit between the two continents (Fig. 9d) given the errors in age and position. A strange dichotomy exists between the geological and paleomagnetic data. The 'low-latitude Laurentia' scenario is in agreement with the geological data, but poles from Laurentia and Baltica used to constrain the paleogeography are similar to younger poles and thus potentially represent remagnetizations. In contrast, the "high latitude scenario" looks less suspicious from a

paleomagnetic perspective (i.e. the poles are unique for the last 600 Ma), but require a wide Iapetus Ocean in direct contrast to the available geological data.

CONCLUSIONS

A paleomagnetic study of gray and maroon sandstone and siltstone of the Basu Formation (Ediacaran-age) from the eastern deformed margin of Baltica was carried out at several localities spread over 150 kilometers. The age of the Basu Formation is constrained to the 570-560 Ma interval by correlation, fossils and isotopic ages of overlying and underlying units. A dual-polarity magnetization was isolated in about two-thirds of a large collection. The HTC is considered primary on the basis of positive reversal, fold and slump tests. A schematic APWP for Baltica reveals motion of this craton with velocities of 7 to 9 cm/year for the 480-615 Ma interval. An attempt to reconstruct the early stages of the Iapetus Ocean between Baltica and Laurentia is problematic. As is well known, late Cryogenian and Ediacaran poles for Laurentia form two distinct groups (a high- and a low-latitude ones) that are difficult to reconcile.

Solutions to this conundrum have been proposed (Mitchell et al., 2011; Abrajevitch and Van der Voo, 2010). Mitchell et al. (2011) propose that the swings from low (615 Ma) to high (580 Ma) back to low (550 Ma) latitudes of Laurentia are the result of true polar wander. Abrajevitch and Van der Voo (2010) argue that the Earth's magnetic field exhibited a strange geometry wherein a long-standing equatorial dipole field alternated with a geocentric axial dipole field to produce the apparent high-low latitude transitions. In our opinion, neither option is wholly satisfying. As noted by Meert (2013), the equatorial dipole argument is a global phenomenon that should be reflected in all paleomagnetic data. In the case of Baltica, a review of existing data (repeated above) along with our new result show no swings between high and low latitude results. The true polar wander hypothesis remains possible, but requires additional data from all continents.

Acknowledgements. We thank all people from Institute of Geology in the Ufa City, who helped to carry out the fieldworks and all researchers who took part in sampling. Special thanks are due to Nina Dvorova and Olga Krezhovskikh for demagnetization and measurements of the collections. The constructive reviews by two anonymous reviewers are appreciated. This paper is a product of coordinated efforts of the Russian scientists from Moscow and Ufa and American researchers from the University of Florida in Gainesville. This study was funded by grant EAR11-19038- from the National Science Foundation (J. Meert), grants 11-05-00037 (N. Levashova) and 11-05-00137 (I. Golovanova) from the Russian Foundation of Basic Research, grant 14.Z50.31.0017 of the Government of the Russian Federation, and by Program #10, Earth Science Division, Russian Academy of Sciences.

- 636 Abrajewitch, A., Van der Voo, R., 2010. Incompatible Ediacaran paleomagnetic directions
637 suggest an equatorial geomagnetic dipole hypothesis, *Earth and Planetary Science*
638 *Letters*, 293, 164-170.
- 639 Aleinikoff, J.N., Zartman, R.E., Walter, M., Rankin, D.W., Lyttle, P.T., and Burton, W.C., 1995.
640 U-Pb ages of metarhyolites of the Catoclin and Mount Rogers formations, central and
641 southern Appalachians: Evidence for two phases of Iapetian rifting: *American Journal of*
642 *Science*, 295, 428–454.
- 643 Bazhenov, M.L., 1981. Inclination shallowing in Paleogene sandstones of the South Darvaz
644 Range, *Doklady Akademii Nauk SSSR*, 260, 1336-1339 (in Russian).
- 645 Bekker, Y.R., 1988. Precambrian molasses, Nedra, Leningrad, 289 p (in Russian).
- 646 Bekker, Y.R., 1992. The oldest Ediacaran biota of the Urals, *Izvestia Acad. Sci. SSSR, ser.*
647 *Geol.*, 6, 16 –24 (in Russian).
- 648 Bekker, Y.R., 1996. The discovery of Ediacaran Fauna in the uppermost Vendian of the South
649 Urals, *Regional Geology and Metallogeny*, Leningrad, VSEGEI, No. 5, 111-131 (in
650 Russian).
- 651 Bingen, B., Demaiffe, D. & van Breemen, O., 1998. The 616 Ma old Egersund basaltic dike
652 swarm, SW Norway, and late Neoproterozoic opening of Iapetus ocean. *The Journal of*
653 *Geology*, v. 106, p. 565-574.
- 654 Bogdanova, S.V., Bingen, B., Gorbatshev, R., Kheraskova, T.N., Kozlov, V.I., Puchkov, V.N.,
655 Volozh, Yu.A., 2008. The East European Craton (Baltica) before and during the assembly
656 of Rodinia, *Precambrian Research*, 160, 23-45.
- 657 Buchan, K.L., Ernst, R.E., Kumarapeli, P.S., 2004. Paleomagnetism of the Grenville diabase
658 dyke swarm and implications for the mid Vendian paleolatitude of Laurentia. *EOS:*
659 *Trans. Am. Geophys. Union* 85 (17) (Abstract GP24A-02, JA166).
- 660 Cawood, P.A., van Gool, J.A.M., Dunning, G.R., 1996. Geological development of eastern
661 Humber and western Dunnage zones: Corner Brook - Glover Island region,
662 Newfoundland: *Canadian Journal of Earth Sciences*, 33, 182–198.

- 663 Cawood, P.A., McCausland, P.J.A., Dunning, G.R., 2001. Opening Iapetus: Constraints from the
664 Laurentian margin in Newfoundland, Geological Society of America Bulletin, 113, 443-
665 453.
- 666 Cogné, J.P., 2003. PaleoMac: a Macintosh application for treating paleomagnetic data and
667 making plate reconstructions, *Geochem. Geophys. Geosyst.*, 4(1), 1007,
668 doi:10.1029/2001GC000227
- 669 Cocks, L.R.M., Torsvik, T.H., 2002. Earth geography from 500 to 400 million years ago: a
670 faunal and paleomagnetic review, *Journal of Geological Society, London*, 159, 631-644.
- 671 Compston, W., Sambridge, M.S., Reinefrank, R.F., Moczydlowska, M., Vidal, G., Claesson, S.,
672 1995. Numerical ages, volcanic rocks and the earliest faunal zone within the Late
673 Precambrian of east Poland, *Journal of the Geological Society, London*, 152, 599–611.
- 674 Dahlgren, S., 1994. 1994, Late Proterozoic and Carboniferous ultramafic magmatism- from the
675 Fen Carbonatite Complex, S. Norway: *Lithos*, v. 31, p. 141–154
- 676 Elming, S.-A., Kravchenko, S., Layer, P., Rusakov, O., Glevasskaya, A., Mikhailova, N.,
677 Bachtadse, V., 2007. Palaeomagnetism and $^{40}\text{Ar}/^{39}\text{Ar}$ age determinations of the
678 Ediacaran traps from the southwestern margin of the East European Craton, Ukraine:
679 relevance to the Rodinia break-up, *Journal of the Geological Society, London*, 164, 969–
680 982.
- 681 Fisher, R.A., 1953. Dispersion on a sphere, *Proceedings of Royal Astronomical Society London*,
682 Ser. A, 217, 295-305.
- 683 Golovanova, I.V., Danukalov, K.N. , Kozlov, V.I., Puchkov, V.N., Pavlov, V.E., Gallet, Y.,
684 Levashova, N.M., Sirota, G.S., Khairullin, R.R., Bazhenov, M.L., 2011. Paleomagnetism
685 of the Upper Vendian Basu Formation of the Bashkirian Meganticlinorium Revisited,
686 *Physics of the Solid Earth*, 47, 623-635.
- 687 Grahdankin, D.V., 2003. The composition and depositional environment of the Vendian
688 complex in the Southeastern White Sea Region, *Stratigraphy and Geological Correlation*,
689 11(4), 3-34.
- 690 Grahdankin, D.V., Marusin, V.V., Meert, J., Krupenin, M.T., Maslov A.V., 2011. Kotlin
691 Regional Stage in the South Urals, *Doklady Earth Sciences*, 440, 1222-1226.

- 692 Higgins, M.D., van Breemen, O., 1998. The age of the Sept Iles layered mafic intrusion, Canada:
693 Implications for the Late Neoproterozoic/Cambrian history of southeastern Canada,
694 Journal of Geology, 106, 421–431.
- 695 Hodych, J.P., Cox, R.A., Kosler, J., 2004. An equatorial Laurentia at 550 Ma confirmed by
696 Grenvillian inherited zircons dated by LAM ICP-MS in the Skinner Cove volcanics of
697 western Newfoundland: implications for inertial interchange true polar wander.
698 Precambrian Research, 129, 93–113.
- 699 Iglesia Llanos, M.P., Tait, J.A., Popov, V., Abalmasova, A., 2005. Palaeomagnetic data from
700 Ediacaran (Vendian) sediments of the Arkhangelsk region, NW Russia: An alternative
701 apparent polar wander path of Baltica for the Late Proterozoic-Early Palaeozoic, Earth
702 and Planetary Science Letters, 240, 732-747.
- 703 Iosifidi, A., Bachtadse, V., Khramov, A., Kuznetsova, A., 2001. Paleomagnetic data for Vendian
704 basalts from Ukraine, 3rd International Conference on Problems of Geocosmos, Abstracts
705 Volume, St. Petersburg, 74-75.
- 706 Iosifidi, A.G., Khramov, A.N., Bachtadse, V., 2005. Multicomponent magnetization of Vendian
707 sedimentary rocks in Podolia, Ukraine, Russian Journal of Earth Sciences, 7(1), 1–14.
- 708 Iosifidi A.G., Mikhailova, V.A., Sal'naya, N.V., Khramov, A.N., 2012. Paleomagnetism of
709 sedimentary rocks of the Asha series, the western slope of the South Ural: new results,
710 Oil and Gas Geology. Theory and Practice, 7(4), http://www.ngtp.ru/rub/2/57_2012.pdf
711 (in Russian).
- 712 Ivanushkin, A.G., Bogoyavlenskaya, O.V., Zenkova, G.G., Breivel, I.A., Kurik, E.Y., 2009.
713 Devonian deposits of western slope of the Urals, Lithosphere, # 1, 3-22 (in Russian)
- 714 Johnson, C.L., Constable C.G., Tauxe L., Barendregt R.W., Brown L.L., Coe R.S., Layer P.W.,
715 Mejia, V., Opdyke, N.D., Singer B.S., Staudigel, H., Stone D.B., 2008. Recent
716 investigations of the 0- to 5-Ma geomagnetic field recorded by lava flows, Geochem.
717 Geophys. Geosyst., 9, Q04032, <http://dx.doi.org/10.1029/2007GC001696>.
- 718 Kamo, S.L., Gower, C.F., Krogh, T.E., 1989. Birthdate for the Iapetus Ocean – a precise U-Pb
719 zircon and baddeleyite age for the Long-Range dykes, southeastern Labrador, Geology,
720 17, 602-605.

- 721 Kamo, S.L., Gower, C.F., 1994. U-Pb baddeleyite dating clarifies age of characteristic
722 paleomagnetic remanence of Long-Range dykes, southeastern Labrador, Atlantic
723 Geology, 30, 259-262.
- 724 Kamo, S.L., Krogh, T.E., Kumarapeli, P.S., 1995. Age of the Grenville dyke swarm, Ontario-
725 Quebec – implications for the timing of Iapetan rifting, Canadian Journal of Earth
726 Sciences, 32, 273-280.
- 727 Kaufman, A.J., Corsetti, F.A., and Varni, M.A. (2007) The effect of rising atmospheric oxygen
728 on carbon and sulfur isotope anomalies in the Neoproterozoic Johnnie Formation, Death
729 Valley, USA. Chemical Geology 237: 47-63.
- 730 Khramov, A.N., Iosifidi, A.G., 2009. Paleomagnetism of the Lower Ordovician and Cambrian
731 sedimentary rocks in the section of the Narva River right bank, for the construction of the
732 Baltica kinematic model in the early Paleozoic. Izvestia, Physics Solid Earth, 45, 465-48.
- 733 King, R.F., 1955. Remanent magnetism of artificially deposited sediments, Monthly Notices of
734 Royal Astronomical Society, Geophysical Supplement, 7, 115-134.
- 735 Kirschvink, J.L., 1980. The least-square line and plane and the analysis of palaeomagnetic data:
736 Geophysical Journal of Royal Astronomical Society, 62, 699-718.
- 737 Kirschvink, J.L., Ripperdan, R.L., Evans, D.A., 1997. Evidence for a large-scale reorganization
738 of Early Cambrian continental masses by inertial interchange true polar wander, Science,
739 277, 541-545.
- 740 Komissarova, R.A., 1963. Paleomagnetic studies of the Asha series from the western slope of the
741 South Urals, in: Paleomagnetic stratigraphic studies, VNIGRI Reports, Leningrad, 204,
742 69-82 (in Russian).
- 743 Komissarova, R.A., 1970. The studies of ancient remanence of some sedimentary rocks in the
744 South Ural and the problem of secondary remagnetization, PhD thesis, IFZ, Moscow, 140
745 p. (in Russian).
- 746 Kozlov, V.I., Ed., 2002. Geological Map of the Russian Federation and Adjacent Territory of the
747 Republic of Kazakhstan. Scale: 1:1 000000 (a New Series). Sheet N-40(41)-Ufa). St.
748 Petersburg: VSEGEI.

- 749 Kozlov, V.I. Upper Riphean and Vendian of the South Urals, Nauka, Moscow, 1982, 128 p. (in
750 Russian)
- 751 Kumarapeli, P.S., Dunnning, G.R., Pintson, H., Shaver, J., 1989. Geochemistry and U-Pb zircon
752 age of comenditic metafelsites of the Tibbit Hill Formation, Quebec Appalachians:
753 Canadian Journal of Earth Sciences, 26, 1374–1383.
- 754 Kuznetsov, N.B., Natapov, L.M., Belousova, E.A., Reilly, S.Y.O', Griffin, W.L., 2010.
755 Geochronological, geochemical and isotopic study of detrital zircon suites from late
756 Neoproterozoic clastic strata along the NE margin of the East European Craton:
757 Implications for plate tectonic models, Gondwana Research, 17, 583-601.
- 758 Kuznetsov, N.B., Romanyuk, T.V., Shatsillo, A.V., Golovanova, I.V., Danukalov, K.N., 2012a.
759 The first results of the bulk U/Pb-dating of detrital zircons (LA-ICP-MS) from Asha
760 Formation, the Southern Urals - the stratigraphic, paleogeographic and paleo-tectonic
761 aspects, Doklady Earth Sciences, 447, 1240-1246.
- 762 Kuznetsov, N.B., Romanyuk, T.V., Shatsillo, A.V., Orlov, S.Y., Golovanova, I.V., Danukalov,
763 K.N., Ipatieva, I.S., 2012b. The Age of the detrital zircons from Asha Group, the
764 Southern Urals - verification of idea about the spatial conjugation of Baltica and Australia
765 within the Rodinia Supercontinent (a positive test of the "Australia Upside Down
766 conception"), Lithosphaera, 4, 59–77 (in Russian).
- 767 Kuznetsov, N.B., Meert, J.G., Romanyuk, T.V. Ages of Detrital Zircons (U/Pb, La-ICP-MS)
768 from the Latest Neoproterozoic - Middle Cambrian(?) Asha Group and Early Devonian
769 Takaty Formation, the South-Western Urals: a Test of an Australia-Baltica connection
770 within Rodinia, Precambrian Research, 2013 doi: 10.1016/j.precamres.2013.09.011).
- 771 Lewandowski, M., Abrahamsen, N., 2003. Paleomagnetic results from the Cambrian and
772 Ordovician sediments of Bornholm (Denmark) and Southern Sweden and
773 paleogeographical implications for Baltica, Journal of Geophysical Research, 108,
774 doi:10.1029/2002JB002281.
- 775 Levashova, N.M., Bazhenov M.L., Joseph G. Meert J.G., Kuznetsov N.B., Golovanova I.V.,
776 Danukalov K.N., Fedorova N.M., 2013. Paleogeography of Baltica in the Ediacaran:
777 Paleomagnetic and geochronological data from the clastic Zigan Formation, South Urals,
778 Precambrian Research, 236, 16-30.

- 779 Martin, M.W., Grazhdankin, D.V., Bowring, S.A., Evans, D.A.D., Fedonkin, M.A., Kirschvink,
780 J.L., 2000. Age of Neoproterozoic bilaterian body and trace fossils, White Sea, Russia:
781 Implications for metazoan evolution, *Science*, 288, 841-845.
- 782 Maslov, A.V., 2004. Riphean and Vendian sedimentary sequences of the Timanides and
783 Uralides, the eastern periphery of the East European Craton, in: Gee D.G., Pease V.
784 (eds.), *The Neoproterozoic Timanides Orogen of Eastern Baltica*, Geological Society of
785 London Memoirs, 30, 19-35.
- 786 Maslov, A.V., Meert, J., Levashova, N.M., Ronkin, Yu. L., Grazhdankin, D.V., Kuznetsov, N.B.,
787 Krupenin, M.T., Federova, N.M., Ipat'eva, I.S., 2013. New constraints for the age of
788 Vendian glacial deposits, *Doklady Earth Sciences*, 449, 303-308.
- 789 McCausland, P.J.A., Hodych, J.P., 1998. Palaeomagnetism of the 550 Ma Skinner Cove
790 volcanics of western Newfoundland and opening of the Iapetus Ocean, *Earth and*
791 *Planetary Science Letters*, 163, 15–29.
- 792 McCausland, P.J.A., Van der Voo, R., Hall, C.M., 2007. Circum-Iapetus paleogeography of the
793 Precambrian–Cambrian transition with a new paleomagnetic constraint from Laurentia,
794 *Precambrian Research*, 156, 125-152.
- 795 McElhinny, M.W., 1964. Statistical significance of the fold test in palaeomagnetism.
796 *Geophysical Journal of Royal Astronomical Society*, 8, 338-340.
- 797 McFadden, P.L., McElhinny, M.W., 1990. Classification of the reversal test in paleomagnetism,
798 *Geophysical Journal International*, 103, 725-729.
- 799 Meert, J.G., Van der Voo, R., Powell, C.M., Li, Z., McElhinny, M.W., Symons, D.T.A., 1993. A
800 plate-tectonic speed limit, *Nature*, 363, 216-217.
- 801 Meert, J.G., Van der Voo, R., Payne, T., 1994. Paleomagnetism of the Catotxin volcanic
802 province: a new Vendian–Cambrian apparent polar wander path for North America,
803 *Journal of Geophysical Research*, 99 (B3), 4625–4641.
- 804 Meert, J.G., Van der Voo, R., 1997. The assembly of Gondwana 800-550 Ma, *Journal of*
805 *Geodynamics*, 23, 223-235.
- 806 Meert, J.G., Torsvik, T.H., Eide, E.A., Dahlgren, S., 1998. Tectonic significance of the Fen

807 Province, S. Norway: constraints from geochronology and paleomagnetism. *Journal of*
808 *Geology*, 106, 553– 564.

809 Meert, J.G., Van der Voo, R., 2001. Comment on "New palaeomagnetic result from Vendian red
810 sediments in Cisbaikalia and the problem of the relationship of Siberia and Laurentia in
811 the Vendian" by S. A. Pisarevsky, R. A. Komissarova and A. N. Khramov, *Geophysical*
812 *Journal International*, 146, 867 – 870.

813 Meert, J.G., Tamrat, E., 2004. A mechanism for explaining rapid continental motion in the Late
814 Neoproterozoic, in: Eriksson, P.G., Altermann, W., Nelson, D.R., Mueller, W.U. and
815 Cantuneanu, O. (eds.) *The Precambrian Earth: Tempos and Events, Developments in*
816 *Precambrian Geology* 12, Elsevier, Amsterdam, pp. 255-267.

817 Meert, J.G., Walderhaug, H.J., Torsvik, T.H., Hendriks, B.W.H., 2007. Age and paleomagnetic
818 signature of the Alnø carbonatite complex (NE Sweden): Additional controversy for the
819 Neoproterozoic paleoposition of Baltica, *Precambrian Research*, 154, 3-4, 159-174.

820 Meert J.G., 2014. Ediacaran–Early Ordovician paleomagnetism of Baltica: A review, *Gondwana*
821 *Research*, 25, 159-169.

822 Mitchell, R.M., Kilian, T.M., Raub, T.D., Evans, D.A.D., Bleeker, W., Maloof, A.C., 2011.
823 Sutton hotspot: Resolving Ediacaran-Cambrian tectonics and true polar wander for
824 Laurentia, *American Journal of Science*, 311, 651-663.

825 Murphy, J.B., Pisarevsky, S.A., Nance, R.D., Keppie, J.D., 2004. Neoproterozoic-Early
826 Paleozoic evolution of peri-Gondwanan terranes: Implications for Laurentia–Gondwana
827 connections. *International Journal of Earth Sciences*, 93, 659–682.

828 Murthy, G.S., 1971. The paleomagnetism of diabase dykes from the Grenville Province.
829 *Canadian Journal of Earth Sciences*, 8, 802–812.

830 Murthy, G.S., Gower, C.F., Tubrett, M., Patzold, R., 1992. Paleomagnetism of Eocambrian Long
831 Range dykes and Double Mer Formation from Labrador, Canada, *Canadian Journal of*
832 *Earth Sciences*, 29, 1224–1234.

833 Nawrocki, J., Boguckij, A., Katinas V., 2004. New Late Vendian palaeogeography of Baltica
834 and the TESZ, *Geological Quarterly*, 48 (4), 309–316.

- 835 O'Brien, T.M., van der Pluijm, B.A., 2012. Timing of Iapetus Ocean rifting from Ar
836 geochronology of pseudotachylytes in the St. Lawrence Rift System of southern Quebec.
837 *Geology*, 40, 443–446.
- 838 Park, J.K., 1992. Did Laurentia pass over the south pole during earliest Cambrian time?
839 *Geological Survey of Canada Paper* 92-1E, 11-22.
- 840 Park, J.K., 1994. Paleomagnetic constraints on the position of Laurentia from middle
841 Neoproterozoic to early Cambrian times, *Precambrian Research*, 69, 95-112.
- 842 Petterson, R.J., Prave, A.R., Wernicke, B.P., Fallick, A.E., 2011. The Neoproterozoic Noonday
843 Formation, Death Valley region, California, *Bulletin of the Geological Society of*
844 *America*, doi: 10.1130/B30281.1.
- 845 Piper, J.D.A., 1988. Paleomagnetism of (late Vendian–earliest Cambrian) minor alkaline
846 intrusions, Fen Complex, southeast Norway, *Earth and Planetary Science Letters*, 90,
847 422–430.
- 848 Pisarevsky, S.A., Komissarova, R.A., Khramov. A.N., 1999. A multicomponent magnetization
849 of the Vendian red sediments from Southern Urals and the palaeoposition of the Baltica,
850 *Aarhus Geoscience*, 8, 95-98.
- 851 Pisarevsky, S.A., Komissarova, R.A., Khramov. A.N., 2000. New palaeomagnetic result from
852 Vendian red sediments in Cisbaikalia and the problem of the relationship of Siberia and
853 Laurentia in the Vendian, *Geophysical Journal International*, 140 (3), 598-610.
- 854 Pisarevsky, S. A., Komissarova, R.A., Khramov. A.N., 2001. Reply to comment by J. G. Meert
855 and R. Van der Voo on "New palaeomagnetic result from Vendian red sediments in
856 Cisbaikalia and the problem of the relationship of Siberia and Laurentia in the Vendian",
857 *Geophysical Journal International*, 146, 871 – 873.
- 858 Pisarevsky, S.A., Murphy, J.B., Cawood, P.A., Collins, A.S., 2008. Late Neoproterozoic and
859 Early Cambrian palaeogeography: models and problems. In: Pankhurst, R.J., Trouw,
860 R.A.J., Brito Neves, B.B., de Wit, M.J. (Eds.), *West Gondwana: pre-Cenozoic*
861 *Correlations across the South Atlantic Region*. Geological Society, Special Publications,
862 294, 9–31.

- 863 Poorter, R.P.E., 1972. Preliminary paleomagnetic results from the Fen carbonatite complex, S.
864 Norway, Earth and Planetary Science Letters, 17, 194-198.
- 865 Popov, V., Iosifidi, A., Khramov, A., Tait, J., Bachtadse, V., 2002. Paleomagnetism of Upper
866 Vendian sediments from the Winter Coast, White Sea region, Russia: Implications for the
867 paleogeography of Baltica during Neoproterozoic times, Journal of Geophysical
868 Research, 107, 10.1029/2001JB001607.
- 869 Popov, V.V., Khramov, A.N., Bachtadse, V., 2005. Palaeomagnetism, magnetic stratigraphy,
870 and petromagnetism of the Upper Vendian sedimentary rocks in the sections of the
871 Zolotitsa River and in the Verkhotina Hole, Winter Coast of the White Sea, Russia,
872 Russian Journal of Earth Sciences, 7, 2, 1–29.
- 873 Powell, C.M., Li, Z.X., McElhinny, M.W., Meert, J.G., and Park, J.K., 1993. Palaeomagnetic
874 constraints on timing of the Neoproterozoic breakup of Rodinia and the Cambrian
875 formation of Gondwana, Geology, 21, 889–892.
- 876 Powell, C.M., 1995. Are Neoproterozoic glacial deposits preserved on the margins of Laurentia
877 related to the fragmentation of two supercontinents?: Comment. Geology, 23, 1053–
878 1054.
- 879 Puchkov, V.N., Svetlakova A.N., Razuvaev V.I., 2001. Geologic interpretation of the
880 URALSEIS seismic profile (western domain), in: A.F. Morozov, ed., Deep structure and
881 geodynamics of the South Urals (URALSEIS Project), GERS Publishers, Tver, 148-154
882 (in Russian).
- 883 Puchkov, V.N., 2003. The Uralides and Timanides: their structural relationship and position in
884 the geologic history of the Ural-Mongolian fold belt, Russian Geology and Geophysics,
885 44, no. 1-2, 28-39.
- 886 Ronkin, Y., Grazhdankin, D.V., Maslov, A.V., Mizens, G.A., Matukov, D.I., Krupenin, M.T.,
887 Petrov, G.A., Lepikhina, O.P., Kornilova, A.Y., 2006. U-Pb (SHRIMP II) age of zircons
888 from ash beds of the Chernokamen Formation, Vendian Sylvitsa Group (Central Urals),
889 Doklady Earth Sciences, 411, 1341-1345.
- 890 Stratotype of the Riphean: Stratigraphy, Geochronology 1983. B.M. Keller and N.M. Chumakov,
891 eds., Nauka, Moscow, 184 p. (in Russian).

- 892 Symons, D.T.A., Chiasson, A.D., 1991. Paleomagnetism of the Callander complex and the
893 Cambrian apparent polar wander path for North America, *Canadian Journal of Earth*
894 *Sciences*, 28, 355–363.
- 895 Sun, W.W., Kodama, K.P., 1992. Magnetic anisotropy, scanning electron microscopy, and x ray
896 pole figure goniometry study of inclination shallowing in compacting clay-rich sediment,
897 *Journal of Geophysical Research*, 97, 19,599-19,616.
- 898 Tan, X., Kodama, K.P., 2002. Magnetic anisotropy and paleomagnetic inclination shallowing in
899 red beds: Evidence from the Mississippian Mauch Chunk Formation, Pennsylvania,
900 *Journal of Geophysical Research*, 107, 2311, doi: 10.1029/2001JB001636.
- 901 Tanczyk, E.I., Lapointe, P., Morris, W.A., Schmidt, P.W., 1987. A paleomagnetic study of the
902 layered mafic intrusion at Sept-Iles, Quebec, *Canadian Journal of Earth Sciences*, 24,
903 1431–1438.
- 904 Tauxe, L., Kylstra, N., Constable, C., 1991. Bootstrap statistics for paleomagnetic data, *Journal*
905 *of Geophysical Research*, 96, 11723-11740
- 906 Tauxe, L., Kent, D.V., 2004. A Simplified Statistical Model for the Geomagnetic Field and the
907 Detection of Shallow Bias in Paleomagnetic Inclinations: Was the Ancient Magnetic
908 Field Dipolar? in: *Timescales of the Paleomagnetic Field*, J.E.T. Channell, D.V. Kent, W.
909 Lowrie, and J.G. Meert, eds., *AGU Geophys. Monogr.*, 135, 101-116.
- 910 Tauxe, L., 2005. Inclination flattening and the geocentric axial dipole hypothesis, *Earth and*
911 *Planetary Science Letters*, 233, 247-261.
- 912 Tohver, E., D'Agrella-Filho, M.S., Trindade, R.I.F., 2006. Paleomagnetic record of Africa and
913 South America for the 1200–500 Ma interval, and evaluation of Rodinia and Gondwana
914 assemblies, *Precambrian Research*, 147, 193–222.
- 915 Torsvik, T.H., Trench, A., 1991. Ordovician magnetostratigraphy: Llanvirn-Caradoc limestones
916 of the Baltic platform, *Geophysical Journal International*, 107, 171-184
- 917 Torsvik, T.H., Rehnström, E.F., 2001. Cambrian palaeomagnetic data from Baltica: implications
918 for true polar wander and Cambrian palaeogeography, *Journal of Geological Society*
919 *London*, 158, 321– 329.

- 920 Torsvik, T.H., Van der Voo, R., Preeden, U., MacNiocaill, C., Steinberger, B., Doubrovine, P.V.,
921 van Hinsbergen, D.J.J., Domeir, M., Gaina, C., Tohver, E., Meert, J.G., McCausland,
922 P.J.A., Cocks, R.M., 2012. Phanerozoic polar wander, palaeogeography and dynamics,
923 Earth Science Reviews, 114, 325–368.
- 924 Trower, E.J. and Grotzinger, J. P., 2010. Sedimentology, diagenesis, and stratigraphic occurrence
925 of giant ooids in the Ediacaran Rainstorm Member, Johnnie Formation, Death Valley
926 region, California. Precambrian Research, 180 (1-2). pp. 113-124
- 927 Van Alstine, D.R., Gillett, S.L., 1979. Paleomagnetism of the upper Precambrian sedimentary
928 rocks from the Desert Range, Nevada, Journal of Geophysical Research, 84, 4490–4500.
- 929 Van der Voo, R., 1990. The reliability of paleomagnetic data, Tectonophysics, 184, 1–9.
- 930 Van Staal, C. R., Dewey, J. E, Mac Niocaill, C., McKerrow, W. S. 1998. The Cambrian-Silurian
931 tectonic evolution of the northern Appalachians and British Caledonides: history of a
932 complex, west and southwest Pacific-type segment of Iapetus. In: Blundell, D. J. & Scott,
933 A. C. (eds) *Lyell: the Past is the Key to the Present*. Geological Society, London, Special
934 Publications, 143, 199-242.
- 935 Velikanov, V.A., 1985. The general Vendian section of Podolia, in: B.S. Sokolov, M.A.
936 Fedonkin (eds.), *The Vendian system*, Nauka, Moscow, pp. 35-67 (in Russian).
- 937 Walderhaug, H.J., Torsvik, T.H., Halvorsen, E., 2007. The Egersund dykes (SW Norway): a
938 robust Early Ediacaran (Vendian) palaeomagnetic pole from Baltica, Geophysical Journal
939 International, 168, 935–948.
- 940 Williams, H., 1995. Geology of the Appalachian-Caledonian orogen in Canada and Greenland:
941 Ottawa, Geological Survey of Canada, Geology of North America, v. F1, 944 p.
- 942 Wingate, M.T.D., Giddings, J.W., 2000. Age and paleomagnetism of the Mundine Well dyke
943 swarm, Western Australia: Implications for an Australia-Laurentia connection at 755 Ma,
944 Precambrian Research, 100, 335–357.
- 945 Zijdeveld, J.D.A., 1967. AC demagnetization of rocks: analysis of results, in: Collinson, D.W.
946 and Creer, K.M., (eds.), *Methods in Paleomagnetism*, Elsevier, Amsterdam, 254-286.
- 947

FIGURE CAPTIONS

Figure 1. (a) Location of the Baltica block with Precambrian basement (shaded), the Urals (brown band), and the study area in the South Urals (SU, red square). (b) Schematic map of the SW Urals with the limits of the Bashkir Anticlinorium (Uplift) shown as thick dotted line (simplified after Kozlov, 2002). The thickest red line denotes the Zilmerdak Fault, to the west of which Ediacaran rocks are overlain by Paleozoic rocks without angular unconformity. Thick black line is the Main Uralian Fault. Red stars denote the sampling localities numbered as in the text and Table 1, where the Upper Ediacaran Basu Formation was studied; white unlabeled stars denote localities, where no consistent results were obtained. (c) Simplified stratigraphic column of the Ediacaran sequence of the SW Urals.

Figure 2. Representative thermal demagnetization plots of the Ediacaran Basu Formation from the western part of the South Urals, in stratigraphic coordinates. Isolated intermediate- (blue) and high-temperature (red) components are denoted by thick dashed lines. Temperature steps are in degrees Celsius. Magnetization intensities are in mA/m. Full (open) circles represent vector endpoints projected onto the horizontal (vertical) plane.

Figure 3. (a-b) Stereoplots of high-temperature component, HTC, site-mean directions in the late Ediacaran clastic rocks of the Basu Fm. in situ (a) and after tilt correction (b). Black (brown) symbols and lines denote the sites with confidence circles of $< 15^\circ$ ($> 15^\circ$). Stars are the overall mean directions with associated confidence circles (thick lines) for the sites with confidence circles of $< 15^\circ$. (c) Stereoplots of HTC sample directions in the same rocks after tilt correction: circles, accepted for calculation of site-means, inverted triangles, discarded as anomalous. Star is mean direction for accepted samples. Diamonds and shaded circles are mean directions and confidence circles for a high-temperature component that is claimed by Iosifidi et al. (2012) to be present in the Basu rocks (see text for detail). Full (open) symbols and solid (dashed and dotted) lines are projected onto lower (upper) hemisphere. (d) Plot of concentration parameter k versus stepwise unfolding for the sites with confidence circles of $< 15^\circ$. Solid vertical line denotes 95% confidence level.

Fig. 4. (a-c) Representative thermal demagnetization plots of samples of the Ediacaran Basu Formation from slump-1 (b-c) and non-deformed rocks (d) in stratigraphic coordinates. Isolated high-temperature components are denoted by thick dashed red lines. Other notation as in Figure 2. (e-f) Stereoplots of high-temperature component, HTC, in the samples from the slump (circles) and non-deformed rocks (inverted triangles) in situ (e) and after tilt correction (d). Star (large inverted triangle) is overall mean direction with associated confidence circle (thick line) for the slump samples (non-deformed rocks). Other notation as in Figure 3. (g) Plot of

concentration parameter k versus degree of unfolding for the slump samples only. Solid vertical line denotes 95% confidence level.

Fig. 5. Picture of the sampled slump-2 and paleomagnetic data. (a) Picture of slump-2. Circles (inverted triangles) denote sample locations in the slumped (non-deformed) rocks. (b-d) Representative thermal demagnetization plots of samples of the Ediacaran Basu Formation from slump-2 (a-b) and non-deformed rocks nearby (c) in stratigraphic coordinates (some samples from non-deformed rocks fall outside the picture). Other notation as in Figures 2 and 4b-d. (e-g) Stereoplots of high-temperature component, HTC, in the samples from the slump (circles) and non-deformed rocks (inverted triangles) in situ (d), after tilt correction (e), and at optimal unfolding of 55% (f). Other notation as in Figures 3 and 4e-f. (g) Plot of concentration parameter k versus degree of unfolding for the slump samples only. Solid vertical line denotes 95% confidence level.

Figure 6. Stereoplots of HTC sample directions (circles) for different values of shallowing factor (see text for detail) transferred to the projections pole. Thin blue circle is shown for scale. Thick dashed red ellipse drawn according to the elongation parameter E is to highlight the form of the distribution, with its long axis shown as solid black line. Solid arrow denotes mean declination of the HTC.

Figure 7. The APWP's for Baltica for the 140 – 480 Ma interval (Torsvik et al., 2012) and the selected Cambrian-Ediacaran paleomagnetic poles from Baltica keyed as in Table 5 and the text. Reference data are shown as north poles (solid dots) with associated confidence circles (thin solid lines) connected with thick solid green line; the ages in Ma are shown for some reference poles. The Ediacaran poles that are used for interpretation are shown as stars with yellow-filled confidence circles and connected with thick dashed lines with the age differences between neighboring data; the poles that are discussed but discarded are shown as squares with gray-filled confidence circles.

Figure 8. (a) The Phanerozoic APWP's for Laurentia (Torsvik et al., 2012) and the selected Ediacaran paleomagnetic poles from Laurentia keyed as in Table 6 and the text. Reference data are shown as north poles (solid dots) with associated confidence circles (thin solid lines; not shown for Mesozoic and Cenozoic for clarity) connected with thick solid green line; the ages in Ma are shown for some reference poles. The Ediacaran poles that infer a low-latitude position of Laurentia are shown as stars with yellow-filled confidence circles; the “high-latitude” poles are squares with non-filled confidence circles. (b) The time succession of the Early Cambrian-Ediacaran paleomagnetic poles from Laurentia. Arrow-heads point from older to younger data. Other notation as in Figure 8a.

Figure 9. The Baltica-Laurentia relationship at different times according to paleomagnetic poles listed in Tables 5 and 6. (a) “Standard” reconstruction for 480 Ma using mean poles for all continents from Torsvik et al. (2012); (b) Reconstruction for 550 Ma using the VWZ pole for Baltica and the Skinner Cove pole for Laurentia; (c) Reconstruction for 565 Ma using the Basu pole for Baltica and the Sept Iles A and B poles for Laurentia; (d) Reconstruction for 615 Ma using the Egersund Dykes pole for Baltica and the Long Range Dykes pole for Laurentia. Longitudinal separation of continents is arbitrary. Baltica is yellow, low-latitude Laurentia is blue, high-latitude Laurentia is gray. Red bands denote the margins of the late Neoproterozoic Iapetus Ocean.

1024 Table 1. Summary of paleomagnetic results on the Asha Series

Formation & datum	n	CS	D°	I°	k	α_{95}°	Ref.
Bakeev	15	No consistent components are isolated					Unpublished data
Uryuk-green	40	No consistent components are isolated					Unpublished data
Uryuk-red	(6)	60%	245	-37	29	13	Levashova et al., 2013
Basu-1	(17)	S	55	-37	39	6	Golovanova et al., 2011
Basu-2	5[?]	S	153	-23	18	18	Iosifidi et al., 2012
Kuk-Karauk-1	?	S	344	-8	18	8	Pisarevsky et al., 1999
Kuk-Karauk-2	?	S	39	84	9	10	Pisarevsky et al., 1999
Kuk-Karauk-3	?	S	245	-7	9	9	Pisarevsky et al., 1999
Kuk-Karauk-4	9[?]	S	46	-40	22	11	Iosifidi et al., 2012
Kuk-Karauk-5	18[?]	S	153	-14	24	7	Iosifidi et al., 2012
Kuk-Karauk-6	35(5)	Very scattered data					Unpublished data
Kuk-Karauk Congl.	29	G	343	68	12	8	Golovanova et al., 2011
Zigan	(37)	S	106	-16	25	5	Levashova et al., 2013

1025 Comments. Formation & datum is the formation name and the label of a paleomagnetic
1026 result as defined in the text. n, is the number of samples(sites)[localities] used; question marks
1027 denote the cases where the level of statistics and/or distribution of samples between sites and
1028 localities are unknown. Unpublished data are obtained by the authors of this paper but are not
1029 published separately. CS, coordinate system, which the results are presented in: S, stratigraphic,
1030 G, geographic, 60%, after 60-percent unfolding. D, declination. I, inclination. k, concentration
1031 parameter. α_{95} , radius of 95% confidence circle (Fisher, 1953).
1032

Table 2. High-temperature component directions in clastic rocks of the Basu Formation

			In situ				Tilt-corrected			
Id	S/d	n	D°	I°	k	α_{95}°	D°	I°	k	α_{95}°

Gabdukovo Village, 54.4°N, 57.2°E

GB3-1	347/18	11/10	232.6	12.6	19.6	11.2	229.5	29.1	21.0	10.8
GB3-2	341/22	4/4	66.0	-22.8	46.5	13.6	64.6	-44.2	59.3	12.0
GB		(6/2)	59.1	-17.8	-	-	56.3	-36.9	-	-

Basu River, eastern limb, 54.3°N, 57.3°E

BSE5	329/35	4/3	231.5	-15.6	8.3	45.9	231.4	18.6	8.2	46.3
BSE6-N	4/35	3/3	50.8	-8.7	19.8	28.5	40.5	-33.5	26.0	24.7
BSE6-R	4/39	6/6	245.8	12.3	40.9	10.6	233.0	45.3	34.8	11.5
BSE7	4/42	6/5	224.2	21.0	14.0	21.2	200.5	42.2	17.7	18.7
BSE9	12/45	4/2	261.0	7.0	-	-	250.1	47.8	-	-
BSE10	357/44	10/9	78.6	26.5	13.3	14.7	79.1	-17.1	12.2	15.4
BSE19	7/64	4/2	42.0	12.1	-	-	35.4	-24.6	-	-
BSE20	2/54	5/5	65.5	-3.5	22.1	16.6	48.8	-49.6	32.1	13.7
BSE		(13/8)	59.8	0.2	13.9	15.4	50.1	-36.1	16.2	14.2

Basu River, western limb, 54.3°N, 57.1°E

BSW10	197/63	12/6	351.6	-35.2	23.0	14.3	42.6	-35.5	27.8	12.9
BSW11	206/68	9/9	358.5	-38.6	39.0	8.4	58.3	-34.3	24.9	10.5
BSW12	210/78	7/7	350.5	-26.2	23.0	13.0	54.3	-40.5	91.1	6.4
BSW13	194/63	10/5	347.7	-39.1	16.7	19.3	44.2	-36.9	14.4	20.8
BSW16-1	194/78	9/7	143.0	40.4	42.7	9.3	240.1	45.6	36.3	10.1
BSW		(10/5)	346.7	-36.5	44.8	11.6	51.7	-38.8	109.2	7.4

Takata River, 54.0°N, 56.9°E

C247	197/67	7/7	358.6	-54.8	31.9	10.8	68.0	-29.4	31.9	10.8
P1644	196/72	8/8	332.9	-45.0	71.2	6.6	61.4	-42.8	43.7	8.5
P1652	196/65	8/8	329.8	-49.7	86.9	6.0	63.3	-48.5	66.1	6.9
TA		(3/3)	339.5	-50.5	54.7	16.8	64.5	-40.3	64.1	15.5

Kuk-Karauk Brook, 53.6°N, 56.7°E

KK1R	212/10	10/10	225.0	29.5	22.1	10.5	230.2	26.7	24.1	10.0
KK1N	215/7	6/6	47.7	-28.6	27.2	13.1	51.3	-26.8	22.5	14.4
KK2	14/10	10/10	225.2	25.0	18.0	11.7	220.8	29.8	16.4	12.3
KK3	214/10	6/5	245.8	50.9	29.3	14.4	254.4	45.3	27.0	15.0
KK4	207/7	6/6	240.3	47.0	11.4	20.7	246.2	42.6	9.2	23.3
KK5	177/6	6/6	237.1	29.5	43.5	10.3	238.6	24.5	34.2	11.6
KK6	190/5	6/5	36.7	-18.8	13.4	21.7	38.1	-16.6	14.4	20.9
KK8R	197/13	7/6	251.0	52.4	15.0	17.9	258.3	41.8	15.0	17.9
KK8N	206/14	7/6	31.7	-40.1	43.1	10.3	42.8	-37.3	50.8	9.5
KK9	195/13	11/11	45.6	-51.1	54.8	6.2	57.0	-43.4	58.5	6.0
KK10	187/18	7/7	218.5	39.3	92.4	6.3	228.3	28.5	209.4	4.2
KK11	190/13	8/7	216.4	45.3	51.0	8.5	226.0	38.8	44.8	9.1
KK13	193/13	6/6	52.7	-22.6	77.6	7.7	55.9	-13.9	68.0	8.2
KK14	187/14	6/5	56.1	-37.8	46.1	11.4	61.6	-26.7	51.8	10.7
C100*	224/9	7/7	60.1	-33.7	85.0	6.6	65.5	-30.8	184.2	4.5
C107*	198/19	8/8	48.2	-50.8	24.0	11.5	63.3	-38.7	25.0	11.3
KK		(17/16)	49.3	-38.1	33.5	6.5	55.5	-32.5	35.9	6.2
ALL**		(49/34)	41.1	-31.9	6.2	10.8	54.8	-34.7	31.1	4.5
Precise**		(24)	36.1	-38.2	7.8	11.4	55.9	-35.3	47.8	4.3
Imprecise**		(10)	51.4	-14.9	5.4	23.0	52.1	-33.3	15.9	12.5
LOCs**		[5]	33.2	-33.3	4.6	40.0	55.7	-36.7	215.8	5.2

Precise –Mean pole: $n=(24)$ Plat = 1.7°N, Plong = 186.1°E, K = 62.5, A₉₅ = 3.8°

- 1033 * - these sites are close to two slump folds that are discussed separately.
- 1034 ** - these means are computed by recalculating all site-means and locality-means to the common
- 1035 point at 54.3°N, 57.3°E.
- 1036 Comments. Localities are labeled as in Figure 1b; ALL is the overall mean for all sites listed in
- 1037 the table; Precise/Imprecise is the mean for all sites with $\alpha_{95} < 15^\circ$ ($\alpha_{95} > 15^\circ$); LOCs is the
- 1038 mean for five locality-means, each of which is calculated for all sites from the locality. S/d are
- 1039 mean strike and dip for the site. n, number of samples (sites) [localities] studied/used. Mean pole
- 1040 is computed by averaging site-mean virtual geomagnetic poles: Plat, pole latitude; Plong, pole
- 1041 longitude; K concentration parameter, A₉₅, radius of 95% confidence circle. Other notation as in
- 1042 Table 1.
- 1043
- 1044

1044 Table 3. Paleomagnetic data on two slumps and adjacent sites from the Basu Fm. at locality KK.

		In situ				Tilt-corrected			
Data	n	D°	I°	k	α_{95}°	D°	I°	k	α_{95}°
Slump1	20/20	57.5	-71.9	5.4	15.5	75.7	-52.4	19.1	7.7
C100	7/7	60.1	-33.7	85.0	6.6	65.5	-30.8	184.2	4.5
Slump2	22/15	135.8	-63.1	27.2	7.5	126.9	-43.3	39.4	6.2
55%	22/15					129.6	-53.0	71.5	4.6
C107	8/8	48.2	-50.8	24.0	11.5	63.3	-38.7	25.0	11.3

1045 Notation as in Tables 1 and 2.

1046

1046 Table 4. Results of distribution-shape analysis for 195 HTC sample directions.

f	D°	I°	k	α_{95}°	ΔGC°	E	E _{pr}
1.0	55.4	-36.4	17.0	2.5	40.5	1.14	2.1
0.9	55.4	-40.3	16.2	2.6	9.5	1.22	2.0
0.8	55.4	-45.2	15.2	2.7	3.7	1.70	1.9
0.7	55.4	-51.3	14.1	2.8	2.8	2.46	1.7

1047 Comments. f, flattening. ΔGC , angular deviation of the great circle fitted to the data from the
1048 stereonet center. E, observed elongation value. E_{pr} , predicted elongation value (interpolated
1049 from Tauxe and Kent (2004)). 95% critical value of E is ~1.3. Other notation as in Table 1.

1050

1050

Table 5. Paleomagnetic poles from Baltica (Late Ediacaran – Early Ordovician)

Key	Rock unit	LLat °N	Llong °E	Age	Pole (north)		A ₉₅ °
					Lat°	Long°	
EO	480 Ma mean pole			480	-27	239	6.6
NS	Narva Sediments	59	28	500	-22	267	6.0
AS	Andrarum Shale	56	14	500	-52	291	8.5
T	Tornetrask Fm.	68	19	535	-56	295	13.2
NE	Nekso Sandstone	55	15	545	-40	357	5.3
VN	Volhynia North	53	28	540-550	-20	184	3.3
VS	Volhynia South	53	28	540-550	-24	103	3.3
ZI	Zigan Fm.	54	57	542-550	-16	138	3.7
Z	Zolotitsa	65	40	550-555	-32	113	2.1
VH	Verkhovina	65	40	550-555	-32	117	2.0
AR	Arkhangelsk	65	40	550-555	-28	110	3.5
WC	Winter Coast	65	40	550-555	-25	132	2.9
VWZ [#]	VS, Z, VH, AR, WC, ZI			540-555	-27	119	11.4
BA	Basu Fm.	54	57	560-570	2	186	3.8
F	Fen Complex	50	9	583	-57	331	8.8
A1	Alnø steep	62	17	584	-63	281	13.3
A2	Alnø shallow	62	17	584	-4	89	24.4
ED	Egersund Dykes	58	6	616	-31	224	15.7

1051

- overall mean of the listed poles.

1052

Comment: References by pole labels: (EO) – Torsvik et al., 2012; (NS) - Khramov and Iosifidi,

1053

2009; (AS, T) - Torsvik and Rehnstrom, 2001; (NE) - Lewandowski and Abrahamsen, 2003;

1054

(VN, VS) - Elming et al., 2007; (ZI) – Levashova et al., 2013; (ZO, VH) - Popov et al., 2005;

1055

(AR) - Iglesia-Llanos et al., 2004; (WC) - Popov et al., 2002; (BA) – this study; (F) - Meert et

1056

al., 1998; (A1,A2) - Meert et al., 2007; (ED) - Walderhaug et al., 2007. Llat, Locality latitude

1057

(°N). Llong, Locality longitude (°E). Plat, Paleomagnetic pole latitude (°N if positive). Plong,

1058

Paleomagnetic pole longitude (°E). A₉₅ (°), radius of confidence circle for paleomagnetic poles

1059

(Fisher, 1953). Age, rock age in Ma.

1060

1061

1061 Table 6. Early Cambrian - Ediacaran paleomagnetic poles from Laurentia

Key	Rock unit				Pole (north)			
		Llat °N	Llong °E	Age	Lat°	Long°	A ₉₅ °	Pos
MR	Mont Rigaud	45.4	285.7	532±2	-12	185	4.8	L
SC	Skinner Cove Fm.	49.5	302.0	550±3	16	157	9.1	L
JR	Johnnie Fm. (unrotated)	36.8	244.7	ca. 555	10	122	8.0	L
SI-A	Sept Iles intrusion 'A'	50.2	295.5	564±4	20	141	7.1	L
SI-B	Sept Iles intrusion 'B'	50.2	295.5	564±4	-44	134	9.8	H
CT-A	Catoctin Basalts 'A'	38.5	281.8	572±5	-42	117	17.5	H
CT-B	Catoctin Basalts 'B'	38.5	281.8	572±5	-4	184	10.0	L
CT-C	Catoctin Basalts 'C'*	38.5	281.8	572±5	19	134	9.4	L
CC	Callander Complex	46.2	280.6	577±1	-46	121	6.0	H
GD-A	Grenville Dykes 'A'	46.0	282.0	590±2	-3	151	12	L
GD-B	Grenville Dykes 'B'	46.0	282.0	590±2	-60	160	14.0	H
LR	Long Range dykes	53.7	303.1	615±2	-19	175	17.4	L

1062 * - identified as a Middle Paleozoic overprint by Meert et al., 1994

1063 Comments: References by pole labels: (MR) - McCausland et al., 2007; (SC) – Cawood et
1064 al., 2001; (JR) – Meert et al., 1994, but see Hodych et al., and discussion in the text; (SI) –
1065 Tanczyk et al., , 1987; (CT) – Meert et al., 1994; Aleinikoff et al., 1995; (CC, GD) – Symons et
1066 al., 1991; Kamo et al., 1995; (LR) - Kamo et al., 1989 and Kamo and Gover, 1994. Pos, inferred
1067 position of Laurentia: H, high-latitude; L, low-latitude. Other notation as in Table 5.

1068

Figure 1. Levashova et al.

Figure 2. Levashova et al.

Figure 3 Levashova et al.

Figure 5. Levashova et al.

Figure 6. Levashova et al.

Figure 7 Levashova et al.

Figure 8. Levashova et al.

Figure 9 Levashova et al

Graphical Abstract