**Provenance of Permian-Triassic Gondwana sequence unit accreted to the Banda Arc:
constraints from zircon U–Pb and Hf isotopes**

Christopher J Spencer^{1,2*}, Ron A Harris¹, Jon R Major^{1,3}

¹ Department of Geological Sciences, Brigham Young University, Provo UT 84602, USA

² Department of Applied Geology, Curtin University, Perth WA 6845, Australia

³ Bureau of Economic Geology, University of Texas at Austin, Austin TX 78758, United States

*Corresponding Author

Dr Christopher J Spencer

Department of Applied Geology,

Curtin University,

Perth WA 6845

Australia

Email: cspencer@curtin.edu.au

Phone: +61 (0)8 9266 1951

Fax: +61 (0)8 9266 3153

Abstract

Analysis of zircons from Australian affinity Permian-Triassic units of the Timor region yield age distributions with large age peaks at 230-400 Ma and 1750-1900 Ma, which are similar to zircon age spectra found in rocks from NE Australia and the eastern Cimmerian block. It is likely that these terranes, which are now widely separated, were once part of the northern edge of Gondwana near what is now the NW margin of Australia. The Cimmerian Block rifted from Gondwana in the Early Permian during the initial formation of the Neo-Tethys Ocean. The zircon age spectra of the Gondwana Sequence of NW Australia is most similar to the terranes of northern Tibet and Malaysia, further substantiating their initial connection. A large 1750-1900 Ma zircon peak is also very common in other terranes from Baoshan (SW China) to Borneo.

Hf analysis of zircon from the Aileu Complex in Timor and Kisar shows a bimodal distribution (both radiogenically enriched and depleted) in the Gondwana Sequence of NW Australia at ~300 Ma. The magmatic event from which these zircons were derived was likely bimodal (i.e. mafic and felsic). This is substantiated by the presence of Permian mafic and felsic rocks in the Banda Arc. Similar rock types and isotopic signatures are also found in Permian-Triassic igneous units throughout the Cimmerian continental block.

The Permian-Triassic rocks of the Timor region fill syn-rift intra-cratonic basins that successfully rifted in the Jurassic to form the NW margin of Australia. This passive continental margin first entered the Sunda Trench in the Timor region at around 7-8 Ma causing the Permo-Triassic rocks to accrete to the edge of the Asian Plate and emerge as a series of mountainous islands in the young Banda collision zone. Eventually, the Australian continental margin will collide with the southern edge of the Asian plate and these Gondwanan terranes will rejoin.

Keywords: Banda arc; Timor; detrital zircons; Hf isotopes; Gondwana; Cimmeria

Introduction

Asia is an evolving supercontinent built mostly of continental fragments rifted from the former supercontinent of Gondwana (i.e. Metcalfe, 2013). Some of these fragments were rifted from what is now the passive margin of northern Australia. Petrographic and paleocurrent studies of the Gondwana affinity sedimentary rocks of northern Australia indicate they were derived mostly from the north where continental fragments of Gondwana were subsequently rifted away (Bird, 1987; Bird and Cook, 1991; Zobel, 2007). Stratigraphic detective work in Asia has not been able to resolve where these continental fragments now reside. Various paleogeographic reconstructions attach the Lhasa, Sibumasu (Siam-Burma-Malaysia-Sumatra), East Java and Borneo terranes to NW Australia at various times. This paper aims to test the veracity of these reconstructions by combining the methods of sandstone petrography and U-Pb analysis of detrital zircons to address the question of what rifted away from the NW margin of Australia.

Large sections of the strata making up the NW passive margin of Australia are accreted to the Banda Arc via Late Miocene to present arc-continent collision (i.e. Harris, 2011). We have analyzed petrological relations and U-Pb ages of detrital zircons from Gondwana affinity sandstones and metamorphic rocks in the Banda Arc in order to determine a provenance and age fingerprint to compare with various terranes in Asia.

Description of the Gondwana Sequence

Three major tectono-lithic units make up the Banda Arc collision. The Banda Terrane forms the upper plate of the collision. It is of Asian affinity and occupies the highest structural level (Harris, 2006; Standley and Harris, 2009). Subducting beneath the Banda Terrane is the Australian passive margin. It consists of two mega-sequences separated by a Late Jurassic breakup unconformity (Fig. 2). Pre- and syn-breakup strata below the unconformity formed during the Pennsylvanian to Jurassic while Australia was part of Gondwana and are known as the Gondwana Mega-sequence. The Australian Passive Margin Mega-sequence overlies the unconformity.

The recent subduction of the Australian passive margin beneath the Banda Terrane caused the two mega-sequences on it to detach and accrete to the upper plate. The Australian passive margin mega-sequence detaches from the subducting Australian Plate at the Jurassic breakup unconformity to form a classic imbricate stack. The underlying Gondwana mega-sequence is carried further down the subduction zone where it detaches to form a duplex system.

The stratigraphy and sedimentology of Gondwana mega-sequence units in the Banda Arc is described by many previous studies dating back to early Dutch expeditions of late 19th and early 20th centuries (Rothpletz, 1891; Wanner, 1913 and 1956). Other definitive studies of these rocks include: Simons (1939), Gageonnet and Lemoine (1957), Audley-Charles (1968), Gianni (1971), Grady and Berry (1977), Charlton (1989), Bird and Cook (1991), Hunter (1993), Barkam (1993), Sawyer et al. (1993), Prasetyadi and Harris (1996), Harris et al. (1998, 2000 and 2009), Charlton (2002), Charlton et al. (2009), Vorkink (2004), McCartain et al. (2006), Zobel (2007), Haig et al. (2008), and Major (2011). From these studies it is evident that Gondwana Mega-sequence units exposed in the Banda Collision and drilled offshore on the Australian margin represent is divided into two series, the Permian-Triassic Aileu-Maubisse Series, and the

Permian-Jurassic Keknenno Series (Lemoine, 1959). The stratigraphic relationship between the two series is ambiguous due to the fact that in most places the Aileu-Maubisse series is thrust over the Keknenno Series.

Aileu Complex

The Aileu Complex is the metamorphosed part of the Aileu-Maubisse Series. It is found in several places along the north coast of Timor as far east as Manatutu (Fig. 1). It is also exposed on the islands of Kisar (Harris, 2006; Major et al., 2013) and likely on other islands to the east (Kanenko et al., 2007), although the age of these occurrences is unknown. The Aileu complex consists of a protolith of Permian-Triassic, and possibly Jurassic, psammite and limestone, and basalt, rhyolite, tuffaceous material and gabbro and granite plutons (Berry and McDougall, 1986; Prasetyadi and Harris, 1996; Harris, 2012). These units are metamorphosed into pelitic schist and gneiss, marble, phyllite and amphibolite. Metamorphic grade varies between lower greenschist to upper amphibolite facies (Berry and Grady, 1981). Some high temperature metamorphism may have occurred during the rifting event that formed the edge of the Australian continental margin. This event is overprinted by medium to high-pressure metamorphism during Late Miocene onset of collision in central Timor (Berry and McDougall, 1986; Harris, 1991; Harris et al., 1998).

Maubisse Formation

The Maubisse Formation consists of a distinct red, crinoidal limestone, shale and volcanic rocks that were deposited mostly during the Permian with ages ranging from latest Pennsylvanian to Triassic, and represents the oldest rocks exposed in the Banda Orogen (de Roever, 1940; Audley-

Charles, 1968; Davydov et al., 2013). Pillow lavas found within the Maubisse Formation have geochemical signatures of within-plate and ocean-ridge basalt, which is interpreted as representing the onset of rifting (Berry and Jenner, 1982). Clastic sedimentary units found in the Maubisse Formation show fining in grain size toward to the south (Carter et al., 1976). Rocks similar to the Maubisse Formation have been documented on the Sahul Shoals of the undeformed Australian continental margin (Grady and Berry, 1977).

Atahoc Formation

The Atahoc Formation is most likely transitional with the Maubisse Formation and is Permian in age. This formation is mainly shale with some fine-grained sandstone and volcanic rocks. The basal contact has not been found and the upper contact is amygdaloidal basalt (Sawyer et al., 1993).

Cribas Formation

The Permian-Triassic? Cribas Formation is dominated by organic shale in the lower section and inter-bedded to massive sandstone units in the upper part. This sandstone is interpreted as being part of a submarine fan complex (Hunter, 1993) deposited on a shallow shelf (Bird, 1987). Another distinguishing characteristic is the presence of ironstone nodules indicative of anoxic conditions (Charlton et al., 2002).

Niof Formation

The Triassic Niof Formation is recognized in West Timor and may comprise the upper part of the Cribas Formation in East Timor. It consists of thin, inter-bedded claystone, brown,

gray and black shale, and sandstone. Bird and Cook (1991) interpreted the deposition of the Niof as turbidites in shallow to deep water. The upper portion of the Niof is interbedded with the Aitututu formation.

Aitututu Formation

The Triassic Aitututu Formation is the most distinctive unit of the Gondwana Sequence. It is a rhythmically bedded white to pink limestone with thin inter-beds of dark grey shale. The Aitututu Formation was deposited on an open marine outer shelf (Sawyer et al., 1993). It is the most lithologically distinct unit of the Kekeno Series and is used as a marker unit for structural reconstructions (Harris, 2012).

Babulu Formation

The Triassic Babulu Formation is only recognized in central Timor (Gianni, 1971; Bird and Cook, 1991) and Savu (Vorkink, 2004), and may comprise the base of the Wai Luli Formation in East Timor. The Babulu Formation consists of sandstone, shale and silts with some massive sandstone beds. Deposition of this unit most likely occurred in a proximal near shore to shelf break (Sawyer et al., 1993) through turbidity currents from a prograding delta (Bird and Cook, 1991).

Wai Luli Formation

The Late Triassic to Jurassic Wai Luli Formation is a thick succession of mostly smectite-rich mudstone (Harris et al., 1998) with some well-bedded marl, calcilutite, micaceous shale and quartz arenite. Towards the top of the formation are conglomerate and red shale units

(Audley-Charles, 1968). Iron stones are very common and commonly accumulate as loag deposits on the surface. In the Banda collision zone the thick, mechanically weak Wai Luli mudstone serves as a major decollement for imbrication of the overlying Australian Passive Margin Sequence; and as a roof thrust for structural duplexes of the Gondwana Sequence (Harris, 1991). It also is the source of much of the *mélange* matrix found along the suture zone between accreted Australian continental material and structurally overlying Asian affinity arc terranes (Harris et al., 1998).

Methods

We investigated possible source regions for Gondwana Sequence sandstones by petrographic studies of grain types and textures, and U–Pb age analyses of detrital zircon grains. The petrographic analysis (reported in Zobell, 2007) includes 16 samples of Triassic Gondwana Sequence sandstone collected from Savu, West Timor and East Timor.

U–Pb zircon age and Hf analyses were conducted on 5 sandstone and 4 metamorphic rock samples from Savu, Timor and and Kisar (Fig. 2). These samples were crushed using a jaw crusher and the < 500 μm size fraction was magnetically separated using the Carpco and Frantz magnetic separators. Heavy minerals were separated using wifely table and tetraboroethelene. Zircon grains were then hand picked for analysis. Cathode luminescence and backscatter images were acquired using a Hitachi 3400N SEM/Catan Chroma CL system and a Cameca SX-50 electron microprobe, respectively.

In situ U–Pb isotope analyses for individual grains were performed using a Nu Plasma HR MC-ICPMS coupled to a New Wave 193 nm ArF and a Photon Machines G3 193 nm ArF laser ablation system at the University of Arizona with ablation done in a He carrier gas using a

40 μm diameter spot size. Laser fluence of $\sim 4 \text{ J/cm}^2$ at 8 Hz for 30 seconds of integration. Detailed U–Pb analytical procedures are described by Gehrels et al. (2008). Samples were analyzed in sets of 9 to 12 analyses, which include 5 to 8 unknown spots, bracketed beginning and end by a pair of analyses of an in house Sri Lanka zircon standard (Gehrels et al., 2008). The R33 zircon standard was also analyzed at the beginning and end of each sample set as an independent control on reproducibility and instrument stability. Concordance is defined for ages above 700 Ma using the ratio of $^{206}\text{Pb}/^{238}\text{U}$ and $^{207}\text{Pb}/^{206}\text{Pb}$ ages, and $^{206}\text{Pb}/^{238}\text{U}$ and $^{207}\text{Pb}/^{235}\text{U}$ ages are used for those younger than 700 Ma. The accepted ages were selected from a 95% concordant subset, wherein the $^{206}\text{Pb}/^{238}\text{U}$ and $^{207}\text{Pb}/^{206}\text{Pb}$ ages are used for zircons younger and older than 700 Ma, respectively; this age was chosen because there is a natural gap in the ages of the zircons in these samples. Visualization of U-Pb concordia and zircon ages is achieved using Isoplot 4.15 (Ludwig, 2003) and densityplotter (Vermeesch, 2012). GPS locations of samples are presented in Table 2.

Hf isotope analyses were performed with a Nu Plasma HR MC-ICPMS coupled to a Photon Machines G3 193 nm ArF laser ablation system. Instrument settings were established first by analysis of 10 ppb solutions of JMC475 and a Spex Hf solution, and then by analysis of 10 ppb solutions containing Spex Hf, Yb, and Lu. The mixtures range in concentration of Yb and Lu, with $^{176}(\text{Yb}+\text{Lu})$ up to 70% of the ^{176}Hf . When all solutions yield an accurate $^{176}\text{Hf}/^{177}\text{Hf}$ ratio, instrument settings are optimized for laser ablation analyses and seven different standard zircons (Mud Tank, 91500, Temora, R33, FC52, Plesovice, and Sri Lanka) are analyzed. These standards are included with unknowns on the same epoxy mounts. Laser ablation analyses were conducted with a laser beam diameter of 50 microns, with ablation pits located on top of the U-Pb analysis pits or on an adjacent spot of similar zonation. CL images were used to ensure that

the ablation pits do not overlap multiple age domains or inclusions (Fig. 3). Each acquisition consisted of one 40-second integration on backgrounds (on peaks with no laser firing) followed by 60 one-second integrations with the laser firing. Using a typical laser fluence of ~ 5 J/cm² and pulse rate of 7 Hz. Sets of each standard was analyzed once for every ~ 15 unknowns.

Isotope fractionation was accounted for using the method of Woodhead et al. (2004): β_{Hf} is determined from the measured $^{179}\text{Hf}/^{177}\text{Hf}$; β_{Yb} is determined from the measured $^{173}\text{Yb}/^{171}\text{Yb}$ (except for very low Yb signals); β_{Lu} is assumed to be the same as β_{Yb} ; and an exponential formula is used for fractionation correction. Yb and Lu interferences were corrected by measurement of $^{176}\text{Yb}/^{171}\text{Yb}$ and $^{176}\text{Lu}/^{175}\text{Lu}$ (respectively), as advocated by Woodhead et al. (2004). Critical isotope ratios are $^{179}\text{Hf}/^{177}\text{Hf} = 0.73250$ (Patchett & Tatsumoto, 1980); $^{173}\text{Yb}/^{171}\text{Yb} = 1.132338$ (Vervoort et al. 2004); $^{176}\text{Yb}/^{171}\text{Yb} = 0.901691$ (Vervoort et al., 2004; Amelin and Davis, 2005); $^{176}\text{Lu}/^{175}\text{Lu} = 0.02653$ (Patchett, 1983). All corrections are done line-by-line. For very low Yb signals, β_{Hf} is used for fractionation of Yb isotopes. The corrected $^{176}\text{Hf}/^{177}\text{Hf}$ values are filtered for outliers (2-sigma filter), and the average and standard error are calculated from the resulting ~ 58 integrations. There is no capability to use only a portion of the acquired data.

The $^{176}\text{Hf}/^{177}\text{Hf}$ at time of crystallization is calculated from measurement of present-day $^{176}\text{Hf}/^{177}\text{Hf}$ and $^{176}\text{Lu}/^{177}\text{Hf}$, using the decay constant of ^{176}Lu ($\lambda = 1.867\text{e-}11$) from Scherer et al. (2001) and Soderlund et al. (2004). The uncertainty propagation of the epsilon notation also includes the uncertainty of the $^{207}\text{Pb}/^{206}\text{Pb}$ crystallization age, as it is time integrated. Although this may be an over propagation of uncertainty, we prefer this conservative approach for the epsilon notation when defining specific fields of similar ϵ_{Hf} compositions. Uncertainties that

incorporate the crystallization age uncertainty are on average 50 % ($1\sigma = 20\%$) larger than uncertainties that do not consider the crystallization age uncertainty.

Results

Petrography

Point count analysis of Gondwana Sequence sandstones from Savu and West and East Timor (initially reported by Zobell, 2007) indicate they are mostly texturally immature, and are characterized by large, sub-angular to sub-rounded framework grains (Fig. 4a). Modal abundances show high percentages of quartz with significant amounts of unaltered twinned feldspar (Fig. 4b), fresh mica (Fig. 4c) and lithic fragments. In combination with high percentages of matrix, or matrix replacing cement show compositional immaturity. These data preclude a far-traveled source for most of the detrital zircons and rather were likely derived from proximal sources. Compositionally, most sandstone samples plot as quartz wackes to lithic wackes (Fig. 4d) on the classification diagram of Williams (1982). On discrimination diagrams of Dickinson et al. (1983) the sandstones plot as recycled orogen provenance (Fig. 4e).

Zircon U-Pb Analysis

Zircons were separated in 5 sandstone and 4 metamorphic rock samples of the Gondwana Sequence in the Timor region. These zircon grains provide the first constraints for depositional provenance and maximum age of deposition for various sections of the Gondwana Sequence (Table 2). Four of these samples are from Triassic units exposed on the island of Savu (Indonesia), one is from Triassic sandstone in East Timor and the other four are from Permian-Jurassic units of the Aileu Complex in East Timor and Kisar Island (see Fig. 2). Most of the grains are amber to clear or pink to mauve in color with variable amounts of abrasion and rounding. Proterozoic zircons vary in color, but Paleozoic zircons are only amber to clear. Both

abraded and pristine zircons are present, although there is no apparent relationship between abrasion and age. Zircons from nine samples are variably concordant (Figs. 5, 6) and those that are < 5% discordant cluster into two main populations at ~300 Ma and ~1875 Ma (Fig. 7). Several other small peaks are found in some grains between these two age groups. The peak at ~300 Ma includes a spread of ages from mid-Carboniferous to Jurassic (~330 to 150 Ma).

Sample TL23b was collected from a leucosome pod within a migmatitic unit in the Aileu Complex. Zircons within this leucosome (17/19 analyses) give a weighted average of 300.8 ± 3.3 Ma (MSWD = 1.5) (Fig. 6). This age implies that there was partial melting associated with the rifting episode in this region, although there is no evidence of partial melting during the Miocene collision of Australia (see also Berry and Grady, 1981).

Two amphibolite samples (TL89, TL88b) were analyzed using both 40 μm and 50 μm diameter spots. Both sets of data reveal a suite of ages between ~300 Ma and ~7 Ma that lie along a poorly defined discordia. The weighted averages of oldest clusters of ages whose $^{206}\text{Pb}/^{238}\text{U}$ and $^{207}\text{Pb}/^{235}\text{U}$ uncertainties overlap with the concordia are 266 ± 4 (n = 5; MSWD = 1.7) and 256 ± 3 (n = 4; MSWD = 0.33) ($^{206}\text{Pb}/^{238}\text{U}$ age) in sample TL88b and 269 ± 4 and 244 ± 13 (n = 3; MSWD = 2.5) in sample TL89 (Fig. 8).

Hf Analysis

Hf isotopic analyses of three detrital zircon samples from Kisar (KIS05 and MT4891) and Timor (09-HS21-4) have initial ϵHf values of 11 to -21 (Fig. 9). The pre-400 Ma zircons have ϵHf values of 5 to -10 whereas post-400 Ma zircons have only subchondritic ϵHf values. Very young zircons (~7 Ma) from amphibolite samples (TL88b, TL89) have initial ϵHf values of 4.2 to -2.3 although this is likely a maximum estimate given the potential for lead loss in the ages

used to calculate the initial ϵ_{Hf} . Hf isotopic analyses of zircon grains from the Aileu Complex migmatite (sample TL23b) have initial ϵ_{Hf} values of 11 to 16.

Discussion

Provenance of the Gondwana Sequence

Samples of the Gondwana Sequence on the islands of Savu, Timor, and Kisar have dominant zircon U-Pb age peaks at ~ 300 and ~ 1875 Ma with minor Neoproterozoic to Archean subpopulations (Fig. 10). The ~ 1875 Ma population was likely derived from the North Australian Craton within the Kimberly Basin, and Halls Creek and Pine Creek orogens (Tyler, 2005; Downes et al., 2007; Worden et al., 2008; Lewis and Sircombe, 2013). Additionally, the few pre-2.4 Ga detrital zircons were also likely derived from the North Australian Craton (Downes et al., 2007; Cawood and Korsch, 2008). Potential sources for the few Meso- to Neoproterozoic detrital zircons have been identified within the Musgrave Province, Albany-Fraser Orogen, and Pinjarra Orogen of central and western Australia. The presence of zircons from Western Australia was likely transported along the west and north coast prior to convergence with the Pacific Plate (Lewis and Sircombe, 2013). However, it should be noted that the proportions of Meso- and Paleo-Proterozoic zircons seen along coastal Western Australia and the northwest Australian Shelf do not match that of the samples from Savu, Timor, and Kisar. This mismatch might reflect a different provenance than proposed or idiosyncrasies that would be resolved with a greater number of analyses. It should be further noted, that despite the inclusion of potentially far travelled detrital zircon populations, the presence of sub-angular grain fragments along with pristine mica flecks indicate that any far-travelled detrital zircon populations were reworked along a rift shoulder and deposited in a proximal basin. Paleoproterozoic zircon grains, with minor modes at 1.8 to 1.9 Ga, are found in nearly every

sample. Smaller population of Neo- and Mesoproterozoic are also present and three of the samples (SV-9, SV28C, 09HS21-4) also contain minor Archean (~ 2.5 Ga) grains. The early Permian maximum depositional age of the Aileu Formation inferred by the youngest detrital zircon age peak is consistent with that proposed by Brunnschweiler et al. (1978), who reported Permian ammonites fossils in phyllitic rocks.

Zircons in meta-gabbro samples TL88b and TL89 show incomplete lead loss from ~ 260 Ma to ~ 7 Ma (Fig. 8). Based upon the mafic nature of these samples and the incomplete replacement of the zircon, we postulate, the cores of these grains were initially baddeleyite hosted in gabbro, which was replaced by zircon during ~ 7 Ma collision-related metamorphism and the influx of Si-rich fluids. Further support to this hypothesis is that the U/Th ratios of these “metamorphic” zircons are < 10 (3.1 average). Alternatively, these Permian-age cores are simply heavily altered zircon with a pristine oscillatory zoned zircon rims. It is likely the mafic protolith for the amphibolite is a gabbro as argued from faint primary igneous textures. We interpret this unit as having been emplaced along the shoulder of a rift resulting from the Permian separation of the Cimmerian block from Gondwana.

Hf isotopes in the detrital zircon samples display variable radiogenic enrichment with average Paleoproterozoic to Neoproterozoic depleted mantle model ages. An exception to this is seen in the apparent bimodal distribution of ϵ_{Hf} values in the ~ 300 Ma dominant age peak. The end members of this bimodal distribution lie at the +16 and -22 , which correspond to the depleted mantle and a depleted mantle model age of ~ 2600 Ma, respectively. This unique distribution implies these zircons were derived from a bimodal magmatic suite, which likely formed during rifting.

Evidence for this bimodal magmatic event is seen in the Gondwana Sequence where minor amounts of rhyolite and granite are accompanied by larger volumes of alkali basalt (Audley-Charles, 1968; Wopfner and Jin, 2009). These compositions typify continental rifting events (McKee et al., 1970). Furthermore, the ~ 260 Ma upper intercept age of the metagabbro samples in this study is similar to the 270 ± 3 Ma zircon U-Pb age of the Maubisse Trachyandesite reported by Kwon et al. (2014) and these probably represent the same suite of bimodal volcanic rocks associated with the rifting of the Cimmerian block.

Our story of the Gondwana Sequence comes full circle when the final remnants of the Tethys Ocean close and the Gondwana Sequence is both subducted beneath and accreted to the Banda Arc (Harris et al., 2000; Harris, 2006; Harris, 2011). Zircons from the meta-gabbros in this study have imprecisely constrained the metamorphism associated with this event at ~ 7 Ma.

Conclusions

From the data presented in this study, we find that:

1. The Permian-Triassic Gondwana Sequence was deposited off the northern margin of the Australian continent and is now present in exhumed thrust sheets accreted to the Banda Arc complex.
2. Sandstones of the Gondwana Sequence are mostly immature and were derived predominantly from rocks of the NW Australia and fragments of the Cimmerian Block found in Tibet and Malaysia.
3. These sediments, with the accompanying bimodal volcanism, represent proximal rift shoulder deposits that were laid down shortly after the initiation of Tethys Ocean rifting in the early Permian era.

4. Zircons in meta-gabbro provide age estimates for both the rifting of Cimmeria from Gondwana in the grain cores (~300 Ma) as well as the collision of the Banda Arc with Australia (~7 Ma).
5. ϵ_{Hf} of detrital zircons provide evidence for bimodal rift-related magmatism with one cluster of data near the depleted mantle and another with Neoproterozoic depleted mantle model ages.

Acknowledgements

This project was funded by the US National Science Foundation (NSF EAR-0337221 to R.H.) and by the College of Physical and Mathematical Sciences at Brigham Young University.

Michael Vorkink, Elizabeth (Zobell) Ruiz and Jon Major provided some samples and analyses for this paper from their MSc. Thesis research in the Banda Arc. Gratitude is also expressed to George Gehrels, Mark Pecha, and Nicky Giesler who facilitated the analytical work at the University of Arizona Laserchron lab (NSF EAR-0929777 and NSF EAR-0443387). We appreciate the help of Carl Hoiland and Josh Flores with the analytical work.

Figure Captions

1. Regional tectonic map of the Banda Arc region showing active faults (black, dashed if poorly defined) and active volcanoes (red triangles) (from Harris, 2011). Stars represent localities from which samples were taken.
2. Stratigraphic column of the Gondwana Sequence.
3. Cathode luminescence (CL) image of zircon with outline of typical U-Pb (40/50 μm), and Hf (50 μm) analytical spots.

4. Column a) Photomicrographs in cross polarized light showing sub-angular to sub-rounded sandstone textures of representative Triassic Gondwana sequence sandstone from East and West Timor and Savu. b) Photomicrographs in cross polarized light of unaltered twinned feldspar grains (circled). c) Photomicrographs in cross polarized light of fresh micas. d) QFL diagram of Williams et al. (1982) for classification of sandstone samples from the Triassic Gondwana Sequence. e) QFL sedimentary provenance based on Dickinson et al. (1983). Photomicrographs and point counting data are from Zobell (2007).
5. Pb/U concordia diagram of zircon analyses from each sample constructed using Isoplot 4.15 (Ludwig, 2003). Uncertainties are shown at the 2σ level.
6. Top: Pb/U concordia diagram of zircon analyses for sample TL23b. Bottom: weighted mean diagram of the same. The youngest two analyses in grey were discarded as they fall outside the 2-sigma envelope and likely represent Pb-loss.
7. Kernel density estimation (solid line) plots of detrital zircon ages from each sample. Only ages that $< 5\%$ discordant are used. Plot is constructed using densityplotter (Vermeesch, 2012).
8. A) Pb/U concordia diagram of ages (Ma) of zircon grains from samples TL89 and TL88b constructed with the use of Isoplot (Ludwig, 2003). Uncertainties are shown at the 2σ level. 40 and 50 μm spots are displayed in gray and black, respectively. B) Pb/U concordia diagram of ages (Ma) of the youngest zircon ages analyzed with 50 μm spots. Concordia ages from the youngest three analyses. Gray analyses are not included in the concordia age calculation. C) Zircon $^{206}\text{Pb}/^{238}\text{U}$ age analyses plotted with 2σ uncertainties. The oldest plateau of ages represents the crystallization of a primary U-bearing phase in

gabbroic rocks (possibly baddelyite) and the youngest plateau represents the age of metamorphic zircon growth whereas the zircon analyses with between these two events represent varying degrees of lead loss. 40 and 50 μm spots are displayed in black and gray, respectively. Analyses within polygons (a) through (e) corresponds to weighted averages of three to five analyses that broadly represent the igneous and metamorphic crystallization events.

9. A) εHf_t vs. U-Pb analyses of detrital zircons from the samples in this study.

Uncertainties are displayed as 2σ . DM = depleted mantle. B) KDE of εHf from the youngest set of zircon ages displaying a bimodal distribution.

10. Kernel density estimation (solid line) plots of composite detrital zircon age spectra from potential source regions. Data compiled from: Cawood and Nemchin, 2000; van Wyck and Williams, 2002; Veevers et al., 2005; Zobeil, 2007; Ely, 2009; Korsch et al., 2009; Sevastjanova et al., 2011; Gehrels et al., 2012; Kwon et al., 2014; this study. Plot is constructed using densityplotter (Vermeesch, 2012).

References:

- Amelin, Y., Davis, W.J., 2005. Geochemical test for branching decay of ^{176}Lu . *Geochim. Cosmochim. Acta* 69, 465–473. doi:10.1016/j.gca.2004.04.028
- Audley-Charles, M., 1968. The Geology of Portuguese Timor. *Geol. Soc. London, Mem.* doi:10.1144/GSL.MEM.1968.004.01.02
- Barkham, S.T., 1993. The structure and stratigraphy of the Permo-Triassic carbonate formations of West Timor, Indonesia. Royal Holloway, University of London.
- Berry, R., Jenner, G., 1982. Basalt geochemistry as a test of the tectonic models of Timor. *J. Geol. Soc. London*. 139, 593–604.
- Berry, R., McDougall, I., 1986. Interpretation of $^{40}\text{Ar}/^{39}\text{Ar}$ and K/Ar dating evidence from the Aileu Formation, East Timor, Indonesia. *Chem. Geol. Isot. Geosci. Sect.* 59, 43–58. doi:10.1016/0168-9622(86)90056-4

- Berry, R.F., Grady, A.E., 1981. Deformation and metamorphism of the Aileu Formation, north coast, East Timor and its tectonic significance. *J. Struct. Geol.* 3, 143–167. doi:10.1016/0191-8141(81)90011-0
- Bird, P.R., 1987. The geology of the Permo-Triassic rocks of Kekeno, West Timor. Royal Holloway, University of London.
- Bird, P.R., Cook, S.E., 1991. Permo-Triassic successions of the Kekeno area, West Timor: implications for palaeogeography and basin evolution. *J. Southeast Asian Earth Sci.* 6, 359–371. doi:10.1016/0743-9547(91)90081-8
- Brunnschweiler, R.O., 1978. Notes on the geology of eastern Timor. *Bull. Bur. Miner. Resour. Geol. Geophys. Aust.* 192, 9–18.
- CARTER, D.J., AUDLEY-CHARLES, M.G., BARBER, A.J., 1976. Stratigraphical analysis of island arc--continental margin collision in eastern Indonesia. *J. Geol. Soc. London.* 132, 179–198. doi:10.1144/gsjgs.132.2.0179
- Cawood, P. a., Korsch, R.J., 2008. Assembling Australia: Proterozoic building of a continent. *Precambrian Res.* 166, 1–35. doi:10.1016/j.precamres.2008.08.006
- Charlton, T., Barber, A., Barkham, S., 1991. The structural evolution of the Timor collision complex, eastern Indonesia. *J. Struct. Geol.* 13, 489–500. doi:10.1016/0191-8141(91)90039-L
- Charlton, T.R., 2002. The structural setting and tectonic significance of the Lolotoi, Laclubar and Aileu metamorphic massifs, East Timor. *J. Asian Earth Sci.* 20, 851–865. doi:10.1016/S1367-9120(01)00075-X
- Charlton, T.R., Barber, A.J., McGowan, A.J., Nicoll, R.S., Roniewicz, E., Cook, S.E., Barkham, S.T., Bird, P.R., 2009. The Triassic of Timor: Lithostratigraphy, chronostratigraphy and palaeogeography. *J. Asian Earth Sci.* 36, 341–363. doi:10.1016/j.jseaes.2009.06.004
- Davydov, V.I., Haig, D.W., McCartain, E., 2013. A latest Carboniferous warming spike recorded by a fusulinid-rich bioherm in Timor Leste: Implications for East Gondwana deglaciation. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 376, 22–38. doi:10.1016/j.palaeo.2013.01.022
- Downes, P.J., Griffin, B.J., Griffin, W.L., 2007. Mineral chemistry and zircon geochronology of xenocrysts and altered mantle and crustal xenoliths from the Aries micaceous kimberlite: Constraints on the composition and age of the central Kimberley Craton, Western Australia. *Lithos* 93, 175–198. doi:10.1016/j.lithos.2006.06.005
- Ely, K.S., Sandiford, M., Phillips, D., Boger, S.D., 2013. Detrital zircon U-Pb and ⁴⁰Ar/³⁹Ar hornblende ages from the Aileu Complex, Timor-Leste: provenance and metamorphic cooling history. *J. Geol. Soc. London.* 171, 299–309. doi:10.1144/jgs2012-065
- Gageonnet, R., Lemoine, M., 1957. Sur la Stratigraphie de L'autochtone au Timor Portugais. *COMPTE* ....
- Gehrels, G.E., Valencia, V. a., Ruiz, J., 2008. Enhanced precision, accuracy, efficiency, and spatial resolution of U-Pb ages by laser ablation-multicollector-inductively coupled plasma-mass spectrometry. *Geochemistry, Geophys. Geosystems* 9, n/a–n/a. doi:10.1029/2007GC001805
- Gianni, L., 1971. The geology of the Belu district of Indonesian Timor. University of London.

- Grady, A.E., Berry, R.F., 2007. Some Palaeozoic- Mesozoic stratigraphic- structural relationships in east timor and their significance in the tectonics of timor. *J. Geol. Soc. Aust.* 24, 203–214. doi:10.1080/00167617708728981
- Haig, D.W., McCartain, E.W., Keep, M., Barber, L., 2008. Re-evaluation of the Cablac Limestone at its type area, East Timor: Revision of the Miocene stratigraphy of Timor. *J. Asian Earth Sci.* 33, 366–378. doi:10.1016/j.jseas.2008.03.002
- Harris, R., 2011. The nature of the Banda Arc–continent collision in the timor region. *Arc-Continent Collis.*
- Harris, R., 2006. Rise and fall of the Eastern Great Indonesian arc recorded by the assembly, dispersion and accretion of the Banda Terrane, Timor. *Gondwana Res.* 10, 207–231. doi:10.1016/j.gr.2006.05.010
- HARRIS, R., LONG, T., n.d. The Timor ophiolite, Indonesia: Model or myth? *Spec. Pap. - Geol. Soc. Am.* 321–330.
- Harris, R., Vorkink, M.W., Prasetyadi, C., Zobell, E., Roosmawati, N., Apthorpe, M., 2009. Transition from subduction to arc-continent collision: Geologic and neotectonic evolution of Savu Island, Indonesia. *Geosphere* 5, 152–171. doi:10.1130/GES00209.1
- Harris, R.A., 1991. Temporal distribution of strain in the active Banda orogen: a reconciliation of rival hypotheses. *J. Southeast Asian Earth Sci.* 6, 373–386. doi:10.1016/0743-9547(91)90082-9
- Harris, R.A., Sawyer, R.K., Audley-Charles, M.G., 1998. Collisional melange development: Geologic associations of active melange-forming processes with exhumed melange facies in the western Banda orogen, Indonesia. *Tectonics* 17, 458–479. doi:10.1029/97TC03083
- Hunter, D., 1993. A stratigraphic and structural study of the Maubisse area, East Timor, Indonesia.
- Kaneko, Y., Maruyama, S., Kadarusman, A., Ota, T., Ishikawa, M., Tsujimori, T., Ishikawa, A., Okamoto, K., 2007. On-going orogeny in the outer-arc of the Timor–Tanimbar region, eastern Indonesia. *Gondwana Res.* 11, 218–233. doi:10.1016/j.gr.2006.04.013
- Kwon, C.W., Kim, S.W., Park, S.-I., Park, J., Oh, J.-H., Kim, B.C., Koh, H.J., Cho, D.-L., 2014. Sedimentological characteristics and new detrital zircon SHRIMP U–Pb ages of the Babulu Formation in the Fohorem area, Timor-Leste. *Aust. J. Earth Sci.* 61, 865–880. doi:10.1080/08120099.2014.924554
- Lemoine, M., 1959. Un exemple de tectonique chaotique: Timor essai de coordination et d’interpretation. *Rev. Geogr. Phys. Geol. Dyn.* 2, 205.
- Lewis, C., Sircombe, K., n.d. Use of U-Pb geochronology to delineate provenance of North West Shelf sediments, Australia. asi-pl.com.au.
- Ludwig, K.R., 2012. User ’s Manual for A Geochronological Toolkit for Microsoft Excel. *Berkeley Geochronol. Cent. Spec. Publ.* 1–75.
- Major, J., 2011. Evolution and Emergence of the Hinterland in the Active Banda Arc-Continent Collision: Insights From the Metamorphic Rocks and Coral Terraces of Kisar, Indonesia. *All Theses Diss.*
- Major, J., Harris, R., Chiang, H.-W., Cox, N., Shen, C.-C., Nelson, S.T., Prasetyadi, C., Rianto, A., 2013. Quaternary hinterland evolution of the active Banda Arc: Surface uplift and neotectonic deformation recorded by coral terraces at Kisar, Indonesia. *J. Asian Earth Sci.* 73, 149–161. doi:10.1016/j.jseas.2013.04.023

- McCartain, E., Backhouse, J., 2006. Gondwana-related Late Permian palynoflora, foraminifers and lithofacies from the Wailuli Valley, Timor Leste. *Neues Jahrb. für* ....
- McKee, E.H., 1970. Fish Creek Mountains Tuff and volcanic center, Lander County, Nevada. Prof. Pap.
- Metcalfe, I., 2013. Gondwana dispersion and Asian accretion: Tectonic and palaeogeographic evolution of eastern Tethys. *J. Asian Earth Sci.* 66, 1–33. doi:10.1016/j.jseas.2012.12.020
- Patchett, P.J., 1983. Importance of the Lu-Hf isotopic system in studies of planetary chronology and chemical evolution. *Geochim. Cosmochim. Acta* 47, 81–91. doi:10.1016/0016-7037(83)90092-3
- Patchett, P.J., Tatsumoto, M., 1980. Hafnium isotope variations in oceanic basalts. *Geophys. Res. Lett.* 7, 1077–1080. doi:10.1029/GL007i012p01077
- Prasetyadi, C., Harris, R., 1996. Hinterland structure of the active Banda arc-continent collision, Indonesia: constraints from the Aileu complex of East Timor. *Proc. 25th Conv. Indones. Assoc. Geol.*
- Roever, W.P. d., 1955. Genesis of jadeite by low-grade metamorphism. *Am. J. Sci.* 253, 283–298. doi:10.2475/ajs.253.5.283
- Rothpletz, A., 1891. The Permian, Triassic, and Jurassic Formations in the East-Indian Archipelago (Timor and Rotti). *Am. Nat.*
- Vermeesch, P., 2012. On the visualisation of detrital age distributions. *Chem. Geol.* 312–313, 190–194. doi:10.1016/j.chemgeo.2012.04.021
- Wanner, J., 1956. Zur stratigraphie von Portugiesisch Timor. *Zeitschrift der Dtsch. Geol. Gesellschaft* 108, 109–140.
- Wanner, J., 1913. Geologie von Westtimor. *Geol. Rundschau* 4, 136–150. doi:10.1007/BF01801936

Figure 1

Figure 3

Figure 4

Figure 5

Figure 6 TL23b

Figure 8

Figure 9

B.

Figure S1