Large Reductions in Urban Black Carbon Concentrations in the United States between 1965 and 2000

Thomas W. Kirchstetter^{1,2}, Chelsea V. Preble², Odelle L. Hadley^{1,a}, Tami C. Bond³, Joshua S. Apte⁴

¹Environmental Energy Technologies, Lawrence Berkeley National Laboratory, California

²Department of Civil and Environmental Engineering, University of California, Berkeley

³Department of Civil and Environmental Engineering, University of Illinois at Urbana-Champaign

⁴Department of Civil, Architectural and Environmental Engineering, University of Texas at Austin

^aCurrent address: Olympic Region Clean Air Agency, Olympia, Washington

ABSTRACT

Long-term pollutant concentration trends can be useful for evaluating air quality effects of emission controls and historical transitions in energy sources. We employed archival records of coefficient of haze (COH), a now-retired measure of light-absorbing particulate matter, to reconstruct historical black carbon (BC) concentrations at urban locations in the United States (U.S.). The following relationship between COH and BC was determined by reinstating into service COH monitors beside aethalometers for two years in Vallejo and one year in San Jose, California: $BC (\mu g m^{-3}) = 6.7COH + 0.1$, $R^2 = 0.9$. Estimated BC concentrations in ten states stretching from the East to West Coast decreased markedly between 1965 and 1980: 5-fold in Illinois, Ohio, and Virginia, 4-fold in Missouri, and 2.5-fold in Pennsylvania. Over the period from the mid-1960s to the early 2000s, annual average BC concentrations in New Jersey and California decreased from 13 to 2 $\mu g m^{-3}$ and 4 to 1 $\mu g m^{-3}$, respectively, despite concurrent increases in fossil fuel consumption from 1.6 to 2.1 exajoules ($EJ = 10^{18} J$) in New Jersey and 4.2 to 6.4 EJ in California. New Jersey's greater reliance on BC-producing heavy fuel oils and coal in the 1960s and early 1970s and subsequent transition to cleaner fuels explains why the decrease was larger in New Jersey than California. Patterns in seasonal and weekly BC concentrations and energy consumption trends together indicate that reducing wintertime emissions – namely substituting natural gas and electricity for heavy fuel oil in the residential sector – and decreasing emissions from diesel vehicles contributed to lower ambient BC concentrations. Over the period of study, declining concentrations of BC, a potent and short-lived climate warming pollutant, contrast increasing fossil fuel carbon dioxide (CO_2) emissions in the U.S. Declining BC emissions may have had the benefit of mitigating some atmospheric warming driven by increased CO_2 emissions with complementary health benefits.

KEY WORDS: black carbon; concentration trends; coefficient of haze; elemental carbon; air pollution; energy consumption; carbon dioxide emissions

INTRODUCTION

Background. Black carbon (BC) is an air pollutant produced during fossil and biomass fuel combustion. It is a major component of diesel engine exhaust particulate matter (PM), which is classified as a toxic air contaminant and linked to significant health impacts including cancer

(IARC, 2012). It also constitutes a significant portion of the PM emitted during biomass fuel combustion, a leading factor in premature mortality worldwide (Lim et al., 2012). In addition to adversely impacting public health, BC is a strong absorber of sunlight and contributes to climate change, alters regional precipitation patterns, and enhances ice melting in the Arctic and Himalayas (Hansen and Nazarenko, 2004; Menon et al., 2002). Recent assessments have found that the positive radiative forcing of BC emissions is comparable to or larger than that of other pollutants except carbon dioxide (CO₂) (Bond et al., 2013). Since BC has a short atmospheric lifetime, controlling PM sources with high BC mass fraction has been proposed as a means of combating near-term global warming in the face of increasing CO₂ emissions, with the co-benefit of improving air quality (Grieshop et al., 2009; Jacobson 2002).

This study presents a historical record of ambient BC concentrations in the United States (U.S.) between the years 1965 and 2000. Such records are useful for illustrating the air quality benefits associated with emission controls and transitions to cleaner energy sources. For example, a retrospective examination of ambient lead concentrations shows the efficacy of unleaded gasoline use (EPA, 2014b; Needleman, 2000). Likewise, long-term carbon monoxide and sulfur dioxide trends demonstrate how automobile catalytic converters, reducing sulfur emissions from power plants, and reducing the sulfur content of fuels improved ambient air quality (EPA, 1998; [Hand et al., 2012](#)). Pollution trends can also support epidemiological studies, emission inventory development, and air quality modeling.

Whereas the abovementioned criteria air pollutants have been monitored for many decades in the U.S., similar long records of BC concentrations do not exist. Historical BC concentrations must be estimated from other PM air pollution measurements. Murphy et al. (2011) examined elemental carbon (EC) concentration trends measured at remote locations across the U.S. by the Interagency Monitoring of Protected Visual Environments (IMPROVE) network of aerosol samplers. Annual and wintertime average EC decreased by 30% and 50%, respectively, between 1990 and 2004. Bahadur et al. (2011) interpreted California IMPROVE EC as BC and indicated a 50% decrease over the 20-year period beginning 1998.

In this study, BC concentrations are estimated using archived records of coefficient of haze (COH). Whereas IMPROVE samplers are remotely located mostly in national parks, COH monitors were primarily operated in urban regulatory monitoring networks. The COH record begins in the early 1960s, which affords an opportunity to look back several decades, as well as over different time scales, to provide some insight about dominant BC sources and other factors affecting concentration trends. This study extends an earlier study of BC concentration trends in the San Francisco Bay Area (Kirchstetter et al., 2008) to other urban air basins in California and nine additional states stretching from the East Coast to the West Coast of the U.S.

Coefficient of Haze. COH was an early air pollution particulate matter measurement method in the U.S. (Hemeon et al., 1953). Air quality agencies in many states deployed COH monitors in urban locations beginning in the mid-1960s. COH has been related to contemporary measures of atmospheric PM, including total suspended particles and coarse and fine particle mass concentrations, but the technique for measuring COH closely resembles that now used for measuring BC rather than PM, [which is sensitive to light-absorbing material and insensitive to nonabsorbing light-scattering material](#) (Wolff et al., 1983). As newer, more specific methods for

measuring PM were adopted by regulatory agencies, COH instruments were retired. For example, California operated COH monitors at more than 50 locations in 1980, but that number had dropped to 15 by 2003. Only one monitor remained in use in California by 2005.

The COH instrument, like the aethalometer used for measuring BC (Hansen et al., 1984), is a filter-based aerosol absorption photometer. In both instruments (shown in Figure S1 of the Supporting Information), air is continuously drawn through a white filter and the intensity of light transmitted through the filter is periodically measured. Transmission decreases as light-absorbing particles are collected on the filter. COH and BC are analogously calculated with equations 1 and 2, respectively:

$$\text{COH}(1000\text{ft}^{-1}) = \log(1/T) \frac{A_{\text{COH}}}{Q_{\text{COH}}\Delta t} \quad (1)$$

$$\text{BC}(\mu\text{g m}^{-3}) = \frac{1}{\sigma} \ln(1/T) \frac{A_{\text{aeth}}}{Q_{\text{aeth}}\Delta t} \quad (2)$$

T is transmission calculated from successive light intensity measurements over the period Δt (s), A is the area of filter used to collect particles (m^2), and Q is the volumetric air sampling rate ($\text{m}^3 \text{s}^{-1}$). In equation 2, σ is an attenuation coefficient ($\text{m}^2 \text{g}^{-1}$) relating particle light attenuation, $\ln(1/T)$, to BC mass. The aethalometer reports BC in units of mass concentration. COH is defined as the aerosol amount that produces an optical density, $\log(1/T)$, of 0.01 and concentrations are reported in terms of COH per 1000 linear feet (305 m) of sampled air.

The COH monitor uses a broadly emitting incandescent lamp to illuminate the particle sample and a photocell to measure the intensity of transmitted light, whereas the contemporary aethalometer uses a discrete light-emitting diode focused at 880 nm and a photodiode sensor. Light scattering within the aethalometer's quartz fiber filter (Pallflex type Q250F) amplifies its response to light-absorbing particles and diminishes its response to light-scattering particles (Hansen et al., 1984; Arnott et al., 2005). Similarly, Hemeon et al. (1953) reported that the COH filter paper (Whatman No. 4) amplified the instrument's response to smoke particles.

METHODS

Sampling. To support constructing a historical record of ambient BC, COH monitors and aethalometers were collocated and operated at two locations. Two COH monitors were restored to working condition with original lamps, photocells, and filter paper. Each unit was upgraded with a mass flow controller (see Figure S1) and connected to a computer for data acquisition and tape advance control. Prior to field deployment, both COH monitors were evaluated in the laboratory using soot generated with a methane-air flame (Kirchstetter and Novakov, 2007). Measured COH concentrations from the two monitors were in good agreement: linearly correlated with a slope of 1.03, zero intercept, and correlation coefficient $R^2 = 0.96$ (Figure S2). These tests established that time-resolved COH concentrations, like time-resolved BC concentrations measured using the aethalometer, are affected by a “filter loading” artifact (Weingartner et al., 2003; Arnott et al., 2005; Kirchstetter and Novakov 2007; Cohen et al., 2010), wherein COH concentrations erroneously decrease as the instrument's particle collection

1 filter becomes increasingly loaded with BC (Figure S3). These tests also established that
2 averaging COH concentrations over one or more sampling cycles largely eliminates the influence
3 of this artifact (Figure S3). Therefore, in this study, we examine trends in concentrations
4 averaged over days, months, and years rather than hours.

5
6 After laboratory evaluation, the COH instruments were collocated with aethalometers in the air-
7 monitoring network of the Bay Area Air Quality Management District (BAAQMD) where COH
8 was historically measured in Vallejo and San Jose. Vallejo's population is 120,000 (in year
9 2010) and the monitoring station sits in a mixed commercial and residential neighborhood one
10 mile east of downtown and 0.5 miles west of a major freeway. San Jose is the largest city in the
11 San Francisco Bay Area with a population of 950,000 (in year 2010). The air monitoring station
12 is located in a commercial and residential part of downtown San Jose encircled by major
13 freeways. The aethalometers and COH monitors were operated for 14 months (March 2010
14 through April 2011) in San Jose and 25 months (March 2010 through March 2012) in Vallejo.
15 One aethalometer had recently been serviced and was active in the monitoring network. The
16 other had been inactive and was serviced by the manufacturer at the start of this study.

17
18 Consistent with historical practice, the COH monitors in this study were installed with 0.25 inch
19 Teflon sample tubing, without an inlet cyclone, and with an airflow rate of 6.25 L min^{-1} .
20 Consistent with contemporary sampling practice, each aethalometer sampled ambient air through
21 a $2.5 \mu\text{m}$ diameter cut-point inlet cyclone (BGI, model SCC1.829) and electrically conductive
22 tubing (0.375 inch aluminum in San Jose and 0.625 inch stainless steel in Vallejo). The
23 aethalometers were operated at an airflow rate of 5 L min^{-1} .

24
25 The COH monitors were initially programmed to advance their filter tapes once every two hours,
26 again consistent with historical practice. However, current ambient BC concentrations were too
27 low to provide sufficient signal after only two hours of sampling. To increase the amount of PM
28 collected on the COH filter and thereby increase the measurement sensitivity, the instruments
29 were reprogrammed for tape advances once every six hours. With each advance, a six-hour
30 average COH concentration was recorded. In effect, the six-hour sampling interval provided the
31 same filter loading that a two-hour interval would have accumulated when COH concentrations
32 were three times higher than they are now. The aethalometer was programmed to advance its
33 filter when measured attenuation reached the factory default value of 75, which occurred about
34 four times every five days.

35
36 **Data Sources.** The California Air Resources Board (CARB) and the New Jersey Department of
37 Environmental Protection (NJDEP) provided COH data collected in California and New Jersey
38 from the early 1960s to 2000. COH monitors were located predominantly in urban areas and the
39 location and number of active monitors varied over time (Figure S4). The Environmental
40 Protection Agency's (EPA) Air Quality System (EPA, 2012) supplied COH data for several
41 other states. The COH records of some states were not preserved as well as those kept by
42 California and New Jersey. As such, the data obtained from the EPA were comparatively limited
43 both temporally and spatially. The following presentation includes the data-rich states of
44 California and New Jersey and eight other states with COH records covering at least ten years:
45 Colorado, Illinois, New York, Missouri, Ohio, Pennsylvania, Virginia, and Washington.

1 Since BC is a primary air pollutant, urban emissions and local concentrations are tightly linked
2 (Apte et al., 2012). Therefore, long-term trends in annual average BC concentrations and weekly
3 cycles in daily average BC concentrations are discussed in relation to energy consumption and
4 CO₂ emissions trends as well as temporal patterns in emission sources. Energy consumption and
5 CO₂ emissions data are from the U.S. Energy Information Administration (EIA, 2012; EIA,
6 2016).

7
8 Meteorological variability over the course of a year moderates air pollution levels. Thus,
9 seasonal trends in BC concentrations are discussed in relation to the local meteorology, namely
10 wind speed (u) and mixing height (H), which govern pollutant dispersion. Meteorological data
11 were acquired with minimal processing as described by Apte et al. (2012) for the 2007-2009
12 period from NASA's Modern-Era Retrospective Analysis for Research and Applications
13 (MERRA) (Rienecker et al., 2011). We assume the meteorological trends are the same for the
14 whole period of COH record. MERRA uses the Goddard Earth Observing System atmospheric
15 model to assimilate meteorological observations into a global data set with a spatial and temporal
16 resolution of 0.5° longitude × 0.76° latitude × 1 hr. At mid-latitudes in the U.S., this corresponds
17 to a spatial resolution of approximately 50 × 75 km. Data for cities was extracted by matching
18 latitude and longitude coordinates to the nearest MERRA grid point. The harmonic means of u,
19 H, and uH were calculated. The product uH is the normalized dilution rate (m² s⁻¹) and provides
20 a measure of atmospheric dilution for an urban area (Apte et al., 2012; Marshall et al. 2005).

21 22 RESULTS AND DISCUSSION

23
24 **Relationship between COH and BC.** Figure 1 shows the relationship between daily averaged
25 concentrations of BC and COH at the Vallejo and San Jose sites. Lowest concentrations were
26 measured in May, June, and July and the highest in December and January. A linear regression
27 of the data from both locations yields the equation that we use in this study to calculate BC
28 concentrations:

$$29 \quad 30 \quad BC (\mu\text{g m}^{-3}) = 6.7COH + 0.1 \quad (3)$$

31
32 The high correlation coefficient ($R^2 = 0.9$) indicates that COH is a good proxy for BC. The linear
33 relationship suggests that the COH-BC relationship is not seasonally dependent. Moreover, since
34 the BC fraction of ambient PM varies throughout the year as a result of meteorological
35 influences on secondary PM, the relationship between COH and BC does not appear to be highly
36 sensitive to ambient PM composition. The intercept of the linear equation is small compared to
37 past BC concentrations (see Figure 2), which means that historical BC and COH are
38 approximately proportional. Analysis of monthly averaged concentrations of BC and COH yields
39 essentially the same result (Figure S5), indicating that the same relationship can effectively be
40 used to convert daily and monthly averaged values of COH to BC.

41
42 Others have previously examined the relationship between COH and EC. While EC and BC are
43 not the same, they are often well correlated (Venkatachari et al., 2006), such that examining the
44 relationship between EC and COH may be instructive. The BAAQMD derived EC ($\mu\text{g m}^{-3}$) =
45 $7.5COH$ from measurements during the period 1991-2004 in San Francisco and San Jose,
46 California (BAAQMD, 2007). A decade prior in the summer of 1981 in Detroit, Michigan, Wolff

et al. (1983) measured $EC (\mu g m^{-3}) = 8.9COH - 0.63$ with $R^2 = 0.9$ for EC concentrations up to $10 \mu g m^{-3}$.

Together, the data collected in this work (Figure 1) and the findings of the studies cited above demonstrate that COH correlates with light-absorbing PM, indicate that the relationship does not change dramatically with location or time, and thus support using COH to estimate past BC concentrations. While uncertainty remains in the use of a constant COH-BC relationship at all sites, over all years, and at concentrations exceeding those in Figure 1 and the above-cited studies, it facilitates a reasonable analysis of the COH database.

Figure 1. Scatter plot of daily averaged BC and COH concentrations measured in San Jose (March 2010 through April 2011) and Vallejo (March 2010 through March 2012). The linear regression applies to the entire data set.

Black Carbon Concentration Trends. Figure 2 shows the trends in annual average urban BC concentrations for the ten states considered in this study, in each case computed as the arithmetic mean of BC concentrations at all monitoring sites in the state. At the start of this record, BC concentrations were highest in New Jersey, New York, and Illinois and lowest in California. Over the 15 year period from the mid-1960s to 1980, BC concentrations decreased by 5-fold in Illinois, Ohio, and Virginia, 4-fold in Missouri, and 2.5-fold in Pennsylvania. From the mid-1960s to 2000, annual average BC concentrations in the urban regions of New Jersey and

1 California decreased from 13 to 2 $\mu\text{g m}^{-3}$ and 4 to 1 $\mu\text{g m}^{-3}$, respectively. Figure S6 shows BC
2 concentrations separately for different air-basins in California and New Jersey.

3
4 New Jersey's greater reliance on BC-producing fuels in the 1960s is most likely why the BC
5 concentration reduction was more dramatic in New Jersey than California. Clean burning natural
6 gas was the single largest energy provider in California from 1960 to 2000, whereas energy
7 production in New Jersey relied more on residual and distillate fuel oils than natural gas until
8 1974 (Figure S7) and coal was consumed in significant amounts by New Jersey's industrial
9 sector up until about 1970 (Figure S8).

12
13
14 **Figure 2.** Annual average urban BC concentrations in ten states. California's BC concentration trend
15 includes the densely populated South Coast air basin only from 1969 to 1976 because that region's COH
16 data were unavailable outside of this period. BC concentrations in the South Coast air basin were higher
17 than in other California air basins (Figure S6), but increased statewide average BC concentrations only by
18 a small amount.

19
20
21 Figures 3 and S9 show that BC concentrations decreased much more in winter than summer in
22 many East Coast states. The transition to lower wintertime BC concentrations was especially
23 quick for Pennsylvania and Illinois, where the seasonal BC cycle essentially disappeared (i.e.,
24 the curve flattened out) between the 1960s and 1970s. Pennsylvania's wintertime BC
25 concentrations decreased from about 10 to 5 $\mu\text{g m}^{-3}$, whereas its summertime concentrations did
26 not decrease much at all. An annual cycle is evident in New York's data in the 1960s, but much
27 less so in the 1990s. Similarly, the seasonality evident in New Jersey's BC concentrations in the

1 1970s steadily diminished to the point where concentrations were relatively uniform throughout
2 the year in the 1990s and 2000s.

3
4 Seasonal variations in BC concentrations reflect the combined influence of emissions and
5 meteorology. Whereas the latter probably has not changed when averaged over many years, a
6 transition to cleaner fuels used for wintertime heating has occurred. For example, beginning in
7 the early 1970s, BC-producing distillate fuel oil in New Jersey's residential sector (EDF, 2009)
8 was substituted with cleaner burning natural gas and electricity (Figure S10). Therefore, reduced
9 wintertime emissions associated with heating likely caused the seasonality in BC concentrations
10 to disappear and drove the major reductions in annual BC concentrations early in the record
11 (Figure 2), as previously discussed.

Figure 3. Monthly average BC concentrations by decade in ten states.

Figure 4. Weekly cycle of BC concentration by decade. The vertical axis reports the ratio of average BC concentrations each day of the week to the average Sunday concentration.

1 In contrast, California's seasonal cycle persisted throughout the entire period of record: BC
2 concentrations are consistently 2 to 3 times higher in the winter than in the summer (Figures 3
3 and S9). This seasonality appears to be driven by meteorology rather than emissions. Whereas
4 California's residential energy sector relied on cleaner energy sources over the entire study
5 period (Figure S10), atmospheric dispersion of pollutants in California is 2 to 3 times stronger in
6 summer than in winter (Figures S11 and S12; Glen et al., 1996). Thus, BC concentrations in
7 urban California are naturally much higher in winter. The seasonal variation in atmospheric
8 dilution in New Jersey and other eastern cities is weaker and opposite that of California,
9 consistent with the absence of a strong seasonal cycle in BC after the wintertime heating
10 emissions were largely reduced. The study by Glen et al. (1996) similarly attributed wintertime
11 maxima and summertime minima in CO concentrations in San Francisco, California to lower
12 wind speeds and mixing heights in the winter, and a weak or nonexistent seasonality in
13 concentrations across several East Coast cities to the lack of a seasonal cycle in pollutant
14 dispersion (Figure S13).

15
16 In addition to the seasonal cycle, all states exhibit a weekly pattern of lower BC concentrations
17 on weekends compared to weekdays and lowest BC concentrations on Sundays (Figure 4). For
18 example, BC concentrations in the 1970s in Missouri and Colorado were approximately 1.5
19 times higher during the week than they were on Sunday. This weekly cycle strengthened in most
20 cases as baseline BC concentrations decreased over time. The persistent weekly pattern points to
21 a source of BC emissions with the same weekly pattern that has been persistently dominant
22 throughout the study period. Emission inventories include diesel trucks as a major BC source
23 (e.g., Bond et al., 2004) and, as previously reported, diesel truck activity is appreciably lower on
24 weekends. For example, on-road diesel truck volume on Sundays is 25% of the weekday average
25 (Dreher and Harley, 1998) and weekend off-road diesel vehicle activity is less than 10% of
26 weekday activity (Coe et al., 2003).

27
28 A number of studies have attributed declining ambient BC (or EC) concentrations to diesel truck
29 emission regulations (Christoforou et al., 2000; Murphy et al., 2011; Bahadur et al., 2011),
30 notably smoke emission limits in 1970 and PM emission standards beginning in 1998. According
31 to Sawyer and Johnson (1995) and Lloyd and Cackette (2001), PM emission factors from new
32 heavy-duty highway diesel engines decreased by a factor of 10 from 1970 to 1990. Changes in
33 in-use emission rates are more pertinent than new truck emission standards when relating to
34 ambient concentration trends. The longest span of measurement of in-use heavy-duty diesel truck
35 fleet PM emissions at one location is 1975 to 1999 in Pennsylvania's Tuscarora Tunnel. Over
36 that 24-year period, the fleet average diesel PM emission factor decreased by a factor of 6
37 (Gertler et al. 2002). This is double the factor of 3 increase in taxable (i.e., highway) diesel fuel
38 consumption in California and triple the factor of 2 increase in New Jersey's distillate
39 consumption (not exclusively used by highway trucks) between 1975 and 2000 (Figure S14).
40 More recently, between 1997 and 2006, the average BC emission factor for in-use heavy-duty
41 diesel trucks in California's Caldecott Tunnel decreased by 61% (Ban-Weiss et al., 2008), about
42 twice as much as consumption of California taxable diesel increased (33%) over the same period
43 (Figure S14). These data suggest that the decrease in on-road diesel truck BC emission rates
44 outpaced increased fuel consumption and thus contributed to the trend of decreasing BC
45 concentrations shown in Figure 2.

1 The recent study of McDonald et al., (2015) highlights the importance of off-road engines when
2 assessing long-term ambient BC trends. Off-road diesel engines were not subjected to EPA
3 emission standards until 1996 and have higher emission rates than highway diesel vehicles. Data
4 available for California indicate that off-road diesel fuel consumption declined by 40% from
5 1970 to 2000 (Figure S14). Thus, BC emissions from off-road diesel engines likely also
6 contributed to the ambient BC concentration decline over the period of record in California, due
7 to decreasing consumption rather than reduced emission factors.
8

9 **Relationship between BC and CO₂.** Over the period from the mid-1960s to the early 2000s,
10 when annual average BC concentrations in New Jersey and California decreased from 13 to 2 μg
11 m^{-3} and 4 to 1 $\mu\text{g} \text{ m}^{-3}$, respectively, fossil fuel consumption increased from 4.2 to 6.4 EJ in
12 California and 1.6 to 2.1 EJ in New Jersey (Figure S15). Figure 5 shows the rise in energy
13 consumption trends in terms of fossil fuel greenhouse gas CO₂ emissions, which strongly
14 contrast with declining BC concentrations. Since BC is a potent but short-lived climate warming
15 pollutant, reducing BC emissions may have had the benefit of mitigating some atmospheric
16 warming driven by increased CO₂ emissions. For example, Bahadur et al. (2011) reported a
17 negative radiative forcing (i.e., climate cooling) over California due to a reduction in BC
18 concentrations. Given that BC is a major component of diesel PM, which is a toxic air pollutant
19 and a human carcinogen, these reductions may have had complementary health benefits
20 (Anenberg et al., 2012).
21
22
23
24
25
26
27
28
29

Figure 5. Major reductions in BC concentrations over time contrast with increases in CO₂ emissions in both California (top) and New Jersey (bottom).

ACKNOWLEDGEMENTS

This work was supported by the National Science Foundation (award number 0852775) and the California Air Resources Board (contract 08-323). The statements and conclusions herein are those of the authors and do not necessarily reflect the views of the project sponsors. We thank the Bay Area and Lake County Air Quality Management Districts and the California Air Resources Board for supplying COH monitors. We also thank the Bay Area Air Quality Management District for supporting us in our efforts to operate COH monitors at their facilities in Vallejo and San Jose.

REFERENCES

- Allen, GA; Lawrence, J; Koutrakis, P (1999) Field validation of a semi-continuous method for aerosol black carbon (aethalometer) and temporal patterns of summertime hourly black carbon measurements in southwestern PA, *Atmos. Environ.* 33, 817–823.
- Anenberg, SC, Schwartz, J, Shindell, D, Amann, M, Faluvegi, G, Klimont, Z, Janssens-Maenhout, G, Pozzoli, L, Van Dingenen, R, Vignati, E, Emberson, L, Muller, NZ, West, JJ, Williams, M, Demkine, V, Hicks, WK, Kuysenstierna, J, Raes, F, Ramanathan, V (2012) Global air quality and health co-benefits of mitigating near-term climate change through methane and black carbon emission controls, *Environ Health Perspect.*, 120, 831–839, doi: 10.1289/ehp.1104301.
- Arnott, WP; Hamasha, K; Moosmuller, H; Sheridan, PJ; Ogren, JA (2005) Towards aerosol light absorption measurements with a 7-wavelength aethalometer: evaluation with a photoacoustic instrument and a 3-wavelength nephelometer, *Aerosol Sci & Technol.*, 39, 17-29.
- Apte, JS; Bombrun, E; Marshall, JD; Nazaroff, WW (2012) Global intraurban intake fractions for primary air pollutants from vehicles and other distributed sources. *Environmental Science & Technology* 46, 3415-3423.
- BAAQMD (2007) Bay Area data on relationship of EC and COH, Bay Area Air Quality Management District, Office memorandum, obtained by personal communication with David Fairley.
- Bahadur, R; Feng, Y; Russell, LM; Ramanathan, V (2011) Impact of California's air pollution laws on black carbon and their implications for direct radiative forcing, *Atmos. Environ.*, 45, 1162-1167.
- Ban-Weiss, GA; McLaughlin, JP; Harley, RA; Lunden, MM; Kirchstetter, TW; Kean, AJ; Strawa, AW; Stevenson, ED; Kendall, GR (2008) Long-term changes in emissions of nitrogen oxides and particulate matter from on-road gasoline and diesel vehicles, *Atmos. Environ.*, 42, 220-232.
- Bond, TC; Doherty, SJ; Fahey, DW; Forster, PM; Berntsen, T; DeAngelo, BJ; Flanner, MG; Ghan, S; Kärcher, B; Koch, D; Kinne, S; Kondo, Y; Quinn, PK; Sarofim, MC; Schultz, MG; Schulz, M; Venkataraman, C; Zhang, H; Zhang, S; Bellouin, N; Guttikunda, SK; Hopke, PK; Jacobson, MZ; Kaiser, JW; Klimont, Z; Lohmann, U; Schwarz, JP; Shindell, D; Storelvmo, T;

Warren, SG; Zender, CS (2013) Bounding the role of black carbon in the climate system: A scientific assessment. *J. Geophys. Res. Atmos.*, 118, 5380–5552.

Bond, TC; Street, DG; Yarber, KF; Nelson, SM; Woo, JH; Klimont, Z (2004) A technology-based global inventory of black and organic carbon emissions from combustion, *J. Geophys. Res.*, 109, D14203, doi: 10/1029/2003JD003697.

Christoforou, CS; Salmon, LG; Hannigan, MP; Solomon, PA; Cass, GR (2000) Trends in fine particle concentration and chemical composition in Southern California, *J. Air & Waste Manage. Assoc.*, 50, 43-53.

Coe, DL; Gorin, CA; Chinkin, LR; Reid, SR (2003) Observations of Weekday–Weekend Activity Patterns for Area Sources in the Los Angeles Area, Presented at the US Environmental Protection Agency 12th International Emission Inventory Conference – “Emission Inventories – Applying New Technologies”, Sonoma Technology, Inc. April 2003.

Coen, MC; Weingartner, E; Apituley, A; Ceburnis, D; Fierz-Schmidhauser, R; Flentje, H; Henzing, JS; Jennings, SG; Moerman, M; Petzold, A; Schmid, O; Baltensperger, U (2010) Minimizing light absorption measurement artifacts of the Aethalometer: evaluation of five correction algorithms, *Atmos. Meas. Tech.*, 3, 457-474, doi:10.5194/amt-3-457-2010.

Dieselnet (2012) Emission standards, United States, Nonroad diesel engines, <http://www.dieselnet.com/standards/us/nonroad.php>

EDF, Environmental Defense Fund (2009) Bottom of the Barrel, available online https://www.edf.org/sites/default/files/10085_EDF_Heating_Oil_Report.pdf

EIA, Energy Information Administration (2012) State Energy Data: Consumption, 1960-2010, California. Available online at <http://www.eia.gov/beta/state/seds/seds-data-complete.cfm?sid=CA#Consumption>.

EIA, Energy Information Administration (2016) State Carbon Dioxide Emissions. Available online at http://www.eia.gov/environment/emissions/state/state_emissions.cfm

EPA (1998) National air quality and emission trends report 1996, U.S. Environmental Protection Agency, 454/R-97-013; <http://www3.epa.gov/airtrends/aqtrnd96/chapter2.pdf>

EPA (2002) Health assessment document for diesel engine exhaust, National Center for Environmental Assessment, U.S. Environmental Protection Agency, EPA/600/8-90/057F; <http://nepis.epa.gov/Exe/ZyPDF.cgi/300055PV.PDF?Dockkey=300055PV.PDF>

EPA (2012) Environmental Protection Agency, Air Quality System [internet database] available at <http://www.epa.gov/ttn/airs/aqsdatamart>. Accessed March 01, 2012.

EPA (2014a) Environmental Protection Agency, Nonroad compression-ignition engines, Exhaust Emission Standards: <http://www.epa.gov/otaq/standards/nonroad/nonroadci.htm>

1
2 EPA (2014b) Environmental Protection Agency, National trends in lead levels:
3 <https://www3.epa.gov/airtrends/lead.html>
4
5 Glen, WG; Zelenka, MP; Graham, RC (1996) Relating meteorological variables and trends in
6 motor vehicle emissions to monthly urban carbon monoxide concentrations, *Atmos. Environ.*, 30,
7 4225-4232.
8
9 Grieshop, AP; Reynolds, C; Kandlikar M; Dowlatbadi, H (2009) A black carbon mitigation
10 wedge, *Nature Geoscience* 2, 533 – 534, doi:10.1038/ngeo595
11
12 Hand, JL; Schichtel, BA; Malm, WC; Pitchford, ML (2012) Particulate sulfate ion concentration
13 and SO₂ emission trends in the United States from the early 1990s through 2010, *Atmos. Chem.*
14 *Phys.*, 12, 10353-10365, doi:10.5194/acp-12-10353-2012.
15
16 Hansen, J; Nazarenko, L (2004) Soot climate forcing via snow and ice albedos. *Proc. Natl Acad.*
17 *Sci.*, 101, 423–428.
18
19 Hansen, ADA; Rosen, H; Novakov, T (1984) The aethalometer – an instrument for the real-time
20 measurement of optical absorption by aerosol particles, *Sci. Tot. Environ.*, 36, 191-196.
21
22 Hemeon, WCL; Haines, GF; Ide, HM (1953) Determination of haze and smoke concentrations
23 by filter paper samples, *Air Repair*, 3, 22–28.
24
25 IARC (2012) IARC: DIESEL ENGINE EXHAUST CARCINOGENIC, Press release No. 213;
26 https://www.iarc.fr/en/media-centre/pr/2012/pdfs/pr213_E.pdf
27
28 Jacobson, MZ (2002) Control of fossil-fuel particulate black carbon and organic matter, possibly
29 the most effective method of slowing global warming, *J. Geophys. Res.*, 107, D19, 4410.
30
31 Jeonge, CH; Hopke, PK; Kim, E; Lee, DW (2004) The comparison between thermal-optical
32 transmittance elemental carbon and Aethalometer black carbon measured at multiple monitoring
33 sites, *Atmos. Environ.*, 38, 5193-5204.
34
35 Kirchstetter, TW; Aguiar, J; Tonse, S; Novakov, T; Fairley, D (2008) Black carbon
36 concentrations and diesel vehicle emission factors derived from coefficient of haze
37 measurements in California: 1967-2003. *Atmos. Environ.*, 42, 480-491.
38
39 Kirchstetter, TW; Novakov, T (2007) Controlled generation of black carbon particles from a
40 diffusion flame and applications in evaluating black carbon measurement methods, *Atmos.*
41 *Environ.*, 41, 1874-1888.
42
43 Kirchstetter, TW; Novakov, T; Hobbs PV; Magi, B (2003) Airborne measurements of
44 carbonaceous aerosols in southern Africa during the dry, biomass burning season, *J. Geophys.*
45 *Res.*, 108, 10.1029/2002JD002171.
46

- Kirchstetter, TW; Harley, RA; Kreisberg, NM; Stolzenburg, MR; Hering, SV (1999) On-road measurement of fine particle and nitrogen oxide emissions from light- and heavy-duty motor vehicles. *Atmos. Environ.*, 33, 2955-2968.
- Lim SS; Vos, T; Flaxman, AD; Danaei, G; Shibuya, K; ...Lopez, AD; Murray, CJL; Ezzati, M (2012) A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: A systematic analysis for the Global Burden of Disease Study 2010. *Lancet*, 380, 2224–2260.
- Lloyd, AC; Cackette, TA (2001) Diesel engines: Environmental impact and control, *J. Air & Waste Manage. Assoc.*, 51, 809-847.
- Marshall, JD; Teoh, SK; Nazaroff, WW (2005) Intake fraction of nonreactive vehicle emissions in US urban areas, *Atmos Environ*, 39, 1363-1371.
- McDonald, BC; Goldstein, AH; Harley, RA (2015) Long-term trends in California mobile source emissions and ambient concentrations of black carbon and organic aerosol, *Environ. Sci. Technol.*, 49, 5178–5188, doi: 10.1021/es505912b
- Menon, S; Hansen, J; Nazarenko, L; Luo, Y (2002) Climate Effects of Black Carbon Aerosols in China and India, *Science*, 297; doi: 10.1126/science.1075159
- Murphy, DM; Chow, JC; Leibensperger, EM; Malm, WC; Pitchford, M; Schichtel, BA; Watson, JG; White, WH (2011) Decreases in elemental carbon and fine particle mass in the United States, *Atmos. Chem. Phys.*, 11, 4679–4686, doi:10.5194/acp-11-4679-2011.
- Needleman, HL (2000) The removal of lead from gasoline: historical and personal reflections, *Environ. Res. Section A*, 84, 20-35.
- Novakov, T; Ramanathan, V; Hansen, J; Kirchstetter, TW; Sinton, JE; Sathaye, JA (2003) Large historical changes of fossil-fuel black carbon aerosols, *Geophys. Res. Letters*, 30, 10.1029/2002GL016345.
- Rienecker, MM, Suarez, MJ, Gelaro, R, Todling, R, Bacmeister, J, Liu, E, Bosilovich, MG, Schubert, SD, Takacs, L, Kim, G-K, Bloom, S, Chen, J, Collins, D, Conaty, A, da Silva, A, Gu, W, Joiner, J, Koster, RD, Lucchesi, R, Molod, A, Owens, T, Pawson, S, Pegion, P, Redder, CR, Reichle, R, Robertson, FR, Ruddick, AG, Sienkiewicz, M, Woollen, J (2011) MERRA: NASA's modern-era retrospective analysis for research and applications. *J Climate* 24, 3624-3648.
- Venkatachari, P; Zhou, L; Hopke, PK; Schwab, JJ; Demerjian, KL; Weimer, S; Hogrefe, O; Felton, D; Rattigan, O (2006) An intercomparison of measurement methods for carbonaceous aerosol in the ambient air in New York City, *Aerosol Sci. Technol.*, 40, 788-795, doi: 10.1080/02786820500380289.
- Watson, JG (2002) Visibility: Science and regulation – 2002 critical review, *J. Air Waste Manage.*, 52, 628-713.

1
2 Weingartner, E; Saathoff, H; Schnaiter, M; Streit, N; Bitnar, B; Baltensperger, U (2003)
3 Absorption of light by soot particles: determination of the absorption coefficient by means of
4 aethalometers, *J Aerosol Sci.*, 34, 1445–1463, doi:10.1016/S0021-8502(03)00359-8.
5
6 Wolff, GT; Stroup, CM; Stroup, DP (1983) The coefficient of haze as a measure of particulate
7 elemental carbon. *JAPCA*, 33, 746-750.
8
9 Yan, F; Winijkul, E; Jung, S; Bond, TC; Streets, DG (2011) Global emission projections of
10 particulate matter (PM): I. Exhaust emissions from on-road vehicles, *Atmos. Environ.*, 45, 4830–
11 4844, <http://dx.doi.org/10.1016/j.atmosenv.2011.06.018>.