Abstract

Despite increasing evidence of the relationship between neighborhood cohesion and depressive mood, little is known about this longitudinal association in old age. This study examined the association between perceived neighborhood cohesion and depressive mood and the stress-buffering effect of perceived neighborhood cohesion on depressive mood among older Japanese people using the 2010 (baseline) and 2012 (follow-up) Hatoyama Cohort Study datasets. We analyzed 655 participants aged 65–84 at baseline. Although perceived neighborhood cohesion at baseline was not associated with depressive mood at follow-up, high neighborhood cohesion partially offset the deleterious effect of anticipated daily stressors on depressive mood. This effect was stronger for long-term residents of the neighborhood. Interventions to strengthen neighborhood cohesion may help reduce the deleterious effect of stressors on older residents' depressive mood.

Keywords: neighborhood cohesion; depressive mood; old age; longitudinal study; Japan

1 **Introduction**

2 Previous research on the global burden of disease in 2010 identified depressive disorders
3 as a leading cause of this burden (Ferrari et al., 2013; Whiteford et al., 2013). The cost of depression
4 is as high as that of other major illnesses, such as cardiovascular diseases or AIDS (Greenberg et al.,
5 1993). The prevention of depression is an important public health issue, particularly in old age.
6 People often experience changes in factors such as social functions, social relations, and physical
7 condition at this stage of life (Müller-Spahn and Hock, 1994; Rowe and Kahn, 1997) and these
8 changes are often related to depression in old age (Emmerson et al., 1989). In fact, the rate of
9 depression increases with age (Davey et al., 2004; Luppá et al., 2012; Stordal et al., 2001).
10 Moreover, depression is a risk factor for functional decline in later life (Stuck et al., 1999).

11 Many individual-level factors increase the risk of depression. However, neighborhood
12 environments can also affect depression. A number of theories have been proposed to explain the
13 association between neighborhood characteristics and mental health. Features of neighborhoods
14 may constrain or enhance health-related behaviors, may function as stressors, and may act as a
15 buffer against stress (Cattell, 2001; Diez Roux and Mair, 2010; Murayama et al., 2012a). Mair et al.
16 (2008) reviewed the literature on neighborhood characteristics and depressive mood and classified
17 neighborhood characteristics into two dimensions: structural features (e.g., socioeconomic and
18 racial composition, stability) and social processes (e.g., social cohesion, ties among neighbors).
19 They found that social processes were associated more consistently with depressive mood than with
20 structural features. However, there has been little research on the association between social
21 processes, particularly social cohesion, and depressive mood in later life.

22 Neighborhood cohesion refers to the extent of connectedness and solidarity in a
23 neighborhood and represents resources that individuals can access via membership (Kawachi and
24 Berkman, 2000); it can be distinguished from social network interaction, social support, and social

1 exchange in that it assesses norms and expectations for behavior rather than actual ties or direct
2 interaction (Kawachi et al., 2008). Because older people are more likely to spend time in their
3 neighborhood of residence and have stronger social networks in the community than younger
4 generations (Cabinet Office, 2007), neighborhood cohesion may play an important role in their
5 mental health. Therefore, a study on the association between neighborhood cohesion and depression
6 in old age is warranted.

7 Neighborhood cohesion can also act as a buffer against sources of stress related to mental
8 health and psychological outcomes (Cattell, 2001; Diez Roux and Mair, 2010). For example, high
9 perceived neighborhood cohesion buffers the effects of stressors on daily negative affect (Robinette
10 et al., 2013) and the effect of area income deprivation on mental health (Fone et al., 2007). Takagi
11 et al. (2013) used cross-sectional data to reveal that higher area-level trust among neighbors and
12 higher social participation buffered the adverse effect of social distance (derived by taking averaged
13 differences in sociodemographic characteristics between individuals and their neighbors) on
14 depressive risk among older Japanese people. Findings on this stress-buffering effect can help to
15 identify the mechanism by which neighborhood cohesion affects people's mental health. However,
16 evidence on this effect in old age is sparse.

17 In addition to the shortcomings noted above, several important gaps in research remain to
18 be addressed. First, most studies on neighborhood cohesion and depression have been conducted in
19 Western countries. In Japan, which has a relatively collectivist society with intense group ties,
20 people feel comfortable living in a community with systems of mutual assurance and monitoring
21 among residents (Nakane, 1970; Yamagishi et al., 1998; Yamagishi and Yamagishi, 1994). Because
22 of the difference in the cultural and historical backgrounds of Japan and Western countries, it is
23 important to explore the association between neighborhood cohesion and depressive mood in Japan.
24 Data from non-Western societies highlights many conditions not observable in Western countries

1 and could offer useful insights about the mechanisms underlying these factors. Second, most
2 previous work on neighborhood cohesion and depressive mood has used cross-sectional designs that
3 failed to identify causality. Longitudinal designs are needed to determine the direction of causality.

4 To address these gaps in this study, we analyzed data from a cohort study of
5 community-dwelling older Japanese people. Our purpose was to examine the longitudinal
6 association between perceived neighborhood cohesion and depressive mood and to test whether
7 perceived neighborhood cohesion buffers the effect of anticipated daily stressors on depressive
8 mood among older Japanese people. We hypothesized that higher perceived neighborhood cohesion
9 is associated with less likelihood of depressive mood (H_1) and that perceived neighborhood
10 cohesion partially offsets the deleterious effect of anticipated daily stressors on depressive mood
11 (H_2). Additionally, as neighborhood cohesion represents resources individuals can access via
12 membership in the neighborhood, the amount of time people have spent there (i.e., age or years of
13 residence in the neighborhood) could influence the linkage of neighborhood cohesion with
14 depressive mood and its stress-buffering effect. However, there is little evidence concerning this.
15 Therefore, we also explored whether the direct and stress-buffering effects of perceived
16 neighborhood cohesion on depressive mood varied by age or years of residence in the neighborhood.
17 We hypothesized that these effects are stronger in those who are older or have lived longer in the
18 neighborhood compared with those who are younger or have lived for a shorter time in the
19 neighborhood (H_3).

22 **Methods**

23 *Study population*

24 The Hatoyama Cohort Study consisted of community-dwelling individuals aged 65 years

1 or older, living in the town of Hatoyama in Saitama, Japan. Hatoyama is a suburban area located 50
2 kilometers northwest of central Tokyo. In June 2010, it had a population of 15,424, and the
3 proportion of people aged 65 years or older was 26.1% (vs. 23.1% nationally). To recruit the study
4 participants, we used stratified sampling of four groups classified by age (65–74 and 75–84 years)
5 and residential area of the town (the traditional area and the newly developed area). People with
6 long-term care certification (levels 1–5) and those admitted to hospitals or residing in nursing
7 homes were excluded. For the group of residents aged 65–74 living in the newly developed area, we
8 used a random sampling strategy because the number of target residents was larger than in the other
9 three groups. For the other three groups, we used a complete census (i.e., we approached all
10 residents aged 65–84 in the traditional area and aged 75–84 in the newly developed area). The
11 participation rates in these four stratified groups were 23.1% (aged 65–74 in the traditional area),
12 18.3% (aged 75–84 in the traditional area), 35.0% (aged 65–74 in the newly developed area), and
13 27.1% (aged 75–84 in the newly developed area). In addition to the sampling recruitment, we
14 recruited study participants using the Hatoyama town bulletin, which was distributed to all
15 households in the town, to permit broader recruitment. Further information on sampling and the
16 participants is included in a previous paper (Murayama et al., 2012b).

17 A total of 742 people participated in the baseline survey in 2010 (722 obtained by the
18 random sampling recruitment, and 20 by the bulletin recruitment). Among the study participants (n
19 = 742), 57.7% were male, the mean age was 71.9 ± 5.2 years, and 65.1% lived in the newly
20 developed area. Conversely, among individuals in the general population of the town aged 65–84
21 (i.e., the target population), 51.2% were male, the mean age was 71.9 ± 5.4 years, and 63.5% lived
22 in the newly developed area. In 2012, a follow-up survey of the participants in the baseline survey
23 was conducted. Of 742 participants, eight had died between the baseline and follow-up surveys, 27
24 had declined further inclusion in this cohort study, and 26 could not participate in the 2012

1 follow-up survey for various reasons (e.g., health-related, schedule conflict). As a result, 681
2 (91.8%) participants completed both baseline and follow-up surveys. In this study, we analyzed the
3 data of those participants from this sample of 681 who answered the question about depressive
4 mood in both baseline and follow-up surveys (n = 655; 638 obtained by the random sampling
5 recruitment, and 17 by the bulletin recruitment). The Ethics Committee of the Tokyo Metropolitan
6 Institute of Gerontology, Japan, reviewed and approved the study protocol. All subjects provided
7 written consent to participate in this study.

8 9 **Measurements**

10 Face-to-face interviews were used to collect data for both baseline and follow-up surveys.

11 *Neighborhood cohesion*

12 We asked three items about neighborhood cohesion at the baseline survey: “Do people in
13 your neighborhood have close relationships with each other?,” “Can people in your neighborhood
14 be trusted?,” and “Do people in your neighborhood usually help each other?” Respondents
15 answered these items using a four-point Likert scale (“agree,” “slightly agree,” “slightly disagree,”
16 or “disagree”). These three questions were aggregated into one neighborhood cohesion scale by the
17 following steps. First, we dichotomized the responses (1 = “agree” or “slightly agree” and 0 =
18 “slightly disagree” and “disagree”) because these items were highly skewed; second, we summed
19 these dichotomized items (range of scores: 0–3); and third, after reviewing the distribution of this
20 scale, we classified responses into three categories: low (0–1), moderate (2), and high (3). This
21 means that those with high neighborhood cohesion (a score of 3) had positive responses (“agree” or
22 “slightly agree”) to all three items, whereas those with low neighborhood cohesion (a score of 0–1)
23 had positive responses to one item at most.

24 Neighborhood cohesion in this study was assessed by the participants’ perceptions. One

1 common way to create neighborhood-level variables is by aggregating individual responses in each
2 neighborhood (Blakely and Subramanian, 2006). However, this study's sample could not be
3 representative of the residents of the study town (e.g., the participants were aged 65–84 at baseline
4 and healthy volunteer bias may exist). Moreover, the number of participants within a district unit
5 was limited. Therefore, in the analysis, the neighborhood cohesion variable was defined at an
6 individual level (i.e., we did not create an aggregated district-level variable of neighborhood
7 cohesion).

8 9 *Anticipated Daily Stressors*

10 Ten items were used to assess psychological daily stressors at baseline. The items assessed
11 negative expectation for the following 10 points: crime; no expectation of help in an emergency
12 situation; natural disaster; living expenses and medical/care costs; loss of home; future long-term
13 care; loss of friends; dementia; burden on the family if respondent needs long-term care; and social
14 isolation. Respondents answered these questions using a four-point Likert scale (1–4). We
15 conducted a factor analysis and chose to use the one factor that had an eigenvalue greater than 1.00.
16 Therefore, the responses for the 10 items were summed (range of scores: 10–40). Cronbach's alpha
17 at baseline was 0.89. Higher scores indicated greater stress.

18 19 *Depressive mood*

20 Depressive mood was assessed using the Geriatric Depression Scale (GDS) short-form
21 (Burke et al., 1991; Schreiner et al., 2003) at baseline and follow-up. Respondents answered
22 dichotomized questions and responses were summed (range of scores: 0–15). Cronbach's alpha at
23 baseline was 0.77. A cutoff point of 5/6 was adopted, and a score of ≥ 6 indicated depressive mood
24 (Schreiner et al., 2003).

Covariates

Baseline data on gender, age, years of residence in the neighborhood, marital status, socioeconomic status (SES), comorbidities, functional capacity, and social isolation were used as covariates in the analysis. Marital status was categorized as “married” or “not married” (never married, divorced, and widowed). SES included educational attainment (“less than 12 years” or “more than 13 years”), financial stability (“lack of financial resources,” “adequate,” or “wealthy”), and current occupation (“employed” or “unemployed [not in paid employment]”). Information on comorbidities was assessed in a medical interview by a doctor or registered nurse and included the following five diseases: hypertension, cardiovascular disease, cerebrovascular diseases, hyperlipidemia, and diabetes mellitus.

Functional capacity is composed of basic activities of daily living (BADL) and higher-order competence of independence. In our sample, almost all participants were independent in BADL (Murayama et al., 2012b). Therefore, we used only higher-order competence of independence in the analysis. This was measured using the Tokyo Metropolitan Institute of Gerontology Index of Competence (TMIG-IC), which consists of a 13-item index (score range: 0–13; Koyano et al., 1991). A higher score indicated greater independence.

Although neighborhood cohesion is a collective dimension of society and distinct from forms of individual-level social network interaction (Kawachi and Berkman, 2000), the two may be linked. Therefore, we adjusted for individual social networks in the analysis. Social isolation, which is often defined by extremely poor social networks, is related to health status, including mental health (e.g., Cacioppo and Hawkley, 2003). Participants who had contact (meetings, telephone calls, and mail) less than once a month with people other than family members living with them were classified as socially isolated.

District-level indicators

The 2010 Japanese census data were used to obtain objective values for the district. Hatoyama town consists of 28 small districts (*chou-chou* in Japanese). However, area units in the Japanese census data do not correspond to these 28 small districts. Therefore, instead of these small districts, we used postal district as an analytic unit in this study. There are 15 postal districts in the town (12 in the traditional area and 3 in the newly developed area).

To control the effect of structural features of the neighborhood, we used district-level indicators of stability and SES. District-level stability was assessed by calculating the proportion of people who had lived in the district for 20 years and more among each district's adult population. With regard to district-level SES, we used two types of indicators: educational level (the proportion of people who had graduated from university/college and higher) and occupational status (the proportion of people with white-collar jobs) in each district. Higher proportions indicated higher district-level SES. Household income data in each district had not been collected in the Japanese census, so these were the only indicators used to measure district-level SES. In addition to district-level stability and SES, population size and the proportion of people aged 65 or older (aging rate) in each postal district were calculated from the same census data.

Statistical analysis

As Hatoyama has 15 postal districts, participants are nested within each district. Therefore, we fitted the data using a multilevel model comprising individuals nested within districts. This model included a random intercept to assess the likelihood of a GDS score of ≥ 6 versus a score of ≤ 5 . The model partitioned the variation of the likelihood of depressive mood into variance within and between the districts. Thus, this modeling strategy accounts for the hierarchical structure of the

1 data. The estimation method was full maximum likelihood procedure.

2 First, to examine the influence of neighborhood cohesion on depressive mood (H_1), we set
3 the baseline neighborhood cohesion as the independent variable and the follow-up status of
4 depressive mood as the dependent variable. Anticipated daily stressors, baseline GDS,
5 individual-level covariates, and district-level indicators were used for the adjustment. Second, to
6 assess the buffering effect of neighborhood cohesion on depressive mood (H_2), the interaction
7 between anticipated daily stressors and neighborhood cohesion was added into the model. To
8 investigate further the direct and the stress-buffering effects of neighborhood cohesion on
9 depressive mood (H_3), we tested the interactions between neighborhood cohesion and
10 age/residential year and the interaction among anticipated daily stressors, neighborhood cohesion,
11 and age/residential year in the model, respectively. Finally, to check the robustness of the effects of
12 neighborhood cohesion on depressive mood, the analysis was performed excluding the participants
13 who reported depressive mood ($GDS \geq 6$) at baseline. The results of the fixed effects are expressed
14 as odds ratios (ORs) with 95% confidence intervals (CIs). Statistical analysis was performed using
15 HLM 6.08 (Scientific Software International, Inc., Skokie, IL).

16 17 18 **Results**

19 Table 1 shows the characteristics of the participants. Females comprised 41.7% and the
20 mean age was 71.9 ± 5.1 years. The mean number of years of residence in the neighborhood was
21 31.8 ± 16.7 years (median: 30.0 years). A total of 12.4% had a GDS score of ≥ 6 at baseline.
22 Regarding perceived neighborhood cohesion, 24.5%, 40.1%, and 35.4% evaluated this as low,
23 moderate, and high, respectively. At the follow-up survey, 15.9% reported depressive mood (GDS

1 ≥6). Between baseline and follow-up, 9.0% and 5.5% changed their GDS status from
2 non-depressive to depressive, and from depressive to non-depressive, respectively.

3 <Insert Table 1 here>

4 Table 2 indicates the characteristics of the study district and the correlations among the
5 district-level variables. District-level population size was highly correlated with district-level
6 educational level and occupational status ($r = 0.86$ and 0.84 , respectively). District-level educational
7 level was also highly correlated with district-level occupational status ($r = 0.80$). Therefore, in the
8 multilevel analysis, we used district-level aging rate, stability and education to avoid
9 multicollinearity.

10 <Insert Table 2 here>

11 Table 3 shows the association between perceived neighborhood cohesion and depressive
12 mood. In Model 1, when adjusting for age, gender, and baseline GDS, people who had low and
13 moderate neighborhood cohesion were more likely to experience depressive mood, compared with
14 those with high neighborhood cohesion (ORs [95% CIs] were 1.91 [1.19–3.08] for the low category
15 and 1.77 [1.05–3.00] for the moderate category). However, after adjusting for other
16 sociodemographic variables, comorbidities, functional capacity, and social isolation at baseline in
17 Models 2 and 3, this association became non-significant. This remained the case after the
18 adjustment of district-level indicators in Model 4. Higher anticipated daily stressor scores were
19 related to a greater likelihood of depressive mood (1.08 [1.03–1.13] in Model 4). Higher
20 district-level education was significantly associated with less likelihood of depressive mood (0.62
21 [0.46–0.86] in Model 4).

22 To test the effects of the interaction between perceived neighborhood cohesion and
23 age/years of residence in the neighborhood on depressive mood, we added the interaction terms into
24 Model 4 (Model 5). We found a marginally significant interaction between low neighborhood

1 cohesion and years of residence (1.33 [0.94–1.90], $p = 0.085$). This implied that the association
2 between neighborhood cohesion and depressive mood was stronger among long-term residents of
3 the neighborhood. Indeed, in the analysis for those who had lived in the neighborhood for 30 years
4 (median in our sample) and longer in Model 4, the OR for the low category was 1.96, though this
5 was not statistically significant (data not shown). In contrast, the interaction between neighborhood
6 cohesion and age on depressive mood was not significant.

7 <Insert Table 3 here>

8 We examined the buffering effect of perceived neighborhood cohesion on the association
9 between anticipated daily stressors and depressive mood (Table 4). In Model 6, there was a
10 significant interaction between anticipated daily stressors and low neighborhood cohesion on
11 depressive mood (1.13 [1.01–1.28]). Figure 1 illustrates this interaction. For ease of presentation,
12 we divided anticipated daily stressors into two categories by median (high and low). The figure
13 indicates that perceived high neighborhood cohesion partially offsets the deleterious effect of
14 anticipated daily stressors on depressive mood. Additionally, we tested whether this buffering effect
15 differed by participants' age or years of residence in the neighborhood. In Model 7, a significant
16 interaction among anticipated daily stressors, low neighborhood cohesion, and years of residence
17 was found (1.01 [1.00–1.02]). This suggested that the stress-buffering effect of neighborhood
18 cohesion on depressive mood was stronger among long-term residents of the neighborhood. We did
19 not find an interaction among anticipated daily stressors, neighborhood cohesion, and age.

20 <Insert Table 4 and Figure 1 here>

21 We performed an analysis with the same models as shown in Tables 3 and 4, but
22 excluding participants who reported depressive mood ($GDS \geq 6$) at baseline. The results of Models
23 3–7 are shown in Table 5. These were consistent with those for the entire sample.

24 <Insert Table 5 here>

Discussion

The present study examined the direct association of perceived neighborhood cohesion with depressive mood and the buffering effect of perceived neighborhood cohesion on the relationship between anticipated daily stressors and depressive mood, using 2-year longitudinal data from older Japanese residents. A major strength of this research lies in its use of longitudinal data from older Japanese people. It is important to examine this association from a longitudinal perspective to identify causality; however, to our knowledge, this is the first attempt to examine the longitudinal association of neighborhood cohesion with depressive mood in old age. Moreover, findings from a non-Western nation that has the most rapidly aging population in the world may stimulate further research and discussion on the link between neighborhood cohesion and mental health, and also provide guidelines for health promotion policies for both current and future aging societies.

Our study setting, Hatoyama, has unique characteristics. Hatoyama was developed as a commuter town for Tokyo at the end of the high-growth period from the mid-1950s to the mid-1970s (Murayama et al., 2012b). Many people from outside Hatoyama moved to the town, especially to the newly developed area. The population size within each postal district in the newly developed area is higher than that in the traditional area (614 ± 483 in the traditional area, and $2,614 \pm 110$ in the newly developed area; data not shown in the table). Moreover, people in the newly developed area have a higher SES (e.g., education, occupation) than those in the traditional area (Murayama et al., 2012b). Therefore, we found strong correlations between population size and SES at the district level. Additionally, the proportion of people aged 65 or older in Hatoyama has been increasing rapidly. From a level of 26.1% in 2010, it is predicted to reach 50.7% by 2035,

1 which will substantially exceed the national estimates of 33.7% (National Institute of Population
2 and Social Security Research, 2007, 2009). However, although Hatoyama has such unique
3 characteristics, there are many commuter towns with similar characteristics, not only in Japan (e.g.,
4 the suburbs of Tokyo and Osaka metropolitan areas; National Institute of Population and Social
5 Security Research, 2009), but also in other countries. For example, in the United States, a rapid
6 growth in senior residents between 2010 and 2030 is predicted, particularly for some states in the
7 West, and for Texas, Georgia, and Florida in the South (Frey, 2010). Although aging suburbs are
8 most numerous and relatively concentrated in Florida (36% of all suburbs in Florida have a rapidly
9 graying population), such aging communities can be found throughout the country (Mikelbank,
10 2010). Therefore, our findings could be generalized to, and used in, these communities when
11 developing health promotion policies.

12 Although the association of perceived neighborhood cohesion with depressive mood was
13 statistically significant in Model 1 (adjusted for age, gender, and baseline GDS), this association
14 became non-significant after the additional adjustment for factors such as anticipated daily stressors
15 and some actual stressors (e.g., financial instability, social isolation, poor functional capacity) in
16 Models 2–4. This meant that our first hypothesis (H_1) was not supported. In a review paper, 11 out
17 of 16 studies reported a protective effect of social interaction between neighbors against depressive
18 mood (Mair et al., 2008). Following this literature review, several studies also demonstrated this
19 protective association (e.g., Echeverria et al., 2008; Mair et al., 2010; Mair et al., 2009). There may
20 be several reasons why neighborhood cohesion was not associated with depressive mood in this
21 study. First, the participants were limited to those aged 65–84 at baseline, while the
22 above-mentioned earlier studies on neighborhood cohesion and depressive mood were not
23 conducted with older people. Our sample might be homogeneous with regard to age, and therefore
24 homogeneity of the sample might make the influence of neighborhood cohesion on the outcome

unclear. Second, our sample had better mental health than is generally found in older people. The proportions of those with depressive mood were 12.4% at baseline and 15.9% at follow-up. These proportions are lower than that reported in another study of community-dwelling older Japanese people (33.5%; Wada et al., 2004). Thus, the influence of neighborhood cohesion on depressive mood might be underestimated in our sample.

We found a stress-buffering effect of perceived neighborhood cohesion on depressive mood, supporting our second hypothesis (H_2). It has been suggested that the social cohesion of communities could influence associations between potentially adverse factors on mental health and psychological outcomes of individuals (Fone et al., 2007; Robinette et al., 2013; Takagi et al., 2013). Neighborhood cohesion is considered distinct from social support and social networks, because it assesses norms and expectations for behavior rather than actual ties or direct interaction (Kawachi et al., 2008). However, it is widely accepted that people who live in a cohesive neighborhood may find it easier to obtain social support from neighbors to cope with daily stress. Being able to depend on neighbors for help may attenuate the adverse effects on depressive mood of living in cohesive neighborhoods. Moreover, as the diffusion of knowledge and information (e.g., about health promotion) is likely to occur faster in such a cohesive neighborhood, knowledge and information about how to deal with stress may be easily available when needed.

Furthermore, we found significant interactions between perceived neighborhood cohesion and years of residence in the neighborhood, and among anticipated daily stressors, neighborhood cohesion, and years of residence, in Models 5 and 7, respectively. However, the direct effect and the stress-buffering effect of perceived neighborhood cohesion on depressive mood did not differ by age. These results partly support our third hypothesis (H_3). Robinette et al. (2013) examined whether the association between perceived neighborhood cohesion and well-being varied by age among adults and reported that neighborhood cohesion buffered the negative affect from daily

1 stressors among younger adults more than among middle-aged and older adults. They argued that
2 older adults might rely less on peripheral network members for support, including neighbors, than
3 do younger adults. Compared with this previous research, our participants were more homogeneous
4 in age (aged 65–84 at baseline), making it unlikely that an age difference would be found for our
5 sample. That is, older people experience a buffering effect of neighborhood cohesion on depressive
6 mood regardless of age.

7 In contrast, among the older population, length of residence in the neighborhood had a
8 stronger impact on the relationship among neighborhood cohesion, anticipated daily stressors, and
9 depressive mood. The uniqueness of the study area background may explain the impact of length of
10 residence. Many people from outside Hatoyama moved to the town when it was developed as a
11 commuter town. Indeed, 85% of the participants were born outside of the town, whereas only 5%
12 were born in their current residential neighborhood (data not shown). That is, community members’
13 length of residence varied and this influenced the degree of exposure to the neighborhood. Because
14 residential length increases the chance to obtain support and knowledge/information from neighbors
15 to cope with stress in this study setting, this might have had a strong impact on the relationship
16 among neighborhood cohesion, anticipated daily stressors, and depressive mood.

17 Additionally, this uniqueness of the study area may have affected the non-significance of
18 the direct association between neighborhood cohesion and depressive mood, discussed above as H_1 .
19 A previous study indicated that people living in a community that was developed in later years tend
20 to have lower general trust (Hanibuchi et al. 2012); therefore, neighborhood cohesion might be
21 relatively lower in this town compared with other settings. This could explain the small variance in
22 neighborhood cohesion in our sample; thus, we might have underestimated the association between
23 neighborhood cohesion and depressive mood. This was also the case for social isolation. We
24 adjusted for social isolation in the model because individual-level social networks are often closely

1 associated with neighborhood cohesion, but we found no association between social isolation and
2 depressive mood. Although 8.7% of our sample was socially isolated, this proportion is lower than
3 that among older Japanese in urban area (11.9%; Murayama et al., 2011). Therefore, social isolation
4 had only a small impact on depressive mood in our sample.

5 Because we measured both neighborhood cohesion and depressive mood at an individual
6 level, there was a concern about reporting bias. That is, people who already experience depressive
7 mood tend to report lower social cohesion. To reduce the effect of this reporting bias, we excluded
8 participants who reported depressive mood ($GDS \geq 6$) at baseline, which strengthened the effects of
9 neighborhood cohesion on depressive mood. This could support the robustness of our findings.

10 Regarding the district-level indicators, we adjusted for district-level aging rate in the
11 model instead of population size. Previous studies have reported that the rate of depression
12 increases with age (Davey et al., 2004; Luppa et al., 2012; Stordal et al., 2001). In addition, it is
13 possible that depression spreads from one individual to another (this is called “emotional
14 contagion”; Howes et al., 1985). Therefore, a higher proportion of people aged 65 or older in the
15 residential neighborhood may affect one’s depressive mood. These factors suggest that the
16 adjustment of aging rate in the model was appropriate. In the multilevel analysis, higher
17 district-level educational level was associated with less likelihood of depressive mood. This pattern
18 is consistent with previous research on elderly people (Aneshensel et al., 2007; Beard et al., 2009).
19 Greater neighborhood SES seems to be associated with higher individual social trust (Murayama et
20 al., 2014). Therefore, those living in a neighborhood characterized by high SES might obtain more
21 support and feel less stress/unease.

22 The current study has some limitations. First, we used the variable of neighborhood
23 cohesion at an individual level, not at a district level. Future work should attempt to replicate the
24 current findings using a more objective measurement of neighborhood cohesion. For example, using

representative survey data or objectively collected data on neighborhood cohesion could possibly avoid same-source bias (Mujahid et al., 2007). We used this method for the district-level indicators, but there was no available data on neighborhood cohesion in the study area. Second, the participants of this cohort study do not necessarily represent the target population because we used stratified sampling of four groups classified by age and residential area (random sampling for one group but used a complete census for the other three groups). Moreover, the response rate was relatively low in the four stratified groups. Although the male-female ratio, mean age, and residential area ratio (participants residing in traditional areas as compared to those living in the newly developed area) of the participants did not substantially differ from those of the overall population of the town aged 65–84, and age and residential area were taken into consideration in the analysis, care is needed in the interpretation of the findings. Furthermore, we included 17 participants in the analysis who were recruited through the town bulletin because these people possibly possessed different characteristics from those recruited through the other sampling methods. Therefore, we undertook a sensitivity analysis by performing the same analyses excluding these 17 participants. This analysis produced similar results to those reported here. Third, the potential for bias due to selection of healthy volunteers must be considered (i.e., healthy volunteer effect). In fact, our study participants tended to be healthy and wealthy compared with the Japanese average at baseline (Murayama et al., 2012b). Moreover, those who participated in both baseline and follow-up surveys ($n = 681$) had better health at baseline, compared with those who did not participate in the follow-up survey ($n = 61$), for example for depressive mood (11.9% vs. 19.7%, respectively, $p = 0.066$) and TMIG-IC score (12.4 ± 1.1 vs. 11.8 ± 2.1 , respectively, $p = 0.001$). These factors might have caused selection bias and an underestimation of the association between neighborhood cohesion and depressive mood. Fourth, data for this study came from a single suburban area. Therefore, the generalizability of these findings should be examined by conducting further studies in different settings.

Conclusions

Although we did not find any direct association between perceived neighborhood cohesion and depressive mood, the present longitudinal study provides evidence that perceived neighborhood cohesion had a stress-buffering effect on depressive mood in old age. Additionally, this effect was stronger for long-term residents of the neighborhood. Therefore, interventions to strengthen neighborhood cohesion may help to prevent depressive mood in older residents.

Funding

This study was supported by research grants from the Research Institute of Science and Technology for Society (RISTEX), the Japan Science and Technology Agency, the Ministry of Education, Culture, Sports, Science and Technology (MEXT) KAKENHI Grant Number 24792611, and the town of Hatoyama.

References

Aneshensel, C.S., Wight, R.G., Miller-Martinez, D., Botticello, A.L., Karlamangla, A.S., Seeman, T.E., 2007. Urban neighborhoods and depressive symptoms among older adults. *J Gerontol*

- 1 B Psychol Sci Soc Sci 62, S52-59.
- 2 Beard, J.R., Cerda, M., Blaney, S., Ahern, J., Vlahov, D., Galea, S., 2009. Neighborhood
- 3 characteristics and change in depressive symptoms among older residents of New York City.
- 4 Am J Public Health 99, 1308-1314. DOI:10.2105/AJPH.2007.125104
- 5 Blakely, T., Subramanian, S.V., 2006. Multilevel studies, In: Oakes, J.M., Kaufman, J.S. (Eds.),
- 6 Methods in social epidemiology. Jossey-Bass, San Francisco, pp. 316-340.
- 7 Burke, W.J., Roccaforte, W.H., Wengel, S.P., 1991. The short form of the Geriatric Depression
- 8 Scale: a comparison with the 30-item form. J Geriatr Psychiatry Neurol 4, 173-178.
- 9 DOI:10.1177/089198879100400310
- 10 Cabinet Office, 2007. White paper on the national lifestyle 2007. Jiji Gahosha, Tokyo.
- 11 Cacioppo, J.T., Hawkley, L.C., 2003. Social isolation and health, with an emphasis on underlying
- 12 mechanisms. Perspect Biol Med 46, S39-52. DOI:10.1353/pbm.2003.0063
- 13 Cattell, V., 2001. Poor people, poor places, and poor health: the mediating role of social networks
- 14 and social capital. Soc Sci Med 52, 1501-1516. DOI:10.1016/S0277-9536(00)00259-8
- 15 Davey, A., Halverson, C.F., Jr., Zonderman, A.B., Costa, P.T., Jr., 2004. Change in depressive
- 16 symptoms in the Baltimore longitudinal study of aging. J Gerontol B Psychol Sci Soc Sci
- 17 59, P270-277. DOI:10.1093/geronb/59.6.P270
- 18 Diez Roux, A.V., Mair, C., 2010. Neighborhoods and health. Ann N Y Acad Sci 1186, 125-145.
- 19 DOI:10.1111/j.1749-6632.2009.05333.x
- 20 Echeverria, S., Diez-Roux, A.V., Shea, S., Borrell, L.N., Jackson, S., 2008. Associations of
- 21 neighborhood problems and neighborhood social cohesion with mental health and health
- 22 behaviors: the Multi-Ethnic Study of Atherosclerosis. Health Place 14, 853-865.
- 23 DOI:10.1016/j.healthplace.2008.01.004
- 24 Emmerson, J. P., Burvill, P. W., Finlay-Jones, R., Hall, W., 1989. Life events, life difficulties and

confiding relationships in the depressed elderly. *Br J Psychiatry* 155, 787-792. DOI: 10.1192/bjp.155.6.787

Ferrari, A.J., Charlson, F.J., Norman, R.E., Patten, S.B., Freedman, G., Murray, C.J., Vos, T., Whiteford, H.A., 2013. Burden of depressive disorders by country, sex, age, and year: findings from the global burden of disease study 2010. *PLoS Med* 10, e1001547. DOI:10.1371/journal.pmed.1001547

Fone, D., Dunstan, F., Lloyd, K., Williams, G., Watkins, J., Palmer, S., 2007. Does social cohesion modify the association between area income deprivation and mental health? a multilevel analysis. *Int J Epidemiol* 36, 338-345. DOI:10.1093/ije/dym004

Frey, W.H., 2010. Baby boomers and the new demographics of America's seniors. *J Am Soc Aging* 34, 28-37.

Greenberg, P.E., Stiglin, L.E., Finkelstein, S.N., Berndt, E.R., 1993. Depression: a neglected major illness. *J Clin Psychiatry* 54, 419-424.

Hanibuchi, T., Kondo, K., Nakaya, T., Shirai, K., Hirai, H., Kawachi, I., 2012. Does walkability mean sociable? Neighborhood determinants of social capital among older adults in Japan. *Health Place* 18, 229-239. DOI:10.1016/j.healthplace.2011.09.015

Howes, M.J., Hokanson, J.E., Lowenstein, D.A., 1985. Induction of depressive affect after prolonged exposure to a mildly depressed individual. *J Pers Soc Psychol* 49, 1110-1113. DOI:10.1037/0022-3514.49.4.1110

Kawachi, I., Berkman, L.F., 2000. Social cohesion, social capital, and health, In: Berkman, L.F., Kawachi, I. (Eds.), *Social epidemiology*. Oxford University Press, New York, pp. 174-190.

Kawachi, I., Subramanian, S.V., Kim, D., 2008. Social capital and health: a decade of progress and beyond, In: Kawachi, I., Subramanian, S.V., Kim, D. (Eds.), *Social capital and health*. Springer, New York, pp. 1-26.

- 1 Koyano, W., Shibata, H., Nakazato, K., Haga, H., Suyama, Y., 1991. Measurement of competence:
2 reliability and validity of the TMIG Index of Competence. Arch Gerontol Geriatr 13,
3 103-116. DOI:10.1016/0167-4943(91)90053-S
- 4 Lupp, M., Sikorski, C., Luck, T., Ehreke, L., Konnopka, A., Wiese, B., Weyerer, S., Konig, H.H.,
5 Riedel-Heller, S.G., 2012. Age- and gender-specific prevalence of depression in latest-life:
6 systematic review and meta-analysis. J Affect Disord 136, 212-221.
7 DOI:10.1016/j.jad.2010.11.033
- 8 Mair, C., Diez Roux, A.V., Galea, S., 2008. Are neighbourhood characteristics associated with
9 depressive symptoms? a review of evidence. J Epidemiol Community Health 62, 940-946.
10 DOI:10.1136/jech.2007.066605
- 11 Mair, C., Diez Roux, A.V., Morenoff, J.D., 2010. Neighborhood stressors and social support as
12 predictors of depressive symptoms in the Chicago Community Adult Health Study. Health
13 Place 16, 811-819. DOI:10.1016/j.healthplace.2010.04.006
- 14 Mair, C., Diez Roux, A.V., Shen, M., Shea, S., Seeman, T., Echeverria, S., O'Meara, E.S., 2009.
15 Cross-sectional and longitudinal associations of neighborhood cohesion and stressors with
16 depressive symptoms in the multiethnic study of atherosclerosis. Ann Epidemiol 19, 49-57.
17 DOI:10.1016/j.annepidem.2008.10.002
- 18 Mikelbank, B.A., 2010. A typology of U.S. suburban places. Hous Policy Debate 15, 935-964.
19 DOI:10.1080/10511482.2004.9521527
- 20 Mujahid, M.S., Diez Roux, A.V., Morenoff, J.D., Raghunathan, T., 2007. Assessing the
21 measurement properties of neighborhood scales: from psychometrics to econometrics. Am J
22 Epidemiol 165, 858-867. DOI:10.1093/aje/kwm040
- 23 Müller-Spahn, F., Hock, C., 1994. Clinical presentation of depression in the elderly. Gerontology
24 40, 10-14. DOI:10.1159/000213615

- 1 Murayama, H., Arami, R., Wakui, T., Sugawara, I., Yoshie, S., 2014. Cross-level interaction
- 2 between individual and neighbourhood socioeconomic status in relation to social trust in a
- 3 Japanese community. *Urban Stud* 51, 2770-2786. DOI:10.1177/0042098013513648
- 4 Murayama, H., Fujiwara, Y., Kawachi, I., 2012a. Social capital and health: a review of prospective
- 5 multi-level studies. *J Epidemiol* 22, 179-187. DOI:10.2188/jea.JE20110128
- 6 Murayama, H., Nishi, M., Shimizu, Y., Kim, M.J., Yoshida, H., Amano, H., Fujiwara, Y., Shinkai, S.,
- 7 2012b. The Hatoyama cohort study: design and profile of participants at baseline. *J*
- 8 *Epidemiol* 22, 551-558. DOI:10.2188/jea.JE20120015
- 9 Murayama, H., Shibui, Y., Fukuda, Y., Murashima, S., 2011. A new crisis in Japan—social isolation
- 10 in old age. *J Am Geriatr Soc* 59, 2160-2162. DOI: 10.1111/j.1532-5415.2011.03640.x.
- 11 Nakane, C., 1970. Japanese society. University of California Press, Los Angeles.
- 12 National Institute of Population and Social Security Research, 2007. Nihon no shorai suike jinko:
- 13 2006–2055 [Population projections for Japan: 2006–2055]. Health and Welfare Statistics
- 14 Association, Tokyo.
- 15 National Institute of Population and Social Security Research, 2009. Nihon no shikuchoson betsu
- 16 shorai suike jinko: 2005–2035 [Population projections by municipality, Japan: 2005–2035].
- 17 Health and Welfare Statistics Association, Tokyo.
- 18 Robinette, J.W., Charles, S.T., Mogle, J.A., Almeida, D.M., 2013. Neighborhood cohesion and
- 19 daily well-being: results from a diary study. *Soc Sci Med* 96, 174-182.
- 20 DOI:10.1016/j.socscimed.2013.07.027
- 21 Rowe, J.W., Kahn, R.L., 1997. Successful aging. *Gerontologist* 37, 433-440.
- 22 DOI: 10.1093/geront/37.4.433
- 23 Schreiner, A.S., Hayakawa, H., Morimoto, T., Kakuma, T., 2003. Screening for late life depression:
- 24 cut-off scores for the Geriatric Depression Scale and the Cornell Scale for Depression in

- 1 Dementia among Japanese subjects. *Int J Geriatr Psychiatry* 18, 498-505.
2
3
4
5
6
7
8
9 DOI:10.1002/gps.880
10
11 3 Stordal, E., Kruger, M.B., Dahl, N.H., Kruger, O., Mykletun, A., Dahl, A.A., 2001. Depression in
12
13 4 relation to age and gender in the general population: the Nord-Trøndelag Health Study
14
15 5 (HUNT). *Acta Psychiatrica Scandinavica* 104, 210-216.
16
17 6 DOI:10.1034/j.1600-0447.2001.00130.x
18
19
20 7 Stuck, A.E., Walthert, J.M., Nikolaus, T., Bula, C.J., Hohmann, C., Beck, J.C., 1999. Risk factors
21
22 8 for functional status decline in community-living elderly people: a systematic literature
23
24 9 review. *Soc Sci Med* 48, 445-469. DOI:10.1016/S0277-9536(98)00370-0
25
26
27 10 Takagi, D., Kondo, K., Kondo, N., Cable, N., Ikeda, K., Kawachi, I., 2013. Social
28
29 11 disorganization/social fragmentation and risk of depression among older people in Japan:
30
31 12 multilevel investigation of indices of social distance. *Soc Sci Med* 83, 81-89.
32
33 13 DOI:10.1016/j.socscimed.2013.01.001
34
35
36 14 Wada, T., Ishine, M., Sakagami, T., Okumiya, K., Fujisawa, M., Murakami, S., Otsuka, K., Yano,
37
38 15 S., Kita, T., Matsubayashi, K., 2004. Depression in Japanese community-dwelling elderly:
39
40 16 prevalence and association with ADL and QOL. *Arch Gerontol Geriatr* 39, 15-23.
41
42
43 17 Whiteford, H.A., Degenhardt, L., Rehm, J., Baxter, A.J., Ferrari, A.J., Erskine, H.E., Charlson, F.J.,
44
45 18 Norman, R.E., Flaxman, A.D., Johns, N., Burstein, R., Murray, C.J., Vos, T., 2013. Global
46
47 19 burden of disease attributable to mental and substance use disorders: findings from the
48
49 20 Global Burden of Disease Study 2010. *Lancet* 382, 1575-1586.
50
51 21 DOI:10.1016/S0140-6736(13)61611-6
52
53
54 22 Yamagishi, T., Cook, K.S., Watabe, M., 1998. Uncertainty, trust, and commitment formation in the
55
56 23 United States and Japan. *Am J Sociol* 104, 165-194. DOI:10.1086/210005
57
58 24 Yamagishi, T., Yamagishi, M., 1994. Trust and commitment in the United States and Japan. *Motiv*
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Emot 18, 129-166. DOI:10.1007/BF02249397

Longitudinal association between neighborhood cohesion and depressive mood in old age: A Japanese prospective study

Hiroshi Murayama*, Mariko Nishi, Yu Nofuji, Eri Matsuo, Yu Taniguchi, Hidenori Amano, Yuri Yokoyama, Yoshinori Fujiwara, and Shoji Shinkai

Research Team for Social Participation and Community Health, Tokyo Metropolitan
Institute of Gerontology, Tokyo, Japan

35-2 Sakae-cho, Itabashi-ku, Tokyo 173-0015, Japan

*Corresponding author: Hiroshi Murayama, PhD, RN, PHN

Research Team for Social Participation and Community Health, Tokyo Metropolitan
Institute of Gerontology

Tel: +81-3-3964-3241 (ext. 4253)

Fax: +81-3-3579-4776;

e-mail: murayama@tmig.or.jp

Table 1. Participant characteristics (n = 655)

		No. of respondents	n (%) or Mean $\pm$ SD ^a	
Baseline (2010)				
Gender	Female	655	273	(41.7)
Age (years)		655	71.9 $\pm$ 5.1	
Residing in neighborhood (years)		655	31.8 $\pm$ 16.7	
Marital status	Not married ^a	653	106	(16.2)
Education	Less than 12 years	655	427	(65.2)
Occupational status	Currently unemployed	654	506	(77.4)
Financial stability		655		
	Lack of financial resources		55	(8.4)
	Adequate		488	(74.5)
	Wealthy		112	(17.1)
Comorbidities		655		
	0		160	(24.4)
	1		242	(36.9)
	≥ 2		253	(38.6)
TMIG-IC (range: 0–13)		654	12.4 $\pm$ 1.1	
Social isolation	Isolated	655	57	(8.7)
Anticipated daily stressors (range: 10–40)		645	24.9 $\pm$ 5.9	
GDS	≥ 6	655	81	(12.4)
Neighborhood cohesion		653		
	Low		160	(24.5)
	Moderate		262	(40.1)
	High		231	(35.4)
Follow-up (2012)				
GDS	≥ 6	655	104	(15.9)
Change in GDS between baseline and follow-up		655		
No change (≤ 5 to ≤ 5)			515	(78.6)
No change (≥ 6 to ≥ 6)			45	(6.9)
Non-depressive to depressive (≤ 5 to ≥ 6)			59	(9.0)
Depressive to non-depressive (≥ 6 to ≤ 5)			36	(5.5)

GDS: Geriatric Depression Scale. SD: standard deviation. TMIG-IC: Tokyo Metropolitan Institute of Gerontology Index of Competence.

^aNot married includes never married, divorced, and widowed.

Table 2. Characteristics of the district-level variables and their correlations (n = 15)

	Mean $\pm$ SD	Min–Max	Correlation coefficient (Pearson's <i>r</i>)				
			a	b	c	d	e
a. Population size	1,020 $\pm$ 944	75–2,762		0.10	-0.35	0.86***	0.84***
b. Aging rate							
% people aged ≥ 65	27.6 $\pm$ 7.6	17.7–49.9			-0.15	-0.03	0.03
c. Stability							
% people living in the district for ≥ 20 years	60.7 $\pm$ 13.5	37.4–81.3				-0.05	-0.43
d. Education							
% people graduating from university/college or higher	13.7 $\pm$ 7.5	4.8–28.7					0.80***
e. Occupation							
% people with white-collar jobs	53.6 $\pm$ 6.7	39.2–66.6					

SD: standard deviation.

 $p < 0.001$.

Table 3. Associations of neighborhood cohesion with depressive mood for the entire sample: multilevel binomial logistic regression model

		Model 1		Model 2		Model 3		Model 4		Model 5	
		OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
Female		0.84	(0.61–1.15)	0.82	(0.54–1.23)	0.77	(0.44–1.37)	0.75	(0.41–1.35)	0.74	(0.41–1.33)
Age	(every 10 years)	1.83	(1.17–2.88)	1.72	(1.13–2.61)	1.46	(0.90–2.38)	1.50	(0.90–2.50)	1.51	(0.91–2.52)
Years of residence	(every 10 years)			0.90	(0.75–1.07)	0.88	(0.73–1.06)	0.88	(0.74–1.06)	0.85	(0.71–1.02)
Not married				1.28	(0.61–2.69)	1.53	(0.79–2.96)	1.54	(0.79–3.04)	1.55	(0.79–3.07)
Less than 12 years of education				0.92	(0.70–1.22)	0.91	(0.52–1.60)	0.88	(0.48–1.61)	0.92	(0.50–1.68)
Currently unemployed				1.39	(0.81–2.36)	1.23	(0.65–2.36)	1.25	(0.64–2.45)	1.20	(0.61–2.36)
Financial stability ^a	Lack of financial resources			7.78	(3.23–18.72)	4.76	(1.44–15.71)	5.92	(1.54–22.76)	5.39	(1.38–21.05)
	Adequate			4.03	(2.76–5.87)	3.05	(1.15–8.12)	3.86	(1.21–12.30)	3.64	(1.13–11.74)
Comorbidities ^b	1					1.19	(0.61–2.33)	1.21	(0.60–2.44)	1.19	(0.59–2.41)
	≥2					1.14	(0.60–2.17)	1.19	(0.61–2.35)	1.17	(0.59–2.32)
Functional capacity	(every 1-point increase in TMIG-IC score)					0.70	(0.58–0.86)	0.71	(0.58–0.87)	0.70	(0.57–0.86)
Socially isolated						1.44	(0.68–3.06)	1.46	(0.67–3.21)	1.45	(0.66–3.20)
Higher stressor scores						1.08	(1.03–1.13)	1.08	(1.03–1.13)	1.08	(1.03–1.13)
Baseline GDS	≥6	9.32	(6.21–13.97)	7.25	(4.49–11.69)	5.98	(3.29–10.89)	6.11	(3.32–11.26)	6.77	(3.63–12.63)
Neighborhood cohesion ^c	Low	1.91	(1.19–3.08)	1.57	(0.96–2.59)	1.19	(0.60–2.37)	1.19	(0.58–2.43)	0.47	(0.12–1.88)
	Moderate	1.77	(1.05–3.00)	1.59	(0.93–2.74)	1.38	(0.75–2.54)	1.38	(0.73–2.62)	2.04	(0.54–7.75)
Low neighborhood cohesion ^c × years of residence										1.33	(0.94–1.90)
Moderate neighborhood cohesion ^c × years of residence										0.87	(0.60–1.26)
District-level aging rate								1.02	(0.98–1.06)	1.02	(0.98–1.06)
District-level stability ^d	(higher quartile)							0.76	(0.53–1.09)	0.75	(0.52–1.07)
District-level education ^e	(higher quartile)							0.62	(0.46–0.86)	0.64	(0.47–0.88)

CI: confidence interval. GDS: Geriatric Depression Scale. OR: odds ratio. TMIG-IC: Tokyo Metropolitan Institute of Gerontology Index of Competence.

^aRelative to wealthy.^bRelative to no disease.^cRelative to high neighborhood cohesion.^dDefined as the proportion of people who had lived in the neighborhood for ≥20 years.^eDefined as the proportion of people who graduated from university/college or higher.

Table 4. Stress-buffering effect of neighborhood cohesion on depressive mood: multilevel binomial logistic regression model

		Model 6		Model 7	
		OR	(95% CI)	OR	(95% CI)
Higher stressor scores		1.07	(1.02–1.13)	1.08	(1.02–1.13)
Neighborhood cohesion ^a	Low	0.04	(0.00–1.31)	0.04	(0.00–1.37)
	Moderate	0.43	(0.02–8.22)	0.28	(0.01–5.71)
Low neighborhood cohesion ^a × higher stressor scores		1.13	(1.01–1.28)	1.09	(0.96–1.25)
Moderate neighborhood cohesion ^a × higher stressor scores		1.04	(0.94–1.17)	1.08	(0.96–1.23)
Low neighborhood cohesion ^a × higher stressor scores × years of residence				1.01	(1.00–1.02)
Moderate neighborhood cohesion ^a × higher stressor scores × years of residence				0.99	(0.98–1.01)

CI: confidence interval. OR: odds ratio.

All other individual-level variables (gender, age, years of residence, marital status, education, current occupation, financial stability, comorbidities, functional capacity, social isolation, and depressive mood at baseline) and district-level variables (aging rate, stability, and education) were adjusted in both models.

^aRelative to high neighborhood cohesion.

Table 5. Associations of neighborhood cohesion with depressive mood excluding those who had depressive mood (GDS ≥ 6) at baseline: multilevel binomial logistic regression model

		Model 3		Model 4		Model 5		Model 6		Model 7	
		OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)	OR	(95% CI)
Higher stressor scores		1.07	(1.01–1.13)	1.07	(1.01–1.13)	1.07	(1.01–1.13)	1.07	(1.01–1.13)	1.07	(1.01–1.14)
Neighborhood cohesion ^a	Low	1.51	(0.67–3.38)	1.50	(0.65–3.47)	0.50	(0.10–2.67)	0.02	(0.00–1.30)	0.02	(0.00–1.59)
	Moderate	1.48	(0.73–3.00)	1.50	(0.72–3.15)	2.40	(0.51–11.42)	0.20	(0.01–5.16)	0.14	(0.01–3.92)
Low neighborhood cohesion ^a $\times$ years of residence						1.39	(0.91–2.12)				
Moderate neighborhood cohesion ^a $\times$ years of residence						0.83	(0.52–1.32)				
Low neighborhood cohesion ^a $\times$ higher stressor scores								1.17	(1.01–1.36)	1.12	(0.95–1.32)
Moderate neighborhood cohesion ^a $\times$ higher stressor scores								1.08	(0.96–1.22)	1.12	(0.97–1.29)
Low neighborhood cohesion ^a $\times$ higher stressor scores $\times$ years of residence										1.02	(1.01–1.03)
Moderate neighborhood cohesion ^a $\times$ higher stressor scores $\times$ years of residence										0.99	(0.97–1.01)

CI: confidence interval. GDS: Geriatric Depression Scale. OR: odds ratio.

All other individual-level variables (gender, age, years of residence, marital status, education, current occupation, financial stability, comorbidities, functional capacity, and social isolation) and district-level variables (aging rate, stability, and education) were adjusted in all models.

^aRelative to high neighborhood cohesion.

Figure(s)

Figure 1.
Predicted probability of depressive mood at follow-up by anticipated daily stressors and neighborhood cohesion at baseline among entire sample (based on Model 1 in Table 2). For ease of presentation, anticipated daily stressor scores were dichotomized: high (above median) and low (below median).