**Remnants of a Late Triassic oceanisland in the Gufeng area,
northern Tibet: implications for the opening and early
evolution of the Bangong–NujiangTethyan Ocean**

Jian-Jun Fan^{1*}, Cai Li¹, Ming Wang¹, Yi-Ming Liu¹, Chao-Ming Xie¹

¹*College of Earth Sciences, Jilin University, Changchun, 130061, P.R. China*

Corresponding author: Tel.: +86-431-88502046; Fax: +86-431-88584422; Email:
fanjj03@163.com (**J.-J. Fan**)

E-mail addresses: fanjj03@163.com (**J.-J. Fan**); licai010@126.com (**C. Li**);
wm609@163.com (**W. Wang**); 992702229@qq.com (**Y.-M. Liu**);
xcmxcm1983@126.com (**C.-M Xie**)

Postal address: 2199 Jianshe Street, College of Earth Sciences, Jilin
University, Changchun 130061, China

ABSTRACT

In this paper we present new major and trace element compositions of basaltic rocks in the Gufeng ocean island (GFOI) area in the western segment of the Bangong–Nujiang Suture Zone, northern Tibet. Our aim was to assess the genesis of these rocks and discuss the implications of this new dataset for the evolution of the Bangong–Nujiang Tethyan Ocean. An ocean-island-type double-layer structure comprising a basaltic basement and an oceanic sedimentary cover sequence found within the GFOI provides direct evidence for the interpretation that the assemblage is a typical ocean island. The basalts in the GFOI can be divided into three types (named G1, G2 and G3 basalts), and these basalts range in composition from MORB to OIB types, which is typical of ocean islands. The G1 basalts have MORB-type affinities, possibly indicating the existence of MORB oceanic crust under the GFOI. The G2 basalts represent the early stage of formation of the GFOI, and are produced by the interaction of rising OIB-type basaltic magma and the existing MORB oceanic crust. The G3 basalts are typical OIB basalts and they are the products of the direct eruption of OIB-type basaltic magmas. The G3 basalts have high $(La/Yb)_N$ (12.3–14.4), $(Ce/Yb)_N$ (10.8–11.8), $(La/Sm)_N$ (2.39–2.76), and $(Sm/Yb)_N$ (4.89–5.23) ratios, indicating the presence of oceanic lithosphere below the GFOI with a thickness of 50–60 km. Geochemical analyses of the GFOI cherts show that they contain terrigenous material, indicating the GFOI formed close to a continental margin. Norian conodont fossils within the GFOI limestones indicate the GFOI formed during the Late Triassic. These data, combined with geological evidence and a half-space model of lithosphere cooling, where the thickness of the oceanic

lithosphere is determined from the age of the lithosphere, indicate that the western segment of the Bangong–Nujiang Tethyan Ocean opened initially in the late Permian, expanded rapidly during the Early–Middle Triassic, and was a mature ocean during the Late Triassic.

Key words: Tibetan Plateau; Bangong–Nujiang Suture Zone; Gufeng ocean island; late Permian; Late Triassic; geochemistry

1. Introduction

Ocean islands and ophiolites, which are remnants of oceanic lithosphere, provide direct evidence of the nature of a pre-existing ocean. Because of their greater elevation and lower density compared with ophiolites, ocean islands might be more easily retained in a suture zone during subduction and the closing of an ocean basin (Dobretsov *et al.*, 2004; Cai *et al.*, 2006; Yuan *et al.*, 2009). Further investigation of ocean islands would therefore be important when investigating the evolution of ancient oceans (Dobretsov *et al.*, 2004; Li, 2004; Yan *et al.*, 2008; Pan *et al.*, 2008, 2012; Fan *et al.*, 2014a, b).

However, the difficulties encountered in determining the ages of ocean islands mean that little work has been done on the ocean islands of most ancient suture zones. The Bangong–Nujiang Suture Zone (BNSZ) in China is a typical example of such an ancient suture zone. It lies between the Lhasa Block to the south and the Southern Qiangtang Block to the north (Fig. 1a), and provides a natural laboratory for studying

the features of the Meso-Tethys Oceans Ocean. Although many scholars have conducted research on the strata, volcanic rocks, and ophiolites in the BNSZ since the 1980s(Pan, 1983; Wang *et al.*, 1987; Dewey *et al.*, 1988; Bureau of Geology and Mineral Exploration of Tibet Province, 1993; Pan *et al.*, 1997, 2006; Yin and Harrison., 2000; Shi *et al.*, 2012; Fan *et al.*, 2014a, b; Zhang *et al.*, 2014), there are many uncertainties regarding the evolution of the suture zone, such as the processes involved in the opening and early evolution of the Bangong–NujiangTethyan Ocean.

Although most scholars have proposed that the Bangong–NujiangTethyan Ocean opened in the Late Triassic(Chang and Cheng, 1973; Pan, 1983; Allègre *et al.*, 1984; Pan *et al.*, 1997; Yin and Harrison, 2000; Kapp *et al.*, 2003; Qiangba *et al.*, 2009), others have suggested the Carboniferous–Permian(Pan *et al.*, 2006; Zhu *et al.*, 2013), late Permian to Early Triassic(Ren and Xiao., 2004; Fan *et al.*, 2014b, 2015b), Early Jurassic(Qiu *et al.*, 2004; Xia *et al.*, 2008; Qiu *et al.*, 2010),and Middle to Late Jurassic(Wang *et al.*, 1987; Zhang, 2007). The oldest reliable ages of oceanic rocks in the BNSZ are Early Jurassic(Qiu *et al.*, 2004; Xia *et al.*, 2008; Qiu *et al.*, 2010), and the general absence of earlier oceanic remnants has left us with little information on when and how the early Bangong–NujiangTethyan Ocean developed.

In this paper, we describe a newly found ocean island in the western segment of the BNSZ. Our research has shown that this ocean island formed during the Late Triassic. The existence of a Late Triassic ocean island indicates that the ocean already existed at that time, suggesting an earlier opening age than the Late Triassic proposed by previous scholars(Chang and Cheng, 1973; Pan, 1983; Allègre *et al.*, 1984; Pan *et*

al., 1997; Yin and Harrison., 2000; Kapp *et al.*, 2003; Qiangba *et al.*, 2009). Here, we focus on the Late Triassic Gufeng ocean island and present the results of detailed petrological and geochemical analyses of its basalts and cherts. We discuss the genesis of these rocks and implications for the opening and early evolution of the Bangong–Nujiang Tethyan Ocean.

2. Geological background

The Tibetan Plateau is located in the eastern section of the Alpine–Himalayan tectonic domain and has a complex geological history that includes the formation and evolution of the Paleo-, Meso- and Neo-Tethys Oceans (Pan *et al.*, 1997; Yin and Harrison, 2000; Kapp *et al.*, 2005; Zhu *et al.*, 2013). The Tethys Ocean closed along four main suture zones that have been identified on the Tibetan Plateau (from south to north): the Indus–Yarlung Zangbo suture zone (IYZSZ), the Bangong–Nujiang suture zone (BNSZ), the Longmuco–Shuanghu–Lancangjiang suture zone (LSLSZ), and the Jinshajiang suture zone (JSSZ) (Fig. 1a). These suture zones divide the plateau from south to north into the Himalayan, Lhasa, Southern Qiangtang, Northern Qiangtang, and Bayan Har–Songpan Garze blocks (Fig. 1a; Yin and Harrison, 2000; Li *et al.*, 2006; Pan *et al.*, 1997, 2006, 2012; Wang *et al.*, 2013; Zhai *et al.*, 2013, 2016; Zhu *et al.*, 2013).

The BNSZ crosses the middle of the Tibetan Plateau in an east–west direction (Fig. 1a). The suture zone starts in Kashmir and extends eastwards for 2400 km through the Bangong Co, Rutog, Gerze, Dongqiao, Dingqing, and Jiayuqiao areas before

continuing into Burma, Thailand, and Malaysia. The BNSZ in China is divided into three segments, which from west to east are the Bangong–Gerze, Gerze–Dingqing, and Dingqing–Nujiang segments(Pan *et al.*, 1997, 2006; Qiu *et al.*, 2004).

The present study area is located in the Gufeng area, 100 km northwest of Gerze County, in the western segment of the BNSZ(Fig. 1b). The area contains ocean island fragments, ophiolites, granitoids, and sediments of the Mugangri Group, Quse Formation, and Riganpeico Formation (Fig. 1b). The Jurassic Mugangri Group consists of bathyal to abyssal flysch deposits (mainly graywacke and shale) that formed within the Bangong–NujiangTethyan Ocean(Bureau of Geology and Mineral Exploration of Tibet Province, 1993; Fan *et al.*, 2015a). The Early Jurassic Quse Formation consists mainly of sandstone deposited on a continental slope. The Late Triassic Riganpeico Formation consists mainly of shallow marine limestones that formed in the Bangong–NujiangTethyan Ocean. Early Cretaceousgranitoids are widespread throughout the study area and intrude the Mugangri Group(Li *et al.*, 2008, 2011; Yu *et al.*, 2011; Li *et al.*, 2012; Chen *et al.*, 2013).

The ophiolites in the study area are dominated by peridotites, gabbros, and basalts, and have not been described previously.The remnant ocean island in the study area is called the Gufeng ocean island (GFOI; Fig. 2a, b), and is dominated by basalt, limestone, chert(Fig. 2c), and conglomerate (Fig. 2d).

3. Field and petrographic observations

The transect across the Gufeng area provides a representative section through the geology of the GFOI(Fig.3). The GFOI was thrust onto the Jurassic Mugangangri Group(Fig. 2aand Fig. 3), forming nappes that are exposed within klippe in the Gufeng area. This tectonic activity resulted in strong deformation, with numerous chevron folds developed in the GFOI(Fig. 2e and Fig. 3), but the island still retains its double-layered structure comprising a basaltic basement and an oceanic sedimentary cover sequence.

The dominant lithologies encountered within the transect across the GFOI are as follows.

(1) Basalt. The basalts within the GFOI are usually gray–green in color, and generally occur as either pillow basalt(Fig. 2f) or massive basalt(Fig. 2c, g). The pillows are generally 60 × 100 cm in size (Fig. 2f)and the massive basalts occur as individual units with thicknesses of generally >2 m(Fig. 2c, g). Both the pillow basalts and the massive basalts are porphyritic, with pyroxene and plagioclase phenocrysts set in an intergranular matrix(Fig. 2h), and they have undergone greenschist-facies metamorphism, resulting in some pyroxene phenocrysts being altered to chlorite (Fig. 2i). In addition, they commonly contain calcite-bearing amygdales that make up ≤10% of the rock(Fig. 2h).

(2) Limestone. The limestones within the Gufeng transect are in conformable contact with basalts(Fig.2g,j) and can be divided into two types. The first type forms thick layers that record deposition in an isolated, clear-water sedimentary environment

associated with a carbonate platform(Fig.2c). The second type forms thin layers (<20 cm) interbedded with layers of basalt(Fig.2j)or chert(Fig.2k) of similar thickness.

(3) **Chert.**The cherts within the Gufeng transect are in conformable contact with basalts(Fig.2c)and limestones (Fig. 2l), and are mostly thin-bedded (<10 cm thick) and gray to gray–black in color (Fig. 2l). They are laterally extensive, which is typical of cherts that form in an ocean basin.

(4) **Conglomerate.** The conglomerates within the Gufeng transect can be divided into basaltic conglomerates, in which both the gravel and matrix are made of basaltic debris, and another conglomerate that contains minor amounts of limestone in addition to the basaltic debris (Fig.2d). The significant deformation in this area means that the limestone clasts within the latter type of conglomerate have been stretched. These conglomerates provide evidence of a sedimentary environment with a restricted provenance, proximal deposition, a far-shore-type setting, and rapid accumulation of the sediments in the ocean surrounding the island.

The double-layer structure of the **GFOI** comprises a basaltic basement and an oceanic sedimentary cover sequence (limestone, chert, and conglomerate; Fig. 2b), interbedded limestone and basalt(Fig.2g, j), limestone with chert (Fig.2k), and two types of conglomerate (Fig.2d), which together provide direct evidence that the **GFOI** was a typical ocean island that formed in an ocean basin.

4. Analytical methods

Nine pillow basalt samples, 15 massive basalt samples and 10 chert samples from

the **GFOI** (Fig. 1b and Fig. 2) were collected for whole-rock geochemical analyses. Before analysis, samples were trimmed to remove weathered surfaces, cleaned with deionised water, crushed, and then powdered in an agate mill. Major element compositions were determined using inductively coupled plasma–optical emission spectroscopy (ICP–OES; Leeman Prodigy with high-dispersion Echelle optics) at China University of Geosciences, Beijing, China. Loss on ignition (LOI) values were determined by heating 1 g of sample in a furnace at 950°C for several hours before being cooled in a desiccator and reweighed. Analytical uncertainties are generally better than 1wt% for all major elements. Trace element analyses were performed using inductively coupled plasma–mass spectrometry (ICP–MS; Agilent-7500a) at China University of Geosciences. Analytical accuracy and precision during trace element analyses were monitored by analyzing the standards AGV-2 (US Geological Survey) and GSR-3 (National Geological Standard Reference Materials of China). Analytical accuracy, as indicated by relative differences between measured and recommended values, is better than 5wt% for most elements, and analysis of the AGV-2 and GSR-3 international standards was in good agreement with recommended values (Govindaraju, 1994). For details of the analytical procedures, see Zhai *et al.* (2013) and Hu *et al.* (2013).

5. Results

5.1. Whole-rock major and trace element geochemistry

5.1.1. Basalts

Whole-rock major and trace element data for the basalts of the GFOI are given in Tables 1–3. The GFOI basalts contain calcite amygdals and have undergone lower-greenschist-facies metamorphism, resulting in a range of loss on ignition (LOI) values and changes in the concentrations of mobile elements (e.g., Na, K, Ca, Cs, Rb, Ba, and Sr) compared with protolith values. However, the concentrations and ratios of immobile elements (e.g., the rare earth elements and high field strength elements) and transition metals (e.g., V, Ni, Cr, and Fe) have not been affected by these processes, and can therefore be used to classify and determine the petrogenesis and tectonic setting of the samples.

The GFOI basalts can be divided into three types based on their geochemical characteristics. Basalts of the first type (named G1 basalts), which consist of eight samples of pillow basalt, have high contents of SiO₂ (46.8–57.3 wt%), Al₂O₃ (12.8–15.3 wt%), and Fe₂O₃ (10.3–13.9 wt%), intermediate contents of Na₂O (0.89–4.20 wt%), MgO (4.16–5.40 wt%), and TiO₂ (1.09–1.30 wt%), low Mg[#] values (44.2–50.7), and very low contents of K₂O (0.03–0.06 wt%). They contain TiO₂ concentrations (1.09–1.30 wt%) that are higher than those of island-arc volcanic rocks (0.58–0.85 wt%), but close to the concentrations in mid-ocean-ridge basalts (MORB; 1.0–1.5 wt%; Hishashi, 1954; Irvine *et al.*, 1971; Winchester *et al.*, 1977; Pearce, 1983;

Li, 1993). The G1 basalts have low total rare earth element (REE) concentrations (33–44 ppm), and they have flat chondrite-normalized REE (Fig. 4a; Sun and McDough, 1989) and primitive-mantle-normalized trace element patterns (Fig. 4b; Sun and McDough, 1989) that are similar to those of N-MORB basalts. The G1 basalts have Nb contents (1.38–2.06 ppm) and Nb/La ratios (0.49–0.78) that are also similar to those of N-MORB basalts (Nb < 3 ppm, Nb/La = 0.68; Pearce, 1983; Irvine *et al.*, 1971; Winchester *et al.*, 1977; Crawford, 1989; Kimura *et al.*, 2002), and all the samples lie in the normal mid-ocean-ridge field on the Ti/100 vs. Zr vs. (Y × 3) and (Nb × 2) vs. Zr/4 vs. Y immobile element discrimination diagrams (Fig. 5a, b). These data indicate that the G1 basalts are MORB-type basalts.

Basalts of the second type (named G2 basalts), which consist of one sample of pillow basalt and eight samples of massive basalt, have variable contents of SiO₂ (42.7–55.1 wt%), Al₂O₃ (11.2–15.3 wt%), Fe₂O₃ (7.55–11.8 wt%), and CaO (6.69–18.8 wt%). They have high contents of TiO₂ (1.22–1.76 wt%), Na₂O (2.82–4.35 wt%), and MgO (4.12–8.24 wt%), high Mg[#] values (55.2–64.9), and low K₂O (0.17–1.53 wt%) contents. They have higher contents of REE (65–87 ppm) than the G1 basalts (33–44 ppm), and have chondrite-normalized REE patterns that are enriched in light rare earth elements (LREEs; La_N/Yb_N = 1.91–4.74; Fig. 4a; Sun and McDough, 1989). In addition, they show weak enrichments in highly incompatible elements (e.g., high field strength elements [HFSEs]: Nb, Ta, and Ti) in primitive-mantle-normalized multi-element variation diagrams (Fig. 4b; Sun and McDough, 1989). The compositions of the G2 basalts fall between those of enriched

MORB (E-MORB) and ocean island basalt (OIB; Fig.4a, b), and lie in the intraplate basalt and E-type MORB field on both the Ti/100 vs. Zr vs. (Y × 3) and (Nb × 2) vs. Zr/4 vs. Y immobile element discrimination diagrams (Fig. 5a, b).

Basalts of the third type (G3 basalts), which consist of seven samples of massive basalt, also have variable contents of SiO₂ (43.6–49.4 wt%), Al₂O₃ (12.5–16.4 wt%), and CaO (6.79–10.8 wt%). They have higher contents of TiO₂ (2.41–2.64 wt%), Na₂O (3.82–5.40 wt%), and Fe₂O₃ (10.4–12.6 wt%), lower contents of MgO (3.95–6.92 wt%), and lower Mg[#] values (46.2–55.7) than the G2 basalts. The G3 basalts have higher concentrations of TiO₂ (2.41–2.64 wt%) than island-arc volcanic rocks (0.58–0.85 wt%) and MORB (1–1.5 wt%), similar to the concentrations in OIB (2.20 wt%; Hishashi, 1954; Irvine *et al.*, 1971; Winchester *et al.*, 1977; Pearce, 1983; Li, 1993). G3 basalts have significantly higher total REE concentrations (149–178 ppm) than the G1 (33–44 ppm) and G2 basalts (65–87 ppm). The G3 basalts also have LREE-enriched chondrite-normalized REE patterns (La_N/Yb_N = 12.3–14.4; Fig.4a; Sun and McDough, 1989), with low concentrations of the heavy REEs (HREEs) and no Eu anomaly, similar to typical OIB compositions. They show enrichment in HFSEs in primitive-mantle-normalized trace element variation diagrams, similar to OIB (Fig.4b; Sun and McDough, 1989). In addition, all the G3 basalt samples plot in the intraplate alkali basalt field on both the Ti/100 vs. Zr vs. (Y × 3) and (Nb × 2) vs. Zr/4 vs. Y immobile element discrimination diagrams (Fig. 5a, b). These data indicate that the G3 basalts represent typical OIB-type magmas.

The G1 basalts plot along the boundary between the sub-alkaline basalt and

basaltic andesite fields on the Zr/TiO₂ vs. Nb/Y immobile element classification diagram (Fig. 6; Winchester *et al.*, 1977), whereas G2 basalts plot on the boundary between the sub-alkali basalt and alkali basalt fields, and G3 basalts plot in the alkali basalt field (Fig. 6).

5.1.2. Cherts

Whole-rock major and trace element data for the GFOI cherts are given in Table 4. The cherts have high contents of SiO₂ (84.9–95.0 wt%), Al₂O₃ (0.58–2.16 wt%), Fe₂O₃ (0.32–1.25 wt%), and CaO (0.42–6.73 wt%), and low contents of TiO₂ (0.04–0.20 wt%), MnO (0.01–0.08 wt%), MgO (0.19–0.88 wt%), Na₂O (0.04–0.62 wt%), and K₂O (0.00–0.32 wt%). They have variable contents of the REEs (20–68 ppm) and they have North American shale (NASC)-normalized REE patterns with weak negative Ce anomalies (Ce* = 0.54–0.98; Fig. 7), similar to the Ce-anomalies of cherts that form on a continental margin or in a pelagic setting (Murray, 1994).

6. Discussion

6.1. Age of the GFOI

Five conodont fossils of *Epigondolella abneptis* subspecies A were found in the GFOI limestone (Fig. 8). This subspecies is typical of the Norian (Michael, 1983), and is widespread in the Lizhou, Yawa, and Dibuco areas of the Gangdese region on the Tibetan Plateau (Jiet *et al.*, 2003, 2006, 2010). The *Epigondolella abneptis* subspecies A

fossils in the **GFOI** indicate, therefore, that the **GFOI** formed during the Norian of the Late Triassic (227–209 Ma).

6.2. Petrogenesis

6.2.1. Crustal contamination

Before discussing magma sources, it is necessary to consider whether the basalts analyzed in this study were affected by crustal contamination. Th and Ta are sensitive indicators of crustal contamination, because such contamination produces a rapid increase in Th/Ta ratios (Condie, 1993). The **G1**, **G2**, and **G3** basalts of the **GFOI** all have relatively low Th/Ta ratios (2.67–5.20, 1.28–1.92, and 1.18–2.17, respectively), similar to those in volcanic rocks derived from primitive mantle (Th/Ta = 2.3; Condie, 1993), and much lower than those in the upper crust (Th/Ta >10), indicating that the **GFOI** basalts did not undergo crustal contamination. Zr/Nb ratios remain almost constant during fractional crystallization, but change significantly during magma mixing or crustal contamination (Weaver *et al.*, 1996). Although the **G1**, **G2**, and **G3** basalts have a wide range of SiO₂ contents (46.8–57.3, 42.7–55.1, and 43.6–49.4 wt%, respectively), each type of basalt has only a small range in Zr/Nb ratios (29.7–37.8, 4.35–8.69, and 2.92–5.70, respectively), and this is consistent with the **GFOI** basalts not being contaminated by the crust.

6.2.2. Magma source

Various studies have shown that basaltic magmas are commonly derived from

the partial melting of mantle peridotite, and that their REE patterns are controlled mainly by the contents of garnet and spinel in the source mantle, rather than by the contents of olivine, clinopyroxene, or orthopyroxene, or by pressure and temperature (Beattie *et al.*, 1994; McKenzie and O’Nions, 1991; Schwandt *et al.*, 1998; Horne *et al.*, 1994). In general, basalts derived from spinel peridotite have flat chondrite-normalized REE patterns with inconspicuous fractionation between the LREEs and HREEs. However, basalts derived from garnet peridotite display obvious fractionation between the LREEs and HREEs, and high $\text{La}_\text{N}/\text{Yb}_\text{N}$ and $\text{Ce}_\text{N}/\text{Yb}_\text{N}$ ratios (Hart *et al.*, 1993; Hauriet *et al.*, 1994).

The chondrite-normalized REE patterns of the **G1** basalts are flat (Fig. 4a), similar to those of basalt derived from spinel peridotite. The low Dy/Yb (1.45–1.57), Sm/Yb (0.78–0.93), and La/Sm (0.84–1.24) ratios of the **G1** basalts, and the fact that they all plot in an area that corresponds to ~30% partial melting of spinel peridotite on the Sm/Yb vs. La/Sm diagram (Fig. 9), further indicate that the magma source of the **G1** basalts was spinel peridotite in the mantle.

The **G3** basalts have LREE-enriched chondrite-normalized REE patterns ($\text{La}_\text{N}/\text{Yb}_\text{N} = 12.3\text{--}14.4$; Fig. 4a), similar to those of basalts derived from garnet peridotite. The high Dy/Yb (2.89–3.18), Sm/Yb (4.40–4.70), and La/Sm (3.70–4.27) ratios of the **G3** basalts, and the fact that they all plot in an area that corresponds to ~5% partial melting of garnet peridotite on the Sm/Yb vs. La/Sm diagram (Fig. 9), further indicate that the magma source of the **G3** basalts was garnet peridotite in the mantle.

The compositions of the **G2** basalts fall between those of the **G1** and **G3** basalts

(Figs 4–6). We infer, therefore, that they formed by the mixing of the G1 and G3 basalts, as also indicated by the fact that the G2 basalts plot in an intermediate position between the G1 basalts and the G3 basalts on the Sm/Yb vs. La/Sm diagram (Fig. 9).

6.2.3. Fractional crystallization

Most of the Cr and Ni concentrations and $Mg^\#$ values in the G1 basalts (Cr = 26.5–68.1 ppm, Ni = 25.0–42.4 ppm, $Mg^\#$ = 44.2–50.7), G2 basalts (Cr = 183–306 ppm, Ni = 25.0–42.4 ppm, $Mg^\#$ = 55.2–64.9), and G3 basalts (Cr = 6.69–245 ppm, Ni = 42.4–130 ppm, $Mg^\#$ = 46.2–55.7) are lower than the values for primary mantle-derived magmas (Cr = 300–500 ppm, Ni = 300–400 ppm, $Mg^\#$ = 68–76; Frey *et al.*, 1978; Hess *et al.*, 1992). This indicates that all the GFOI basalts underwent fractional crystallization of olivine, chromite, and pyroxene (Wilson *et al.*, 1989; Jung *et al.*, 1998). The absence of negative Eu anomalies in the chondrite-normalized REE patterns of the GFOI basalts (Fig. 4a) argues against the fractionation of plagioclase.

In summary, we conclude that the G1 basalts were derived from spinel peridotite, whereas the G3 basalts were derived from garnet peridotite. The G2 basalts were formed by the mixing of the G1 basalts and the G3 magmas. The ascending G1, G2, and G3 magmas underwent varying degrees of fractional crystallization of olivine, chromite, and pyroxene, but the magmas did not experience crustal contamination.

6.3. Tectonic setting

The ocean-island-type double-layered structure comprising a basaltic basement and an oceanic sedimentary cover sequence (Fig. 2b and Fig. 3), the presence of limestones interbedded with basalts (Fig. 2g, j) and limestones interbedded with cherts (Fig. 2k), and the types of conglomerate found within the GFOI (Fig. 2d) provide direct evidence that the GFOI is a typical ocean island.

As discussed in section 5.1.1, the GFOI basalts are divided into the G1, G2, and G3 basalts, and these basalts range in composition from MORB to OIB types, which is a common phenomenon in most modern and ancient ocean islands (Wu *et al.*, 2006; Fan *et al.*, 2014a). The G1 basalts have MORB-type affinities, possibly indicating the existence of a MORB oceanic crust under the GFOI. The G2 basalts, which formed from the mixing of MORB- and OIB-type basaltic material, may represent the early stage of formation of the GFOI, and could be the products of interaction between new OIB-type basaltic magmas and existing MORB oceanic crust. The G3 basalts are typical OIB basalts that represent the mature stage of the GFOI, at which time the OIB magmas were erupted through well-established channels with little further interaction between the rising OIB magmas and the existing MORB oceanic crust.

In conclusion, our petrological and geochemical analyses indicate that the GFOI was a typical ocean island that formed in an ocean basin.

6.3.1. Thickness of the oceanic lithosphere beneath the GFOI

Although previous studies have shown that the geochemistry of OIB-type

magmas can be influenced by various factors, such as compositional variations in the mantle, variations in the mantle potential temperature, and variations in the thickness of the underlying oceanic lithosphere, the factor that exerts a first-order control is the thickness of the oceanic lithosphere, with the other factors having only secondary effects(Niu and Hékinian, 1997; Niu and O'Hara, 2007, 2008; Humphreys and Niu, 2009; Niu, 2013). Although all OIB basalts have garnet signatures with enrichments in LREEs and depletion in HREEs, the intensity of the garnet signature depends on the relative proportion of melt that is produced in the garnet (vs. spinel) peridotitefacies. With decreasing thickness of oceanic lithosphere, the extent of melting increases, with more melt produced by decompression in the spinel peridotitefacies. As a result, the intensity of the garnet signature in an OIB melt is inversely related to the extent of dilution, with less dilution in melts produced by low degrees of melting beneath a thick lithosphere and more dilution in melts produced by high degrees of melting beneath a thin oceanic lithosphere(Niu *et al.*, 1999; Niu and O'Hara, 2008; Humphreys and Niu, 2009; Niu, 2013).

In the case of the **G3** basalts, which represent the primary OIB-type basaltic magmas of the **GFOI**, there is an intense garnet signature, as reflected by the high $(\text{La/Yb})_{\text{N}}$ (12.3–14.4) and $(\text{Ce/Yb})_{\text{N}}$ (10.8–11.8) ratios. In general, $(\text{Sm/Yb})_{\text{N}}$ ratios provide important information on the source of a basaltic magma, and values are as high as 5 in OIB basalts that are derived solely from the garnet peridotitefacies, but as low as 1.7 in OIB basalts if they are diluted by melts produced by decompression in the spinel peridotitefacies(Humphreys and Niu, 2009; Niu, 2013). As for the **G3**

basalts, they have high $(\text{Sm}/\text{Yb})_{\text{N}}$ values (4.89–5.23) that are similar to those of OIB basalts that are fully derived from the garnet peridotite facies, and which are less diluted by melts produced by decompression in the spinel peridotite facies. These features indicate that the underlying oceanic lithosphere of the GFOI was very thick.

Humphrey and Niu (2009) examined island-averaged geochemical data for 115 volcanic islands in the Pacific, Atlantic, and Indian Oceans that have known eruption ages and known ages for the underlying lithosphere (Humphreys and Niu, 2009). These age data allow one to calculate the thickness of the oceanic lithosphere at the time of volcanism. Humphrey and Niu (2009) concluded that the island-averaged $(\text{La}/\text{Sm})_{\text{N}}$ and $(\text{Sm}/\text{Yb})_{\text{N}}$ ratios increase with increasing thickness of oceanic lithosphere (Humphreys and Niu, 2009). The G3 basalts of GFOI have high $(\text{La}/\text{Sm})_{\text{N}}$ (2.39–2.75) and $(\text{Sm}/\text{Yb})_{\text{N}}$ (4.89–5.23) values that are similar to those of modern OIB basalts that formed on an underlying oceanic lithosphere with a thickness of 50–60 km.

6.3.2. Tectonic environment

A chert is a chemically precipitated sedimentary rock, essentially monomineralic, and composed chiefly of microcrystalline and/or chalcedonic quartz, with subordinate coarse-grained quartz (Folk, 1980). Cherts may contain up to 95% or more silica (Hesse, 1988). Cherts are chemically inert and markedly resistant to weathering and alteration, making them suitable for geochemical studies. Their elemental compositions include simple, rare, and trace elements that reflect minor constituents

such as volcanic detritus, terrigenous clasts, and hydrothermal precipitates (Kato and Nakamura, 2003). Consequently, cherts have been used as provenance indicators on the basis of their geochemical characters. Rare earth elements (REEs) are mostly immobile in cherts, and their REE patterns therefore provide evidence of tectonic environment and depositional location (Murray, 1994). Cherts are common in the GFOI, and their geochemical analysis could further enhance our understanding of the tectonic environment of the GFOI.

Murray *et al.* (1990) studied the chert and shale sequences of the Franciscan assemblage in the Claremont and Monterey formations of coastal California, and they identified three depositional regimes: spreading ridge proximal, ocean-basin floor, and continental margin. Back-arc basins, marginal seas, epicontinental seas, and open continental shelves are all listed under the heading of continental margin (Murray, 1994). The chemical compositions of bedded cherts are generally determined by biogenic silica derived from radiolarians and sponge spicules, and modified by clastic material and hydrogenous (authigenic) components. In addition, cherts deposited near spreading ridges are influenced by metalliferous activity. Modification during diagenesis of chemical and isotopic systems is also conceivable.

The high SiO₂ content of bedded radiolarian cherts indicates a greater contribution of biogenic silica whereas high MnO contents indicate greater contributions of hydrogenous components (Sugisaki *et al.*, 1982; Kunimaru *et al.*, 1998). Al₂O₃ is a component derived from terrigenous clastic rocks, and a high Al₂O₃ content in a bedded chert indicates, therefore, a greater contribution of terrigenous material.

Murray (1994) proposed a diagram using shale-normalized (La/Ce) ratios versus the $\text{Al}_2\text{O}_3/(\text{Al}_2\text{O}_3 + \text{Fe}_2\text{O}_3)$ ratio as a chemical depositional criterion that is independent of diagenetic modification. In his (La/Ce) vs. $\text{Al}_2\text{O}_3/(\text{Al}_2\text{O}_3 + \text{Fe}_2\text{O}_3)$ diagram, cherts that formed in ridge-proximal, pelagic, and continental margin environments have (La/Ce)_{SN} (shale-normalized ratios) of 3–4, 1–2.5, and 0.5–1.5, and $\text{Al}_2\text{O}_3/(\text{Al}_2\text{O}_3 + \text{Fe}_2\text{O}_3)$ ratios of 0.05–0.4, 0.4–0.7, and 0.55–0.9, respectively.

Among the REE features of cherts, a negative Ce-anomaly usually implies a pelagic character. Strongly negative Ce-anomalies are commonly observed for deep-ocean water, but not for clastic rocks. Murray *et al.* (1990) and Murray (1994) noted that the Ce-anomaly reflects the depositional location of chert, so that areas proximal to spreading ridges have values of $\text{Ce}/\text{Ce}^* = \text{ca. } 0.3$ and (La/Ce)_{SN} = 3–4, ocean-basin floor or pelagic areas have values of $\text{Ce}/\text{Ce}^* = \text{ca. } 0.5$ and (La/Ce)_{SN} = 1–2.5, and continental margins have values of $\text{Ce}/\text{Ce}^* = \text{ca. } 0.9\text{--}1.3$ and (La/Ce)_{SN} = 0.5–1.4.

In the case of the GFOI cherts, they are mostly thin-bedded (<20 cm bed thickness; Fig. 2c, l) and laterally extensive in the field, which is typical of cherts formed in an ocean basin. The GFOI chert samples have high contents of SiO_2 (84.9–95.0 wt%) and low contents of MnO (0.01–0.08 wt%), indicating a large contribution of biogenic silica, and a small contribution of hydrogenous components. This inference is further supported by the fact that all GFOI chert samples plot in the non-hydrothermal field on the Fe–Al–Mn diagram (Fig. 10a).

The GFOI cherts have high contents of Al_2O_3 (0.58–2.16 wt%), indicating a

significant contribution of terrigenouslastic rock material during their deposition. Slightly negative Ce anomalies (Fig. 7; $Ce^* = 0.54\text{--}0.98$) are similar to the Ce-anomalies of cherts that formed on a continental margin or in pelagic areas, and exclude the possibility of deposition proximal to a spreading ridge. High $Al_2O_3/(Al_2O_3 + Fe_2O_3)$ ratios (0.58–0.76) and low $(La/Ce)_{SN}$ (0.90–1.77) ratios indicate that these cherts were deposited close to a continental margin, and this inference is also supported by the fact that all the GFOI chert samples plot in the continental margin fields of the $100 \times (Fe_2O_3/SiO_2)$ vs $100 \times (Al_2O_3/SiO_2)$, $Fe_2O_3/(100 - SiO_2)$ vs. $Al_2O_3/(100 - SiO_2)$, and Fe_2O_3/TiO_2 vs. $Al_2O_3/(Al_2O_3 + Fe_2O_3)$ diagrams (Fig. 10b–d).

In summary, we conclude that the GFOI was located on an oceanic lithosphere that had a thickness of 50–60 km. This ocean island was probably formed close to a continental margin.

6.4.A tectonic model: implications for the opening and early evolution of the Bangong–NujiangTethyan Ocean

As mentioned above, most scholars have proposed that the Bangong–NujiangTethyan Ocean opened during the Late Triassic (Chang and Cheng, 1973; Pan, 1983; Allègre *et al.*, 1984; Pan *et al.*, 1997; Yin and Harrison., 2000; Kapp *et al.*, 2003; Qiangba *et al.*, 2009), but other possible opening times have been suggested including Carboniferous–Permian (Pan *et al.*, 2006; Zhu *et al.*, 2013), late Permian to Early Triassic (Ren and Xiao., 2004; Fan *et al.*, 2014b, 2015b), Early

Jurassic(Qiuet *et al.*, 2004; Xia *et al.*, 2008; Quet *et al.*, 2010),and Middle to Late Jurassic(Wang *et al.*, 1987; Zhang, 2007). And until recently, the oldest reliable age of oceanic remains in the BNSZ was Early Jurassic(Qiuet *et al.*, 2004; Xia *et al.*, 2008; Quet *et al.*, 2010), which meant we had been left in a state of great uncertainty about how the Bangong–NujiangTethyan Ocean actually developed in its early stages of opening.

Ocean islands are commonly found in both ancient and modern ocean basins, and they generally form in oceanic basins with a mature oceanic crust(Cai *et al.*, 2006; Lai *et al.*, 2010; Fan *et al.*, 2014a). Therefore, the discovery of the Late Triassic GFOI in the BNSZ, suggests that at least by the Late Triassic, the Bangong–NujiangTethyan Ocean had developed as a mature ocean. This implies that the ocean had already opened well before the Late Triassic, which is earlier than the timing of opening proposed by most previous researchers(Chang and Cheng, 1973; Pan, 1983; Allègre *et al.*, 1984; Pan *et al.*, 1997; Yin and Harrison., 2000; Kapp *et al.*, 2003; Qiangba *et al.*, 2009).

The Late Triassic NorianGFOI (227–209 Ma) is located near the northern margin of the BNSZ (Fig. 1a), and it formed close to a continental margin, and from this we infer that it formed near the northern margin of the Bangong–NujiangTethyan Ocean. Numerous studies have shown that northwards subduction of the Bangong–NujiangTethyan oceanic lithosphere did not occur during the Late Triassic(Quet *et al.*, 2009; Kapp *et al.*, 2005; Pullen *et al.*, 2011; Du *et al.*, 2011; Li *et al.*, 2014; Liu *et al.*, 2014, 2016; Fan *et al.*, 2014a, 2015b). The oceanic lithosphere that

underlies the **GFOI** could therefore represent an early stage within the Bangong–Nujiang Tethyan Ocean, with an age that is younger than but close to the age of the initial opening of the ocean.

According to the theory of plate tectonics, oceanic lithosphere initially forms at a mid-ocean ridge and becomes cooler and thicker as it moves away from the ridge (Oxburgh and Parmentier, 1977). The thickness of the oceanic lithosphere can be determined from the age of the lithosphere using the half-space lithosphere cooling model; i.e., $T = 11 \times t^{1/2}$ (where T is lithosphere thickness in km, and t is the age in Ma). The model is reliable for lithosphere younger than ~70 Myr (Parsons and Sclater, 1977; Phipps Morgan and Smith, 1992; Stein and Stein, 1992). Because oceanic lithosphere reaches its full thickness at an age of ~70 Myr, we assume a constant thickness of ~90 km (i.e., $11 \times 70^{1/2} = 92$ km) for older lithosphere (Humphreys and Niu, 2009).

Because the oceanic lithosphere below an ocean island existed before the eruption of the OIB magma and the thickness of the regional oceanic lithosphere does not change with local OIB volcanism (Parsons and Sclater, 1977; Phipps Morgan and Smith, 1992; Stein and Stein, 1992; Niu and Hékinian, 1997; Niu and O'Hara, 2007, 2008; Humphreys and Niu, 2009; Niu, 2013), the half-space lithosphere cooling model can be used to calculate the age or thickness of oceanic lithosphere below an ocean island at the time of OIB volcanism (Humphreys and Niu, 2009; Niu, 2013). Applying this concept to the **GFOI**, we obtain a thickness of 50–60 km for the oceanic lithosphere that underlies the **GFOI**. We then used the average thickness of the

oceanic lithosphere below the **GFOI** (55 km) to calculate the age of the oceanic lithosphere below the **GFOI** at the time the **GFOI** formed in the Late Triassic, and we obtained an age of 25 Ma. Given that the present-day age of the Late Triassic–Norian is 227–209 Ma we can propose that the initial opening of the Bangong–Nujiang Tethyan Ocean was earlier than the Early Triassic (252–233 Ma, i.e., 227–209 Ma + 25 Ma), though probably not a lot earlier. We infer, therefore, that the western segment of the Bangong–Nujiang Tethyan Ocean opened initially in the late Permian. This inference is further supported by the following lines of evidence. (i) The angular unconformity between the late Permian and middle Permian strata formed in both the Lhasa block (i.e., the angular unconformity between the middle Permian Xiala Formation and the overlying upper Permian Jianzhanong and Dibucuo formations) and the Southern Qiangtang block (i.e., the angular unconformity between the middle Permian Longge Formation and the overlying upper Permian Jipuria Formation; [Xia and Liu., 1997](#); [Zhou et al., 2002](#); [Wang et al., 2013](#); [Zhang et al., 2014](#)). This indicates the occurrence of a tectonic event that resulted in intense uplift of the surrounding continental regions of the BNSZ during the late Permian. We infer, therefore, that this tectonic event corresponds to the initial opening of the Bangong–Nujiang Tethyan Ocean ([Fan et al., 2015b](#)). (ii) Detrital zircons from deposits in the Early Cretaceous Zhaga Formation, as well as inherited zircons from basalts and gabbros of the Zhonggang ocean island and Kangqiong ophiolites, yield ages that peak at 247–255 Ma ([Fan et al., 2015a, b](#)), also indicating an intense volcanic event in the areas surrounding the BNSZ during the late Permian–Early Triassic.

In summary, we conclude that the western segment of the Bangong–NujiangTethyan ocean opened initially in the late Permian(Fig. 11a)and had opened substantially by the Early Triassic. It then expanded rapidly during the Early–Middle Triassic and developed into a mature ocean by the Late Triassic (Fig. 11b).

7. Conclusions

(1) The Late Triassic Gufeng ocean island (GFOI) is a typical ocean island that formed within the Bangong–NujiangTethyan ocean basin.

(2) The GFOI basalts can be divided into G1 basalts with MORB-type characteristics, G2 basalts that formed by the mixing of G1basalts and G3basaltic magma, and G3 basalts with OIB-type characteristics. The G1 basalts represent the pre-existing MORB oceanic crust that underlies the GFOI, and the G2 basalts represent the early stage of GFOI formation and were produced by the interaction of rising OIB-type basaltic magma and the existing oceanic crust. The G3 basalts represent the mature stage of GFOI formation and they are the products of the direct eruption of OIB-type basaltic magmas.

(3) The GFOI was located on an oceanic lithosphere that had a thickness of 50–60 km.

(4) The GFOI chertswere significantly affected by terrigenous material during their deposition, indicating that GFOIprobably formed close to a continental margin.

(5) The western segment of the Bangong–NujiangTethyan Ocean opened in the

late Permian, and expanded rapidly during the Early–Middle Triassic. The ocean continued to develop as a mature ocean during the Late Triassic.

Acknowledgments

We thank MrHao Wu, MrTianyu Zhang, MrQiangyuan Jiang, and MrJianxinXu for their help in the field. This research was supported by the China Postdoctoral Innovative Talent Support Program (Grant No. BX201600061), the National Science Foundation of China (Grant Nos. 41272240 and 41402190) and the China Geological Survey project (Grant Nos. DD20160026 and 121201010000150014).

References

- Allègre, C.J., Courtillot, V., Tapponnez, P., other 32 co-authors., 1984. Structure and evolution of the Himalaya–Tibet orogenic belt. *Nature* 307, 17–22.
- Beattie, P., 1994. Systematics and energetics of trace—element partitioning between olivine and silicate melts: Implications for the nature of mineral/melt partitioning. *Chemical Geology* 117, 57–71.
- Bureau of Geology and Mineral Exploration of Tibet Province, 1993. Regional geology of Tibet Province. Beijing: Geological Publishing House, 1–707 (in Chinese with English abstract).
- Cai, X.F., Cai, H.L. and Liu, D.M., 2006, Stratigraphic sequences of mixtite series in oceanic island of orogenic belts and the significance of the study. *Marine Geology Letters* 22, 9–11 (in Chinese with English abstract).
- Chang, C.F., Cheng, H.L., 1973. Some tectonic features of the Mt. JolmoLungma area, southern Tibet, China. *Scientia Sinica* 16, 257.
- Chen, H.A., Zhu, X.P., Ma, D.F., Huang, H.X., Li, G.M., Li, Y.B., Li, Y.C., Wei, L.J.,

592 Liu, C.Q., 2013. Geochronology and geochemistry of the Bolong porphyry
 593 Cu-Au deposit, Tibet and its mineralizing significance. *ActaGeologicaSinica*
 594 87, 1593–1611 (in Chinese with English abstract).
 595 Condie, K.C., 1993. Chemical composition and evolution of the upper continental
 596 crust: Contrasting results from surface samples and shales. *Chemical*
 597 *Geology* 104, 1–37.
 598 Crawford, A.J., Falloon, T.J., Green, D.H., 1989. Classification, progress and
 599 tectonics setting of Boninites. In: Crawford A J, ed. *Boninites and Related*
 600 *Rocks*. London: Unwyn Hyman, 1–49.
 601 Dewey, J.F., Shackelton, R.M., Chang, C., Sun, Y., 1988. The tectonic evolution of the
 602 Tibetan plateau. *Philosophical Transactions of the Royal Society of London*
 603 A327, 379–413.
 604 Dobretsov, N.L., Buslov, M.M., Yu, U., 2004. Fragments of oceanic islands in
 605 accretion–collision areas of Gorny Altai and Salair Southern Siberia Russia:
 606 Early stages of continental crustal growth of the Siberian continent in
 607 Vendian–Early Cambrian time. *Journal of Asian Earth Sciences*, 23, 673–
 608 690.
 609 Du, D.D., Qu, X.M., Wang, G.H., Xin, H.B., Liu, Z.B., 2011. Bidirectional
 610 subduction of the Middle Tethys oceanic basin in the west segment of
 611 Bangonghu–Nujiang suture, Tibet: evidence from zircon U–Pb LA-ICP-MS
 612 dating and petrogeochemistry of arc granites. *ActaPetrologicaSinica* 27,
 613 1993–2002 (in Chinese with English abstract).
 614 Fan, J.J., Li, C., Liu, Y.M., Xu, J.X., 2015a. Age and nature of the late Early
 615 Cretaceous Zhaga Formation, northern Tibet: constraints on when the
 616 Bangong–Nujiang Neo–Tethys Ocean closed. *International Geology Review*
 617 57, 342–353.
 618 Fan, J.J., Li, C., Xie, C.M., Wang, M., 2014b. Petrology, geochemistry, and
 619 geochronology of the Zhonggang ocean island, northern Tiber: implications
 620 for the evolution of the Bangongco–Nujiang oceanic arm of Neo–Tethys.
 621 *International Geology Review* 56, 1504–1520.

622 Fan, J.J., Li, C., Xie, C.M., Wang, M., Chen, J.W., 2015b. The evolution of the
 623 Bangong–Nujiang Neo–Tethys ocean: evidence from zircon U–Pb and
 624 Lu–Hf isotopic analyses of Early Cretaceous oceanic islands and ophiolites.
 625 *Tectonophysics* 655, 27–41.

626 Fan, J.J., Li, C., Xu J.X., Wang, M., 2014a. Petrology, geochemistry and geological
 627 significance of the Nadongoccean island, Banggongco–Nujiang suture,
 628 Tibetan Plateau. *International Geology Review* 56, 915–928.

629 Folk, R.L., 1980. *Petrology of Sedimentary Rocks*. Hemphill, Austin, TX, 79.

630 Frey, F.A., Green, D.H., Roy, S.D., 1978. Integrated models of basalt petrogenesis: a
 631 study of quartz tholeiites to olivine melilitites from South Eastern Australia
 632 utilizing geochemical and experimental petrological data. *Journal of*
 633 *Petrology* 19, 463–513.

634 Govindaraju, K., 1994. Compilation of working values and sample description for 383
 635 geostandards. *Geostandards Newsletter* 18, 1–158.

636 Hart, S.R., Dunn, T., 1993. Experimental cpx/melt partitioning of 24 trace elements.
 637 *Contributions to Mineralogy and Petrology* 113, 1–8.

638 Hauri, E.H., Wagner, T.P., Grove, T.L., 1994. Experimental and natural partitioning of
 639 Th, U, Pb and other trace elements between garnet, clinopyroxene and
 640 basaltic melts. *Chemical Geology* 117, 149–166.

641 Hess, P.C., 1992. Phase equilibria constraints on the origin of ocean floor basalts.
 642 *Geophysical Monograph* 71, 67–102.

643 Hesse, R., 1988. Origin of chert: diagenesis of biogenic siliceous sediments.
 644 *Geoscience Canada* 15, 171–192.

645 Hishashi, K., 1954. Geology and petrology of O–Shima Volcano. *International*
 646 *Geology Review* 1, 48–59.

647 Horn, I., Foley, S.F., Jackson, S.E., Jenner, G.A., 1994. Experimentally determined
 648 partitioning of high field strength and selected transition elements between
 649 spinel and basaltic melt. *Chemical Geology* 117, 193–218.

650 Hu, P.Y., Li, C., Wang, M., Xie C.M., Wu, Y.W., 2013. Cambrian volcanism in the
 651 Lhasa terrane, southern Tibet: Record of an early Paleozoic Andean-type

652 magmatic arc along the Gondwana proto-Tethyan margin. *Journal of Asian*
653 *Earth Sciences* 77, 91–107.

654 Humphreys, E.R., Niu, Y.L., 2009. On the composition of ocean island basalts (OIB):
655 The effects of lithospheric thickness variation and mantle
656 metasomatism. *Lithos* 112, 118–136.

657 Irvine, T.N., Baragar, W.A.R., 1971. A guide to the chemical classification of the
658 common volcanic rocks. *Canadian Journal of Microbiology* 8, 523–548.

659 Ji, Z.S., Yao, J.X., Wu, G.C., 2010. Revision on the Age of the Mamuxia Formation at
660 the Yawa Area, Coqin County, Tibet and Its Geological
661 Significances. *Acta Geologica Sinica* 84, 1379–1388 (in Chinese with English
662 abstract).

663 Ji, Z.S., Yao, J.X., Wu, G.C., Liu, G.Z., 2006. Discovery of Late Triassic
664 Norian Epigondolella in the “Xiala Formation” in the Dibu Co area, Coqin
665 County, Tibet, China. *Geological Bulletin of China* 25, 138–141 (in Chinese
666 with English abstract).

667 Ji, Z.S., Yao, J.X., Yang, X.D., Zang, W.S., Wu, G.C., 2003. Conodont zonation of
668 Norian in Lhasa area, Xizang (Tibet) and their global
669 correlation. *Acta Palaeontologica Sinica* 42, 382–392 (in Chinese with English
670 abstract).

671 Jung, S., Mesberg, P., 1998. Major and trace-element systematics and isotope
672 geochemistry of Cenozoic mafic volcanic rocks from the Vogelsberg (central
673 Germany) Constraints on the origin of continental alkaline and tholeiitic
674 basalts and their mantle sources. *Journal of Volcanology and Geothermal*
675 *Research* 86, 151–177.

676 Kapp, P., Murphy, M.A., Yin, A., Harrison, T.M., Ding, L. and Guo, J.H., 2003,
677 Mesozoic and Cenozoic tectonic evolution of the Shiquanhe area of western
678 Tibet. *Tectonics* 22, 3–23.

679 Kapp, P., Yin, A., Harrison, T.M., Ding, L., 2005. Cretaceous–Tertiary shortening,
680 basin development, and volcanism in central Tibet. *Geological Society of*
681 *America Bulletin* 117, 865–878.

682 Kato, Y., Nakamura, K., 2003. Origin and global tectonic significance of
683 Early Archean cherts from the Marble Bar greenstone belt, Pilbara Craton,
684 Western Australia. *Precambrian Research* 125, 191–242.

685 Kimura, J.I., Manton, W.I., Sun, C.H., Iizumi, S., Yoshida, T., J. stern, R., 2002.
686 Chemical diversity of the Ueno BASALTS, Central Japan: identification of
687 mantle and crustal contributions to arc basalts. *Journal of Petrology* 43, 1923
688 —1946.

689 Kunimaru, T., Shimizu, H., Takahashi, K., Yabuki, S., 1998. Differences in
690 geochemical features between Permian and Triassic cherts from the Southern
691 Chichibu terrane, southwest Japan: REE abundances, major element
692 compositions and Sr isotopic ratios. *Sedimentary Geology* 119, 195–217.

693 Lai, S.C., Qin, J.F., Li, X.J and Zang, W.J., 2010, Geochemistry of the OIB-type
694 basalt from the Wumulong-Tongchangjie area, Changning-Menglian suture
695 zone: its petrogenesis and tectonic implications. *Earth Science Frontiers* 17, 44
696 —52 (in Chinese with English abstract).

697 Li, C., Huang, X.P., Zhai, Q.G., Zhu, T.X., Yu Y.S., Wang G.H., Zeng Q.G., 2006. The
698 Longmu Co–Shuanghu–Jitang plate suture and the northern boundary of
699 Gondwanaland in the Qinghai–Tibet plateau. *Earth Science Frontiers* 13,
700 136–147 (in Chinese with English Abstract).

701 Li, J.L., 2004. Basic characteristics of accretion-type orogens. *Geological Bulletin of*
702 *China* 23, 947–951 (in Chinese with English abstract).

703 Li, J.X., Li, G.M., Qin, K.Z., Xiao, B., 2008. Geochemistry of porphyries and
704 volcanic rocks and ore-forming geochronology of Duobuza gold-rich
705 porphyry copper deposit in Bangonghu belt, Tibet: Constraints on
706 metallogenic tectonic settings. *Acta Petrologica Sinica* 24, 531–543 (in
707 Chinese with English abstract).

708 Li, J.X., Qin, K.Z., Li, G.M., Richards, J.P., Zhao, J.X., Cao, M.J., 2014.
709 Geochronology, geochemistry, and zircon Hf isotopic compositions of Mesozoic
710 intermediate-felsic intrusions in central Tibet: Petrogenetic and tectonic

711 implications. *Lithos* 198–199, 77–91.

712 Li, J.X., Qin, K.Z., Li, G.M., Xiao, B., Zhao, J.Z., Chen, L., 2011.
713 Magmatic-hydrothermal evolution of Cretaceous Duolong gold-rich
714 porphyry copper deposit in the Bangongcometallogenic belt, Tibet: Evidence
715 from U-Pb and $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology. *Journal of Asian Earth Sciences* 41,
716 525–536.

717 Li, S.G., 1993. Ba–Nb–Th–La diagrams user to identify tectonic environments of
718 ophiolite. *Acta Petrologica Sinica* 9, 146–157 (in Chinese with English
719 abstract).

720 Li, Y.B., Zhong, W.T., Zhang, T.P., Chen, H.A., Li, Y.C., Chen, H.Q., Fan, A.H., 2012.
721 Geochemical characteristics and genesis of the Bolong porphyry copper-gold
722 deposit in Gerze County, Tibet. *Acta Geoscientica Sinica* 33, 579–587 (in
723 Chinese with English abstract).

724 Liu, D.L., Huang, Q.S., Fan, S.Q., Zhang, L.Y., Shi, R.D., Ding, L., 2014. Subduction
725 of the Bangong–Nujiang Ocean: constraints from granites in the Bangong Co
726 area, Tibet. *Geological Journal* 49, 188–206.

727 Liu, D.L., Shi, R.D., Ding, L., Huang, Q.S., Zhang, X.R., Yue, Y.H., Zhang, L.Y.,
728 2016. Zircon U–Pb age and Hf isotopic compositions of Mesozoic granitoids
729 in southern Qiangtang, Tibet: Implications for the subduction of the
730 Bangong–Nujiang Tethyan Ocean. *Gondwana Research*, doi:
731 10.1016/j.gr.2015.04.007.

732 McKenzie, D., OpNions, R.K., 1991. Partial melt distributions from inversion of rare
733 Earth element concentrations. *Journal of Petrology* 32, 1021–1091.

734 Michael, J.O., 1983. Epigondolella populations and their phylogeny and zonation in
735 the Upper Triassic. *Fossil and Strata* 11, 177–192.

736 Murray, R.W., 1994. Chemical criteria to identify the depositional environment of
737 chert: general principles and applications. *Sedimentary Geology* 90,
738 213–232.

739 Murray, R.W., Buchholtz ten Brink, M.R., Jones, D.L., Gerlac, D.C., Russ III, G.P.,

740 1990. Rare earth elements as indicators of different marine depositional
 741 environments in chert and shale. *Geology* 18, 268–271.

742 Niu, Y.L., 2013. Global tectonics and geodynamics—A petrological and geochemical
 743 approach. Science Press, 1–307 (In China)

744 Niu, Y.L., Collerson, K.D., Batiza, R., Wendt, J.I., Regelous, M., 1999. Origin of
 745 enriched, type mid-ocean ridge basalt at ridges far from mantle plumes: the
 746 East Pacific Rise at 11°20'N. *Journal of Geophysical Research* 104,
 747 7067–7087

748 Niu, Y.L., Hékinian, R., 1997. Spreading rate dependence of the extent of mantle
 749 melting beneath ocean ridges. *Nature* 385, 326–329.

750 Niu, Y.L., O'Hara, M.J., 2008. Global correlations of ocean ridge basalt chemistry
 751 with axial depth: A new perspective. *Journal of Petrology* 49, 633–664.

752 Niu, Y.L., O'Hara, M.J., 2007. Varying Ni in OIB olivines—product of process not
 753 source. *Geochimica et Cosmochimica Acta* 71, a721–a721.

754 Oxburgh, E.R., Pamentier, E.M., 1977. Compositional and density stratification in
 755 oceanic lithosphere-causes and consequences. *Journal of the Geological*
 756 *Society of London* 133, 343–355.

757 Pan, G.T., 1983. An evolution of Tethys in global ocean–continent
 758 transGroup. *Tethyan Geology* 18, 23–40 (in Chinese with English abstract).

759 Pan, G.T., Chen, Z.L., Li, X.Z., Yang, Y.J., Xu, X.S., Xu, Q., Jiang, X.S., Wu, Y.L.,
 760 Luo, J.N., Zhu, T.X and Peng, Y.M., 1997. Geological–tectonic evolution in
 761 the eastern Tethys. Beijing: Geological Publishing House, 1–218 (in Chinese
 762 with English abstract).

763 Pan, G.T., Mo, X.X., Hou, Z.Q., Zhu, D.C., Wang, L.Q., Li, G.M., Zhao, Z.D., Geng,
 764 Q.R., Liao, Z.L., 2006. Spatial–temporal framework of the Lhasa Orogenic
 765 Belt and its evolution. *Acta Petrologica Sinica* 22, 521–533 (in Chinese with
 766 English abstract).

767 Pan, G.T., Wang, L.Q., Li, R.S., Yin, F.G., Zhu, D.C., 2012. Tectonic model of
 768 archipelagic arc-basin systems: The key to the continental geology.
 769 *Sedimentary Geology and Tethyan Geology* 32, 1–20 (in Chinese with

English abstract).

Pan, G.T., Xiao, Q.H., Lu, S.N., Deng, J.F., Feng, Y.M., Zhang, K.X., Zhang, Z.Y., Wang, F.G., Xing, G.J., Feng, Y.F., 2008. Definition, classification, characteristics and diagnostic indications of tectonic facies. *Geological Bulletin of China* 27, 1613–1637 (in Chinese with English abstract).

Parsons, B., Sclater, J.G., 1977. An analysis of the variation of ocean floor bathymetry and heat flow with age. *Journal of Geophysical Research* 82, 803–827.

Pearce, J.A., 1983. The role of sub–continental lithosphere in magmagenesis at destructive plate margins. In: Hawkesworth C J, et al. eds, *Continental Basalts and Mantle Xenoliths*. Nantwich Shiva, 230–249.

Phipps Morgan, J., Smith, W.F.S., 1992. Flattening of the sea-floor depth-age curve as a response to asthenospheric flow. *Nature* 359, 524–527.

Pullen, A., Kapp, P., Gehrels, G.E., Ding, L., Zhang, Q.H., 2011. Metamorphic rocks in central Tibet: lateral variations and implications for crustal structure. *Geological Society of America Bulletin* 123, 585–600.

Qiangba, Z.X., Xie, Y.W., Wu, Y.W., Xie, C.M., Li, Q.L., Qiu, J.Q., 2009. Zircon SIMS U–Pb dating and its significance of cumulate gabbro from Dengqenophiolite, eastern Tibet, China. *Geological Bulletin of China* 28, 1253–1258 (in Chinese with English abstract).

Qiu, R.Z., Zhou, S., Deng, J.F., Li, J.F., Xiao, Q.H., Cai, Z.Y., 2004. Dating of gabbro in the Shemalagouophiolite in the western segment of the Bangong Co–Nujiangophiolite belt, Tibet–With a discussion of the age of the Bangong Co–Nujiangophiolite belt. *Geological China* 31, 262–268 (in Chinese with English abstract).

Qu, X.M., Wang, R.J., Xin, H.B., Zhao, Y.Y and Fan, X.R., 2009. Geochronology and geochemistry of igneous rocks related to the subduction of the Tethys oceanic plate along the Bangong Lake arc zone, the western Tibetan Plateau. *Geochimica* 38, 523–535 (in Chinese with English abstract).

Qu, X.M., Xin, H.B., Zhao, Y.Y., Wang, R.J., Fan, H.T., 2010. Opening time of Bangong Lake Middle Tethys oceanic basin of the Tibet Plateau: Constraints

800 from petro–geochemistry and zircon U–Pb LAICPMS dating of mafic
801 ophiolites. *Earth Science Frontiers* 17, 53–63 (in Chinese with English
802 abstract).

803 Ren, J.S., Xiao, L.W., 2004. Lifting the mysterious veil of the tectonics of the
804 Qinghai-Tiber Plateau by 1:250000 geological mapping. *Geological Bull of*
805 *Chian* 23, 1–11(in Chinese with English abstract).

806 Schwandt, C.S., McKay, G.A., 1998. Rare earth element partition coefficients from
807 enstatite/melt synthesis experiments. *Geochimica et Cosmochimica Acta* 62,
808 2845–2848.

809 Shi, R.D., Griffin, W.L., Reilly, S.Y., Huang, Q.S., Zhang, X.R., Liu, D.L., Zhi, X.C.,
810 Xia, Q.X., Ding, L., 2012. Melt/mantle mixing products podiform chromite
811 deposits in ophiolites: Implication of Re–Os systematics in the Dongqiao
812 Neo-tethyan ophiolite, northern Tibet. *Gondwana Research* 21, 194–206.

813 Stein, C.A., Stein, S., 1992. A model for the global variation in oceanic depth and
814 heatflow with lithospheric age. *Nature* 359, 123–129 1992.

815 Sugisaki, R., Yamamoto, K., Adachi, M., 1982. Triassic bedded cherts in central Japan
816 are not pelagic. *Nature* 298, 644–647.

817 Sun, S.S., McDough, W.F., 1989. Chemical and isotope systematics of oceanic basalts:
818 implications for mantle composition and processes. In: Saunders A D (eds.),
819 *Magmatism in ocean Basins*. Geological Society Publication 42, 313–345.

820 Wang, L.Q., Pan, G.T., Ding, J., Yao, D.S., Luo, J.N., 2013. The geological map and
821 instruction of the Tibetan Plateau and its adjacent areas. Geological Publishing
822 House, 1–288.

823 Wang, X.B., Bao, P.S., Deng, W.M., 1987. Tibetan ophiolite. Beijing: Geological
824 Publishing House, 1–124 (in Chinese with English abstract).

825 Weaver, B., Kar, A., Davidson, J., Colucci, M., 1996. Geochemical characteristics of
826 volcanic rocks from Ascension Island South Atlantic Ocean. *Geothermics* 25,
827 449–470.

828 Wilson, M., 1989. *Igneous Petrogenesis*. London: Unwin Hyman, 1–466.

829 Winchester, J.A., Flody, P.A., 1977. Geochemical magma type discrimination:
 830 application to altered and metamorphosed basic igneous rocks. *Earth and*
 831 *Planetary Science Letters* 28, 459–469.

832 Wu, G.Y., 2006, Azores-type oceanic island basalts in the Paleo-Tethyan major
 833 oceanic basin in the Bitu area, eastern Tibet, China. *Geological Bulletin of*
 834 *China* 25, 772–781 (in Chinese with English abstract).

835 Xia, B., Xu, L.F., Wei Z.Q., Zhang, Y.Q., Wang, R., Li, J.F and Wang, Y.B., 2008,
 836 SHRIMP zircon dating of gabbro from the Donqiao Ophiolite in Tibet and its
 837 geological implications. *Acta Geological Sinica* 82, 528–531 (in Chinese
 838 with English abstract).

839 Xia, D.X., Liu, S.S., 1997. The lithostratigraphy of the Tibet. Publishing House of
 840 China University of Geosciences, 1–155 (in Chinese with English abstract).

841 Yan, Z., Wang, Z.Q., Li, J.L., Yan, Q.R., Wang, T., Chen, J.L., Xu, X.Y.,
 842 2008. Restoring the original tectonic types of sedimentary basins in the
 843 orogenic belts. *Geological Bulletin of China* 27, 2001–2013 (in Chinese with
 844 English abstract).

845 Yin, A., Harrison, T.M., 2000. Geologic evolution of the Himalayan–Tibetan orogen.
 846 *Annual Review of Earth and Planetary Sciences* 28, 211–280.

847 Yu, H.Q., Li, J.W., Ma, D.F., Li, G.M., Zhang, D.Q., Feng, C.W., Qu, W.J., Pan, G.T.,
 848 2009. Molybdenite Re–Os and SHRIMP zircon U–Pb dating of Duobuza
 849 porphyry copper deposit in Tibet and its geological implications. *Mineral*
 850 *Deposits* 28, 737–746 (in Chinese with English abstract).

851 Yuan, S.H., Wang, L.Q., Jiang, X.S., Yin, F.G., 2009. The retrospect and prospect on
 852 Archipelago Model and Composite Arc–Basin System Model of
 853 orogenesis. *Geological Science and Technology Information* 28, 1–11 (in
 854 Chinese with English abstract).

855 Zhai, Q.G., Jahn, B.M., Wang, J., Hu, P.Y., Chung, S.L., Lee, H.Y., Tang, S.H., Tang,
 856 Y., 2016. Oldest Paleo–Tethyan ophiolitic mélange in the Tibetan Plateau.

Geological Society of America Bulletin, doi:10.1130/B31296.1.

Zhai, Q.G., Jahn, B.M., Wang, J., Su, L., Mo, X.X., Wang, K.L., Tang, S.H., Lee, H.Y.,
2013. The Carboniferous ophiolite in the middle of the Qiangtangterrane,
Northern Tibet: SHRIMP U–Pb dating, geochemical and Sr–Nd–Hf isotopic
characteristics. *Lithos* 168–169, 186–199.

Zhang, K.J., Xia, B., Zhang, Y.X., Liu, W.L., Zeng, L., Li, J.F., Xu, L.F., 2014.
Central Tibetan Meso–Tethyan oceanic plateau. *Lithos* 210–211, 278–288.

Zhang, Y.C., Shi, G.R., Shen, S.Z., Yuan, D.X., 2014, Permian Fusuline Fauna from
the Lower Part of the Lugu Formation in the Central Qiangtang Block and its
Geological Implications: *ActaGeologicaSinica* (English Edition) 88, 365–379.

Zhang, Y.X., 2007, Tectonic evolution of the middle–western Bangong–Nujiang
suture, Tibet. A thesis submitted for the degree of doctorate at the Graduate
School of The Chinese Academy of Sciences, 1–256 (in Chinese with
English abstract).

Zhou, Y.Y., Jiang, Y.S., Wang, M.G., 2002. Permian Dibuco formation in the
Coqen-Xainza stratigraphic area, Tibet. *Geological Bulletin of China* 21, 79–82
(in Chinese with English abstract).

Zhu, D.C., Zhao, Z.D., Niu, Y.L., Dilek, Y., Hou, Z.Q., Mo, X.X., 2013. The origin
and pre–Cenozoic evolution of the Tibetan Plateau. *Gondwana Research* 23,
1429–1454.

Figure Captions

Fig. 1. (a) Tectonic framework of the Tibetan Plateau. (b) Geological maps of the
study area. Abbreviations are as follows: JSSZ, Jinshajiang Suture Zone; LSLSZ,

Longmuco–Shuanghu–Lancangjiang Suture Zone; BNSZ, Bangong–Nujiang Suture Zone; IYZSZ, Indus–Yarlung Zangbo Suture Zone.

Fig. 2. (a) and (b) Field photograph of the Gufeng ocean island (GFOI). (c) Field photograph showing basalt, chert, and limestone of the GFOI; (d) Field photograph of the GFOI conglomerate; (e) Chevron folds in GFOI limestones; (f) Field photograph of pillow basalt of the GFOI; (g) Field photograph of massive basalt of the GFOI; (h) Photomicrograph showing the texture and variable amounts of carbonate alteration of GFOI basalt; (i) Photomicrograph showing features indicative of greenschist-facies metamorphism of the GFOI basalt; (j) Field photograph showing interbedded GFOI basalts and GFOI limestones; (k) Field photograph showing interbedded GFOI cherts and GFOI limestones; (l) Field photograph of the conformable contact between GFOI limestone and GFOI chert.

Fig. 3. Geological transect across the Gufeng ocean island (GFOI) in the Gufeng area of Gerze County, Tibet.

Fig. 4. (a) Chondrite-normalized REE diagram for the basalts analyzed in this study. (b) Primitive-mantle-normalized trace element variation diagram for the basalt samples.

902

903 Fig.5.(a) Ti vs. Zr vs. Y diagram for samples analyzed in this study: A, island-arc
904 tholeiite; B, mid-ocean ridge basalt (MORB) + island-arc tholeiites and calc-alkali
905 basalt; C, calc-alkaline basalt; D, intraplate basalt. (b) Zr vs. Nb vs. Y diagram for
906 samples analyzed in this study: AI, intraplate alkali basalts; AII, intraplate alkali
907 basalt and tholeiite; B, E-type MORB; C, intraplate tholeiite and volcanic arc basalt; D,
908 N-type MORB and volcanic arc basalt.

909

910 Fig. 6. Nb/Y vs. Zr/TiO₂ diagram for the samples analyzed in this study.

911

912 Fig.7. NASC-normalized REE diagram for the **GFOI** cherts analyzed during this study.
913 NASC is North American Shale.

914

915 Fig. 8. Conodont fossils within the Gufeng ocean island (**GFOI**). (a–e)
916 Photomicrographs of the conodont fossils. (f–h) CL images of the conodont fossils.

917

918 Fig. 9. Sm/Yb vs. La/Sm diagram (after [Zhao and Zhou, 2007](#)) showing data of the
919 basalts analyzed in this study. The mantle array (heavy line) is defined by depleted
920 MORB mantle (DM; [McKenzie and O’Nions, 1991](#)) and primitive mantle (PM; [Sun](#)
921 [and McDonough, 1989](#)). Melting curves for spinel peridotite and garnet peridotite

with both DM and PM compositions are after [Aldanmaz et al. \(2000\)](#). Numbers along lines indicate the degree of partial melting.

Fig. 10. Al vs. Fe vs. Mn, $100 \cdot (\text{Fe}_2\text{O}_3/\text{SiO}_2)$ vs. $100 \cdot (\text{Al}_2\text{O}_3/\text{SiO}_2)$, $\text{Fe}_2\text{O}_3/(100 - \text{SiO}_2)$ vs. $\text{Al}_2\text{O}_3/(100 - \text{SiO}_2)$, and $\text{Fe}_2\text{O}_3/\text{TiO}_2$ vs. $\text{Al}_2\text{O}_3/(\text{Al}_2\text{O}_3 + \text{Fe}_2\text{O}_3)$ diagrams showing data of the cherts analyzed in this study ([Murray, 1994](#)). A, non-hydrothermal chert; B, hydrothermal chert.

Fig. 11. Schematic model showing the opening and early evolution of the Bangong–Nujiang Tethyan Ocean.

Fig.1

Fig.2

Fig.3

Fig.5

Fig.5

Fig.6

Fig.7

Fig.8

Fig.9

Fig.10

Fig.11

Graphical Abstract

Table 1

Table 1. Major (wt.%) and trace element (ppm) data for the G1 basalts in the Gufeng ocean island.

Sample	G1H1	G1H2	G1H3	G1H4	G1H5	G1H6	G1H7	G1H8
SiO ₂	53.7	57.3	49.5	53.5	52.8	53.7	46.8	54.2
TiO ₂	1.20	1.13	1.30	1.28	1.17	1.09	1.25	1.25
Al ₂ O ₃	14.5	12.8	14.7	13.1	13.3	13.2	15.3	13.8
Fe ₂ O ₃	11.9	10.3	13.2	11.5	11.7	11.7	13.9	12.2
MnO	0.17	0.14	0.15	0.15	0.16	0.13	0.19	0.15
MgO	5.26	4.32	4.50	4.78	4.89	4.16	5.40	4.97
CaO	7.22	6.77	10.08	6.08	9.29	10.69	9.31	7.59
Na ₂ O	4.20	3.35	1.29	3.61	2.64	0.89	2.72	1.40
K ₂ O	0.03	0.03	0.03	0.03	0.03	0.03	0.05	0.06
P ₂ O ₅	0.09	0.08	0.10	0.10	0.10	0.07	0.12	0.09
LOI	0.93	2.93	4.33	5.01	3.07	3.49	4.28	3.46
Li	43.8	46.0	53.2	56.1	41.2	42.3	57.7	72.2
Sc	34.5	31.7	30.4	33.9	32.5	31.3	35.6	33.2
V	435	372	485	388	429	452	561	446
Cr	68.1	44.7	26.5	48.3	49.9	54.8	54.1	60.4
Mn	1341	1073	1140	1148	1229	1050	1530	1204
Co	32.4	28.7	31.9	27.1	29.5	29.1	31.3	31.9
Ni	42.4	30.1	25.0	31.8	32.0	31.4	35.1	37.7
Cu	114	127	97	116	110	88.7	85.4	104
Zn	70.0	62.3	83.8	80.0	69.0	74.1	86.7	82.2
Ga	15.5	14.9	21.4	14.0	17.2	21.8	21.2	19.2
Rb	0.83	0.58	0.51	0.39	0.47	0.28	0.51	1.34
Sr	31.5	44.5	30.8	55.0	23.5	20.4	33.1	18.9
Y	28.2	22.5	32.8	26.4	27.8	28.6	35.1	29.7
Zr	56.1	53.6	61.1	59.5	55.5	52.1	60.0	59.2
Nb	1.74	1.51	2.06	1.78	1.53	1.38	1.76	1.80
Cs	0.29	0.22	0.22	0.19	0.16	0.22	0.49	0.42
Ba	8.58	7.52	39.5	18.0	8.79	6.75	16.8	12.2
La	2.68	2.64	3.07	2.67	2.52	2.82	2.24	3.39
Ce	7.70	7.23	9.07	7.71	7.31	7.78	6.91	9.24
Pr	1.29	1.17	1.48	1.30	1.22	1.24	1.21	1.49
Nd	6.81	5.98	7.85	6.92	6.56	6.51	6.78	7.82
Sm	2.47	2.13	2.80	2.55	2.42	2.32	2.66	2.74
Eu	0.86	0.69	1.02	0.88	0.83	0.86	0.97	1.01
Gd	3.49	2.87	4.08	3.44	3.45	3.48	4.09	3.77
Tb	0.66	0.55	0.76	0.66	0.65	0.65	0.78	0.70
Dy	4.36	3.62	4.93	4.31	4.31	4.31	5.20	4.53
Ho	1.00	0.82	1.12	0.96	0.97	0.98	1.19	1.02
Er	2.99	2.46	3.24	2.89	2.88	2.88	3.53	3.00
Tm	0.44	0.37	0.47	0.43	0.42	0.42	0.51	0.44
Yb	2.96	2.49	3.14	2.90	2.83	2.80	3.41	2.95
Lu	0.45	0.38	0.48	0.44	0.43	0.42	0.51	0.45

Hf	1.56	1.43	1.62	1.65	1.52	1.40	1.67	1.57
Ta	0.17	0.09	0.22	0.10	0.09	0.09	0.10	0.18
Pb	0.94	0.83	1.23	0.36	1.00	1.39	0.59	0.86
Th	0.50	0.45	0.63	0.49	0.46	0.42	0.52	0.48
U	0.13	0.13	0.18	0.16	0.12	0.12	0.15	0.17

Table 2

Table 2. Major (wt.%) and trace element (ppm) data for the G2 basalts in the Gufeng ocean island.

Sample	G2H1	G2H2	G2H3	G2H4	G2H5	G2H6	G2H7	G2H8	G2H9
SiO ₂	52.6	42.7	45.1	55.0	48.2	52.5	52.4	47.6	52.6
TiO ₂	1.62	1.76	1.22	1.60	1.67	1.68	1.51	1.72	1.62
Al ₂ O ₃	13.7	15.3	11.2	13.0	14.1	13.1	13.0	14.3	13.7
Fe ₂ O ₃	11.8	10.4	7.55	8.68	10.0	7.81	9.67	9.90	11.8
MnO	0.13	0.15	0.11	0.14	0.15	0.13	0.15	0.15	0.13
MgO	4.48	8.24	4.12	4.59	7.50	4.15	7.28	7.32	4.48
CaO	7.90	7.59	18.8	6.69	8.14	7.84	6.92	8.04	7.90
Na ₂ O	2.82	3.30	3.53	4.24	3.77	4.35	3.60	3.80	2.82
K ₂ O	0.56	1.53	0.23	0.17	0.62	0.28	0.61	0.60	0.56
P ₂ O ₅	0.23	0.22	0.17	0.19	0.20	0.21	0.19	0.21	0.23
LOI	3.36	7.21	6.21	5.02	4.31	7.13	3.94	4.99	3.36
Li	69.5	69.7	27.4	39.3	62.1	38.3	67.5	59.8	69.5
Sc	27.1	32.6	22.2	27.1	32.3	27.8	36.0	31.3	27.1
V	410	269	192	218	271	218	310	262	410
Cr	41.0	294	221	194	290	183	306	222	41.038
Mn	1084	1052	800	1034	1185	934	1255	1074	1084
Co	32.5	47.5	29.9	35.0	44.6	33.7	48.0	41.6	32.5
Ni	41.4	158	86.3	81.0	162	79.2	173	130	41.4
Cu	93.6	109	18.3	53.5	95.4	46.1	110	114	93.6
Zn	87.6	84.2	75.1	90.2	79.6	77.8	82.8	76.2	87.6
Ga	21.3	19.0	14.8	18.6	18.9	17.5	19.6	18.7	21.3
Rb	31.8	41.5	7.6	5.4	18.5	7.8	19.3	18.4	31.8
Sr	163	258	234	376	334	444	310	323	163
Y	25.4	21.1	16.4	20.1	21.2	20.9	22.1	20.6	25.4
Zr	96.1	122	87.2	97.1	120	107	134	113	96.1
Nb	22.1	14.8	10.9	11.2	14.6	12.8	18.5	14.5	22.1
Cs	3.77	2.45	3.39	2.18	2.33	1.26	2.11	2.41	3.77
Ba	214	302	71.0	205	247	153	221	243	214
La	11.7	13.0	10.6	8.49	12.3	9.39	13.0	12.9	11.7
Ce	26.3	30.2	22.8	20.5	28.8	22.6	29.4	29.8	26.3
Pr	3.49	3.91	2.94	2.78	3.80	3.03	3.89	3.85	3.49
Nd	15.0	17.6	13.1	13.1	17.4	14.2	17.3	17.3	15.0
Sm	3.79	4.46	3.22	3.66	4.44	3.89	4.39	4.35	3.79
Eu	1.32	1.59	1.15	1.40	1.59	1.65	1.56	1.57	1.32
Gd	4.35	4.96	3.66	4.45	4.98	4.57	4.79	4.86	4.35
Tb	0.71	0.77	0.56	0.71	0.77	0.74	0.72	0.75	0.71
Dy	4.24	4.62	3.48	4.40	4.63	4.59	4.44	4.52	4.24
Ho	0.89	0.90	0.67	0.85	0.89	0.89	0.85	0.88	0.89
Er	2.50	2.53	1.91	2.34	2.50	2.49	2.35	2.46	2.50
Tm	0.35	0.34	0.26	0.31	0.34	0.33	0.31	0.33	0.35
Yb	2.31	2.16	1.63	1.93	2.13	2.09	1.98	2.11	2.31
Lu	0.34	0.31	0.23	0.27	0.30	0.29	0.28	0.30	0.34

Hf	2.32	2.89	2.09	2.41	2.92	2.66	3.44	2.81	2.32
Ta	1.42	0.87	0.69	0.66	0.99	0.74	1.10	0.84	1.42
Pb	1.66	1.25	2.44	1.37	1.13	1.04	1.18	2.70	1.66
Th	1.82	1.62	1.28	1.14	1.62	1.29	1.57	1.61	1.82
U	0.54	0.39	1.79	0.46	0.39	0.64	0.36	0.39	0.54

Table 3

Table 3. Major (wt.%) and trace element (ppm) data for the G3 basalts in the Gufeng ocean island.

Sample	G3H1	G3H2	G3H3	G3H4	G3H5	G3H6	G3H7
SiO ₂	47.0	49.4	44.3	43.6	45.5	44.9	45.2
TiO ₂	2.46	2.44	2.52	2.54	2.45	2.64	2.41
Al ₂ O ₃	14.8	14.0	15.1	13.6	12.5	14.6	16.4
Fe ₂ O ₃	11.6	10.7	10.4	12.3	11.8	11.9	12.6
MnO	0.11	0.10	0.14	0.17	0.18	0.15	0.13
MgO	4.28	3.95	4.07	6.67	6.92	5.63	4.66
CaO	8.90	8.09	10.0	10.4	10.8	8.89	6.79
Na ₂ O	4.83	5.24	5.15	3.92	3.82	4.28	5.40
K ₂ O	1.08	1.24	0.34	0.37	0.32	0.74	1.44
P ₂ O ₅	0.46	0.44	0.56	0.48	0.46	0.52	0.56
LOI	3.65	3.52	5.75	4.28	3.86	4.30	3.82
Li	60.1	48.2	62.1	59.0	52.3	72.8	108
Sc	15.7	17.7	16.9	30.8	26.9	26.2	14.1
V	276	288	294	271	305	322	330
Cr	10.3	11.4	14.6	245	172	38.2	6.69
Mn	919	843	1066	1201	1545	1157	1019
Co	34.4	30.1	35.9	40.2	44.9	37.7	34.2
Ni	55.9	53.1	42.4	107	130	69.0	54.5
Cu	67.7	57.4	80.3	24.7	51.8	70.0	78.0
Zn	112	97.1	113	107	120	111	111
Ga	23.1	22.4	23.2	22.7	24.8	26.2	26.1
Rb	49.0	69.7	9.48	9.52	12.5	29.2	81.9
Sr	314	356	517	290	322	278	374
Y	19.8	19.9	21.5	19.5	22.2	21.8	21.6
Zr	204	210	187	179	197	198	186
Nb	44.0	43.2	37.6	31.4	35.3	35.0	63.6
Cs	6.65	8.54	0.96	1.06	1.87	3.16	8.61
Ba	558	576	476	360	559	828	571
La	28.3	29.7	32.7	26.0	32.6	28.7	27.0
Ce	61.6	62.7	71.4	58.8	71.1	65.4	59.1
Pr	7.72	7.80	8.67	7.31	8.61	8.16	7.53
Nd	31.8	32.4	36.0	31.4	36.3	35.0	30.5
Sm	6.80	6.96	7.71	7.03	7.91	7.75	6.60
Eu	2.25	2.31	2.56	2.30	2.67	2.57	2.15
Gd	6.38	6.51	7.28	6.74	7.47	7.44	6.10
Tb	0.83	0.85	0.97	0.91	1.01	1.00	0.85
Dy	4.51	4.58	5.10	4.80	5.33	5.28	4.33
Ho	0.77	0.79	0.88	0.81	0.90	0.91	0.78
Er	1.94	1.97	2.22	2.02	2.29	2.29	1.94
Tm	0.24	0.24	0.28	0.25	0.28	0.28	0.25
Yb	1.46	1.48	1.66	1.51	1.68	1.68	1.50
Lu	0.20	0.20	0.22	0.20	0.23	0.23	0.21

Hf	5.12	5.26	4.49	4.45	4.80	4.82	3.93
Ta	2.57	2.52	2.10	2.03	2.36	2.36	3.95
Pb	7.06	1.82	3.44	2.21	1.52	8.47	1.69
Th	4.10	4.09	4.56	3.70	4.21	4.14	4.66
U	0.81	0.92	1.60	0.91	0.85	0.72	1.05

Table 4

Table 4. Major (wt.%) and trace element (ppm) data for the chert in the Gufeng ocean island.

Sample	GH1	GH2	GH3	GH4	GH5	GH6	GH7	GH8	GH9	GH10
SiO ₂	92.2	93.0	94.2	94.3	94.3	95.0	84.9	91.0	94.1	93.4
TiO ₂	0.07	0.20	0.14	0.05	0.05	0.04	0.13	0.08	0.12	0.10
Al ₂ O ₃	0.76	2.16	1.94	0.94	0.71	0.58	1.69	1.04	1.55	1.54
Fe ₂ O ₃	0.34	1.07	1.25	0.56	0.51	0.41	1.09	0.32	1.01	0.82
MnO	0.03	0.01	0.02	0.03	0.03	0.03	0.08	0.03	0.01	0.02
MgO	0.24	0.71	0.77	0.36	0.29	0.19	0.61	0.38	0.55	0.88
CaO	2.88	0.42	0.81	2.54	2.92	2.73	6.73	4.49	0.18	1.93
Na ₂ O	0.29	0.81	0.62	0.29	0.10	0.04	0.32	0.04	0.54	0.04
K ₂ O	0.00	0.03	0.04	0.09	0.14	0.13	0.23	0.32	0.02	0.31
P ₂ O ₅	0.01	0.04	0.06	0.01	0.02	0.01	0.03	0.02	0.03	0.05
LOI	2.58	0.73	0.93	1.64	1.73	1.67	4.71	2.95	0.49	1.71
Li	3.46	9.99	10.76	4.94	4.79	3.59	8.33	5.99	6.08	14.94
Sc	1.90	2.99	2.58	1.24	1.12	0.81	2.88	2.05	3.04	2.18
V	5.99	13.6	10.4	4.82	4.29	3.84	9.52	6.52	9.90	8.92
Cr	7.77	8.48	9.26	24.8	7.88	4.24	7.20	19.4	10.5	7.53
Co	3.35	4.79	3.94	1.53	1.31	0.69	3.64	1.68	2.97	3.69
Ni	7.49	25.7	23.4	18.7	9.18	2.58	21.1	13.4	17.1	14.8
Cu	6.83	14.8	22.0	10.1	14.2	7.14	109	11.9	20.1	6.52
Zn	5.13	18.9	16.4	37.9	6.65	4.62	15.1	6.73	13.3	18.1
Ga	1.24	3.26	2.70	1.35	1.11	0.85	2.25	1.46	2.73	2.25
Rb	0.50	1.61	1.53	3.30	6.00	5.00	7.56	13.05	0.70	12.5
Sr	98.9	11.3	13.5	47.0	59.8	64.9	116	71.4	5.9	27.3
Y	7.32	6.61	4.29	5.47	4.77	4.00	17.3	9.78	5.69	5.69
Zr	12.4	43.1	32.7	13.5	15.9	11.1	22.8	15.8	22.9	18.6
Nb	1.94	7.62	4.11	1.45	4.10	1.18	2.90	1.85	4.07	2.08
Cs	0.12	0.17	0.15	0.16	0.24	0.19	0.35	0.38	0.13	0.40
Ba	8.68	22.7	15.7	21.3	26.6	19.5	33.6	63.0	9.24	53.0
La	6.36	5.33	6.73	5.82	5.32	3.72	13.6	8.05	8.92	4.62
Ce	10.1	12.5	13.9	9.1	10.3	7.7	16.3	10.8	12.3	10.1
Pr	1.74	1.90	1.91	1.50	1.39	1.02	3.79	2.14	2.45	1.29
Nd	7.64	8.70	7.72	6.20	5.57	4.14	16.67	8.84	11.14	5.28
Sm	1.68	2.09	1.62	1.38	1.28	0.99	4.13	2.06	2.53	1.23
Eu	0.36	0.46	0.35	0.32	0.30	0.23	0.97	0.44	0.50	0.28
Gd	1.63	1.88	1.40	1.35	1.24	0.97	4.47	2.06	2.29	1.22
Tb	0.22	0.26	0.18	0.19	0.17	0.14	0.65	0.30	0.30	0.17
Dy	1.26	1.45	0.94	1.08	0.92	0.71	3.70	1.71	1.54	0.98
Ho	0.24	0.27	0.17	0.19	0.16	0.12	0.65	0.32	0.27	0.18
Er	0.66	0.75	0.46	0.50	0.40	0.31	1.59	0.85	0.72	0.47
Tm	0.09	0.10	0.06	0.06	0.05	0.04	0.19	0.10	0.10	0.06
Yb	0.53	0.59	0.39	0.41	0.28	0.24	1.16	0.65	0.61	0.37
Lu	0.07	0.08	0.05	0.05	0.04	0.03	0.15	0.08	0.08	0.05
Hf	0.30	1.09	0.82	0.33	0.53	0.34	0.52	0.38	0.55	0.43

Ta	0.13	0.47	0.25	0.08	0.31	0.08	0.16	0.11	0.24	0.15
Pb	5.24	3.91	4.76	2.47	2.68	1.77	3.53	3.53	2.34	1.97
Th	0.62	1.36	1.31	0.92	0.72	0.57	1.20	0.82	1.22	0.83
U	0.16	0.33	0.29	0.94	0.68	0.66	0.80	0.23	0.47	0.20