ABA-dependent and -independent signaling in response to osmotic stress in plants

Takuya Yoshida, Junro Mogami and Kazuko Yamaguchi-Shinozaki

5

Laboratory of Plant Molecular Physiology, Graduate School of Agricultural and Life Sciences, The University of Tokyo, Tokyo 113-8657, Japan

Corresponding author: Kazuko Yamaguchi-Shinozaki

10 Tel: +81-3-5841-8137

E-mail: akys@mail.ecc.u-tokyo.ac.jp

1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan

Abstract

Plants have adaptive robustness to osmotic stresses such as drought and high salinity. Numerous genes functioning in stress response and tolerance are induced under osmotic
5 conditions in diverse plants. Various signaling proteins, such as transcription factors, protein kinases and phosphatases, play signal transduction roles during plant adaptation to osmotic stress, with involvement ranging from stress signal perception to stress-responsive gene expression. Recent progress has been made in analyzing the complex cascades of gene expression during osmotic stress response, and especially in identifying specificity and
10 crosstalk in abscisic acid (ABA)-dependent and -independent signaling pathways. In this review, we highlight transcriptional regulation of gene expression governed by two key transcription factors: AREB/ABFs and DREB2A operating respectively in ABA-dependent and -independent signaling pathways.

Introduction

As sessile organisms facing various environmental challenges, higher plants have adaptive robustness at molecular, cellular and physiological levels to environmental stress. Water deficit is a major factor restricting plant growth, survival, yield and distribution. Under osmotic stress conditions such as drought and high salinity, numerous genes functioning in stress response and tolerance are induced, and abscisic acid (ABA), a key plant stress-signaling hormone, is accumulated [1–3]. Osmotic stress-responsive gene expression is regulated by ABA-dependent and -independent pathways [2]. The *cis*-acting element, ABA-responsive element (ABRE), and a group of transcription factors, ABRE-binding protein/ABRE-binding factors (AREB/ABFs), have pivotal functions in ABA-dependent gene expression. Similarly, a *cis*-element, dehydration-responsive element/C-repeat (DRE/CRT), and DRE-/CRT-binding protein 2 (DREB2) transcription factors play key roles in ABA-independent gene expression in response to osmotic stress. In this review, we focus on recent progress in the study of transcriptional regulatory networks governed by AREB/ABFs and DREB2A transcription factors in osmotic stress signaling, with particular emphasis on the functional interaction of ABA-dependent and -independent pathways. Various signaling proteins, such as transcription factors, kinases and phosphatases, are involved in expression of osmotic stress-responsive genes; their roles have been extensively reviewed elsewhere [4,5].

The pivotal role of AREB/ABF transcription factors in ABA-dependent gene expression

Among group-A bZIP transcription factors in *Arabidopsis thaliana* (*Arabidopsis*), nine AREB/ABF transcription factors have a bZIP domain and four conserved domains containing Ser/Thr kinase phosphorylation sites [6]. Consistent with increased *AREB1/ABF2*, *AREB2/ABF4* and *ABF3* expression induced by drought, high salinity and ABA in vegetative tissues [7], overexpression studies have shown that these three AREB/ABFs are positive regulators of ABA signaling under drought stress conditions [7–10]. These studies

and investigation of the *areb1 areb2 abf3* triple mutant [11] have jointly demonstrated that the three AREB/ABFs function as master transcription factors cooperatively regulating ABRE-dependent gene expression in ABA signaling during response to drought stress.

AREB/ABF transcription factors are activated in an ABA-dependent manner
5 through multiple-site phosphorylation of their conserved domains by SNF1-related kinase 2s (SnRK2s) [10,12,13]. Nine of 10 *Arabidopsis* SnRK2s are activated by osmotic stress, with 3 subclass III SnRK2s, SRK2D/SnRK2.2, SRK2E/SnRK2.6/OST1 and SRK2I/SnRK2.3, also strongly activated by ABA [14]. The three SnRK2s co-localize and interact with AREB/ABFs in plant cell nuclei [11,15]. Furthermore, expression of most downstream
10 genes of AREB1/ABF2, AREB2/ABF4 and ABF3 is substantially impaired in the *srk2d/e/i* triple mutant [15], and ABA-dependent phosphorylation of AREB/ABFs is completely eliminated [16,17]. These findings support the view that the three subclass III SnRK2s regulate ABA-responsive gene expression through phosphorylation of AREB/ABFs under osmotic stress conditions (Figure 1).

15 The *srk2d/e/i* triple mutant displays an extreme ABA-insensitive phenotype in regard to seed germination, seedling growth, stomatal regulation and osmotic stress-responsive gene expression [15,16,18]. AREB1/ABF2, AREB2/ABF4 and ABF3 regulate expression of one-third of SnRK2 downstream genes [15]; other than these three AREB/ABFs, however, little is known about the transcription factors functioning
20 downstream of SnRK2s. To gain a comprehensive understanding of signaling networks governed by the three subclass III SnRK2s, two phosphoproteome studies using the *srk2d/e/i* triple mutant have been performed [19,20]. A comparative phosphoproteome analysis of the *srk2d/e/i* mutant and wild-type plants after dehydration or ABA treatment identified 32 phosphopeptides as candidates for SnRK2 substrates, including two
25 phosphopeptides corresponding to AREB1/ABF2, AREB2/ABF4, ABF3 or ABF1 [19]. Although phosphopeptides corresponding to other transcription factors were undetected, SNRK2-SUBSTRATE 1 (SNS1) was identified as a novel SnRK2 substrate (Figure 1). The *sns1* mutant shows increased sensitivity to ABA in post-germination stages, with the expression of ABA-responsive genes such as *RD29B* and *RAB18* being slightly increased.
30 Given that SNS1 is of unknown function, further analyses are required to reveal its precise

roles in ABA signaling.

Another phosphoproteome study using the *srk2d/e/i* mutant identified 84 phosphopeptides as possible SnRK2 substrates, including eight proteins classified as transcription factors and DNA-binding proteins [20]. Results of previous studies imply that physiological roles of the remaining proteins, including AREB3, ENHANCED EM LELVEL (EEL), FLOWERING BHLH 3 (FBH3) and TATA BINDING PROTEIN-ASSOCIATED FACTOR 5 (TAF5), but not AREB1/ABF2, are associated with developmental processes rather than osmotic stress responses [6,21,22] (Figure 1). Although RNA-binding proteins were also identified as candidate SnRK2 substrates, their roles in ABA signaling are still undetermined. Collectively, these phosphoproteome analyses suggest that AREB/ABFs are predominant transcription factors functioning downstream of the three subclass III SnRK2s during ABA signaling in response to osmotic stress. The view derived from the phosphoproteome studies has been further supported by a reverse genetic approach showing that ABF1 is a transcription factor that functions downstream of SnRK2s [23]. Despite lower expression levels of *ABF1* compared with *AREB1/ABF2*, *AREB2/ABF4* and *ABF3*, even under stress conditions, the *areb1 areb2 abf3 abf1* quadruple mutant displays increased sensitivity to drought, decreased sensitivity to ABA in primary root growth, and impaired expression of ABA- and osmotic stress-responsive genes compared with the *areb1 areb2 abf3* triple mutant. Transcriptome analyses have revealed that 70% of downstream genes of the three subclass III SnRK2s are down-regulated in the *areb1 areb2 abf3 abf1* quadruple mutant. These results favor the view that these SnRK2s regulate ABA-responsive gene expression under osmotic stress primarily through the four AREB/ABF transcription factors (Figure 1).

Phosphoproteome and transcriptome approaches have provided further new insights into the role of the three subclass III SnRK2s in ABA signaling. FBH3, identified as a possible SnRK2 substrate, and its homologs are bHLH transcription factors activating *CONSTANS (CO)* expression during floral regulation [21]. Interestingly, *FBH3* and *CO* expression is impaired in both late-flowering *areb1 areb2 abf3 abf1* and early-flowering *srk2d/e/i* mutants [20,23], although their phenotypes are contradictory. Given that other flowering-associated proteins, such as *MOS3* and *XRN3*, are also possible SnRK2

substrates [20], SnRK2s may intricately and finely regulate the floral transition in ABA signaling via a variety of substrates, including AREB/ABFs and FBH3. Surprisingly, FBH3 and CO are involved in stomatal opening [24,25]. FBH3 is identical to ABA-responsive kinase substrate 1 (AKS1), which is negatively regulated in guard cells by SRK2E/SnRK2.6/OST1-dependent phosphorylation [24] (Figure 1). FBH3/AKS1 is a transcriptional activator of *KAT1*, which encodes the major inward-rectifying K⁺ channel involved in stomatal opening. Because *KAT1* is also negatively regulated by SnRK2E/SnRK2.6/OST1-dependent phosphorylation [26], SnRK2-mediated ABA signaling might inhibit stomatal opening through transcriptional and posttranscriptional mechanisms. Studies exploiting these recently identified putative SnRK2 substrates are needed to understand the global regulatory roles of subclass III SnRK2s in ABA signaling.

Negative regulatory mechanisms of DREB2A activity in ABA-independent gene expression

DREB2 proteins are members of the AP2/ERF family of plant-specific transcription factors. Among eight *DREB2s* in *Arabidopsis*, *DREB2A* and *DREB2B* are highly induced by drought, high salinity and heat stress [27,28]. Despite the instability of DREB2A proteins *in planta*, studies of *Arabidopsis* overexpressing a constitutively active form of DREB2A, which harbors an internal deletion of the domain involved in protein stability, have revealed that DREB2A plays pivotal roles in ABA-independent gene expression under osmotic and heat stress conditions [28,29].

Because DREB2A, which induces many genes encoding proteins involved in stress response and tolerance, has adverse effects on plant growth, its transcript and protein levels are tightly regulated. An attempt to elucidate transcriptional regulatory mechanisms of *DREB2A* has identified GROWTH-REGULATING FACTOR7 (GRF7) as a transcriptional repressor of osmotic-stress responsive genes including *DREB2A* [30] (Figure 2). GRF proteins are putative transcription factors harboring two conserved domains. Among nine GRFs in *Arabidopsis*, GRF7 interacts with a short *DREB2A* promoter region that suppresses its expression under unstressed growth conditions. *GRF7* knockout and knockdown mutants,

showing increased *DREB2A* expression, display enhanced salinity tolerance as well as growth retardation. In addition, transcriptome analysis has revealed that hundreds of osmotic stress-responsive genes are up-regulated in the *grf7* knockout mutant compared with wild-type plants grown under normal conditions, suggesting that GRF7 is a repressor regulating a wide range of osmotic-stress responsive genes during unstressed growth conditions.

In addition to transcriptional repression under normal growth conditions, a ubiquitin-proteasome pathway is proposed to degrade leaky expression of the DREB2A protein. DREB2A-INTERACTING PROTEIN1 (DRIP1), a ubiquitin E3 ligase harboring a C3HC4-type RING domain, has been identified as an interacting protein of DREB2A by yeast two-hybrid screening [31]. Whereas DRIP1 overexpression delays the expression of DREB2A downstream genes in response to dehydration, their expression is increased in double knockout mutants of *DRIP1* and its homolog *DRIP2*. Furthermore, green fluorescent protein (GFP)-DREB2A fusion proteins are stably expressed in the *drip1 drip2* mutant, which displays growth retardation under normal growth conditions and enhanced drought stress tolerance. These results suggest that DRIP1 and DRIP2 negatively regulate drought stress-responsive gene expression by targeting DREB2A to 26S proteasome proteolysis (Figure 2). Despite the crucial role of DRIPs in DREB2A degradation, a recent study has shown that DREB2A protein levels, which are rapidly increased in response to heat stress, are reduced by prolonged heat stress even in the *drip1 drip2* mutant [32]. Given that stabilization of DREB2A is required, but not sufficient for substantial induction of its downstream genes [32], DREB2A degradation and activation might be regulated by many yet-to-be-identified proteins in addition to DRIPs.

DREB2A and DRIPs are a well-characterized transcription factor/ubiquitin E3 ligase combination that regulates osmotic stress-responsive genes. A similar regulatory interaction has recently been identified between the transcription factor AtERF53 and two homologous C3HC4-type RING E3 ligases, RING domain ligase 2 (RGLG2) and RGLG1 [33]. The expression of *AtERF53*, an AP2/ERF transcription factor belonging to a non-DREB2 subfamily [34], is induced significantly by drought and high salinity but only slightly by ABA treatment [33,35]. AtERF53 interacts and co-localizes with RGLG2 and

RGLG1, both of which mediate AtERF53 ubiquitination *in vitro*. In contrast to wild-type plants, AtERF53-GFP fusion proteins are stably expressed in the *rglg1 rglg2* mutant that displays enhanced drought stress tolerance. Because AtERF53 is involved in induction of stress-responsive genes such as *COR15A* and *COR15B* [35], AtERF53 and RGLGs may function as positive and negative regulators, respectively, of osmotic stress-responsive gene expression. As each group of AP2/ERF transcription factors and RING-type ubiquitin ligases comprises many genes [34,36], comprehensive analyses of these genes may provide further insights into transcriptional networks in response to osmotic stress.

10 **Crosstalk between ABA-dependent and -independent pathways**

As evidenced by the fact that multiple dehydration-inducible genes are induced by exogenous ABA treatment [2], the global transcriptional network activated in response to osmotic stress is cooperatively but not exclusively regulated by ABA-dependent and -independent pathways. Despite the importance of crosstalk between ABA-dependent and -independent pathways, knowledge of how the two signaling pathways regulate each other has been limited until recently. The three subclass III SnRK2s likely participate in the convergence of ABA-dependent and -independent signaling [6] (Figure 1). Activated by osmotic stress as well as by ABA, these SnRK2s substantially regulate ABA-dependent and -independent gene expression [15]. Analysis of multiple knockout mutants of all 10 *SnRK2s* in *Arabidopsis* has further implied that the three SnRK2s are essential to both osmotic stress response and ABA signaling [37]. In-gel kinase assay results, however, indicate that osmotic stress activates SnRK2s in an ABA-independent manner [38,39]. Consistent with these results, a recent phosphoproteome study has revealed that phosphopeptides corresponding to the three subclass III SnRK2s are phosphorylated after 5 min of treatment with ABA but not by osmotic stresses such as NaCl and mannitol [40]. On the other hand, short-term osmotic stress has been found to phosphorylate a phosphopeptide corresponding to SRK2A/SnRK2.4, SRK2B/SnRK2.10, SRK2G/SnRK2.1 or SRK2H/SnRK2.5, which are subclass I SnRK2s not activated by ABA, and phosphopeptides corresponding to MAP3K and MAP4K. Although subclass III SnRK2s appear to be activated by autophosphorylation during ABA

signaling, novel proteins not yet characterized may be responsible for SnRK2 activation in response to osmotic stress.

Analyses of *DREB2A* promoters have provided new insights into crosstalk between ABA-dependent and -independent signaling (Figure 2). Given that an ABRE motif-containing short promoter region is required for dehydration-responsive *DREB2A* expression, transient expression and chromatin immunoprecipitation (ChIP) analyses have demonstrated that *DREB2A* is regulated by AREB1/ABF2, AREB2/ABF4 and ABF3 during ABA signaling under osmotic stress [41]. Interestingly, several AP2/ERF transcription factors including *DREB2A* interact with AREB1/ABF2, AREB2/ABF4 and ABF3 [42] (Figure 1), but it is unclear whether these interactions have functional significance in gene expression. It is noteworthy that expression of ABA-inducible genes (including *AREB2/ABF4*) and osmotic stress-inducible genes is increased in the *grf7* mutant [30]. This finding implies that GRF7 has roles in repressing both ABA-dependent and -independent gene expression (Figure 2). In contrast to *DREB2A*, transcriptional regulatory mechanisms of *AREB/ABFs* are poorly understood. Further studies of *AREB/ABF* promoters may help to elucidate whole transcriptional networks governed by ABA-dependent and -independent signaling.

Transcriptional networks under osmotic stress have been recently investigated using large-scale data analyses. A differential network analysis based on machine learning, an intelligent data mining technique, was used to reanalyze an abiotic stress-responsive gene expression data set. The analysis identified two novel genes whose mutants are sensitive to salt stress [43]. Integrated transcriptome and metabolome analyses have revealed the roles of ABA-dependent and -independent signaling in metabolic alterations under osmotic stress conditions in *Arabidopsis* and rice [44,45]. Cytokinin signaling is also reportedly decreased by dehydration stress in these two species [45]. This finding supports the idea that crosstalk between ABA and cytokinin plays key roles in osmotic stress signaling [46]. Furthermore, transcriptome and metabolome profiling of *Arabidopsis* grown under mild osmotic stress conditions has shown that cell proliferation and expansion are mainly regulated by ethylene and gibberellin signaling in developing leaves [47]. Importantly, a large-scale phenotyping study, which measured the rosette size of *Arabidopsis* grown in soil using an automated

platform, has suggested that improved drought stress tolerance under lethal conditions is not well correlated with superior growth under moderate drought conditions [48]. Thus, more attention must be focused on the roles of other plant hormones besides ABA to understand the molecular mechanisms used by plants to rigorously control their tolerance and growth under osmotic stress conditions.

Conclusions

In this review, we demonstrated that four AREB/ABF transcription factors, AREB1/ABF2, AREB2/ABF4, ABF3 and ABF1, regulate most downstream genes of three subclass III SnRK2s in ABA-dependent gene expression. Under unstressed conditions, transcriptional and protein levels of DREB2A, a key transcription factor in the ABA-independent pathway, are properly regulated by GRF7 and DRIPs, respectively. Given that *AREB/ABF* and *DREB2A* expression is regulated by the three subclass III SnRK2s, these kinases might serve as a convergence point in the crosstalk between ABA-dependent and -independent gene expression. Because AREB/ABF and DREB2A transcription factors are conserved in land plants [6,34], these transcription factors are available for applied research. Their orthologous genes have indeed been shown to participate in stress-responsive gene expression in crops such as rice and soybean as well as *Arabidopsis* [6,49]. For future application of these transcription factors to various crop, timber and other plants, research involving large-scale analyses, such as transcriptome and phosphoproteome analyses, should provide further insights into transcriptional networks under osmotic stress conditions.

ACKNOWLEDGMENTS

We thank E. Toma for skillful editorial assistance. This work was supported by a Grant-in-Aid for Scientific Research on Innovative Areas (no. 22119004) from the Ministry of Education, Culture, Sports, Science and Technology of Japan (MEXT), the Program for the Promotion of Basic Research Activities for Innovative Biosciences (BRAIN) of Japan,

and the Science and Technology Research Partnership for Sustainable Development (SATREPS) of the Japan Science and Technology Agency (JST)/Japan International Cooperation Agency (JICA).

5

References

1. Bartels D, Sunkar R: **Drought and salt tolerance in plants.** *Critical Reviews in Plant Sciences* 2005, **24**:23-58.
 - 10 2. Yamaguchi-Shinozaki K, Shinozaki K: **Transcriptional regulatory networks in cellular responses and tolerance to dehydration and cold stresses.** *Annu Rev Plant Biol* 2006, **57**:781-803.
 3. Finkelstein R: **Abscisic Acid synthesis and response.** *The Arabidopsis Book* 2013, **11**:e0166. doi: <http://dx.doi.org/10.1199/tab.0166>
 - 15 4. Fujita Y, Fujita M, Shinozaki K, Yamaguchi-Shinozaki K: **ABA-mediated transcriptional regulation in response to osmotic stress in plants.** *J Plant Res* 2011, **124**:509-525.
 5. Qin F, Shinozaki K, Yamaguchi-Shinozaki K: **Achievements and challenges in understanding plant abiotic stress responses and tolerance.** *Plant Cell Physiol* 20 2011, **52**:1569-1582.
 - * 6. Fujita Y, Yoshida T, Yamaguchi-Shinozaki K: **Pivotal role of the AREB/ABF-SnRK2 pathway in ABRE-mediated transcription in response to osmotic stress in plants.** *Physiol Plant* 2013, **147**:15-27.
- This current review focuses on the roles of AREB/ABF transcription factors and SnRK2
- 25 kinases in ABA signaling. This paper also summarizes the nomenclature and phylogenetic relationships of the AREB/ABFs and SnRK2s to help promote the link between basic and applied studies.
7. Fujita Y, Fujita M, Satoh R, Maruyama K, Parvez MM, Seki M, Hiratsu K, Ohme-Takagi M, Shinozaki K, Yamaguchi-Shinozaki K: **AREB1 is a transcription activator of**
 - 30 **novel ABRE-dependent ABA signaling that enhances drought stress tolerance in**

- Arabidopsis**. *Plant Cell* 2005, **17**:3470-3488.
8. Kang JY, Choi HI, Im MY, Kim SY: **Arabidopsis basic leucine zipper proteins that mediate stress-responsive abscisic acid signaling**. *Plant Cell* 2002, **14**:343-357.
 9. Kim S, Kang JY, Cho DI, Park JH, Kim SY: **ABF2, an ABRE-binding bZIP factor, is an essential component of glucose signaling and its overexpression affects multiple stress tolerance**. *Plant J* 2004, **40**:75-87.
 10. Furihata T, Maruyama K, Fujita Y, Umezawa T, Yoshida R, Shinozaki K, Yamaguchi-Shinozaki K: **Abscisic acid-dependent multisite phosphorylation regulates the activity of a transcription activator AREB1**. *Proc Natl Acad Sci U S A* 2006, **103**:1988-1993.
 11. Yoshida T, Fujita Y, Sayama H, Kidokoro S, Maruyama K, Mizoi J, Shinozaki K, Yamaguchi-Shinozaki K: **AREB1, AREB2, and ABF3 are master transcription factors that cooperatively regulate ABRE-dependent ABA signaling involved in drought stress tolerance and require ABA for full activation**. *Plant J* 2010, **61**:672-685.
 12. Uno Y, Furihata T, Abe H, Yoshida R, Shinozaki K, Yamaguchi-Shinozaki K: **Arabidopsis basic leucine zipper transcription factors involved in an abscisic acid-dependent signal transduction pathway under drought and high-salinity conditions**. *Proc Natl Acad Sci U S A* 2000, **97**:11632-11637.
 13. Fujii H, Verslues PE, Zhu JK: **Identification of two protein kinases required for abscisic acid regulation of seed germination, root growth, and gene expression in Arabidopsis**. *Plant Cell* 2007, **19**:485-494.
 14. Boudsocq M, Barbier-Brygoo H, Laurière C: **Identification of nine sucrose nonfermenting 1-related protein kinases 2 activated by hyperosmotic and saline stresses in Arabidopsis thaliana**. *J Biol Chem* 2004, **279**:41758-41766.
 15. Fujita Y, Nakashima K, Yoshida T, Katagiri T, Kidokoro S, Kanamori N, Umezawa T, Fujita M, Maruyama K, Ishiyama K, et al.: **Three SnRK2 protein kinases are the main positive regulators of abscisic acid signaling in response to water stress in Arabidopsis**. *Plant Cell Physiol* 2009, **50**:2123-2132.
 16. Fujii H, Zhu JK: **Arabidopsis mutant deficient in 3 abscisic acid-activated protein**

- kinases reveals critical roles in growth, reproduction, and stress. *Proc Natl Acad Sci U S A* 2009, **106**:8380-8385.
17. Fujii H, Chinnusamy V, Rodrigues A, Rubio S, Antoni R, Park SY, Cutler SR, Sheen J, Rodriguez PL, Zhu JK: **In vitro reconstitution of an abscisic acid signalling pathway**. *Nature* 2009, **462**:660-664.
18. Nakashima K, Fujita Y, Kanamori N, Katagiri T, Umezawa T, Kidokoro S, Maruyama K, Yoshida T, Ishiyama K, Kobayashi M, et al.: **Three Arabidopsis SnRK2 protein kinases, SRK2D/SnRK2.2, SRK2E/SnRK2.6/OST1 and SRK2I/SnRK2.3, involved in ABA signaling are essential for the control of seed development and dormancy**. *Plant Cell Physiol* 2009, **50**:1345-1363.
- ** 19. Umezawa T, Sugiyama N, Takahashi F, Anderson JC, Ishihama Y, Peck SC, Shinozaki K: **Genetics and phosphoproteomics reveal a protein phosphorylation network in the abscisic acid signaling pathway in Arabidopsis thaliana**. *Sci Signal* 2013, **6**:rs8.
- 15 This is the first study integrating genetics with phosphoproteomics to investigate protein phosphorylation networks in *Arabidopsis*. By applying a phosphoproteomic approach to the triple knockout mutant of the three subclass III SnRK2 kinases, the authors identifies multiple components of the ABA-responsive protein phosphorylation network governed by the SnRK2s, including AREB/ABFs, AtMPK1, AtMPK2 and SNS1.
- 20 ** 20. Wang P, Xue L, Batelli G, Lee S, Hou YJ, Van Oosten MJ, Zhang H, Tao WA, Zhu JK: **Quantitative phosphoproteomics identifies SnRK2 protein kinase substrates and reveals the effectors of abscisic acid action**. *Proc Natl Acad Sci U S A* 2013, **110**:11205-11210.
- 25 This phosphoproteomic study using the triple knockout mutant of the three subclass III SnRK2s also identifies candidate substrates of the SnRK2s in ABA signaling, including proteins involved in flowering time regulation, RNA and DNA binding, miRNA and epigenetic regulation, signal transduction, chloroplast function and many other cellular processes. Given that the *srk2d/e/i* triple mutant shows the early
- 30 flowering phenotype, the authors suggest that the SnRK2s are important for

flowering time regulation.

21. Ito S, Song YH, Josephson-Day AR, Miller RJ, Breton G, Olmstead RG, Imaizumi T:
**FLOWERING BHLH transcriptional activators control expression of the
photoperiodic flowering regulator CONSTANS in Arabidopsis.** *Proc Natl Acad
Sci U S A* 2012, **109**:3582-3587.
22. Mougou N, Poulis S, Kaldis A, Vlachonasios K: **Arabidopsis thaliana
TBP-associated factor 5 is essential for plant growth and development.**
Molecular Breeding 2012, **30**:355-366.
- * 23. Yoshida T, Fujita Y, Maruyama K, Mogami J, Todaka D, Shinozaki K,
Yamaguchi-Shinozaki K: **Four Arabidopsis AREB/ABF transcription factors
function predominantly in gene expression downstream of SnRK2 kinases in
abscisic-acid signaling in response to osmotic stress.** *Plant Cell Environ* 2014.
- Based on microarray data from the *areb1 areb2 abf3 abf1*, *areb1 areb2 abf3* and *srk2d/e/i*
mutants, this work demonstrates that most of the downstream genes of the three
subclass III SnRK2s are regulated by four AREB/ABF transcription factors in ABA
signaling during vegetative growth.
24. Takahashi Y, Ebisu Y, Kinoshita T, Doi M, Okuma E, Murata Y, Shimazaki K: **bHLH
Transcription Factors That Facilitate K⁺ Uptake During Stomatal Opening Are
Repressed by Absciscic Acid Through Phosphorylation.** *Sci Signal* 2013, **6**:ra48.
25. Ando E, Ohnishi M, Wang Y, Matsushita T, Watanabe A, Hayashi Y, Fujii M, Ma JF,
Inoue S, Kinoshita T: **TWIN SISTER OF FT, GIGANTEA, and CONSTANS
have a positive but indirect effect on blue light-induced stomatal opening in
Arabidopsis.** *Plant Physiol* 2013, **162**:1529-1538.
26. Sato A, Sato Y, Fukao Y, Fujiwara M, Umezawa T, Shinozaki K, Hibi T, Taniguchi M,
Miyake H, Goto DB, et al.: **Threonine at position 306 of the KAT1 potassium
channel is essential for channel activity and is a target site for ABA-activated
SnRK2/OST1/SnRK2.6 protein kinase.** *Biochem J* 2009, **424**:439-448.
27. Sakuma Y, Liu Q, Dubouzet JG, Abe H, Shinozaki K, Yamaguchi-Shinozaki K:
**DNA-binding specificity of the ERF/AP2 domain of Arabidopsis DREBs,
transcription factors involved in dehydration- and cold-inducible gene**

- expression.** *Biochem Biophys Res Commun* 2002, **290**:998-1009.
28. Sakuma Y, Maruyama K, Qin F, Osakabe Y, Shinozaki K, Yamaguchi-Shinozaki K: **Dual function of an Arabidopsis transcription factor DREB2A in water-stress-responsive and heat-stress-responsive gene expression.** *Proc Natl Acad Sci U S A* 2006, **103**:18822-18827.
29. Sakuma Y, Maruyama K, Osakabe Y, Qin F, Seki M, Shinozaki K, Yamaguchi-Shinozaki K: **Functional analysis of an Arabidopsis transcription factor, DREB2A, involved in drought-responsive gene expression.** *Plant Cell* 2006, **18**:1292-1309.
- ** 30. Kim JS, Mizoi J, Kidokoro S, Maruyama K, Nakajima J, Nakashima K, Mitsuda N, Takiguchi Y, Ohme-Takagi M, Kondou Y, et al.: **Arabidopsis growth-regulating factor7 functions as a transcriptional repressor of abscisic acid- and osmotic stress-responsive genes, including DREB2A.** *Plant Cell* 2012, **24**:3393-3405.
- This work identifies a transcriptional repressor of ABA- and osmotic stress-responsive genes, providing a new insight into how key transcription factors in osmotic stress signaling such as DREB2A are negatively regulated under unstressed conditions.
31. Qin F, Sakuma Y, Tran LS, Maruyama K, Kidokoro S, Fujita Y, Fujita M, Umezawa T, Sawano Y, Miyazono K, et al.: **Arabidopsis DREB2A-interacting proteins function as RING E3 ligases and negatively regulate plant drought stress-responsive gene expression.** *Plant Cell* 2008, **20**:1693-1707.
32. Morimoto K, Mizoi J, Qin F, Kim JS, Sato H, Osakabe Y, Shinozaki K, Yamaguchi-Shinozaki K: **Stabilization of Arabidopsis DREB2A is required but not sufficient for the induction of target genes under conditions of stress.** *PLoS One* 2013, **8**:e80457.
33. Cheng MC, Hsieh EJ, Chen JH, Chen HY, Lin TP: **Arabidopsis RGLG2, functioning as a RING E3 ligase, interacts with AtERF53 and negatively regulates the plant drought stress response.** *Plant Physiol* 2012, **158**:363-375.
34. Mizoi J, Shinozaki K, Yamaguchi-Shinozaki K: **AP2/ERF family transcription factors in plant abiotic stress responses.** *Biochim Biophys Acta* 2012, **1819**:86-96.
35. Hsieh EJ, Cheng MC, Lin TP: **Functional characterization of an abiotic stress-inducible transcription factor AtERF53 in Arabidopsis thaliana.** *Plant*

Mol Biol 2013, **82**:223-237.

36. Stone SL, Hauksdóttir H, Troy A, Herschleb J, Kraft E, Callis J: **Functional analysis of the RING-type ubiquitin ligase family of Arabidopsis.** *Plant Physiol* 2005, **137**:13-30.
- 5 37. Fujii H, Verslues PE, Zhu JK: **Arabidopsis decuple mutant reveals the importance of SnRK2 kinases in osmotic stress responses in vivo.** *Proc Natl Acad Sci U S A* 2011, **108**:1717-1722.
38. Yoshida R, Umezawa T, Mizoguchi T, Takahashi S, Takahashi F, Shinozaki K: **The regulatory domain of SRK2E/OST1/SnRK2.6 interacts with ABI1 and integrates abscisic acid (ABA) and osmotic stress signals controlling stomatal closure in Arabidopsis.** *J Biol Chem* 2006, **281**:5310-5318.
- 10 39. Boudsocq M, Droillard MJ, Barbier-Brygoo H, Laurière C: **Different phosphorylation mechanisms are involved in the activation of sucrose non-fermenting 1 related protein kinases 2 by osmotic stresses and abscisic acid.** *Plant Mol Biol* 2007, **63**:491-503.
- 15 * 40. E Stecker K, Minkoff BB, Sussman MR: **Phosphoproteomic Analyses Reveal Early Signaling Events in the Osmotic Stress Response.** *Plant Physiol* 2014, **165**:1171-1187.

This work shows untargeted and targeted quantitative phosphoproteomic approaches are powerful tools for studying protein phosphorylation events in response to short-term stimuli. The authors also identify a couple of proteins rapidly phosphorylated by osmotic stresses, which may be involved in the initial perception and response to dehydration.

20

41. Kim JS, Mizoi J, Yoshida T, Fujita Y, Nakajima J, Ohori T, Todaka D, Nakashima K, Hirayama T, Shinozaki K, et al.: **An ABRE promoter sequence is involved in osmotic stress-responsive expression of the DREB2A gene, which encodes a transcription factor regulating drought-inducible genes in Arabidopsis.** *Plant Cell Physiol* 2011, **52**:2136-2146.
- 25
42. Lee SJ, Kang JY, Park HJ, Kim MD, Bae MS, Choi HI, Kim SY: **DREB2C interacts with ABF2, a bZIP protein regulating abscisic acid-responsive gene expression,**
- 30

- and its overexpression affects abscisic acid sensitivity. *Plant Physiol* 2010, **153**:716-727.
43. Ma C, Xin M, Feldmann KA, Wang X: **Machine learning-based differential network analysis: a study of stress-responsive transcriptomes in Arabidopsis.** *Plant Cell* 2014, **26**:520-537.
44. Urano K, Maruyama K, Ogata Y, Morishita Y, Takeda M, Sakurai N, Suzuki H, Saito K, Shibata D, Kobayashi M, et al.: **Characterization of the ABA-regulated global responses to dehydration in Arabidopsis by metabolomics.** *Plant J* 2009, **57**:1065-1078.
45. Maruyama K, Urano K, Yoshiwara K, Morishita Y, Sakurai N, Suzuki H, Kojima M, Sakakibara H, Shibata D, Saito K, et al.: **Integrated analysis of the effects of cold and dehydration on rice metabolites, phytohormones, and gene transcripts.** *Plant Physiol* 2014, **164**:1759-1771.
46. Ha S, Vankova R, Yamaguchi-Shinozaki K, Shinozaki K, Tran LS: **Cytokinins: metabolism and function in plant adaptation to environmental stresses.** *Trends Plant Sci* 2012, **17**:172-179.
47. Skirycz A, De Bodt S, Obata T, De Clercq I, Claeys H, De Rycke R, Andriankaja M, Van Aken O, Van Breusegem F, Fernie AR, et al.: **Developmental stage specificity and the role of mitochondrial metabolism in the response of Arabidopsis leaves to prolonged mild osmotic stress.** *Plant Physiol* 2010, **152**:226-244.
48. Skirycz A, Vandenbroucke K, Clauw P, Maleux K, De Meyer B, Dhondt S, Pucci A, Gonzalez N, Hoeberichts F, Tognetti VB, et al.: **Survival and growth of Arabidopsis plants given limited water are not equal.** *Nat Biotechnol* 2011, **29**:212-214.
49. Mizoi J, Ohori T, Moriwaki T, Kidokoro S, Todaka D, Maruyama K, Kusakabe K, Osakabe Y, Shinozaki K, Yamaguchi-Shinozaki K: **GmDREB2A;2, a canonical DEHYDRATION-RESPONSIVE ELEMENT-BINDING PROTEIN2-type transcription factor in soybean, is posttranslationally regulated and mediates dehydration-responsive element-dependent gene expression.** *Plant Physiol* 2013, **161**:346-361.

Figure Legends

5

Figure 1. ABA-dependent signaling pathway and crosstalk with ABA-independent signaling in response to osmotic stress. The four AREB/ABF transcription factors, AREB1, AREB2, ABF3, and ABF1, play key roles in ABA-dependent gene expression. As pivotal positive regulators downstream of PP2C-PYR/PYL/RCAR ABA receptor complexes in ABA signaling, subclass III SnRK2s phosphorylate a variety of substrates, including AREB/ABFs, FBH3/AKS1, and SNS1, and modulate their activities. Recent studies suggest that crosstalk occurs between the AREB/ABF-SnRK2 pathway in ABA signaling and the ABA-independent pathway. Transcription factors and DNA-binding proteins are shown in colored ellipses. Dashed lines indicate possible although unconfirmed routes.

10

15 PYR/PYL/RCAR, pyrabactin resistance1/PYR1-like/regulatory components of ABA receptor; PP2C, protein phosphatase 2C; CE, coupling element.

Figure 2. ABA-independent signaling pathway and crosstalk with ABA-dependent signaling in response to osmotic stress. DREB2A is the key transcription factor in ABA-independent gene expression in response to osmotic stress. DREB2A protein levels are tightly regulated by ubiquitin E3 ligases such as DRIP1 and DRIP2. In addition, *DREB2A* expression is repressed by GRF7 under unstressed growth conditions. Crosstalk between ABA-independent and -dependent pathways is suggested on the basis of recent findings that GRF7 is involved in repression of ABA- and osmotic stress-inducible genes and that *DREB2A* is induced by AREB/ABFs. Transcription factors and DNA-binding proteins are shown in colored ellipses. Dashed lines indicate possible although unconfirmed routes.

20

25

PYR/PYL/RCAR, pyrabactin resistance1/PYR1-like/regulatory components of ABA receptor; PP2C, protein phosphatase 2C; GTE, GRF7-targeting *cis*-element; HSE, heat shock element; Ub, ubiquitin.

Figure 1.

Osmotic stress

ABA-independent
pathway

ABA-dependent
pathway

Figure 2.

Osmotic stress