Root signals that mediate mutualistic interactions in the rhizosphere

Sergio Rasmann* & Ted C. J. Turlings

Abstract

A recent boom in research on belowground ecology is rapidly revealing a multitude of fascinating interactions, in particular in the rhizosphere. Many of these interactions are mediated by photo-assimilates that are excreted by plant roots. Root exudates are not mere waste products, but serve numerous functions to control abiotic and biotic processes. These functions range from changing the chemical and physical properties of the soil, inhibiting the growth of competing plants, combating herbivores, and regulating the microbial community. Particularly intriguing are root-released compounds that have evolved to serve mutualistic interactions with soil-dwelling organisms. These mutually beneficial plant-mediated signals are not only of fundamental ecological interest, but also exceedingly important from an agronomical perspective. Here, we attempt to provide an overview of the plant-produced compounds that have so far been implicated in mutualistic interactions. We propose that these mutualistic signals may have evolved from chemical defenses and we point out that they can be (mis)used by specialized pathogens and herbivores. We speculate that many more signals and interactions remain to be uncovered and that a good understanding of the mechanisms and ecological implications can be the basis for exploitation and manipulation of the signals for crop improvement and protection.

Keywords – Benzoxazinoids, biological control, entomopathogenic nematodes, flavonoids, insect pests, root exudates, root herbivores, soil microbes, terpenes.

Introduction

It has been estimated that between 20 to 80% of the total photosynthates enters the rhizosphere as root exudates [1]. These exudates are primarily composed of carbon-containing compounds, including the respired CO₂, but also of non-carbon-containing compounds including ubiquitous H⁺, inorganic ions, water, and electrons [2]. A large fraction of the rhizodeposition also involves compounds such as phytosiderophores for improving nutrient acquisition [3,4], or polysaccharides, that, in association with rhizospheric microbial cells and clay particles, form a mucigel [5,6]. This mucigel provides protection and lubrication, and can create a suitable environment for symbiotic microorganisms. In this environment other exudates can operate as signals to mediate the interactions with plant-beneficial organisms [7-9].

Indeed, a rapidly increasing number of studies is revealing that plant roots exude a variety of organic compounds that are not directly associated with plant nutrition or growth [9-11]. These may include sugars and simple polysaccharides (such as arabinose, fructose, glucose, maltose, mannose, oligosaccharides), amino acids (such as arginine, asparagine, aspartic, cysteine, cystine, glutamine), organic acids (such as acetic, ascorbic, benzoic, ferulic, malic acids), and phenolic compounds [2]. In addition, roots may exude higher-molecular-weight compounds such as flavonoids, enzymes, fatty acids, growth regulators, nucleotides, tannins, carbohydrates, steroids, terpenoids, alkaloids, polyacetylenes, and vitamins [12-14]. Many of these compounds are thought to mediate interactions with other organisms in the rhizosphere.

Below, we review how root-produced signals may mediate mutualistic interactions (Fig 1.), but also stress that the same signals may be opportunistically used by plant antagonists. We point out gaps in the literature and reiterate that root-

emitted signals might be exploited for application, for instance in crop protection [15].

Plants send chemical cues to other organisms

Plant-plant interaction – Chemically mediated plant-plant interactions have been extensively studied for aboveground tissues, particularly in the context of herbivore-induced volatiles that can be perceived by neighbouring plants and “warn” them for imminent attack [16,17]. Specific volatiles are essential for this plant-to-plant signalling [18,19] and may be exploitable for crop protection [20,21]. This should also be explored for belowground plant-plant signalling.

Indeed, it has been shown that when *Vicia faba* (broad bean) plants are subjected to aphid attack, they release root exudates that induce the release of volatiles in undamaged neighbouring plants, which in turn attracts aphid parasitoids [22]. Similarly, *Phaseolus lunatus* (lima bean) plants under attack by spider mites produce root exudates that induce the release of volatiles in other plants that attract predatory mites [23]. For now, the only other known mutually beneficial underground interaction involves fungal networks via which neighbouring plants can be informed of impending insect attack [24]. The specific signals that are involved in these interactions remain to be elucidated. Although for one type of plant-produced secondary metabolites (iridoid glycosides from *Plantago lanceolata*) it is known that they can be transferred to the fungal hyphae, protecting them against fungivore attack [25], and potentially affecting soil-dwelling herbivores too.

Considerably more is known about the chemicals involved in belowground plant-plant interactions that have negative, allelopathic, effects. Allelopathy involves

the release of chemicals by a plant that inhibits seed germination or growth in other plants. Allelopathic root exudates are worth mentioning here because they are also available as signals that may be exploited by plant mutualists. This appears to be the case for the benzoxazinoids, like DIMBOA, released by maize, wheat and other grasses roots. Like many other plant produced toxins, DIMBOA has allelopathic properties [12], but it is also induced upon root feeding by insects [26], and may have several signalling functions (Box 1).

Plant-arthropod interaction – Soil arthropod communities are largely composed of immature stage of insects (particularly in the case of Coleoptera and Diptera), collembolan, and mites. They can be involved in positive interactions (such as through the indirect positive effects of decomposers), but also in negative interactions, since many of the soil insect larvae feed on roots [11].

In the context of mutualistic root-produced signals, decomposers have not yet been considered. Decomposition by arthropods and other invertebrates, such as earthworms, have profound effects on nutrient cycling, which in turn affects plant performance [27], and defence [28]. It might therefore be highly advantageous for plants to send cues into the soil to attract specific decomposers. This remains to our knowledge unexplored and merits further investigation.

Chemically-mediated host location by root-feeding insects has been extensively reviewed by Johnson and Nielsen [29], showing that several dozens of compounds in root exudates can serve as foraging cues [30]. Such negative consequences of releasing these particular exudates are likely to be counter-balanced by interactions with positive effects for the plant. For instance, adult soil-dwelling arthropods are often predatory and may use plant exudates to find herbivore preys

[31]. Therefore, similar to what has been described for aboveground plant-herbivore interactions [32], co-evolutionary adaptations of defence and counter-defence between plants and their herbivores appear to also take place belowground. An illustrative example are the highly specialized larvae of *Diabrotica virgifera virgifera* that exploit maize-produced benzoxazinoids, that are toxic to most herbivores, to identify the most suitable roots [33] (Box 1). The same larvae also use an inducible root volatiles involved in indirect (see below) to aggregate, as they develop better when multiple larvae feed on the same root system [34]. But this same compound, the sesquiterpene (*E*)- β -caryophyllene, also benefits the plant by attracting entomopathogenic nematodes (see below and Box 2).

Signalling between plants and microbes (bacteria and fungi)- Plant associations with soil mutualistic microbes are largely mediated by root exudation. Three specific groups of organisms can be identified: mycorrhizal fungi, nitrogen fixing bacteria, and growth promoting bacteria [35].

Arbuscular mycorrhizal fungi (AMF) and plant roots form associations in more than 80% of terrestrial plants [36]. These symbiotic relationships facilitate nutrient uptake by the plant, and in turn, the associated fungi extract lipids and carbohydrates from the host root. AMF have also been shown to indirectly benefit plants, by increasing plant resistance against herbivores and pathogens in several systems [37-40], but not all [41]. As obligate biotrophs, the survival of these fungi depends on their ability to rapidly establish a symbiotic relationship with plant roots. The growth and branching of germinating hyphae prior to root infection require the presence of compounds released by roots [42]. Among the diversity of molecules present in root exudates and active in hyphal branching, flavonoids have often been

proposed to switch AMF from non- to pre-symbiotic growth [43]. Strigolactones, which are carotenoid-derived terpenoids, have been found to stimulate cell proliferation in the AM fungus *Gigaspora rosea*, and spore germination in *Glomus intraradices* and *Gl. claroideum* [43], and they induce branching in *G. margarita* [44]. Strigolactones are also involved in several other belowground interactions, again showing complex ecological and evolutionary dynamics at play (see Box 2).

Plants may also release root exudates to recruit nitrogen fixing and growth promoting bacteria (e.g. Rhizobia). A large number of plant species, mainly in the Fabaceae family profit from the intimate association with nitrogen-fixing bacteria and again plant-produced flavonoids are involved in the establishment of these associations [45] (Fig. 1). Interestingly, because most legume species mainly produce their own specific isoflavonoids, it enables the Rhizobia to selectively colonize their specific host plants [10]. Finally, plants may produce exudates to attract plant-growth-promoting bacteria into their rhizosphere [46]. These free-living bacteria have been shown to be attracted to root-produced carbohydrates and amino-acids, including the primary metabolite tryptophan [10]. Recent evidence shows that benzoxazinoids may act similarly [47] (Box 1).

Whether root exudates can also attract the highly ubiquitous soil protists and how this may affect plant-microbe interactions is to our knowledge unknown. Protists have obvious associations with plants [48]. Several guilds of protists are microbial feeders, and should be attracted to locations with high-energy sources, generating a microbial loop that affects plant performance [49]. It is therefore highly likely that root exudates also mediate protist orientation.

156 *Interaction between plants and herbivore-killing predators and pathogens –*

157 Plants can benefit from attracting predators and pathogens to the sites where they are
158 attacked by herbivores. Indeed, root-feeding insects have plenty natural enemies,
159 including arthropods such as predatory mites (Acari), ants (Formicidae), carabid
160 beetles (Carabidae), staphilinid beetles (Staphylinidae), and spiders (Aranea),
161 predatory nematodes, entomopathogenic nematodes (EPNs), entomopathogenic fungi,
162 and insect-pathogenic bacteria [50].

163 As yet, there is limited evidence that root exudates can attract soil-dwelling
164 predatory arthropods, but clearly this has potential to improve biological control
165 [31,50]. For example, it was found that the predatory mite *Neoseiulus cucumeris* is
166 attracted to tulip bulbs infested with rust mites (*Aceria tulipae*). No such attraction
167 was found for mechanically damaged bulbs, strongly suggesting that the bulbs
168 produce a specific odor in response to rust mite infestation [51], but this remains to be
169 elucidated. Similarly, we are not aware of evidence for chemically-mediated
170 mutualistic interactions between entomopathogenic microbes and plants [11]. Besides
171 releasing attractants, it is plausible that plants have evolved mechanisms, such as
172 maintaining a nutrient-rich rhizosphere, in order to retain mutualistic microbes
173 in the rhizosphere, which would represent an evolutionary advantage for both partners
174 [52].

175 Much more is known about the chemically-mediated interactions between
176 roots and entomopathogenic nematodes (EPN). The underlining idea is that EPN use
177 root exudates in their efforts to find hosts, which are predominantly arthropod root
178 herbivores [53,54]. Once found and penetrated, EPNs rapidly kill their hosts, and
179 thereby ultimately increase plant fitness [55]. Indeed, a combination of behavioural
180 assays and metabolomics analyses has revealed the role for root-produced volatiles in

the foraging success of EPNs [56-61]. To our knowledge the successful manipulation of the release of such an EPN attractant represents the only attempt so far to enhance a mutualistic rhizosphere interaction (Box 2). The manipulation greatly improved the protection that maize roots receive from EPN [62], but, as mentioned above, follow-up work also revealed potential negative effects of such an approach, as the root signal is also used by herbivores to locate host plants [33,34]. This attraction may be dose-dependent [33], implying that the intensity of the root signals can be a key factor in determining whether the produced compound benefit the plant or not. This example illustrates the importance of studying all consequences of the signal production in order to optimize their possible application in crop improvement.

Conclusions and perspectives

We here presented the highlights of what is currently known about how root exudates can benefit plants via interaction with other soil-dwelling organisms. Methodological limitations to studying belowground processes are slowly vanishing, opening the way for many new chemically-mediated interactions to be discovered [8,9,11]. With these new discoveries come new opportunities to study plant exudates and gain further insights into their importance in structuring ecological communities, and, from an applied perspective, to exploit the interactions that they mediate to enhance crop yields.

To achieve this we need to understand the relative magnitude of the effects of root exudates on plant fitness across a wide range of natural conditions [63]. This relates to the new frontier in ecological research, which focuses on assessing habitat-mediated intra- and inter-specific variation of root exudation, and subsequent soil ecology effects.

206 From an applied perspective further insight into the signals that are needed to
207 attract soil-dwelling beneficial organisms is badly needed. Specifically, unravelling
208 the mechanisms that are involved in the synthesis and exudation of these signals
209 should make it possible to breed for crops with enhanced capacities to exploit the soil
210 biota for optimal plant health and yields.

211

212 Acknowledgments – We are grateful to Consuelo De Moraes and Mark Mescher for
213 inviting us to write this review. We thank Thomas Degen (www.thomas-degen.ch)
214 for designing the figure. This review was inspired by several rhizosphere-related
215 research projects that are funded by the Swiss National Science Foundation (Sinergia
216 CRSII3_160786, NRP68 406840_161904 to TCJT and PBNE33-114692,
217 PA0033_121483, and 31003A_159869 to SR)

218

Figure 1. Examples of the diverse range of root exudates that are implicated in mutualistic rhizospheric interactions as outlined in the text.

Box 1. *Benzoxazinoids* - Benzoxazinoids (BXs) are natural hydroxamic acids that that are best known as the main chemical defence in many grasses, including important cereals like maize, rye and wheat. In the plants, the BXs are typically glycosidically bound and stored in the vacuole. The toxic and protective aglycones are only liberated upon exposure to β -glucosidases during tissue damage and/or ingestion by herbivores [64-66]. Specialized herbivores may detoxify the aglycones [67-69] and even use them for their own benefit [33]. The best known of the BXs is DIMBOA (2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one), which has been implicated in resistance against insect pests, as well as a number of pathogens [70]. The concentrations of BXs are typically higher in younger plants [71], and can be released in the rhizosphere [72]. The implications of this are still largely unknown, but it is increasingly clear that BXs have an effect on microbiota in the soil. For example, it was shown that DIMBOA and MBOA affected the soil microbial community structure near wheat plants to their competitive advantage through changes in fungal community composition [73]. Similarly, DIMBOA, the dominant BX in the exudates of maize roots, provokes chemotactic responses in *Pseudomonas putida* KT2440, a bacterium with plant-beneficial traits. The role of BXs in the recruitment of *P. putida* was elegantly confirmed with the use of maize mutants impaired in the production of BXs [47]. A complete understanding of these interactions will greatly facilitate their conceivable application in agriculture.

Box 2 *Terpenoids* - Several root-produced compounds in the chemical class of the terpenoids are involved in numerous belowground mutualistic processes, such as the

sesquiterpenes (*E*)- β -caryophyllene and pregeijerene, that attracts entomopathogenic nematodes (EPN) [56,58], and strigolactones that facilitate root associations with arbuscular mycorrhizal fungi (AMF) [44].

The role of terpenoids in EPN attraction was first shown when we investigated EPNs as potential biological control agents against the western corn rootworm (WCR), *Diabrotica virgifera virgifera*, a major pest of maize roots. In response to WCR feeding maize roots release (*E*)- β -caryophyllene [58]. The importance of (*E*)- β -caryophyllene as an EPN attractant was confirmed by restoring its release in a maize variety that has lost the ability to produce it [62,74]. It should be noted that in combination with CO₂, (*E*)- β -caryophyllene is considerably more attractive [75]. This strong synergistic effect may also be true for other interactions, or other exudates that are commonly released from roots. Similarly, EPNs have been considered for the biological control of weevil larvae (Coleoptera: Curculionidae) attacking *Citrus* trees. Upon weevil damage the tree roots release several terpenes that recruit EPN to the site of wounding [56,57]. Enhancing the production of these root exudates can improve crop protection by EPNs (Degenhardt et al., 2009), but also the EPNs themselves can be selected to respond better to the root-produced attractants [60].

Strigolactones have been particularly well studied in the context of plant-beneficial interactions, but also in the recruitment of antagonists. These terpenoid lactones are carotenoid-derived plant hormones that regulate developmental processes, but they are also excreted into the rhizosphere where they recruit symbiotic fungi, but also root parasitic plants [76,77]. Strigolactones are a wonderful example of how mutualistic signalling can be abused by the plants' enemies, in this specific case they serve as germination stimulants for parasitic plants of the genus *Striga* from which they derive their name [e.g. 78,79]. In an evolutionary context, belowground

strigolactone production is likely to be subject to diffuse selection, as is the case for aboveground emitted compounds [80]. Therefore, their production can only be maintained if other beneficial functions outweigh the detrimental effects. We argue that the fact that strigolactones trigger seed germination in parasitic *Striga* plants [76] is a case of “eavesdropping” on a beneficial interaction of the plants with mutualistic fungi. If these negative effects could be minimized, strigolactones represent a wonderful example of how root-produced chemical cues could be enhanced to increase plant production in concomitance with the application mutualists. Strigolactones may also have a function as direct chemical defences against pathogens and herbivores, and efforts are underway to explore their overall potential for crop protection [81].

References

1. Coleman DC: **A review of root production processes and their influence on soil biota in terrestrial ecosystems.** In *The Role of Terrestrial and Aquatic Organisms in Decomposition Processes*. Edited by Anderson JM, Macfadyen A: Blackwell Scientific Publications; 1976:pp. 417 - 434.
2. Inderjit, Weston LA: **Root Exudates: an Overview.** In *Root Ecology*. Edited by Kroon H, Visser EJW: Springer Berlin Heidelberg; 2003:235-255.
3. Hopkins BG, Whitney DA, Lamond RE, Jolley VD: **Phytosiderophore release by sorghum, wheat, and corn under zinc deficiency.** *Journal of Plant Nutrition* 1998, **21**:2623-2637.
4. Romheld V, Marschner H: **Genotypical differences among gramineous species in release of phytosiderophores and uptake of iron.** *Plant and Soil* 1990, **123**:147-153.
5. Guckert A, Breisch H, Reisinger O: **Electron-microscopic study on interrelation of mucigel, clay and microorganisms.** *Soil Biology & Biochemistry* 1975, **7**:241-250.
6. Rovira AD: **Plant root exudates.** *Botanical Review* 1969, **35**:35-&.
7. Kiers ET, Denison RF: **Sanctions, cooperation, and the stability of plant-rhizosphere mutualisms.** *Annual Review of Ecology, Evolution, and Systematics* 2008, **39**:215-236.
8. Venturi V, Keel C: **Signaling in the Rhizosphere.** *Trends in Plant Science* 2016, **21**:187-198.

* Excellent companion paper to this one

9. van Dam NM, Bouwmeester HJ: **Metabolomics in the rhizosphere: Tapping into belowground chemical communication.** *Trends in plant science* 2016, **21**:256-265.
- * Excellent companion paper to this one
10. Bais HP, Weir TL, Perry LG, Gilroy S, Vivanco JM: **The role of root exudates in rhizosphere interactions with plants and other organisms.** *Annual Review of Plant Biology* 2006, **57**:233-266.
11. Johnson SN, Rasmann S: **Root-feeding insects and their interactions with organisms in the rhizosphere.** *Annual Review of Entomology* 2015, **60**:517-535.
- * Review on how root herbivores interact with other soil organisms
12. Bertin C, Yang XH, Weston LA: **The role of root exudates and allelochemicals in the rhizosphere.** *Plant and Soil* 2003, **256**:67-83.
13. Walker TS, Bais HP, Grotewold E, Vivanco JM: **Root exudation and rhizosphere biology.** *Plant Physiology* 2003, **132**:44-51.
14. Bais HP, Loyola-Vargas VM, Flores HE, Vivanco JM: **Root-specific metabolism: The biology and biochemistry of underground organs.** *In Vitro Cellular & Developmental Biology-Plant* 2001, **37**:730-741.
15. Hiltbold I, Turlings TC: **Manipulation of chemically mediated interactions in agricultural soils to enhance the control of crop pests and to improve crop yield.** *J Chem Ecol* 2012, **38**:641-650.
16. Karban R, Yang LH, Edwards KF: **Volatile communication between plants that affects herbivory: a meta-analysis.** *Ecology Letters* 2014, **17**:44-52.
- * A robust confirmation that damaged plants can increase resistance of their neighbours even when highly taxonomically different.
17. Heil M, Karban R: **Explaining evolution of plant communication by airborne signals.** *Trends in Ecology & Evolution* 2010, **25**:137-144.
18. Oluwafemi S, Dewhurst SY, Veyrat N, Powers S, Bruce TJA, Caulfield JC, Pickett JA, Birkett MA: **Priming of production in maize of volatile organic defence compounds by the natural plant activator cis Jasmon.** *PLoS ONE* 2013, **8**:e62299.
19. Erb M, Veyrat N, Robert CAM, Xu H, Frey M, Ton J, Turlings TCJ: **Indole is an essential herbivore-induced volatile priming signal in maize.** *Nature Communications* 2015, **6**.
- * Identification of a common plant volatile that mediates plant-plant communication
20. Ninkovic V, Glinwood R, Dahlin I: **Weed-barley interactions affect plant acceptance by aphids in laboratory and field experiments.** *Entomologia Experimentalis et Applicata* 2009, **133**:38-45.
21. Dewhurst SY, Birkett MA, Loza-Reyes E, Martin JL, Pye BJ, Smart LE, Hardie J, Pickett JA: **Activation of defence in sweet pepper, *Capsicum annum*, by cis-jasmone, and its impact on aphid and aphid parasitoid behaviour.** *Pest Manag Sci* 2012, **68**:1419-1429.
22. Chamberlain K, Guerrieri E, Pennacchio F, Pettersson J, Pickett JA, Poppy GM, Powell W, Wadhams LJ, Woodcock CM: **Can aphid-induced plant signals be transmitted aurally and through the rhizosphere?** *Biochemical Systematics and Ecology* 2001, **29**:1063-1074.
23. Dicke M, Dijkman H: **Within-plant circulation of systemic elicitor of induced defence and release from roots of elicitor that affects**

354 **neighbouring plants.** *Biochemical Systematics and Ecology* 2001,
355 **29:**1075-1087.

356 24. Babikova Z, Gilbert L, Bruce TJA, Birkett M, Caulfield JC, Woodcock C, Pickett
357 JA, Johnson D: **Underground signals carried through common**
358 **mycelial networks warn neighbouring plants of aphid attack.** *Ecology*
359 *Letters* 2013, **16:**835-843.

360 *Nice example of how fungi can mediate plant-plant interaction

361 25. Duhamel M, Pel R, Ooms A, Bucking H, Jansa J, Ellers J, van Straalen NM,
362 Wouda T, Vandenkoornhuyse P, Kiers ET: **Do fungivores trigger the**
363 **transfer of protective metabolites from host plants to arbuscular**
364 **mycorrhizal hyphae?** *Ecology* 2013, **94:**2019-2029.

365 * Perhaps the first example of an underground defense-mediated mutualism
366 between plants and fungi

367 26. Glauser G, Marti G, Villard N, Doyen GA, Wolfender J-L, Turlings TCJ, Erb M:
368 **Induction and detoxification of maize 1,4-benzoxazin-3-ones by**
369 **insect herbivores.** *Plant Journal* 2011, **68:**901-911.

370 27. Hättenschwiler S, Tiunov AV, Scheu S: **Biodiversity and litter**
371 **decomposition in terrestrial ecosystems.** *Annual Review of Ecology,*
372 *Evolution, and Systematics* 2005, **36:**191-218.

373 28. Wurst S: **Plant-mediated links between detritivores and aboveground**
374 **herbivores.** *Frontiers in Plant Science* 2013, **4.**

375 29. Johnson SN, Nielsen UN: **Foraging in the dark - chemically mediated host**
376 **plant location by belowground insect herbivores.** *Journal of Chemical*
377 *Ecology* 2012, **38:**604-614.

378 30. Erb M, Huber M, Robert CAM, Ferrieri AP, Machado RAR, Arce CCM: **The role**
379 **of plant primary and secondary metabolites in root-herbivore**
380 **behaviour, nutrition and physiology.** In *Advances in Insect Physiology.*
381 Edited by Scott N. Johnson IH, Ted CJT: Academic Press; 2013:53-95. vol
382 Volume 45.]

383 31. Kergunteuil A, Dugravot S, Mortreuil A, Le Ralec A, Cortesero AM: **Selecting**
384 **volatiles to protect brassicaceous crops against the cabbage root fly,**
385 **Delia radicum.** *Entomologia Experimentalis et Applicata* 2012, **144:**69-
386 77.

387 32. Pierik R, Ballare CL, Dicke M: **Ecology of plant volatiles: taking a plant**
388 **community perspective.** *Plant Cell and Environment* 2014, **37:**1845-
389 1853.

390 33. Robert CAM, Veyrat N, Glauser G, Marti G, Doyen GR, Villard N, Gaillard MDP,
391 Köllner TG, Giron D, Body M, et al.: **A specialist root herbivore exploits**
392 **defensive metabolites to locate nutritious tissues.** *Ecology Letters*
393 2012, **15:**55-64.

394 * Very elegant study of how underground specialized herbivores can hijack
395 plant defenses for their own benefits

396 34. Robert CAM, Erb M, Hiltpold I, Hibbard BE, Gaillard MDP, Bilat J, Degenhardt
397 J, Cambet-Petit-Jean X, Turlings TCJ, Zwahlen C: **Genetically engineered**
398 **maize plants reveal distinct costs and benefits of constitutive**
399 **volatile emissions in the field.** *Plant Biotechnology Journal* 2013,
400 **11:**628-639.

35. Azcón-Aguilar C, Barea JM: **Arbuscular mycorrhizas and biological control of soil-borne plant pathogens - An overview of the mechanisms involved.** *Mycorrhiza* 1996, **6**:457-464.
36. Van der Heijden MGA, Sanders IR (Ed): *Mycorrhizal Ecology* Berlin Heidelberg: Springer-Verlag; 2002.
37. Pozo MJ, Azcon-Aguilar C: **Unraveling mycorrhiza-induced resistance.** *Current Opinion in Plant Biology* 2007, **10**:393-398.
38. Cameron DD, Neal AL, van Wees SCM, Ton J: **Mycorrhiza-induced resistance: more than the sum of its parts?** *Trends in plant science* 2013, **18**:539-545.
- * Theoretical framework of how mycorrhizae can increase plant defenses
39. Bennett AE, Alers-Garcia J, Bever JD: **Three-way interactions among mutualistic mycorrhizal fungi, plants, and plant enemies: Hypotheses and synthesis.** *American Naturalist* 2006, **167**:141-152.
40. Vannette RL, Rasmann S: **Arbuscular mycorrhizal fungi mediate below-ground plant-herbivore interactions: a phylogenetic study.** *Functional Ecology* 2012, **26**:1033-1042.
41. Koricheva J, Gange AC, Jones T: **Effects of mycorrhizal fungi on insect herbivores: a meta-analysis.** *Ecology* 2009, **90**:2088-2097.
42. Giovannetti M, Sbrana C, Citernesi AS, Avio L: **Analysis of factors involved in fungal recognition responses to host-derived signals by arbuscular mycorrhizal fungi.** *New Phytologist* 1996, **133**:65-71.
43. Besserer A, Puech-Pagès V, Kiefer P, Gomez-Roldan V, Jauneau A, Roy S, Portais J-C, Roux C, Bécard G, Séjalon-Delmas N: **Strigolactones stimulate arbuscular mycorrhizal fungi by activating mitochondria.** *PLoS Biol* 2006, **4**:e226.
44. Akiyama K, Matsuzaki K-i, Hayashi H: **Plant sesquiterpenes induce hyphal branching in arbuscular mycorrhizal fungi.** *Nature* 2005, **435**:824-827.
45. Peters NK, Frost JW, Long SR: **A plant flavone, luteolin, induces expression of Rhizobium meliloti nodulation genes.** *Science* 1986, **233**:977-980.
46. Gray EJ, Smith DL: **Intracellular and extracellular PGPR: commonalities and distinctions in the plant-bacterium signaling processes.** *Soil Biology and Biochemistry* 2005, **37**:395-412.
47. Neal AL, Ahmad S, Gordon-Weeks R, Ton J: **Benzoxazinoids in root exudates of maize attract Pseudomonas putida to the rhizosphere.** *PLoS ONE* 2012, **7**:e35498.
48. Bragazza L, Bardgett RD, Mitchell EAD, Buttler A: **Linking soil microbial communities to vascular plant abundance along a climate gradient.** *New Phytologist* 2015, **205**:1175-1182.
49. Coleman DC: **The microbial loop concept as used in terrestrial soil ecology studies.** *Microbial Ecology* 1994, **28**:245-250.
50. Toepfer S, Haye T, Erlandson M, Goettel M, Lundgren JG, Kleespies RG, Weber DC, Walsh GC, Peters A, Ehlers RU, et al.: **A review of the natural enemies of beetles in the subtribe Diabroticina (Coleoptera: Chrysomelidae): implications for sustainable pest management.** *Biocontrol Science and Technology* 2009, **19**:1-65.
51. Aratchige NS, Lesna I, Sabelis MW: **Below-ground plant parts emit herbivore-induced volatiles: olfactory responses of a predatory mite**

- 450 **to tulip bulbs infested by rust mites.** *Experimental and Applied*
 451 *Acarology* 2004, **33**:21-30.
- 452 52. Vega FE, Goettel MS, Blackwell M, Chandler D, Jackson MA, Keller S, Koike M,
 453 Maniania NK, Monzón A, Ownley BH, et al.: **Fungal entomopathogens:**
 454 **new insights on their ecology.** *Fungal Ecology* 2009, **2**:149-159.
- 455 53. Kanagy JMN, Kaya HK: **The possible role of marigold roots and alpha-**
 456 **terthienyl in mediating host-finding by steinernematid nematodes.**
 457 *Nematologica* 1996, **42**:220-231.
- 458 54. Choo HY, Kaya HK, Burlando TM, Gaugler R: **Entomopathogenic nematodes**
 459 **- Host-finding ability in the presence of plant-roots.** *Environmental*
 460 *Entomology* 1989, **18**:1136-1140.
- 461 55. Strong DR, Kaya HK, Whipple AV, Child AL, Kraig S, Bondonno M, Dyer K,
 462 Maron JL: **Entomopathogenic nematodes: natural enemies of root-**
 463 **feeding caterpillars on bush lupine.** *Oecologia* 1996, **108**:167-173.
- 464 56. Ali J, Alborn H, Stelinski L: **Subterranean herbivore-induced volatiles**
 465 **released by Citrus roots upon feeding by *Diaprepes abbreviatus***
 466 **recruit entomopathogenic nematodes.** *Journal of Chemical Ecology*
 467 2010, **36**:361-368.
- 468 57. Ali JG, Alborn HT, Stelinski LL: **Constitutive and induced subterranean**
 469 **plant volatiles attract both entomopathogenic and plant parasitic**
 470 **nematodes.** *Journal of Ecology* 2011, **99**:26-35.
- 471 58. Rasmann S, Kollner TG, Degenhardt J, Hiltpold I, Toepfer S, Kuhlmann U,
 472 Gershenzon J, Turlings TCJ: **Recruitment of entomopathogenic**
 473 **nematodes by insect-damaged maize roots.** *Nature* 2005, **434**:732-
 474 737.
- 475 59. van Tol R, van der Sommen ATC, Boff MIC, van Bezooijen J, Sabelis MW, Smits
 476 PH: **Plants protect their roots by alerting the enemies of grubs.**
 477 *Ecology Letters* 2001, **4**:292-294.
- 478 60. Hiltpold I, Baroni M, Toepfer S, Kuhlmann U, Turlings TCJ: **Selection of**
 479 **entomopathogenic nematodes for enhanced responsiveness to a**
 480 **volatile root signal helps to control a major root pest.** *J Exp Biol* 2010,
 481 **213**:2417-2423.
- 482 61. Tonelli M, Peñaflor MFGV, Leite LG, Silva WD, Martins F, Bento JMS:
 483 **Attraction of entomopathogenic nematodes to sugarcane root**
 484 **volatiles under herbivory by a sap-sucking insect.** *Chemoecology* 2016,
 485 **26**:59-66.
- 486 62. Degenhardt J, Hiltpold I, Kollner TG, Frey M, Gierl A, Gershenzon J, Hibbard
 487 BE, Ellersieck MR, Turlings TCJ: **Restoring a maize root signal that**
 488 **attracts insect-killing nematodes to control a major pest.** *Proceedings*
 489 *of the National Academy of Sciences of the United States of America* 2009,
 490 **106**:13213-13218.
- 491 63. Hunter MD: **Out of sight, out of mind: the impacts of root-feeding insects**
 492 **in natural and managed systems.** *Agricultural and Forest Entomology*
 493 2001, **3**:3-9.
- 494 64. Czjzek M, Cicek M, Zamboni V, Bevan DR, Henrissat B, Esen A: **The**
 495 **mechanism of substrate (aglycone) specificity in beta-glucosidases is**
 496 **revealed by crystal structures of mutant maize beta-glucosidase-**
 497 **DIMBOA, -DIMBOAGlc, and -dhurrin complexes.** *Proceedings of the*

- 498 *National Academy of Sciences of the United States of America* 2000,
 499 **97**:13555-13560.
- 500 65. Morant AV, Jorgensen K, Jorgensen C, Paquette SM, Sanchez-Perez R, Moller
 501 BL, Bak S: **beta-glucosidases as detonators of plant chemical defense.**
 502 *Phytochemistry* 2008, **69**:1795-1813.
- 503 66. Koehler A, Maag D, Veyrat N, Glauser G, Wolfender J-L, Turlings TCJ, Erb M:
 504 **Within-plant distribution of 1,4-benzoxazin-3-ones contributes to**
 505 **herbivore niche differentiation in maize.** *Plant Cell and Environment*
 506 2015, **38**:1081-1093.
- 507 67. Wouters FC, Reichelt M, Glauser G, Bauer E, Erb M, Gershenzon J, Vassao DG:
 508 **Reglucosylation of the benzoxazinoid DIMBOA with inversion of**
 509 **stereochemical configuration is a detoxification strategy in**
 510 **lepidopteran herbivores.** *Angew Chem Int Ed Engl* 2014, **53**:11320-
 511 11324.
- 512 68. Glauser G, Marti G, Villard N, Doyen GA, Wolfender JL, Turlings TC, Erb M:
 513 **Induction and detoxification of maize 1,4-benzoxazin-3-ones by**
 514 **insect herbivores.** *Plant J* 2011, **68**:901-911.
- 515 69. Maag D, Erb M, Koellner TG, Gershenzon J: **Defensive weapons and defense**
 516 **signals in plants: Some metabolites serve both roles.** *Bioessays* 2015,
 517 **37**:167-174.
- 518 70. Poloni A, Schirawski J: **Red card for pathogens: phytoalexins in sorghum**
 519 **and maize.** *Molecules* 2014, **19**:9114-9133.
- 520 71. Cambier V, Hance T, de Hoffmann E: **Variation of DIMBOA and related**
 521 **compounds content in relation to the age and plant organ in maize.**
 522 *Phytochemistry* 2000, **53**:223-229.
- 523 72. Baetz U, Martinoia E: **Root exudates: the hidden part of plant defense.**
 524 *Trends in Plant Science* 2014, **19**:90-98.
- 525 73. Chen K-J, Zheng Y-Q, Kong C-H, Zhang S-Z, Li J, Liu X-G: **2,4-dihydroxy-7-**
 526 **methoxy-1,4-benzoxazin-3-one (DIMBOA) and 6-methoxy-**
 527 **benzoxazolin-2-one (MBOA) levels in the wheat rhizosphere and**
 528 **their effect on the soil microbial community structure.** *Journal of*
 529 *Agricultural and Food Chemistry* 2010, **58** 12710-12716.
- 530 74. Kollner TG, Held M, Lenk C, Hiltpold I, Turlings TCJ, Gershenzon J, Degenhardt
 531 J: **A maize (*E*)-beta-caryophyllene synthase implicated in indirect**
 532 **defense responses against herbivores is not expressed in most**
 533 **American maize varieties.** *Plant Cell* 2008, **20**:482-494.
- 534 75. Turlings T, Hiltpold I, Rasman S: **The importance of root-produced**
 535 **volatiles as foraging cues for entomopathogenic nematodes.** *Plant*
 536 *and Soil* 2012:1-10.
- 537 76. Bouwmeester HJ, Roux C, Lopez-Raez JA, Becard G: **Rhizosphere**
 538 **communication of plants, parasitic plants and AM fungi.** *Trends in*
 539 *Plant Science* 2007, **12**:224-230.
- 540 77. Seto Y, Kameoka H, Yamaguchi S, Kyoizuka J: **Recent advances in**
 541 **strigolactone research: chemical and biological aspects.** *Plant and Cell*
 542 *Physiology* 2012, **53**:1843-1853.
- 543 78. Bouwmeester HJ, Matusova R, Zhongkui S, Beale MH: **Secondary metabolite**
 544 **signalling in host-parasitic plant interactions.** *Curr Opin Plant Biol*
 545 2003, **6**:358-364.

- 546 79. Yoder JI: **Host-plant recognition by parasitic Scrophulariaceae.** *Curr Opin*
547 *Plant Biol* 2001, **4**:359-365.
- 548 80. Gershenzon J, Dudareva N: **The function of terpene natural products in**
549 **the natural world.** *Nature Chemical Biology* 2007, **3**:408-414.
- 550 81. Screpanti C, Fonne-Pfister R, Lumbroso A, Rendine S, Lachia M, De
551 Mesmaeker A: **Strigolactone derivatives for potential crop**
552 **enhancement applications.** *Bioorg Med Chem Lett* 2016.
- 553 * Nice synthesis of the advancement in different areas of the strigolactone
554 domain and the implications for agronomical applications

Graphical Abstract

Figure 1